

# Sign and signed rank test

STAT 401 - Statistical Methods for Research Workers

Jarad Niemi

Iowa State University

7 September 2013

## Rusty leaves data

	year1	year2	diff	>0
1	38	32	6	1
2	10	16	-6	0
3	84	57	27	1
4	36	28	8	1
5	50	55	-5	0
6	35	12	23	1
7	73	61	12	1
8	48	29	19	1

# Rusty leaves data

	year1	year2	diff	>0
1	38	32	6	1
2	10	16	-6	0
3	84	57	27	1
4	36	28	8	1
5	50	55	-5	0
6	35	12	23	1
7	73	61	12	1
8	48	29	19	1

If the null hypothesis is true, then the final column should have 50/50 change of being 0 or 1.

# Rusty leaves data

	year1	year2	diff	>0
1	38	32	6	1
2	10	16	-6	0
3	84	57	27	1
4	36	28	8	1
5	50	55	-5	0
6	35	12	23	1
7	73	61	12	1
8	48	29	19	1

If the null hypothesis is true, then the final column should have 50/50 change of being 0 or 1. In this example, we have 6 ones out of 8.

# Sign test

The sign test calculates the probability of observing this many ones (or more extreme) if the null hypothesis is true.

# Sign test

The sign test calculates the probability of observing this many ones (or more extreme) if the null hypothesis is true. For our one-sided hypothesis, that is the probability of observing 6, 7, or 8 ones.

# Sign test

The sign test calculates the probability of observing this many ones (or more extreme) if the null hypothesis is true. For our one-sided hypothesis, that is the probability of observing 6, 7, or 8 ones. This is

$$\binom{8}{6}0.5^8 + \binom{8}{7}0.5^8 + \binom{8}{8}0.5^8 = 0.14$$

# Sign test

The sign test calculates the probability of observing this many ones (or more extreme) if the null hypothesis is true. For our one-sided hypothesis, that is the probability of observing 6, 7, or 8 ones. This is

$$\binom{8}{6}0.5^8 + \binom{8}{7}0.5^8 + \binom{8}{8}0.5^8 = 0.14$$

Although conceptually simple, this test is not very powerful and therefore is used rarely.



# Signed rank test

Also known as the Wilcoxon signed rank test:

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.
- Order the absolute differences from smallest to largest and assign them their ranks.

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.
- Order the absolute differences from smallest to largest and assign them their ranks.
- Calculate  $S$ : the sum of the ranks from the pairs for which the difference is positive.

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.
- Order the absolute differences from smallest to largest and assign them their ranks.
- Calculate  $S$ : the sum of the ranks from the pairs for which the difference is positive.
- Calculate  $E[S] = n(n+1)/4$  where  $n$  is the number of pairs.

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.
- Order the absolute differences from smallest to largest and assign them their ranks.
- Calculate  $S$ : the sum of the ranks from the pairs for which the difference is positive.
- Calculate  $E[S] = n(n+1)/4$  where  $n$  is the number of pairs.
- Calculate  $SD[S] = [n(n+1)(2n+1)/24]^{1/2}$ .

# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.
- Order the absolute differences from smallest to largest and assign them their ranks.
- Calculate  $S$ : the sum of the ranks from the pairs for which the difference is positive.
- Calculate  $E[S] = n(n+1)/4$  where  $n$  is the number of pairs.
- Calculate  $SD[S] = [n(n+1)(2n+1)/24]^{1/2}$ .
- Calculate  $(S + c - E[S])/SD[S]$  where  $c$  is the appropriate continuity correction.



# Signed rank test

Also known as the Wilcoxon signed rank test:

- Compute the difference in each pair.
- Drop zeros from the list.
- Order the absolute differences from smallest to largest and assign them their ranks.
- Calculate  $S$ : the sum of the ranks from the pairs for which the difference is positive.
- Calculate  $E[S] = n(n+1)/4$  where  $n$  is the number of pairs.
- Calculate  $SD[S] = [n(n+1)(2n+1)/24]^{1/2}$ .
- Calculate  $(S + c - E[S])/SD[S]$  where  $c$  is the appropriate continuity correction.
- Calculate the pvalue from a standard normal table.

# Signed rank test

	year1	year2	diff	>0	absdiff	rank
5	50	55	-5	0	5	1
1	38	32	6	1	6	2.5
2	10	16	-6	0	6	2.5
4	36	28	8	1	8	4
7	73	61	12	1	12	5
8	48	29	19	1	19	6
6	35	12	23	1	23	7
3	84	57	27	1	27	8

# Signed rank test

	year1	year2	diff	>0	absdiff	rank
5	50	55	-5	0	5	1
1	38	32	6	1	6	2.5
2	10	16	-6	0	6	2.5
4	36	28	8	1	8	4
7	73	61	12	1	12	5
8	48	29	19	1	19	6
6	35	12	23	1	23	7
3	84	57	27	1	27	8

- $S = 32.5$

# Signed rank test

	year1	year2	diff	>0	absdiff	rank
5	50	55	-5	0	5	1
1	38	32	6	1	6	2.5
2	10	16	-6	0	6	2.5
4	36	28	8	1	8	4
7	73	61	12	1	12	5
8	48	29	19	1	19	6
6	35	12	23	1	23	7
3	84	57	27	1	27	8

- $S = 32.5$
- $E[S] = 18$

# Signed rank test

	year1	year2	diff	>0	absdiff	rank
5	50	55	-5	0	5	1
1	38	32	6	1	6	2.5
2	10	16	-6	0	6	2.5
4	36	28	8	1	8	4
7	73	61	12	1	12	5
8	48	29	19	1	19	6
6	35	12	23	1	23	7
3	84	57	27	1	27	8

- $S = 32.5$
- $E[S] = 18$
- $SD[S] = 7.14$

# Signed rank test

	year1	year2	diff	>0	absdiff	rank
5	50	55	-5	0	5	1
1	38	32	6	1	6	2.5
2	10	16	-6	0	6	2.5
4	36	28	8	1	8	4
7	73	61	12	1	12	5
8	48	29	19	1	19	6
6	35	12	23	1	23	7
3	84	57	27	1	27	8

- $S = 32.5$
- $E[S] = 18$
- $SD[S] = 7.14$
- $Z = 1.96$  (with continuity correction of -0.5)

# Signed rank test

	year1	year2	diff	>0	absdiff	rank
5	50	55	-5	0	5	1
1	38	32	6	1	6	2.5
2	10	16	-6	0	6	2.5
4	36	28	8	1	8	4
7	73	61	12	1	12	5
8	48	29	19	1	19	6
6	35	12	23	1	23	7
3	84	57	27	1	27	8

- $S = 32.5$
- $E[S] = 18$
- $SD[S] = 7.14$
- $Z = 1.96$  (with continuity correction of -0.5)
- $p = 0.02$

# SAS code for paired nonparametric test

```
DATA leaves;  
  INPUT tree year1 year2;  
  diff = year1-year2;  
  DATALINES;  
1 38 32  
2 10 16  
3 84 57  
4 36 28  
5 50 55  
6 35 12  
7 73 61  
8 48 29  
;  
  
PROC UNIVARIATE DATA=leaves;  
  VAR diff;  
  RUN;
```



# SAS code for paired nonparametric tests

## The UNIVARIATE Procedure

Variable: diff

### Moments

N	8	Sum Weights	8
Mean	10.5	Sum Observations	84
Std Deviation	12.2007026	Variance	148.857143
Skewness	-0.1321468	Kurtosis	-1.2476273
Uncorrected SS	1924	Corrected SS	1042
Coeff Variation	116.197167	Std Error Mean	4.31359976

### Basic Statistical Measures

Location		Variability	
Mean	10.50000	Std Deviation	12.20070
Median	10.00000	Variance	148.85714
Mode	.	Range	33.00000
		Interquartile Range	20.50000

### Tests for Location: Mu0=0

Test	-Statistic-	-----p Value-----
Student's t	t 2.434162	Pr >  t  0.0451
Sign	M 2	Pr >=  M  0.2891
Signed Rank	S 14.5	Pr >=  S  0.0469

# Conclusion

Removal of red cedar trees within 100 yards is associated with a significant reduction in rusty apple leaves (Wilcoxon signed rank test,  $p=0.023$ ).