

Contents

Part 1	1.1	Co-ordinate Free Vectors	2
Prerequisites			
Chapter 1		Vectors	2

Part 1



Prerequisites

Vectors

In this chapter we'll give an introduction to vectors, and some results in geometry that you may find useful. Vectors are not introduced rigorously here, but we do use some facts from linear algebra.

1.1 Co-ordinate Free Vectors

We first describe vectors as being co-ordinate free. And really, even though we will introduce co-ordinates later, you should still think of vectors as just existing out somewhere, with or without co-ordinates.

You can think of a vector as a *directed line segment*. Vectors are usually denoted using boldface, \mathbf{A} or with an arrow over its name, \vec{A} . We'll use the former, though the latter is often used when writing by hand.

Because they're directed line segments, only their length matters and we're not concerned with the location of their end points. This means that a vector can be freely translated, without altering it.

The magnitude of the vector is its length, which we call it's *norm*, and write as $\|\mathbf{A}\|$ or simply A . You may also see it being written as $|\mathbf{A}|$.

A unit vector, $\hat{\gamma}$ ("gamma hat") is a vector whose magnitude is unity. We use the unit vector to often denote the direction of a vector by multiplying the unit vector with its magnitude. For instance, the unit vector that point in the direction of \mathbf{A} , $\hat{\mathbf{A}}$ can be calculated as,

$$\hat{\mathbf{A}} = \frac{\mathbf{A}}{A}.$$

Further, vectors have certain operations defined on them. The base operations are *Scalar Multiplication* and *Vector Addition*.

Scalar Multiplication

Scalar Multiplication refers to multiplying a vector by a *scalar*. A scalar for our purposes refers to a real number. Consider a vector \mathbf{v} . Multiplying it by a scalar, $\alpha \in \mathbb{R}$ produces a vector $\alpha\mathbf{v}$ parallel to the original vector and changes the magnitude of the vector so that $\|\alpha\mathbf{v}\|$ is $|\alpha|$ times greater than $\|\mathbf{v}\|$, thus $\|\alpha\mathbf{v}\| = |\alpha|\|\mathbf{v}\|$. $|\alpha|$ can be smaller or greater than 1 which accordingly increases/decreases the length of \mathbf{v} .




Figure 1.1: A directed line segment, the vector \mathbf{A} .




Figure 1.2: Two identical vectors.




Figure 1.3: Scalar multiplication of a vector \mathbf{A} by $c > 1$ and -1 .


If $\alpha > 0$, the vector produced is in the same direction as the original vector. If $\alpha < 0$, the direction of the vector is reversed.

Vector Addition


Adding two vectors, \mathbf{A} and \mathbf{B} produces another vector $\mathbf{A} + \mathbf{B}$. Geometrically, vector addition is done by placing the tail of \mathbf{B} on the head of \mathbf{A} , and then the vector joining the tail of \mathbf{A} and head of \mathbf{B} is the vector $\mathbf{A} + \mathbf{B}$ as in fig. 1.3a.

It can also be interpreted as the diagonal of the parallelogram made by placing the tails of the two vectors together, and producing two sides parallel to them as show in fig. 1.3b.

Subtraction of vectors, $\mathbf{A} - \mathbf{B}$ is equivalent to multiplying \mathbf{B} by -1 and then adding it with \mathbf{A} .



(a) Adding vectors, parallelogram interpretation.



(b) Adding vectors, triangle interpretation.

Figure 1.4: Addition of two vectors, \mathbf{A} and \mathbf{B} produces another vector, $\mathbf{A} + \mathbf{B}$.