

RWorksheet_Basa#3b

Jared J Basa

2024-12-17

1. Create a data frame using the table below.

a. Write the codes.

```
Table <- data.frame(  
  Respondents = c(1:20),  
  Sex = c(2,2,1,2,2,2,2,2,2,2,1,2,2,2,2,2, 2, 1,2),  
  Fathers_Occupation = c(1, 3, 3, 3, 1, 2, 3, 1, 1, 1, 3, 2, 1, 3, 3, 1, 3, 1, 2, 1),  
  Persons_at_Home = c(5, 7, 3, 8, 5, 9, 6, 7, 8, 4, 7, 5, 4, 7, 8, 8, 3, 11, 7, 6),  
  Siblings_at_School = c(6, 4, 4, 1, 2, 1, 5, 3, 1, 2, 3, 2, 5, 5, 2, 1, 2, 5, 3, 2),  
  Type_of_House = c(1, 2, 3, 1, 1, 3, 3, 1, 2, 3, 2, 3, 2, 2, 3, 3, 3, 3, 3, 2)  
)
```

Table

##	Respondents	Sex	Fathers_Occupation	Persons_at_Home	Siblings_at_School
## 1	1	2	1	5	6
## 2	2	2	3	7	4
## 3	3	1	3	3	4
## 4	4	2	3	8	1
## 5	5	2	1	5	2
## 6	6	2	2	9	1
## 7	7	2	3	6	5
## 8	8	2	1	7	3
## 9	9	2	1	8	1
## 10	10	2	1	4	2
## 11	11	1	3	7	3
## 12	12	2	2	5	2
## 13	13	2	1	4	5
## 14	14	2	3	7	5
## 15	15	2	3	8	2
## 16	16	2	1	8	1
## 17	17	2	3	3	2
## 18	18	2	1	11	5
## 19	19	1	2	7	3
## 20	20	2	1	6	2
##	Type_of_House				
## 1	1				
## 2	2				
## 3	3				
## 4	1				
## 5	1				
## 6	3				
## 7	3				

```
## 8      1
## 9      2
## 10     3
## 11     2
## 12     3
## 13     2
## 14     2
## 15     3
## 16     3
## 17     3
## 18     3
## 19     3
## 20     2
```

b. Describe the data. Get the structure or the summary of the data

```
str(Table)
```

```
## 'data.frame': 20 obs. of 6 variables:
## $ Respondents : int 1 2 3 4 5 6 7 8 9 10 ...
## $ Sex : num 2 2 1 2 2 2 2 2 2 2 ...
## $ Fathers_Occupation: num 1 3 3 3 1 2 3 1 1 1 ...
## $ Persons_at_Home : num 5 7 3 8 5 9 6 7 8 4 ...
## $ Siblings_at_School: num 6 4 4 1 2 1 5 3 1 2 ...
## $ Type_of_House : num 1 2 3 1 1 3 3 1 2 3 ...
```

```
summary(Table)
```

```
## Respondents Sex Fathers_Occupation Persons_at_Home
## Min. : 1.00 Min. :1.00 Min. :1.00 Min. : 3.0
## 1st Qu.: 5.75 1st Qu.:2.00 1st Qu.:1.00 1st Qu.: 5.0
## Median :10.50 Median :2.00 Median :2.00 Median : 7.0
## Mean :10.50 Mean :1.85 Mean :1.95 Mean : 6.4
## 3rd Qu.:15.25 3rd Qu.:2.00 3rd Qu.:3.00 3rd Qu.: 8.0
## Max. :20.00 Max. :2.00 Max. :3.00 Max. :11.0
## Siblings_at_School Type_of_House
## Min. :1.00 Min. :1.0
## 1st Qu.:2.00 1st Qu.:2.0
## Median :2.50 Median :2.5
## Mean :2.95 Mean :2.3
## 3rd Qu.:4.25 3rd Qu.:3.0
## Max. :6.00 Max. :3.0
```

c. Is the mean number of siblings attending is 5?

```
meanSiblings_at_School <- mean(Table$Siblings_at_School)
meanSiblings_at_School
```

```
## [1] 2.95
```

no

d. Extract the 1st two rows and then all the columns using the subsetting functions. Write the codes and its output.

```
subsetRespondents <- subset(Table, Respondents <= 2)
subsetRespondents
```

```
## Respondents Sex Fathers_Occupation Persons_at_Home Siblings_at_School
```

```
## 1      1  2      1      5      6
## 2      2  2      3      7      4
##   Type_of_House
## 1      1
## 2      2
```

e. Extract 3rd and 5th row with 2nd and 4th column. Write the codes and its result.

```
rowCols <- Table[c(3, 5), c("Sex", "Persons_at_Home")]
rowCols
```

```
##   Sex Persons_at_Home
## 3   1              3
## 5   2              5
```

f. Select the variable types of houses then store the vector that results as types_houses. Write the codes.

```
Type_of_House <- Table$Type_of_House
Type_of_House
```

```
## [1] 1 2 3 1 1 3 3 1 2 3 2 3 2 2 3 3 3 3 3 2
```

g. Select only all Males respondent that their father occupation was farmer. Write the codes and its output.