

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction  
Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications  
Conclusions

# Modeling the Construction and Evolution of Distributed Volcanic Fields on Earth and Mars

Jacob A. Richardson

School of Geosciences  
University of South Florida

19 February 2016

# Acknowledgements

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows

Arsia Mons

Volcanic Field

Methods

Results

Implications

Conclusions

## Some Collaborators

Chuck Connor

Laura Connor

Sylvain Charbonnier

Judy McIlrath

Paul Wetmore

James Wilson

Lis Gallant

Julia Kubanek

Jake Bleacher

Lori Glaze

## Funding Agencies

NASA Mars Data Analysis Program

NSF SSI

# Introduction

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field


Methods  
Results  
Implications

Conclusions

## Distributed-style Volcanism

### Characteristics

- Clusters of volcanoes are formed, sometimes associated with large volcanoes
- New eruptions form new vents
- Eruptions are fed by small volume batches of magma
- Long periods of quiescence



# Outline of Talk

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview


Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions

- Volcanic fields, from the inside out
  - The role of sills in the formation of volcanic fields (Richardson et al., *Geology*, 2015)
  - The spatial organization of vents in volcanic fields (Richardson et al., *LPSC*, 2012)
  - Simulating lava flow emplacement  
(Kabanek et al., *Bull. Volc.*, 2015)
- Evolution of volcanism at Arsia Mons
  - Can its recent rate of volcanism be determined by studying a volcanic field with only satellite data?
- Conclusions



# The Igneous Plumbing System

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions

## San Rafael Volcanic Field, Utah

- Pliocene volcanic activity
- Now eroded to depth of ~1 km
- Sills and Dikes exposed



Chuck Connor with a Terrestrial Lidar (L. Connor)




Photo Credit: Judy McIlrath

# The Igneous Plumbing System

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction


Overview  
Sills

Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications  
Conclusions

## Results of lidar survey

- Sill volume comparable to volume thought to have erupted at surface
- Sills had ability to modulate eruption style by interacting with volcanic conduits
- Conduits deliver magma from depth to distributed volcanoes



*Richardson et al., Geology, 2015*

# Volcano organization in fields

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows

Arsia Mons

Volcanic Field


Methods

Results

Implications

Conclusions

The spatial organization of volcanoes is modeled as a density function



- Area of volcanic field defined as 95<sup>th</sup> percentile of density function

$$\text{Average vent intensity} = \frac{\# \text{ volcanic vents}}{\text{field area}}$$

- This model is applied to fields on Earth, Mars, and Venus

# Vent Spatial Intensity across the Solar System

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills



Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field


Methods  
Results  
Implications

Conclusions

## Average Vent Intensity, Colored by Planet



# The 2012-3 Tolbachik Lava Flow



*NASA Earth Observing-1 Mission, 1 February 2013*

# Lava Flows/Simulators

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview


Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications


Conclusions

- MOLASSES developed after Connor et al., *JAV*, 2012
- Spreads lava over a grid according to universal rules

## Optional Spreading Rules




Using TanDEM-X satellite data, flow simulations match the 2012-3 Tolbachik flow between 70-85%.



Kubanek et al., *Bull. Volc.*, 2015

# Distributed Volcanism of the Tharsis Volcanic Province



## Tharsis Vent Catalog

- >1,000 small volcanic vents cataloged  
(Richardson et al., JVGR, 2013, Bleacher et al., JVGR, 2009)
- Groups of vents form isolated clusters

## Research Questions

- How does distributed-style volcanism occur over time and space in Tharsis?
- How do volcanic fields relate to the larger volcanoes on Mars?

# Arsia Mons Overview

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows

**Arsia Mons**  
Volcanic Field

Methods

Results

Implications

Conclusions

## Arsia Mons

- Large ( $1.5 \cdot 10^6 \text{ km}^3$ ) shield volcano with 110 km diameter caldera
- A cluster of volcanic vents lay in the caldera!

## Motivation

What are the recurrence rate of  
volcanism and delivery rate of  
magma to the surface?



# Arsia Mons Overview

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows

Arsia Mons  
Volcanic Field

Methods

Results

Implications

Conclusions

## Recurrence Rate and Magma Delivery Rate

$$\text{Recurrence Rate} = \frac{\text{Number of Events} - 1}{\text{Time elapsed}}$$

$$\text{Delivery Rate} = \frac{\text{Total Volume}}{\text{Number of Events}} \times \text{Recurrence Rate}$$

- Lavas from these vents can be mapped to estimate volume and timing of emplacement



# Mapping Volcanic Vents

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction


Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions



CTX Image: G10\_022160\_1710\_XN\_09S120W (NASA/JPL-Caltech/MSSS)

# Mapping Lava Flows

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction


Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions



CTX Image: G10\_022160\_1710\_XN\_09S120W (NASA/JPL-Caltech/MSSS)



# Lava Flow Map of Arsia Mons' Caldera

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson


Introduction

Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications

Conclusions



## Mapping results

- 29 vents are cataloged, each with long lava flows
- Lava flow areas are 10s–100s km<sup>2</sup>
- Flow thicknesses assumed to be 10–80 m (Mouginis-Mark & Rowland, *Icarus*, 2008)
- From this, volumes estimates range from 10<sup>-2</sup>–70 km<sup>3</sup>

# Ages: Crater Counting

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction


Overview

Sills  
Vent Intensity  
Lava Flows

Arisia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions



CTX Image: G10\_022160\_1710\_XN\_09S120W (NASA/JPL-Caltech/MSSS)

Impact craters on  
each flow are  
cataloged

More craters  
generally indicates  
older age

Ages and  
uncertainties  
modeled in  
*craterstats2* (Michael,  
*Icarus*, 2013)

# Ages: Crater Counting

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills  
Vent Intensity  
Lava Flows



Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions

Ages of flow emplacement are modeled as Normal Distributions

- Estimated age from *craterstats2* is used as the Mean Value
- Age Uncertainty is Standard Deviation



# Ages: Stratigraphy

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction


Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions




CTX Image: G10\_022160\_1710\_XN\_09S120W (NASA/JPL-Caltech/MSSS)

Stratigraphic  
relationships  
relatively date  
events.



Lava from v19  
overlies v23 lavas

*Graphical Form:*



## Combined Age Information

## Stratigraphy “Web”



# Ages: Information Conflicts

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview



Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions

Mean crater ages can agree stratigraphy...



# Ages: Information Conflicts

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview


Sills  
Vent Intensity  
Lava Flows

Arisia Mons  
Volcanic Field


Methods  
Results  
Implications

Conclusions

...or mean age can disagree with strat.



But by probabilistically modeling age,  
possible ages can still be identified



# Volcanic Event Recurrence Rate Model (VERRM)

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview


Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications  
Conclusions



## Step 1: Find potential ages of all events in the field (Monte Carlo)

### Input Files

Age Model Database			Stratigraphic Relationship Database	
EventID	Mean_age	Uncertainty	Older	Younger
A	50	30	B	A
B	80	20	D	B
C	30	05	D	C
D	80	50	E	D
E	150	75		



For each vent:



Output

Sampled\_Age [A]  
Sampled\_Age [B]  
...  
Sampled\_Age [E]

# Volcanic Event Recurrence Rate Model (VERRM)

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows


Arsia Mons  
Volcanic Field

Methods

Results

Implications

Conclusions



## Step 2: Calculate Recurrence Rate of Volcanism

For each Monte Carlo simulation (i.e. each set of potential ages), model Recurrence Rate (RR) through time:

$$RR(t) = \frac{3}{T_{e-1} - T_{e+2}}$$

## Step 3: Calculate Magma Delivery Rate

Model Magma Delivery Rate (Flux) through time:

$$\text{Flux}(t) = \frac{\text{Volume}_e}{T_e - T_{e+1}}$$

# Results

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction


Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications

Conclusions

## Recurrence Rate through time



- All 10,000 Recurrence Rate functions are combined
- Rate peaks at 150 Ma, producing 1 vent per 1-10 Myr (Median value:  $0.25 \text{ events Myr}^{-1}$ )
- Rate has decreased monotonically:  $\text{RR}(t = 0) \rightarrow 0$

# Volume Flux

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills  
Vent Intensity  
Lava Flows


Arsia Mons  
Volcanic Field

Methods  
Results  
Implications

Conclusions

## Magma Delivery Rate through time

- All 10,000 Magma Delivery Rate functions are combined
- Median rate plateaus at at 150 Ma ( $0.4\text{--}3 \text{ km}^3 \text{ Myr}^{-1}$ ), remains steady until 100 Ma before rapidly waning.
- Uncertainty in magma delivery rate is about 1 order of magnitude to either side of the median



# Tie-in with ashes and glaciers

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows


Arsia Mons  
Volcanic Field

Methods

Results

Implications

Conclusions



- Extant glaciers are preserved on Western flank of Arsia
- Preserved for ~200 Ma by ashes (Kadish et al., *P&SS*, 2014)
- If ashes were sourced near-summit (Mouginis-Mark, *GRL*, 2002), our effusive volcanism might be predated by explosive activity

← from Shean et al., *JGR Planets*, 2007

# Model of waning volcanism of Arsia

Volcanic  
Fields on  
Earth & Mars



Jacob  
Richardson

Introduction

Overview

Sills  
Vent Intensity  
Lava Flows

Arsia Mons  
Volcanic Field  
Methods  
Results  
Implications  
Conclusions



- ① Large magma chamber fed edifice-building, sometimes explosive eruptions (Wilson et al., *JGR*, 2001)
- ② Ashes coated ice-rich deposits before 200 Ma
- ③ The chamber cooled from waning magma flux
- ④ Volcanism evolved from explosive to effusive as magma waned
- ⑤ Volcanism is now in hiatus ( $RR(t = 0) \rightarrow 0$ )

# Conclusions

Volcanic  
Fields on  
Earth & Mars

Jacob  
Richardson

Introduction

Overview

Sills

Vent Intensity

Lava Flows

Arsia Mons

Volcanic Field

Methods

Results

Implications

Conclusions

## Arsia Mons

- ① Late volcanism at Arsia Mons emplaced a volcanic field in its caldera
- ② Multiple dating methods used on lavas can be combined to estimate rate of volcanism, which peaked 150 million years ago
- ③ Volcanism waned and likely ceased 10-90 million years ago
- ④ This field might be related to a larger waning of volcanism at Arsia
- ⑤ Vents in this field are less concentrated than vents in terrestrial volcanic fields
- ⑥ Lava flows are an order of magnitude larger in volume than flows in most terrestrial fields

# Conclusions

## Overall Conclusions

- Volcanic fields have complex, voluminous roots
- Emplacement of lavas in these fields can be modeled
- Volcanic vent distribution in fields has a wide variation across the solar system
- These tools can be used to gain knowledge about volcanic fields on Mars with spaceborne instruments

