

HITTING TIME AND THE KEMENY CONSTANT

SONGTAO MAO

ABSTRACT. In this paper, we prove that the hitting time cannot be controlled by the Kemeny constant.

Key words. Markov chain, hitting time, Kemeny constant

First, Construct a complete graph K_n with n vertices $\{a_1, a_2, \dots, a_n\}$, with each edge having the weight 1, and let a_0 be an added vertices such that it is only connected with a_1 and the edge connecting a_0 and a_1 has the weight w .

This weighted graph can naturally induce a Markov chain with each vertex as the state, where

$$p_{ij} = \frac{w(i, j)}{\sum_i w(i, j)}$$

We will prove that the hitting time from $a_i (i \geq 1)$ cannot be controlled by the Kemeny constant by the following two claim:

Denote $E_i(\tau_j) := E[\inf\{t \geq 0 : X(t) = a_j | X(0) = a_i\}]$, $E_i(\tau_j^+) := E[\inf\{t > 0 : X(t) = a_j | X(0) = a_i\}]$ First:

$$\lim_{w \rightarrow 0} E_1(\tau_0) = \infty$$

It is clear since $E_0(\tau_0^+) = E_1(\tau_0) + 1$ and $E_0(\tau_0^+) = \frac{1}{\pi(0)} \rightarrow 0$ as $w \rightarrow 0$.

However, the kemeny constant can be controlled by a constant when $w < 1$:

Denote

$$\begin{aligned} K &= \sum_{i \neq 0} \pi(i) E_0(\tau_i) = \sum_{i \neq 0} \pi(i) E_1(\tau_i) + 1 \\ &= \sum_{i \neq 0, 1} \pi(i) E_1(\tau_i) + 1 < E_1(\tau_2) + 1 \end{aligned}$$

The last inequality holds since the symmetry of the states a_2, \dots, a_n and $\sum_{i \neq 0,1} \pi(i) < 1$

However, $E_1(\tau_2)$ is an increasing function of t , and thus is controlled by the case when $w < 1$.

So in this example, the hitting time cannot be controlled by the Kemeny constant.

However, when we fix $t = 1$, we find that the hitting time can be controlled by the product of $n + 1$ and Kemeny constant by the following data, and further, we find that

$$\lim K(n+1) - n^2 = 2$$

and this formula is verified by progressive analysis in the paper submitted by Qi Zhou.

n	K	K(n+1)	t_{hit}
1	0.5	1	1
2	1.5	4.5	5
3	2.5417	10.167	9
4	3.5	17.5	16
5	4.4455	26.673	25
10	9.2652	101.92	100
20	19.142	401.98	400
50	49.059	2502	2500
100	99.028	10002	10000
300	299.01	90002	90000

Conjecture: In an unweighted graph with n vertices, the following relationship exists:

$$\max_{i,j} E_i(\tau_j^+) \leq nK$$

where K is the kemeny constant of this graph. This result can be obtained through a lot of computer practice, but the author has not given proof here.

However, I think it can be divided into the following steps to prove the conclusion:

In a graph with n vertices, fix one vertex V_0 and look into the following function $f(w(i, j) : 1 \leq i \neq j \leq n)$:

$$f = \frac{\max_{i,j} E_i(\tau_j^+)}{K}$$

this is a function of all weight of edges, which is 0 or 1, and to prove that this function monotonically increases as $w(0, i)$, and monotonously decreases as $w(i, j)(i, j \neq 0)$.

SCHOOL OF MATHEMATICAL SCIENCES, SHANGHAI JIAO TONG UNIVERSITY, 800 DONGCHUAN ROAD, 200240 SHANGHAI, CHINA

Email address: jarlly678@126.com