

Progress Report

Gillian Queisser (PI), Stephan Grein, James Rosado

Synaptic Spine Calcium Dynamics

We would like to note that the manuscript *Calcium modeling of spine apparatus containing human dendritic spines demonstrates all-or-nothing communication switch between the spine head and dendrite* which was supported by the current funding period has been submitted to a peer-reviewed journal and in this manuscript we state and explain the results summarized in this progress report.

The system of diffusion equations [2] for modeling calcium dynamics is given by:

$$\frac{\partial c_c}{\partial t} = D_c \Delta c_c + (\kappa_b^- (b^{tot} - b) - \kappa_b^+ b c_c), \quad (1)$$

$$\frac{\partial b}{\partial t} = D_b \Delta b + (\kappa_b^- (b^{tot} - b) - \kappa_b^+ b c_c), \quad (2)$$

$$\frac{\partial c_e}{\partial t} = D_e \Delta c_e, \quad (3)$$

where c_c is the cytosolic calcium and c_e is calcium in the ER, (b^{tot}) is the total concentration of CalB present in the cytosol, and κ_b^- , κ_b^+ are rate constants. We also take into account influxes at the PSD, calcium exchange mechanisms located on the ER membrane and PM, the complete set of equations is described in (cite Breit, Queisser Paper). We first established a baseline for the level of calcium concentration in the cytosol when there is no ER inside the cytosol region of the geometry. The baseline simulations were executed using the 9 spine-dendrite geometries, Figure 1a, row A, with no endoplasmic reticular (ER) and with passive ER, and a calcium influx was specified at the PSD for each spine. Both sets of baseline simulations revealed that passive/absent spine apparatus organelles (the ER) have no major impact on spine-to-dendrite calcium signaling. In all cases less than 3% of the initial calcium concentration amplitude in the head was detectable in the dendrite. When the ER was added as a passive compartment (i.e., when it was present as a geometric obstacle, but without any calcium exchange mechanisms), similar and near-zero dendritic calcium concentration profiles are observed in the respective spines Figure 1a, row B. Therefore, calcium influx at the PSD alone without intracellular calcium dynamics from the ER is not enough to induce spine-to-dendrite calcium coupling; in particular, only a small fraction of released Ca^{2+} ions reaches the dendritic compartment. However, with active ER there exists a critical RyR density where by we have significant calcium signaling in the dendritic region Figure 1a, row C.

We determined that when the ER is active with RyRs and SERCA pumps there is a critical RyR density $\rho_{RyRcrit}$, Figure 1b-A, such that if $\rho_{RyR} < \rho_{RyRcrit}$ then there is no spine-dendrite calcium signaling and if




Figure 1: In (a) we show the types of spine geometries and the calcium profiles. (b) shows a vertical *jump* in the concentration of the dendritic cytosol. (b) demonstrates that the critical RyR density is invariant under changes in synaptic calcium influx and calcium buffering. Figure (b)-A is full buffering of calcium and Figure (b)-B is half buffering of calcium.

$\rho_{RyR} \geq \rho_{RyRcrit}$ there is spine-to-dendrite calcium signaling. This was carried out by running a sequence of simulations where the RyR density parameters was increased starting from $0.00 \mu m^{-2}$ to $5.00 \mu m^{-2}$ at increments of $0.01 \mu m^{-2}$. We tested whether changes in Ca^{2+} entry at the synapse affect the critical RyR density by successively increasing the Ca^{2+} flux density at the PSD, changes in the level of calcium buffering present in the cytosol (Figure 1b-A is full buffering, Figure 1b-B is half buffering) and the simulations demonstrate that calcium coupling occurs at the same RyR density, regardless of the initial Ca^{2+} influx and changes in calcium buffering, Figure 1b-B. Therefore, ryanodine receptors of spine apparatus control spine-to-dendrite Ca^{2+} signaling. We also carried out simulations by which we varied the neck diameter of the spine morphologies, and we observed a positive correlation between spine and SA cross-sectional area of the spine neck. This suggests that the ER adjusts to dendritic spine growth and reorganizes in such a way to support spine-to-dendrite Ca^{2+} communication. In fact, previous work has demonstrated that changes in spine neck diameters accompany the induction of excitatory synaptic plasticity, reflected by an increase in spine head size [1, 5, 4]. Simulations are ongoing and below in Table 1 we have listed our current usage (as of submission of the renewal request) and tentative usage for the remaining 20% of project duration.

Synaptic Spine Calcium Dynamics		Full Cell Calcium Dynamics	
Simulation	SU	Simulation	SU
Baseline Runs	13,188	Synaptic strength Runs	128,052
RyR Runs	20,300	Synapse loss Runs	13,426
Pulse Runs	12,706	Stimulation Region Runs	28,196
VDCC Runs	207,710		
Tentative Usage			
Simulation	SU	Simulation	SU
Pulse Runs	100,000	Synapse Loss	120,000
VDCC Runs	80,000	Stimulation Regions	120,000

Table 1: SU usage according to project simulations.

We are currently running simulations, **TMS runs**, incorporating voltage dependent calcium channels (VDCCs) to determine under what conditions we have strong spine-to-dendrite calcium signalling and calcium wave propagation through the dendrite. Current simulation results demonstrate that having either PSD calcium influx or VDCC release is sufficient for calcium signaling to the dendrite; however, having each alone is not enough to produce stable calcium wave propagation.

Whole Cell Dynamics Progress

For the subproject **Whole Cell Dynamics** under investigation, the authors would like to note that two manuscript are currently in preparation resulting from a recently completed PhD dissertation [3] and supported by simulations on the XSEDE SDSC Comet cluster discussing the meshing pipeline and neurobiological implications of intracellular calcium waves - an excerpt of the results to be published will be given in the report below.

To determine a threshold for the number of synapses required to generate stable intracellular calcium waves as a communication device from distal (neurite tips) to proximal regions (soma) a parameter variation of synaptic activation strength was conducted as proposed in the original allocation request, see Tab. 1. Specifically, a variation of synaptic activation strength has been conducted, in which the low and high synaptic strength has been used for the activation of synapses. While strong synapses reliably trigger intracellular calcium waves, we determined a threshold for the number of weak synapses with a fully synchronous activation pattern as depicted in Fig. 2 and proposed previously. Based on this calibration, synapse loss studies are currently being conducted on 6 representative cells, as well as a comparison of 10 morphologically distinct neurons, to investigate the effects of a variation of spatial stimulation as presented in the proposal. For this, the remaining budget of the current allocation will be used (see Tab. 1 for a summary of the estimated number of SUs in the remaining compute period).

Preliminary studies on synaptic strength are being combined with synapse loss studies and a variation of stimulation regions on various neurons. For this the remaining budget of service units on SDSC Comet and Expanse will be used to enable and support the finalizing of the two manuscripts in preparation.




Figure 2: Synapse density threshold experiment: A stable intracellular calcium wave could be elicited by stimulation of the input region (distal dendrites) with $n = 100$ synapses on a cell from the Rice-Baylor archive (entorhinal cortex). Vertical green bar indicates end of simulation time (50 ms).

Explanation for under utilization of research allocation

In the original proposal, the Synaptic Spine calcium subproject requested 270,000 SUs for RyR runs, and 607,500 SUs for the Calcium pulse runs; however, we decreased the number of runs in order to accommodate the 800,000 SUs we were granted. Overall, the international consortium for TMS-research was slowed down, due to COVID-related restrictions on travel and joint in-person research. We also observed that not all spine simulations required equal amounts of simulation wall-time, this is determined by the complexity of the Spine-ER geometry and the density of RyRs on the ER membrane. In order to study the full level of calcium dynamics in the cytosol, we incorporated the VDCC simulations (**TMS runs** referenced in the main document) prior to continuing our runs of Calcium Pulse simulations. We anticipate completing our study of the **TMS** simulation results first, which will use most of the remaining SUs. Once the **TMS** simulations are complete, then we will execute our Calcium Pulse simulations which we anticipate using a minimum of 100,000 SUs, either from the new allocation or use any remaining SUs. Likewise the Whole Cell Calcium Dynamics project used nearly the 150,000 SUs which have been requested in the original proposal for a systematic evaluation of the consequences of synaptic strength on intracellular calcium waves and calcium signaling. During these studies it has been recognized that runtime for simulations can be reduced depending on the stimulation setup, thus less SUs had been required contrary to our expectation. Still the remaining 350,000 service units will be required to complete the tasks of systematic studies on synapse location on realistic cell geometries and synapse loss studies in this subproject. The available SUs will be manually load-balanced in the next two month between the subprojects depending on the priority of simulations for the progression of the projects' research, see tentative usage in Tab. 1 above.

References

- [1] R. Araya, T. P. Vogels, and R. Yuste. Activity-dependent dendritic spine neck changes are correlated with synaptic strength. *Proceedings of the National Academy of Sciences*, 111, 7 2014.
- [2] Markus Breit, Marcus Kessler, Martin Stepniewski, Andreas Vlachos, and Gillian Queisser. Spine-to-dendrite calcium modeling discloses relevance for precise positioning of ryanodine receptor-containing spine endoplasmic reticulum. *Scientific Reports*, 8, 12 2018.
- [3] S. Grein. *A novel mesh generator for the numerical simulation of multi-scale physics in neurons*. PhD thesis, Temple University, 2020.
- [4] Heinz Steffens, Alexander C. Mott, Siyuan Li, Waja Wegner, Pavel Švehla, Vanessa W. Y. Kan, Fred Wolf, Sabine Liebscher, and Katrin I. Willig. Stable but not rigid: Chronic in vivo sted nanoscopy reveals extensive remodeling of spines, indicating multiple drivers of plasticity. *Science Advances*, 7, 6 2021.
- [5] Jan Tønnesen, Gergely Katona, Balázs Rózsa, and U Valentin Nägerl. Spine neck plasticity regulates compartmentalization of synapses. *Nature Neuroscience*, 17, 5 2014.