

Two newly recorded hypogeous fungi in Taiwan—*Endogone flammicorona* and *Sclerogaster compactus*

King-Fai Wong^{1,2}, Ming-Jer Tsai^{3,4}, Chuen-Hsu Fu⁵, Chieh-Lung Lin⁶, Hoi-Tung Li^{1,7*}

¹Department of Forestry and Resource Conservation, National Taiwan University. No.1, Sec.4, Roosevelt Rd., Taipei 10617, Taiwan

²Tianroei limited company. No.151, Jingshan Rd., Shihlin Dist., Taipei City 11192, Taiwan

³Department of Forestry and Resource Conservation, National Taiwan University. No.1, Sec.4, Roosevelt Rd., Taipei 10617, Taiwan

⁴The Experimental Forest, College of Bio-Resources and Agriculture, National Taiwan University. No.12, Sec. 1, Qianshan Rd., Zhushan Township, Nantou County 55750, Taiwan

⁵Division of Forest Protection, Taiwan Forestry Research Institute, COA. No.67, Sanyuan St., Zhongzheng Dist., Taipei City 10079, Taiwan

⁶Watershed Management Division, Taiwan Forestry Research Institute, COA. No.53, Nan-Hai Rd., Taipei 10066, Taiwan

⁷Advance plant protection limited company. No.21, Ln. 230, Sec. 1, Wufu Rd., Siangshan Dist., Hsinchu City 30095, Taiwan

(Accepted: December 30, 2017)

ABSTRACT

Two newly recorded species of hypogeous fungi, *Endogone flammicorona* (Zygomycota) and *Sclerogaster compactus* (Basidiomycota), were collected from plantation of *Picea morrisonicola* along Dasyueshan Trail in central Taiwan, at the altitude of 2500 m.

Key words: Basidiomycota, false truffles, pea truffles, *Picea morrisonicola*, Zygomycota

Introduction

Hypogeous fungi produce fruiting bodies among layers of leaf-litters underground and do not actively disperse spores, being spread passively by their animal vectors. They are classified in three phyla, with true truffles belonging to Ascomycota, false truffles to Basidiomycota, and pea truffles to Zygomycota. *Tuber* species in the Ascomycota are well-known as gourmet hypogeous fungi, while other hypogeous fungi remain

unknown for their edibility. In Taiwan, four species of basidiomycete and zygomycete hypogeous truffles have been reported previously: *Rhizopogon rubescens* (Tul. & C. Tul.) Tul. & C. Tul., *Hymenogaster arenarius* Tul. & C. Tul., *Descomyces albellus* (Masse & Rodway) Bougher & Castellano, and *Gymnoglossum connectens* (Bucholtz) Zeller (Hu, 2002; Chou, 2011, 2016). In this study, we reported two newly recorded hypogeous fungi collected from plantation of *Picea morrisonicola* Hayata in Taiwan; *Endogone flammicorona* Trappe & Gerdem. is a

*Corresponding author, e-mail: doralie614@gmail.com




Fig. 1. *Endogone flammicorona*. a. Sporocarp. b. Yellowish-orange gleba. c. Zygospores surrounded by agglutinated hyphae. d. Fingerprint-like ornamentation on zygospores.

pea truffle and *Sclerogaster compactus* (Tul. & C. Tul.) Sacc. is a false truffle.

Taxonomy

Endogone flammicorona Trappe & Gerdem.,
Trans. Brit. Mycol. Soc. 59: 405 (1972) Fig. 1

Sporocarp hypogeous, irregularly subglobose, ca 1 cm diam. Peridium smooth, thin, pale-brown. Gleba yellowish-orange, densely packed. Zygosporangia not observed. Zygospores subglobose or ovoid, yellowish-orange, $75\text{--}90 \times 87.5\text{--}97.5$ μm , surrounded by agglutinated and condensed

hyphal layer 15–17.5 µm thick, ornamented with conspicuous looped and pseudohelical ridges.

Specimen examined: TAIWAN. Taichung County, Dasyueshan Trail, 2500 m elev., beneath piled leaves under plantation of *Picea morrissonicola*, 8 Aug 2016, Lin, C.-L.

Distribution: Italy, Taiwan, UK, and USA.

Remarks: *Endogone flammicorona* is similar to *E. lactiflua* Berk. & Br., from which it differs by the significant fingerprint-like spore ornamentation with the appearance of tongues of flame




Fig.2. *Sclerogaster compactus*. a. Gastrocarp. b, c. Yellowish-brown to yellowish-green gleba with white veins. d. Tiny verrucous basidiospores.

(Trappe and Gerdemann, 1972). This species has been reported to form an ectomycorrhizal association with Pinaceae (Gerdemann and Trappe, 1974), and were found in humus associated with both coniferous and deciduous trees in the UK (Pegler et al., 1993).

Sclerogaster compactus (Tul. & C. Tul.) Sacc., Syll. Fung. 11: 170 (1895) Fig. 2
 ≡ *Octaviania compacta* Tul. & C. Tul., Giorn. Bot. Ital., Ann. 1, 2(1): 56 (1844)

Gastrocarp hypogeous, irregularly globose, small, less than 1 cm diam. Peridium smooth,

white to milky yellow. Gleba dense, yellowish-brown to yellowish-green, with white veins. Basidia not observed. Basidiospores pale green to pale yellow, $7.5 \times 5 \mu\text{m}$, mostly ovoid, with thickened wall ornamented with tiny verrucae.

Specimen examined: TAIWAN. Taichung County, Dasyueshan Trail, 2500 m elev., beneath piled leaves under plantation of *Picea morrissonicola*, 8 Aug 2016, Lin, C.-L.

Distribution: France, Germany, Italy, Taiwan, and UK.

Remarks: *Sclerogaster compactus* is distinct in having a small and compact gastrocarp and tiny verrucous basidiospores. It was found in leaf litter or decomposing organic layer, likely not forming a mycorrhizal association with plant roots.

References

- Chou WN. 2011. *Hymenogaster arenarius* (Basidiomycota) newly recorded to Taiwan. Taiwan Journal of Biodiversity 13(3): 265–268.
- Chou WN. 2016. Two species of hypogeous gasteroid fungi (Basidiomycota) new to Taiwan. Taiwan Journal of Biodiversity 18(3): 169–172.
- Gerdemann JW, Trappe JM. 1974. The Endogonaceae in the Pacific Northwest. Mycologia Memior 5: 1–76.
- Hu HT. 2002. *Glomus spinosum* sp. nov. in the Glomaceae from Taiwan. Mycotaxon 83: 159–164.
- Korf RP. 1972. Synoptic key to the genera of the Pezizales. Mycologia 64(5): 937–994.
- Montecchi A, Sarasini M. 2000. Funghi ipogei d'Europa. Associazione Micologica Bresadola, Italy. 714 p.
- Pegler DN, Spooner BM, Young TWK. 1993. British truffles: A revision of British hypogeous fungi. The Royal Botanic Gardens, Kew. 216 p.
- Trappe JM, Gerdemann JW. 1972. *Endogone flammicorona* sp. nov., a distinctive segregate from *Endogone lactiflua*. Transactions of the British Mycological Society 59(3): 403–407.

兩種臺灣新記錄地下真菌—火冠豆松露與緻密硬皮腹菌

黃勁暉^{1,2}、蔡明哲^{3,4}、傅春旭⁵、林介龍⁶、李鎧彤^{1,7*}

¹ 國立臺灣大學森林環境暨資源學研究所

² 天蕊股份有限公司

³ 國立臺灣大學森林環境暨資源學系

⁴ 國立臺灣大學實驗林管理處

⁵ 行政院農業委員會林業試驗所森林保護組

⁶ 行政院農業委員會林業試驗所集水區經營組

⁷ 前衛植保有限公司

摘要

本文描述兩種臺灣新記錄地下真菌，分別為接合菌門的火冠豆松露（*Endogone flammicorona*）與擔子菌門的緻密硬皮腹菌（*Sclerogaster compactus*），皆採自臺灣中部大雪山林道、海拔 2500 m 的臺灣雲杉人工林。

關鍵詞： 擔子菌、假松露、接合菌、豆松露、臺灣雲杉