

JANGAON, ESTELLA

DOB: 04/09/1963
Sex: F
Phone: (713) 294-2551
Patient ID: 19288

Age: 59
Fasting: Y

Specimen: HZ663130C
Requisition: 0019758
Lab Reference ID: 330150
Report Status: FINAL / SEE REPORT

Collected: 11/30/2022 09:34
Received: 11/30/2022 09:36
Reported: 12/02/2022 15:36

Client #: 4463400
KUMAR, ARUN
KUMAR, ARUN MD
10726 HUFFMEISTER RD STE 100
HOUSTON, TX 77065-3181
Phone: (281) 477-0666
Fax: (281) 477-0577

FASTING: YES

▲ COMPREHENSIVE METABOLIC PANEL

Analyte	Value	
▲ GLUCOSE	101 H	Reference Range: 65-99 mg/dL
Fasting reference interval		
For someone without known diabetes, a glucose value between 100 and 125 mg/dL is consistent with prediabetes and should be confirmed with a follow-up test.		
UREA NITROGEN (BUN)	12	Reference Range: 7-25 mg/dL
CREATININE	0.65	Reference Range: 0.50-1.03 mg/dL
EGFR	101	Reference Range: > OR = 60 mL/min/1.73m2
The eGFR is based on the CKD-EPI 2021 equation. To calculate the new eGFR from a previous Creatinine or Cystatin C result, go to https://www.kidney.org/professionals/kdoqi/gfr%5Fcalculator		
BUN/CREATININE RATIO	NOT APPLICABLE	Reference Range: 6-22 (calc)
SODIUM	142	Reference Range: 135-146 mmol/L
POTASSIUM	4.6	Reference Range: 3.5-5.3 mmol/L
CHLORIDE	104	Reference Range: 98-110 mmol/L
CARBON DIOXIDE	28	Reference Range: 20-32 mmol/L
CALCIUM	9.6	Reference Range: 8.6-10.4 mg/dL
PROTEIN, TOTAL	7.1	Reference Range: 6.1-8.1 g/dL
ALBUMIN	4.2	Reference Range: 3.6-5.1 g/dL
GLOBULIN	2.9	Reference Range: 1.9-3.7 g/dL (calc)
ALBUMIN/GLOBULIN RATIO	1.4	Reference Range: 1.0-2.5 (calc)
BILIRUBIN, TOTAL	0.8	Reference Range: 0.2-1.2 mg/dL
ALKALINE PHOSPHATASE	60	Reference Range: 37-153 U/L
AST	14	Reference Range: 10-35 U/L
ALT	11	Reference Range: 6-29 U/L

▲ URINALYSIS, COMPLETE

Analyte	Value	
COLOR	YELLOW	Reference Range: YELLOW
APPEARANCE	CLEAR	Reference Range: CLEAR
SPECIFIC GRAVITY	1.012	Reference Range: 1.001-1.035

PH	6.0	Reference Range: 5.0-8.0
GLUCOSE	NEGATIVE	Reference Range: NEGATIVE
BILIRUBIN	NEGATIVE	Reference Range: NEGATIVE
KETONES	NEGATIVE	Reference Range: NEGATIVE
▲ OCCULT BLOOD	TRACE	Reference Range: NEGATIVE
PROTEIN	NEGATIVE	Reference Range: NEGATIVE
NITRITE	NEGATIVE	Reference Range: NEGATIVE
▲ LEUKOCYTE ESTERASE	TRACE	Reference Range: NEGATIVE
WBC	0-5	Reference Range: < OR = 5 /HPF
RBC	NONE SEEN	Reference Range: < OR = 2 /HPF
SQUAMOUS EPITHELIAL CELLS	NONE SEEN	Reference Range: < OR = 5 /HPF
BACTERIA	NONE SEEN	Reference Range: NONE SEEN /HPF
HYALINE CAST	NONE SEEN	Reference Range: NONE SEEN /LPF

NOTE

This urine was analyzed for the presence of WBC, RBC, bacteria, casts, and other formed elements. Only those elements seen were reported.

▲ HEMOGLOBIN A1c

Analyte	Value
▲ HEMOGLOBIN A1c For someone without known diabetes, a hemoglobin A1c value between 5.7% and 6.4% is consistent with prediabetes and should be confirmed with a follow-up test. For someone with known diabetes, a value <7% indicates that their diabetes is well controlled. A1c targets should be individualized based on duration of diabetes, age, comorbid conditions, and other considerations. This assay result is consistent with an increased risk of diabetes. Currently, no consensus exists regarding use of hemoglobin A1c for diagnosis of diabetes for children.	5.9 H Reference Range: <5.7 % of total Hgb

CARDIO IQ® VITAMIN D, 25 HYDROXY

Analyte	Value
VITAMIN D, 25-OH, TOTAL (Note) Vitamin D, 25-Hydroxy reports concentrations of two common forms, 25-OHD2 and 25-OHD3. 25-OHD3 indicates both endogenous production and supplementation. 25-OHD2 is an indicator of exogenous sources such as diet or supplementation. Therapy is based on measurement of Total 25-OHD, with levels <20 ng/mL indicative of Vitamin D deficiency, while levels between 20 ng/mL and 30 ng/mL suggest insufficiency. Optimal levels are > or = 30 ng/mL. Vitamin D is fat-soluble and therefore inadvertent or intentional ingestion of excessively high amounts could be toxic. Studies in children and adults suggest blood levels would need to exceed 150 ng/mL before there is any concern. Holick MF, Binkley NC, Bischoff-ferrari HA, et al. Evaluation, treatment and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. J Clin Endocrinol Metab. 2011;96(7):1911-30. For additional information, please refer to http://education.QuestDiagnostics.com/faq/FAQ199	24 ng/mL

VITAMIN D, 25-OH, D3	16 ng/mL
Reference range: Not established	

VITAMIN D, 25-OH, D2	8 ng/mL
(Note) Reference range: Not established	

This test was developed and its analytical performance characteristics have been determined by medfusion. It has not been cleared or approved by the US Food and Drug Administration. This assay has been validated pursuant to the CLIA regulation and is used for Clinical purposes. MDF med fusion 2501 South State Highway 121, Suite 1100 Lewisville TX 75067 972-966-7300 Michael Chaump, MD

COMMENT

See Note 1

Note 1

For additional information, please refer to <http://education.QuestDiagnostics.com/faq/FAQ199> (This link is being provided for informational/ educational purposes only.)

IRON, TIBC AND FERRITIN PANEL

Analyte	Value
FERRITIN	148 Reference Range: 16-232 ng/mL

IRON AND TOTAL IRON BINDING CAPACITY

Analyte	Value
IRON, TOTAL	82 Reference Range: 45-160 mcg/dL
IRON BINDING CAPACITY	277 Reference Range: 250-450 mcg/dL (calc)
% SATURATION	30 Reference Range: 16-45 % (calc)

LIPID PANEL, STANDARD

Analyte	Value
CHOLESTEROL, TOTAL	151 Reference Range: <200 mg/dL
HDL CHOLESTEROL	53 Reference Range: > OR = 50 mg/dL
TRIGLYCERIDES	72 Reference Range: <150 mg/dL
LDL-CHOLESTEROL	83 mg/dL (calc)
Reference range: <100	
Desirable range <100 mg/dL for primary prevention; <70 mg/dL for patients with CHD or diabetic patients with > or = 2 CHD risk factors.	
LDL-C is now calculated using the Martin-Hopkins calculation, which is a validated novel method providing better accuracy than the Friedewald equation in the estimation of LDL-C. Martin SS et al. JAMA. 2013;310(19): 2061-2068 (http://education.QuestDiagnostics.com/faq/FAQ164)	
CHOL/HDL-C RATIO	2.8 Reference Range: <5.0 (calc)
NON HDL CHOLESTEROL	98 Reference Range: <130 mg/dL (calc)
For patients with diabetes plus 1 major ASCVD risk factor, treating to a non-HDL-C goal of <100 mg/dL (LDL-C of <70 mg/dL) is considered a therapeutic option.	

ALBUMIN, RANDOM URINE W/CREATININE

Analyte	Value
JANGAON, ESTELLA (HZ663130C)	3 / 5

12/2/22

CREATININE, RANDOM URINE**79** Reference Range: 20-275 mg/dL**MICROALBUMIN, RANDOM URINE (W/CREATININE)**

Analyte	Value
ALBUMIN, URINE	1.0 Reference Range: Reference Range mg/dL Not established mg/dL
ALBUMIN/CREATININE RATIO, RANDOM URINE	13 Reference Range: <30 mcg/mg creat
The ADA defines abnormalities in albumin excretion as follows:	
Albuminuria Category	Result (mcg/mg creatinine)
Normal to Mildly increased	<30
Moderately increased	30-299
Severely increased	> OR = 300
The ADA recommends that at least two of three specimens collected within a 3-6 month period be abnormal before considering a patient to be within a diagnostic category.	

URIC ACID

Analyte	Value
URIC ACID	4.2 Reference Range: 2.5-7.0 mg/dL
Therapeutic target for gout patients: <6.0 mg/dL	

CBC (INCLUDES DIFF/PLT)

Analyte	Value
WHITE BLOOD CELL COUNT	5.6 Reference Range: 3.8-10.8 Thousand/uL
RED BLOOD CELL COUNT	4.04 Reference Range: 3.80-5.10 Million/uL
HEMOGLOBIN	12.2 Reference Range: 11.7-15.5 g/dL
HEMATOCRIT	36.8 Reference Range: 35.0-45.0 %
MCV	91.1 Reference Range: 80.0-100.0 fL
MCH	30.2 Reference Range: 27.0-33.0 pg
MCHC	33.2 Reference Range: 32.0-36.0 g/dL
RDW	13.3 Reference Range: 11.0-15.0 %
PLATELET COUNT	271 Reference Range: 140-400 Thousand/uL
MPV	11.1 Reference Range: 7.5-12.5 fL
ABSOLUTE NEUTROPHILS	3315 Reference Range: 1500-7800 cells/uL
ABSOLUTE LYMPHOCYTES	1730 Reference Range: 850-3900 cells/uL
ABSOLUTE MONOCYTES	403 Reference Range: 200-950 cells/uL
ABSOLUTE EOSINOPHILS	101 Reference Range: 15-500 cells/uL
ABSOLUTE BASOPHILS	50 Reference Range: 0-200 cells/uL
NEUTROPHILS	59.2 %
LYMPHOCYTES	30.9 %
MONOCYTES	7.2 %
EOSINOPHILS	1.8 %

BASOPHILS**0.9** %**T4, FREE****Analyte****Value****T4, FREE****1.3** Reference Range: 0.8-1.8 ng/dL**TSH****Analyte****Value****TSH****1.20** Reference Range: 0.40-4.50 mIU/L**VITAMIN B12/FOLATE, SERUM PANEL****Analyte****Value****VITAMIN B12****287** Reference Range: 200-1100 pg/mL

Please Note: Although the reference range for vitamin B12 is 200-1100 pg/mL, it has been reported that between 5 and 10% of patients with values between 200 and 400 pg/mL may experience neuropsychiatric and hematologic abnormalities due to occult B12 deficiency; less than 1% of patients with values above 400 pg/mL will have symptoms.

FOLATE, SERUM**8.5** ng/mL

Reference Range

Low: <3.4

Borderline: 3.4-5.4

Normal: >5.4

Performing Sites

RGA Quest Diagnostics-Houston Lab, 5850 Rogerdale Road, Houston, TX 77072-1602 Laboratory Director: Robert L Breckenridge

Z3E MedFusion-MedFusion, 2501 South State Highway 121, Suite 1100, Lewisville, TX 75067-8188 Laboratory Director: Michael Chaump MD

Key Priority Out of Range  Out of Range

These results have been sent to the person who ordered the tests. Your receipt of these results should not be viewed as medical advice and is not meant to replace discussion with your doctor or other healthcare professional.

Quest, Quest Diagnostics, the associated logo, Nichols Institute, Interactive Insights and all associated Quest Diagnostics marks are the registered trademarks of Quest Diagnostics. All third party marks - '®' and '™' - are the property of their respective owners. Privacy policy can be found at: <http://questdiagnostics.com/home/privacy-policy/online-privacy.html>. © 2022 Quest Diagnostics Incorporated. All rights reserved.



These results have been sent to the person that ordered the tests. Your receipt of these results should not be viewed as medical advice and is not meant to replace discussion with your doctor or other healthcare professional.