

JANGAON, ESTELLA

DOB: 04/09/1963
Sex: F
Phone: (713) 294-2551
Patient ID: 85233759

Age: 62
Fasting: Y

Specimen: HZ287940Q
Requisition: 0029122
Lab Reference ID: 278562731Q
Report Status: FINAL / SEE REPORT

Collected: 09/22/2025 10:51
Received: 09/22/2025 10:52
Reported: 09/27/2025 20:44

Client #: 4463400
HOANG, RACHEL
KUMAR, ARUN MD
10726 HUFFMEISTER RD STE 100
HOUSTON, TX 77065-3181
Phone: (281) 477-0666
Fax: (281) 477-0577

FASTING: YES

▲ COMPREHENSIVE METABOLIC PANEL

Analyte	Value	
▲ GLUCOSE	106 H	Reference Range: 65-99 mg/dL
Fasting reference interval		
For someone without known diabetes, a glucose value between 100 and 125 mg/dL is consistent with prediabetes and should be confirmed with a follow-up test.		
UREA NITROGEN (BUN)	17	Reference Range: 7-25 mg/dL
CREATININE	0.73	Reference Range: 0.50-1.05 mg/dL
EGFR	93	Reference Range: > OR = 60 mL/min/1.73m2
BUN/CREATININE RATIO	SEE NOTE:	Reference Range: 6-22 (calc)
Not Reported: BUN and Creatinine are within reference range.		
SODIUM	139	Reference Range: 135-146 mmol/L
POTASSIUM	4.0	Reference Range: 3.5-5.3 mmol/L
CHLORIDE	103	Reference Range: 98-110 mmol/L
CARBON DIOXIDE	29	Reference Range: 20-32 mmol/L
CALCIUM	9.5	Reference Range: 8.6-10.4 mg/dL
PROTEIN, TOTAL	7.3	Reference Range: 6.1-8.1 g/dL
ALBUMIN	4.3	Reference Range: 3.6-5.1 g/dL
GLOBULIN	3.0	Reference Range: 1.9-3.7 g/dL (calc)
ALBUMIN/GLOBULIN RATIO	1.4	Reference Range: 1.0-2.5 (calc)
BILIRUBIN, TOTAL	0.5	Reference Range: 0.2-1.2 mg/dL
ALKALINE PHOSPHATASE	61	Reference Range: 37-153 U/L
AST	16	Reference Range: 10-35 U/L
ALT	11	Reference Range: 6-29 U/L

▲ ANALYZER(TM)ANA, IFA WITH REFLEX TITER/PATTERN, SYSTEMIC AUTOIMMUNE PANEL 1

Analyte	Value
---------	-------

▲ ANA SCREEN, IFA**POSITIVE**

Reference Range: NEGATIVE

ANA IFA is a first line screen for detecting the presence of up to approximately 150 autoantibodies in various autoimmune diseases. A positive ANA IFA result is suggestive of autoimmune disease and reflexes to titer and pattern. Further laboratory testing may be considered if clinically indicated.

For additional information, please refer to <http://education.questdiagnostics.com/faq/FAQ177> (This link is being provided for informational/educational purposes only.)

DNA AB (DS) CRITHIDIA, IFA	NEGATIVE	Reference Range: NEGATIVE
CHROMATIN (NUCLEOSOMAL) ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
SM ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
SM/RNP ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
RNP ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
SJOGREN'S ANTIBODY (SS-A)	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
SJOGREN'S ANTIBODY (SS-B)	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
SCL-70 ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
JO-1 ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
CENTROMERE B ANTIBODY	<1.0 NEG	Reference Range: <1.0 NEGATIVE AI
COMPLEMENT COMPONENT C3C	149	Reference Range: 83-193 mg/dL
COMPLEMENT COMPONENT C4C	30	Reference Range: 15-57 mg/dL
CARDIOLIPIN AB (IGA)	<2.0	APL-U/mL See Note 1
CARDIOLIPIN AB (IGG)	<2.0	GPL-U/mL See Note 1
CARDIOLIPIN AB (IGM)	<2.0	MPL-U/mL See Note 1
B2 GLYCOPROTEIN I (IGA)AB	<2.0	U/mL See Note 2
B2 GLYCOPROTEIN I (IGG)AB	<2.0	U/mL See Note 2
B2 GLYCOPROTEIN I (IGM)AB	<2.0	U/mL

The antiphospholipid antibody syndrome (APS) is a clinical-pathologic correlation that includes a clinical event (e.g. arterial or venous thrombosis, pregnancy morbidity) and persistent positive antiphospholipid antibodies (IgM, IgG Cardiolipin or b2GPI antibodies greater than the 99th percentile; or a lupus anticoagulant). International consensus guidelines for APS suggest waiting at least 12 weeks before retesting to confirm antibody persistence. The Systemic Lupus International Collaborating Clinics immunological classification criteria for systemic lupus erythematosus (SLE) include testing for isotype IgA, which has yet to be incorporated into APS criteria. Low level antiphospholipid antibodies may sometimes be detected in the setting of infection, drug therapy or aging.

For additional information, please refer to <http://education.questdiagnostics.com/faq/FAQ109> (This link is being provided for informational/educational purposes only.)

Value	Interpretation
-----	-----
<20.0	Antibody not detected
> or = 20.0	Antibody detected

RHEUMATOID FACTOR (IGA)	<5	U See Note 3
RHEUMATOID FACTOR (IGG)	<5	U See Note 3
RHEUMATOID FACTOR (IGM)	<5	U

Reference Range:

<=6 NEGATIVE

>6 POSITIVE

CYCLIC CITRULLINATED PEPTIDE (CCP) AB (IGG)

<16 Units

Reference Range:

NEGATIVE:

<20

WEAK POSITIVE:

20-39

MODERATE POSITIVE:

40-59

STRONG POSITIVE

>59

MUTATED CITRULLINATED VIMENTIN (MCV) AB

<20 Reference Range: <20 U/mL

Anti-mutated citrullinated vimentin antibody may be used as a second-line marker of rheumatoid arthritis, in addition to rheumatoid factor and anti-cyclic citrullinated peptide (CCP).

THYROID PEROXIDASE ANTIBODIES

<1 Reference Range: <9 IU/mL

▲ ANA, TITER AND PATTERN**Analyte****Value****▲ ANA TITER****1:1280 H** titer

Reference Ranges for Anti-Nuclear Ab Titer:

<1:40 Negative

1:40-1:80 Low Antibody Level

>1:80 Elevated Antibody Level

ANA PATTERN**NUCLEAR, DENSE FINE
SPECKLED**

Dense fine speckled pattern is seen in normal individuals and rarely associated with systemic lupus erythematosus (SLE), Sjogren's syndrome and systemic sclerosis.

AC-2: Dense Fine Speckled International Consensus on ANA Patterns <https://doi.org/10.1515/cclm-2018-0052>

▲ HEMOGLOBIN A1c**Analyte****Value****▲ HEMOGLOBIN A1c****6.1 H** Reference Range: <5.7 %

For someone without known diabetes, a hemoglobin A1c value between 5.7% and 6.4% is consistent with prediabetes and should be confirmed with a follow-up test.

For someone with known diabetes, a value <7% indicates that their diabetes is well controlled. A1c targets should be individualized based on duration of diabetes, age, comorbid conditions, and other considerations.

This assay result is consistent with an increased risk of diabetes.

Currently, no consensus exists regarding use of hemoglobin A1c for diagnosis of diabetes for children.

LIPID PANEL WITH REFLEX TO DIRECT LDL**Analyte****Value****CHOLESTEROL, TOTAL****130** Reference Range: <200 mg/dL**HDL CHOLESTEROL****50** Reference Range: > OR = 50 mg/dL**TRIGLYCERIDES****131** Reference Range: <150 mg/dL

LDL-CHOLESTEROL**59** mg/dL (calc)

Reference range: <100

Desirable range <100 mg/dL for primary prevention;
<70 mg/dL for patients with CHD or diabetic patients
with > or = 2 CHD risk factors.

LDL-C is now calculated using the Martin-Hopkins
calculation, which is a validated novel method providing
better accuracy than the Friedewald equation in the
estimation of LDL-C.

Martin SS et al. JAMA. 2013;310(19): 2061-2068
(<http://education.QuestDiagnostics.com/faq/FAQ164>)

CHOL/HDLC RATIO**2.6** Reference Range: <5.0 (calc)**NON HDL CHOLESTEROL****80** Reference Range: <130 mg/dL (calc)

For patients with diabetes plus 1 major ASCVD risk
factor, treating to a non-HDL-C goal of <100 mg/dL
(LDL-C of <70 mg/dL) is considered a therapeutic
option.

CREATINE KINASE, TOTAL**Analyte****Value****CREATINE KINASE, TOTAL****51** Reference Range: 20-243 U/L

Note 1	Value	Interpretation
	-----	-----
	<20.0	Antibody not detected
	> or = 20.0	Antibody detected

Note 2	Value	Interpretation
	-----	-----
	<20.0	Antibody not detected
	> or = 20.0	Antibody detected

Note 3	Reference Range:
	<=6 NEGATIVE
	>6 POSITIVE

Performing Sites

EZ Quest Diagnostics/Nichols SJC-San Juan Capistrano,, 33608 Ortega Hwy, San Juan Capistrano, CA 92675-2042 Laboratory Director: Irina Maramica MD,PhD,MBA

RGA Quest Diagnostics-Houston Lab, 5850 Rogerdale Road, Houston, TX 77072-1602 Laboratory Director: Meredith A Reyes

Key

 Priority Out of Range  Out of Range

These results have been sent to the person who ordered the tests. Your receipt of these results should not be viewed as medical advice and is not meant to replace discussion with your doctor or other healthcare professional.

Quest, Quest Diagnostics, the associated logo, Nichols Institute, Interactive Insights and all associated Quest Diagnostics marks are the registered trademarks of Quest Diagnostics. All third party marks - '®' and '™' - are the property of their respective owners. Privacy policy can be found at: <http://questdiagnostics.com/home/privacy-policy/online-privacy.html>. © 2022 Quest Diagnostics Incorporated. All rights reserved.



These results have been sent to the person that ordered the tests. Your receipt of these results should not be viewed as medical advice and is not meant to replace discussion with your doctor or other healthcare professional.