

# A Short, Quantitative, Summary of “What Everyone Needs to Know about Climate Change”

Juvid Aryaman, Oct 2018

## 1. Introduction

I am writing this document because it is difficult to find a concise synthesis of evidence supporting the claim that climate change poses (one of) the most important existential risk(s) to humanity. There also exists an abundance of disinformation on the topic, so it is important to find credible sources.

The evidence I have chosen to cite is largely based upon Joe Romm’s recent book [1]. Ref. [1] is a credible source of information on climate change because:

- Joe Romm is a credible expert.
  - PhD in oceanography of the Greenland Sea from MIT, working with Walter Munk.
  - Worked for the U.S. Department of Energy as Acting Assistant Secretary for Energy Efficiency and Renewable Energy for 5 years.
  - Founder of Center for Energy and Climate Solutions, working with corporations to lower greenhouse gas emissions.
  - Founding editor of [ClimateProgress.org](http://ClimateProgress.org).
  - In 2009, *Time* magazine named him “Hero of the Environment” and “The Web’s most influential climate-change blogger”.
  - In 2016, Paul Krugman wrote in the *New York Times* “I’ve learnt a lot of what I know about energy economics from Joe Romm”.
- Oxford University Press is a credible publisher.
- Ref. [1] is essentially an extended review article, and cites primary literature.
- The references in Ref. [1] are almost entirely from well-reputed scientific journals and reports, and are therefore relatively likely to reflect the consensus view of the scientific community.

Ref. [1] is, however, 352 pages long, which presents a barrier to a layperson quickly understanding the importance of climate change. Generation of a set of quantitative claims, with corresponding citations, removes subjectivity from the discussion. There is therefore a need for a set of key, quantitative, claims to be made available which unambiguously state why climate change is of the utmost importance to humanity.

Here, I will attempt to provide such a set of key, quantitative, claims. I will cite primary literature where it was obvious in Ref. [1]. This document is an on-going effort, and suggestions for improvement are most welcome.

## 2. Contributors to climate change

### 2.1. Greenhouse gases

- Satellite data show the Earth emitting less infrared radiation (comparing 1970 to 1997 levels) at the wavenumbers at which greenhouse gases such as CO<sub>2</sub> and CH<sub>4</sub> absorb energy [2]. This results in a significant increase in the amount of longwave downward radiation [3], which has been linked to various anthropogenic greenhouse gases [4]. These studies provide *direct evidence* for the role of anthropogenic greenhouse gases in contributing to climate change (see [5]).

- >90% of all anthropogenic CO<sub>2</sub> comes from burning fossil fuels.
- In 2012, CH<sub>4</sub> contributed up to 9% of greenhouse gases. Major sources of CH<sub>4</sub> are: leaks in fossil fuel extraction, livestock, decaying organic waste, and some agricultural practices.
- Brazil reduced its annual rate of Amazon deforestation by 80% between 2004 and 2013. Deforestation is responsible for ~ 8% of all greenhouse gas emissions [Global Carbon Project].

## 2.2. Permafrost


- Permafrost is soil that stays below freezing for at least 2 years, sinking CO<sub>2</sub> from the carbon cycle. It locks up 1.7 trillion tonnes of carbon [6], which is twice of that in the atmosphere.
- None of the IPCC's climate models include CO<sub>2</sub> or CH<sub>4</sub> emissions from permafrost as a feedback.
- Anthropogenic CH<sub>4</sub> emissions are 0.5 bn tonnes per year, whereas the Siberian permafrost contains 70 bn tonnes. It is uncertain whether this carbon will be released as CH<sub>4</sub> or CO<sub>2</sub>.
- Assuming all permafrost emission as CO<sub>2</sub>, and IPCC scenario A1B [7] (atmospheric CO<sub>2</sub> increases to 700 ppm by 2100, then stays constant at 700 ppm after 2100), permafrost carbon feedback is predicted to change the arctic from a carbon sink to a carbon source after the mid-2020's, and is strong enough to cancel 42-88% of the total global land sink by 2100 [8]
- By 2100, permafrost could add 0.25°C, and possibly as much as 0.8°C [9].

## 2.3. Wildfires

- The “boreal” (subarctic) forests rest on permafrost and peatland which release massive amounts of CO<sub>2</sub> when burned. They store > 30% of all carbon stored on land. Boreal forests now have twice as many wildfires as 500-1000 years ago [10].
- Peat is one of the earliest stages in the process of forming coal and burns easily. The peatland fires in Indonesia during 1997-1998 contributed 13-40% of total CO<sub>2</sub> emissions for that year [11]. Therefore peatland fires have a massive potential to contribute to carbon emissions.
- When peatland dries out, wildfires become more severe [12]. Indonesia has drained a great deal of its peatlands, and burned forested areas, to create palm oil plantations.
- Assuming the Amazonian dry season length (DSL) were to increase at half of the rates we observed during 1979–2011, the DSL would be about  $1 \pm 1/3$  mo longer by 2090 than that in the 2000s [13], potentially making the Amazon rainforest an important accelerating feedback for climate change [1].

## 2.4. Oceans

- Ocean warming could cause a reduction in uptake of atmospheric CO<sub>2</sub> by 14–67 billion tonnes of carbon per year per °C of warming [14]. Some models suggest this corresponds to as much as 30% reduction in uptake by 2100, although most models suggest a more modest reduction [14].
- As oceans acidify, phytoplankton appear to produce less dimethylsulphide, which plays a role in cloud formation, meaning more radiative forcing. This effect alone is estimated to contribute 0.23–0.48°C warming [15].



**Figure 1. Human civilization has never seen CO<sub>2</sub> levels as high as they are today.** Image from Ref. [20].

### 3. CO<sub>2</sub> levels

- CO<sub>2</sub> levels are currently at 400 ppm and rising at more than 2 ppm a year. Pre-industrial levels were 280 ppm (see Fig. 1).
- CO<sub>2</sub> levels were approximately 440 ppm 15-20 million years ago, where the earth was 3-6°C warmer and sea levels 25-40 metres higher [16].
- The last time CO<sub>2</sub> levels were comparable to today's levels was 3.6–2.2 million years ago (mid-Pliocene), where summer temperatures were approximately 8°C warmer than today, the WAIS did not exist, and sea levels were 25 metres higher than today [17].
- Currently operating power generation infrastructure will commit us to 300 GtCO<sub>2</sub> whereas the budget for a 1.5°C-2°C world is 240 GtCO<sub>2</sub>. Current pipeline power plants would add a further 270 GtCO<sub>2</sub>. [18]
- Climate change that takes place due to increases in carbon dioxide concentration is largely irreversible for 1,000 years after emissions stop [19]

### 4. Temperature levels

- If, as a whole, the planet warms by 4°C, much of the global population (which lives in the mid-latitudes: US and Europe) will face warming of 5 °C or more.


### 5. Sea levels

- If Greenland completely melts it will raise sea levels by >6 metres (20 ft).

- If the Antarctic ice sheet completely melts it will raise sea levels by 60 metres (200 feet) [90% of all Earth's ice].
- The Antarctic ice sheet is losing 0.44 billion tonnes of ice per day (159 billion tonnes per year). [21]
- One sector of the West Antarctic ice shelf (WAIS) seems to have passed a tipping point [22]. The collapse of this sector would raise sea levels by 1.2 metres (4 ft) in the coming centuries. The sector acts as a linchpin for the stability of the WAIS.
- The WAIS contains enough ice to induce 3.7–4.6 metres of sea level rise (12–15 ft).
- Business as usual: 0.3 m (1 ft) rise by 2050, >1.2–1.8 m (4–6 ft) by 2100, rising 0.3 m per decade thereafter
- Seas will likely rise by around 80 cm (2.6 ft), with the 95th percentile at 180 cm (5.9 ft) [23].
- A glacier in the East Antarctic Ice Sheet also appears to be highly unstable, and has the potential to raise sea levels by 3.5 m (11.5 ft) [24].
- Antarctica has the potential to contribute more than 1 m (3.3 ft) of sea-level rise by 2100 [25]
- “Recent findings have led top climatologists to conclude that we are likely headed toward what used to 3–5 ft of global sea level rise by 2100, with worst-case scenarios being much worse” [1]
- Between 315–411 million people may be living in the 100-year flood plain by 2060, compared to 189 million in the year 2000 [26]. Subsidence in deltaic areas and from groundwater pumping could further enhance these numbers. Such storm-surges are expected to become far more common – see Section 8.
- Global mean sea level rise of 2.4 m (8 ft) by 2100 is physically possible, although the probability of such an outcome is hard to assess [27].
- Projections of salinization in Bangladesh suggest that climate change will cause significant changes in river salinity, likely leading to shortages in drinking and irrigation water [28]. This may reduce rice yields by ~15% [29].
- If CO<sub>2</sub> peaks at 600 ppm, irreversible global average sea level rise of at least 0.4–1.0 m is predicted from thermal expansion alone [19].

## 6. Heat waves, drought & dustbowlification

- In 2010, Russia suffered the most lethal heat wave in human history killing 55,000 people. Russia lost 40% of its wheat crop, and banned grain exports for 18 months. These factors have been suggested to be implicated in the Syrian conflict [30].
- The risk of a decade-scale megadrought in the U.S. southwest in the coming century is at least 80%, and may be higher than 90% in certain areas [31]
- Severe and widespread droughts are predicted in the next 30–90 years over many land areas [32], see Fig. 2.
- Drought conditions like the Dust Bowl of the 1930's are predicted to become normal in the U.S. Southwest and in other subtropical dry zones (reviewed in [33, 31]), see Fig. 2 [32].



**Figure 2. Severe and widespread droughts are predicted over the next century.** Percentage changes from 1980–1999 to 2080–2099 in the multimodel ensemble mean soil-moisture content in the top 10cm layer under the RCP4.5 emissions scenario. Image from Ref. [32] (Fig. 2a therein).

- Drought-affected areas are predicted to increase from 15.4% of global cropland today, to around 44% by 2100 [34]. The most severely affected regions in the next 30 to 90 years will likely be in southern Africa, the United States, southern Europe and Southeast Asia, says the report. In Africa, the report predicts 35% of cropland will become unsuitable for cultivation in a 5°C world [35].
- The growing season temperature at the end of the 21st century will likely exceed the hottest growing season ever observed in regions affecting half the world's population[36].
- Under a scenario where greenhouse gas emissions continue to grow, by 2100, ~74% of the Earth's population are forecast to be exposed to a mixture of mean surface air temperature and humidity that is deadly, for at least 20 days per year [37]. This is reduced to ~48% under a scenario of drastic greenhouse gas emissions, and is currently ~30%.
- Crop yields (mostly wheat, maize, rice, and soy) are forecast to likely reduce on the order of 10% from around 2040 onwards, with ~30% of projections predicting a decrease in yield of ~25% or more by 2100 (Fig. 2.7, [38]).

## 7. Ocean acidification

- Ocean acidification is causing many parts of the ocean to become undersaturated with calcium carbonate, which affects the ability of many oceanic organisms to form shells [39] [CHECK SOURCE LATER].
- Oceans are currently acidifying 10 times faster today than 55 million years ago when a mass extinction of marine species occurred [40, 1]
- Ocean acidification is irreversible on timescales of at least tens of thousands of years [41].

- Marine food supplies are likely to be reduced with significant implications for food production and security in regions dependent on fish protein [41]. The fish that grow and live on coral reefs are a significant food source for half a billion people worldwide [1].

## **8. Coastal flooding & storm surges**

- 2-4 feet sea-level rise by 2100 results in hurricane Sandy-level storm surge events recurring about once per year or more across the US east-coast [42].
- A 1°C rise in global temperature corresponds to 2–7 times increase in the frequency of Katrina magnitude storm surge events (which were 1 in 20 year events since 1923) [43].

## **9. Polar warming**

- Paleoclimate data shows that Arctic temperature change consistently exceeds the Northern Hemisphere average by a factor of 3-4 [44].
- The Arctic has warmed by 2°C since the 1970's [45]. Over this period, the Arctic grew 8% darker; the extra energy absorbed is equal to 25% of the entire heat-trapping effect of CO<sub>2</sub> in that time [45].
- 

## **10. Economic impacts**

- Non-agricultural economic productivity reduces by 2.4% per degree rise in temperature above 25°C[46].
- Productivity impacts alone might reduce per capita output by ~9% in 2080-2099 (in the absence of strong adaptation). This cost exceeds the combined cost of all other projected economic losses combined [1, 47] (see e.g. [48, 49], which do not include temperature-related loss in productivity, and may underestimate economic costs by more than a factor of two).
- CO<sub>2</sub> is a direct pollutant: at 930 ppm, six of nine decision-making performance domains were found to be impacted [50]. For most of human evolution and modern history, CO<sub>2</sub> levels in the atmosphere were between 180–280 ppm [1] (see Fig 1). Surveys of elementary schools in the US have reported CO<sub>2</sub> concentrations >1000 ppm in 45% of 435 classrooms investigated, and was associated with student absence [51, 50]. Peak CO<sub>2</sub> concentrations have been found to exceed 3000 ppm in 21% of classrooms in a Texas survey [51].

## **11. Biodiversity**

- There are very strong indications that the current rate of species extinctions far exceeds anything in the fossil record [52], being about 1000 times the background rate of extinction [53].
- By 2100, under business-as-usual projections, it is predicted that there will be a 50% increase in the suboxic water volume ("dead zones"), in response to the respiration of excess organic carbon formed at higher CO<sub>2</sub> levels [54]. Intermittent periods of low oxygen levels resulting in marine suffocation have been recorded since 2002 [55].

## 12. Basic science

- Global warming potential (GWP): Amount of heat trapped by a gas compared to the same mass of CO<sub>2</sub>, over a given period of time. 20 year GWP is more important due to the pressing nature of climate change, although 100 year GWP is more widely used.
- CH<sub>4</sub> GWP = 34 over 100 years. CH<sub>4</sub> GWP = 86 over 20 years.
- If sea surface temperatures are below 26.5°C, tropical cyclones and hurricanes do not form
- Polar amplification: where poles warm faster than the rest of the planet. This can weaken the jet stream, making weather last longer.

## 13. Further reading

- <https://www.skepticalscience.com/>

## References

- [1] **Romm J.** 2018. *Climate Change: What Everyone Needs to Know®*. What Everyone Needs To Know®, Oxford University Press, 2nd edition.
- [2] **Harries JE, Brindley HE, Sagoo PJ, Bantges RJ.** 2001. Increases in greenhouse forcing inferred from the outgoing longwave radiation spectra of the Earth in 1970 and 1997. *Nature* **410**: 355.
- [3] **Philipona R, Dürr B, Marty C, Ohmura A,** et al. 2004. Radiative forcing-measured at Earth's surface-corroborate the increasing greenhouse effect. *Geophysical Research Letters* **31**.
- [4] **Evans W, Puckrin E.** 2006, Measurements of the Radiative Surface Forcing of Climate.
- [5] **Cook J.** 2009, How do we know CO<sub>2</sub> is causing warming? <https://www.skepticalscience.com/print.php?n=73>.
- [6] **Tarnocai C, Canadell J, Schuur E, Kuhry P,** et al. 2009. Soil organic carbon pools in the northern circumpolar permafrost region. *Global biogeochemical cycles* **23**.
- [7] **IPCC.** 2007. Summary for Policymakers. In S Solomon, D Qin, M Manning, Z Chen, M Marquis, et al., eds., *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, Cambridge University Press, 1–18.
- [8] **Schaefer K, Zhang T, Bruhwiler L, Barrett AP.** 2011. Amount and timing of permafrost carbon release in response to climate warming. *Tellus B: Chemical and Physical Meteorology* **63**: 168–180.
- [9] **MacDougall AH, Avis CA, Weaver AJ.** 2012. Significant contribution to climate warming from the permafrost carbon feedback. *Nature Geoscience* **5**: 719.
- [10] **Kelly R, Chipman ML, Higuera PE, Stefanova I,** et al. 2013. Recent burning of boreal forests exceeds fire regime limits of the past 10,000 years. *Proceedings of the National Academy of Sciences* **110**: 13055–13060.
- [11] **Page SE, Siegert F, Rieley JO, Boehm HDV,** et al. 2002. The amount of carbon released from peat and forest fires in Indonesia during 1997. *Nature* **420**: 61.
- [12] **Turetsky M, Donahue W, Benscoter B.** 2011. Experimental drying intensifies burning and carbon losses in a northern peatland. *Nature Communications* **2**: 514.
- [13] **Fu R, Yin L, Li W, Arias PA,** et al. 2013. Increased dry-season length over southern Amazonia in recent decades and its implication for future climate projection. *Proceedings of the National Academy of Sciences* : 201302584.
- [14] **Gruber N.** 2011. Warming up, turning sour, losing breath: ocean biogeochemistry under global change. *Philosophical Transactions of the Royal Society of London A: Mathematical, Physical and Engineering Sciences* **369**: 1980–1996.

- [15] **Six KD, Kloster S, Ilyina T, Archer SD**, et al. 2013. Global warming amplified by reduced sulphur fluxes as a result of ocean acidification. *Nature Climate Change* **3**: 975.
- [16] **Tripathi AK, Roberts CD, Eagle RA**. 2009. Coupling of CO<sub>2</sub> and ice sheet stability over major climate transitions of the last 20 million years. *Science* **326**: 1394–1397.
- [17] **Brigham-Grette J, Melles M, Minyuk P, Andreev A**, et al. 2013. Pliocene warmth, polar amplification, and stepped Pleistocene cooling recorded in NE Arctic Russia. *Science* **340**: 1421–1427.
- [18] **Pfeiffer A, Hepburn C, Vogt-Schilb A, Caldecott B**. 2018. Committed emissions from existing and planned power plants and asset stranding required to meet the Paris Agreement. *Environmental Research Letters* **13**: 054019.
- [19] **Solomon S, Plattner GK, Knutti R, Friedlingstein P**. 2009. Irreversible climate change due to carbon dioxide emissions. *Proceedings of the national academy of sciences* **106**: 1704–1709.
- [20] The relentless rise of carbon dioxide. [https://climate.nasa.gov/climate\\_resources/24/graphic-the-relentless-rise-of-carbon-dioxide/](https://climate.nasa.gov/climate_resources/24/graphic-the-relentless-rise-of-carbon-dioxide/).
- [21] **McMillan M, Shepherd A, Sundal A, Briggs K**, et al. 2014. Increased ice losses from Antarctica detected by CryoSat-2. *Geophysical Research Letters* **41**: 3899–3905.
- [22] **Rignot E, Mouginot J, Morlighem M, Seroussi H**, et al. 2014. Widespread, rapid grounding line retreat of Pine Island, Thwaites, Smith, and Kohler glaciers, West Antarctica, from 1992 to 2011. *Geophysical Research Letters* **41**: 3502–3509.
- [23] **Jevrejeva S, Grinsted A, Moore JC**. 2014. Upper limit for sea level projections by 2100. *Environmental Research Letters* **9**: 104008.
- [24] **Greenbaum J, Blankenship D, Young D, Richter T**, et al. 2015. Ocean access to a cavity beneath Totten Glacier in East Antarctica. *Nature Geoscience* **8**: 294.
- [25] **DeConto RM, Pollard D**. 2016. Contribution of Antarctica to past and future sea-level rise. *Nature* **531**: 591.
- [26] **Neumann B, Vafeidis AT, Zimmermann J, Nicholls RJ**. 2015. Future coastal population growth and exposure to sea-level rise and coastal flooding—a global assessment. *PloS one* **10**: e0118571.
- [27] **Wuebbles D, Fahey D, Hibbard K, DeAngelo B**, et al. 2017. Executive summary. In D Wuebbles, D Fahey, K Hibbard, D Dokken, B Stewart, T Maycock, eds., *Climate Science Special Report: Fourth National Climate Assessment*, U.S. Global Change Research Program, Washington, DC, USA, volume I, 12–34.
- [28] **Dasgupta S, Akhter Kamal F, Huque Khan Z, Choudhury S**, et al. 2015. River salinity and climate change: evidence from coastal Bangladesh. In *World Scientific Reference on Asia and the World Economy*, World Scientific, 205–242.
- [29] **Dasgupta S, Hossain MM, Huq M, Wheeler D**. 2014. *Climate change, soil salinity, and the economics of high-yield rice production in coastal Bangladesh*. The World Bank.
- [30] **Kelley CP, Mohtadi S, Cane MA, Seager R**, et al. 2015. Climate change in the Fertile Crescent and implications of the recent Syrian drought. *Proceedings of the National Academy of Sciences* : 201421533.
- [31] **Ault TR, Cole JE, Overpeck JT, Pederson GT**, et al. 2014. Assessing the risk of persistent drought using climate model simulations and paleoclimate data. *Journal of Climate* **27**: 7529–7549.
- [32] **Dai A**. 2013. Increasing drought under global warming in observations and models. *Nature Climate Change* **3**: 52.
- [33] **Seager R, Ting M, Held I, Kushnir Y**, et al. 2007. Model projections of an imminent transition to a more arid climate in southwestern North America. *Science* **316**: 1181–1184.
- [34] **Li Y, Ye W, Wang M, Yan X**. 2009. Climate change and drought: a risk assessment of crop-yield impacts. *Climate research* **39**: 31–46.



- [35] Climate Change Report Warns of Dramatically Warmer World This Century (2012). <http://www.worldbank.org/en/news/feature/2012/11/18/Climate-change-report-warns-dramatically-warmer-world-this-century>.
- [36] **Battisti DS, Naylor RL**. 2009. Historical warnings of future food insecurity with unprecedented seasonal heat. *Science* **323**: 240–244.
- [37] **Mora C, Dousset B, Caldwell IR, Powell FE**, et al. 2017. Global risk of deadly heat. *Nature Climate Change* **7**: 501.
- [38] **Allen MR, Barros VR, Broome J, Cramer W**, et al. 2014. IPCC fifth assessment synthesis report-climate change 2014 synthesis report. *IPCC* .
- [39] Pacific Marine Environmental Laboratory: What Is Ocean Acidification? <https://www.pmel.noaa.gov/co2/story/What+is+Ocean+Acidification>
- [40] **Ridgwell A, Schmidt DN**. 2010. Past constraints on the vulnerability of marine calcifiers to massive carbon dioxide release. *Nature Geoscience* **3**: 196.
- [41] **Zhe C, Alper H**. 2009. IAP Statement on Ocean Acidification. *The Interacademy Panel on International Issues* .
- [42] **Sweet W, Zervas C, Gill S, Park J**. 2013. Hurricane Sandy inundation probabilities today and tomorrow. *Bulletin of the American Meteorological Society* **94**: S17–S20.
- [43] **Grinsted A, Moore JC, Jevrejeva S**. 2013. Projected Atlantic hurricane surge threat from rising temperatures. *Proceedings of the National Academy of Sciences* **110**: 5369–5373.
- [44] **Miller GH, Alley RB, Brigham-Grette J, Fitzpatrick JJ**, et al. 2010. Arctic amplification: can the past constrain the future? *Quaternary Science Reviews* **29**: 1779–1790.
- [45] **Pistone K, Eisenman I, Ramanathan V**. 2014. Observational determination of albedo decrease caused by vanishing Arctic sea ice. *Proceedings of the National Academy of Sciences* **111**: 3322–3326.
- [46] **Hsiang SM**. 2010. Temperatures and cyclones strongly associated with economic production in the Caribbean and Central America. *Proceedings of the National Academy of sciences* : 201009510.
- [47] Temperature and worker output (2011). <http://www.fight-entropy.com/2011/08/temperature-and-worker-output.html>.
- [48] **Stern N, Peters S, Bakhshi V, Bowen A**, et al. 2006. *Stern Review: The economics of climate change*, volume 30. HM treasury London.
- [49] **Tol RS**. 2009. The economic effects of climate change. *Journal of economic perspectives* **23**: 29–51.
- [50] **Allen JG, MacNaughton P, Satish U, Santanam S**, et al. 2015. Associations of cognitive function scores with carbon dioxide, ventilation, and volatile organic compound exposures in office workers: a controlled exposure study of green and conventional office environments. *Environmental health perspectives* **124**: 805–812.
- [51] **Corsi R, Torres V, Sanders M, Kinney K**. 2002. Carbon dioxide levels and dynamics in elementary schools: results of the TESIAs Study. *Indoor Air* **2**: 74–79.
- [52] **Magurran AE, Dornelas M**. 2010, Biological diversity in a changing world.
- [53] **Pimm SL, Jenkins CN, Abell R, Brooks TM**, et al. 2014. The biodiversity of species and their rates of extinction, distribution, and protection. *Science* **344**: 1246752.
- [54] **Oschlies A, Schulz KG, Riebesell U, Schmittner A**. 2008. Simulated 21st century's increase in oceanic suboxia by CO<sub>2</sub>-enhanced biotic carbon export. *Global Biogeochemical Cycles* **22**.
- [55] Oceans Are Losing Oxygen and Becoming More Hostile to Life (2015). <https://news.nationalgeographic.com/2015/03/150313-oceans-marine-life-climate-change-acidification-oxygen-fish/>.