Financial Networks Project: working notes

Jash Dalal, last updated 8 September 2026. Newest entries at the bottom.

5 July: Paper 1, learning a Laplacian from returns

Builds a network of stocks straight from price movements instead of building a correlation table and cutting it up. Watches one number, λ_2, which says how tightly the network holds together; small means close to breaking apart, and the rule is to stay invested below a set level and sit in cash above it.

The method, the month-to-month movement of λ_2, and the COVID dodge all reproduce. The money does not. Here λ_2 stays high through April 2020 where the paper's drops, so the strategy sits in cash through the recovery and ends at -0.011 where simply holding ends at $+0.078$. No threshold was moved to fix it. Takeaway: λ_2 is the incumbent indicator on the Laplacian side, and it is fragile.

22 to 28 July: Paper 2, market regimes with TICC

Cuts time into stretches, each with its own pattern of which series depend on which, and the same pattern can come back later. Headline accuracy reproduces. The side experiments that switch parts of the method off come out weaker than reported, so this rebuild makes the method look *more* necessary than the paper's own numbers do. Peripheral to the current question; it is the only one of the three that models a state stretched over time, so keep it as the comparator if a network indicator ever needs checking against a plain regime model.

9 to 10 August: Paper 3, modularity of market networks

Each day, keep the strongest 10% of pairwise correlations as connections, run Louvain, record one number Q for how cleanly the stocks split into groups. Rebuilt at full size: 360 stocks, 25 years, 6,315 daily networks. It holds up on its own terms. The trouble is what the number means.

- **Q does not show the market has groups.** Pure random data through the same pipeline gives 0.2246; the real market gives 0.2215. Noise comes out marginally *more* grouped than the stock market. Cutting up a correlation table always yields a network full of triangles, and those always look grouped.
- **Louvain disagrees with itself.** Run-to-run spread on the same day is 2.09× a normal day-to-day move in Q, on 78% of days.
- **The discarded number works better.** $\tau(t)$, the cut-off needed each day to keep 10% of pairs, is computed, used, then thrown away. It scores 0.807 against Q's 0.538.
- **Compression fails where the story is.** The group summary is cheaper than the network on 69% of ordinary days but only 34% of crisis days, because in a crisis the network shatters and the summary outgrows the thing it was summarising.

Re-audit the same day: the paper's four non-reproducing numbers are real and survive nine implementation variants, but the cause is that these networks fall into disconnected pieces on 63% of days (99.3% after 2002). Eight defects were found in this rebuild's own criticisms and fixed.

12 August: the two matrices behind the two indicators

The prompt. Paper 1 reads λ_2 off the graph Laplacian. Paper 3 reads Q off a grouping of stocks. Modularity has a matrix underneath it that nobody in this literature reports. Is there a study in comparing the two?

Laplacian $\mathbf{L} = \mathbf{D} - \mathbf{A}$: how well connected the network is and where it would break. Number of zero eigenvalues is how many separate pieces it has fallen into; λ_2 is zero when it is already in pieces and large when solidly joined.

Modularity matrix $\mathbf{B} = \mathbf{A} - \mathbf{d}\mathbf{d}^\top/2m$: what was seen minus what was expected. The baseline $d_i d_j/2m$ answers "how many connections would these two have by accident, given how popular each is". Two typical stocks (36 connections each) give 0.10, exactly the network's density. Two heavily connected stocks (100 each) give 0.77, so a link between them is not news. Every entry is a surprise score:

seen	expected	entry	reading
linked	0.10	+0.90	together far more than chance
linked	0.77	+0.23	two popular stocks linked; barely news
not linked	0.10	−0.10	ordinary; most pairs look like this
not linked	0.77	−0.77	two popular stocks *avoiding* each other

That last row is the point: **B** has a meaningful negative end, while the Laplacian, built only from connections that exist, cannot say "should be there and is missing". And Q is just a sum of selected entries of **B**, so the chain runs from a 360×360 matrix, through one grouping picked by an unstable search, down to a single number. Paper 3 reports the last box.

Reading **B** directly instead: μ_1 measures the strongest split (same job as Q, but deterministic), $\#\{\mu_i > 0\}$ caps the number of separable groups so it is a group count with no algorithm involved, and μ_N picks out *anti*-groups, sets of stocks avoiding each other. None need Louvain, and none appear anywhere in finance.

12 August: spectral probe over the full 25 years

Full spectrum of **L**, **B** and the raw correlation table **C** for all 6,315 days, every statistic through the same detector: a 250-day trailing z-score using only prior days, scored against crisis dates fixed before any of this work. Score is AUC, the chance a random crisis day ranks above a random calm day, so 0.50 is a coin flip. Six minutes on one core.

Signal	**What it measures**	**AUC**
spectral entropy of **C**	how spread out the correlation structure is	**0.822**
$\lambda_1(\mathbf{C})/N$	share of movement in the biggest common factor	0.812
$\tau(t)$	cut-off needed that day to keep 10% of pairs	0.807
$\mu_N(\mathbf{B})$	**strongest anti-grouping**	**0.760**
isolated-node count	stocks left with no connections	0.756
$\sum_{\mu_i>0} \mu_i$	**total group mass**	0.750
$\lambda_N(\mathbf{L})$	largest Laplacian eigenvalue	0.731
$\#\{\mu_i > 0\}$	**group count, no algorithm**	0.722
$\lambda_2(\mathbf{L})$, largest piece	Fiedler value, breakage removed	0.652
$\mu_1 - \mu_2$	modularity spectral gap	0.636
# modes above the noise band	how many groups are even resolvable	0.619
$\lambda_2(\mathbf{L})$, whole network	Fiedler value, **Paper 1**	0.617
$\mu_1(\mathbf{B})$	leading modularity eigenvalue	0.607
$Q(t)$	Louvain modularity, **Paper 3**	**0.538**
$\max_k(\lambda_{k+1} - \lambda_k)$	largest Laplacian gap	0.526

1. The matrix beats the single number. Q at 0.538 means it ranks a crash day above a calm day 53.8% of the time; a coin manages 50%. Three separate readings of the matrix Q comes from reach 0.72 to 0.76 on the same networks under the same test, so the information was in **B** all along and the Louvain step is where it is lost. The best modularity signal is the *negative* end, which thresholding structurally cannot see. And μ_1 tracks Q at $r = 0.761$ while being fully deterministic, so it is a drop-in with none of the wobble. Paper 1's 0.617 is not a refutation of Paper 1, which learns a regularised Laplacian on 7 stocks, a different object entirely.

2. The identity breaks, and that is the good news. On a network where every stock has the same number of connections d, it is exactly true that $\lambda_2(\mathbf{L}) + \mu_1(\mathbf{B}) = d$: the Fiedler

value and the leading modularity eigenvalue are the same number read backwards. If that held here, comparing Paper 1's indicator with Paper 3's would be comparing a thing to itself. Paper 3's construction pins the average degree at exactly 35.90 every day, so it nearly should hold. Measured over all days the sum falls short by $\mathbf{9.49 \pm 4.63}$, i.e. 26% of the average degree, and the ± 4.63 says it is not a fixed offset that could be subtracted, it moves by a third of its own size day to day. Restricting to the 2,342 one-piece days gives 9.21 ± 3.60; the largest piece only gives 11.11 ± 5.31. Cleaner still: two quantities summing to a constant would be perfectly anti-correlated at -1, unrelated ones give 0, and the measured value is $\mathbf{-0.187}$. Dual in theory, unrelated in practice. So the question stops being "which indicator wins" and becomes *what is that* 9.49 *made of*: uneven degrees, breakage, or the market-wide component.

3. There may be nothing to detect. 64,620 correlations are estimated from 30 days on 360 stocks, six times more output than input, and at most 29 of the 360 eigenvalues can be non-zero by arithmetic alone. Random-matrix theory puts the noise band's upper edge at 19.93, so anything below that is indistinguishable from what pure noise produces. Counting eigenvalues that escape it, excluding the market-wide component: **mean** 0.70**, max** 3**, often zero**. Less than one resolvable group per day, while Louvain confidently reports a median of nine and the paper narrates their 25-year dynamics. This indicts the whole literature, not just Paper 3; a 30-day window on a few hundred stocks is standard.

4. The strong signals are one signal, and it does not generalise. Spectral entropy, the absorption ratio and $\tau(t)$ move together at $|r| > 0.99$, so they are the same series with different labels, all measuring the market-wide component. That is not four discoveries but one, made in 1999. Training before 2002 and testing after (27 crisis days in the test period):

	training years	unseen years
market-wide component alone	0.812	**0.730**
+ $\mu_1(\mathbf{B})$	0.822	0.629
+ $\mu_N(\mathbf{B})$	0.810	0.719
+ $Q(t)$	0.831	0.699
+ $\lambda_2(\mathbf{L})$, largest piece	0.810	0.700

Every addition helps on the training years and hurts on the unseen ones, the textbook overfitting fingerprint. 27 events is too few to settle it either way, but "too weak to settle" is not a defence. One bright spot: μ_1 tracks the market-wide component at only 0.025 while tracking Q at 0.761, so it really is a new axis. μ_N tracks it at 0.68, so much of its 0.760 is borrowed, which is exactly the D3 gate below.

Code and outputs in `reports/probe/`. It imports the rebuild's own crisis dates, network construction and scoring unchanged; as a check, Q and τ come out at 0.5380684370 and 0.8071316467, matching the stored values to ten places.

12 August: proposed directions and open questions

D1, what breaks the identity. Decompose the 9.49 into uneven degrees, breakage, and the market-wide component, and see which carries the stress information. Same spectrum over a family of matrices ($\mathbf{A}$, $\mathbf{L}$, normalised $\mathbf{L}$, $\mathbf{B}$, normalised $\mathbf{B}$, and the threshold-free correlation modularity of MacMahon & Garlaschelli) under one shared test. Not a bake-off but a structural result with a bake-off attached. Risk: the gap may be dominated by breakage, an artifact of thresholding. The largest-piece variant already survives, but that needs establishing across constructions.

D2, is there anything to detect. Map, over window length, universe size and density, where the market sits relative to the noise band and the Nadakuditi and Newman detectability threshold, so anyone in this literature can check whether their settings are inside the region where grouping means anything. Ship a deterministic replacement for the group count. Risk: if essentially no configuration in the literature is inside that region, the contribution is purely negative. Still publishable, harder to place.

D3, the negative end (gated). Paper 3 discards 4,246,798 pair-days of strong *anti*-correlation, recorded identically to "no relationship", and that series collapses to near zero from about 2003 and never recovers, right beside the paper's own unexplained 2002 break which it lists as future work. μ_N at 0.760 is the best new number here. Gate before committing: does μ_N survive removing the market-wide component? Two-day check.

Recommendation: D1 and D2 as one project, D3 second if the gate passes. D1 alone risks being a bake-off, D2 alone risks being purely negative. Together they are one claim: here is the right matrix to read market group structure from, here is where reading it means anything at all, and here is what the field's two incumbent indicators actually measure.

Open questions.

Q1. Is the identity the right spine, or would you prefer the plainer empirical comparison?

Q2. Keep Paper 3's thresholded networks (comparable to the rebuild) or move to threshold-free correlation modularity (more defensible, breaks comparability)? Inclination is both, and report the difference, since that difference is the "beyond thresholding" argument made concrete.

Q3. 27 unseen crisis days is far too few. Extend to 2026, add exchanges, or go index-level?

Q4. Venue: a physics venue wants the spectral and detectability result, a finance venue wants the indicator to beat the absorption ratio on unseen data, which on current evidence it does not.

Q5. Carry TICC forward as a comparator, or park it?

19 August: the exploration written up properly

The 12 August entries above are too compressed to follow, so the whole exploration is now written out in `reports/2026-08-19_discrepancy_explained.tex` and shared for comment. It builds every result up from the papers' own definitions: the three replications side by side, each observed discrepancy beside what would explain it, the one scoring apparatus everything is judged on, and the three directions in a single symmetric frame. 17 pages, eight diagrams, self-contained, Overleaf-ready; every number in it is re-derived from the stored artifacts by `reports/probe/check_numbers.py`.

Three figures quoted above are stated more carefully there. The 0.68 and 0.025 market-mode correlations hold on the z-scored detector series, not in levels (where they are -0.819 and $+0.294$); the AUC-0.756 row is the count of *connected components*, not of isolated nodes; and the noise-control Q values come from two different runs and are not interchangeable.

31 August: the meeting, and the question set

Asked to take the modularity matrix seriously as a source of crisis indicators: find a quantity derived from $\mathbf{B}$ that carries the same information as $\lambda_2(\mathbf{L})$, tested on Paper 1's own FAAMUNG data, so the two can be declared equivalent and a theory framework built on top. Not necessarily a second-smallest eigenvalue, anything at all. Scope agreed: an empirical menu, Paper 1's dataset only, no Paper 3 scoring.

Also raised at the meeting: the $\lambda_2(\mathbf{L}) + \mu_1(\mathbf{B}) = d$ spine proposed as D1/D2 on 12 August holds only when *every* degree equals d. Substituting a mean degree is not a weakened hypothesis, it is no hypothesis. **The 12 August framing of D2 is retracted**; the quantitative confirmation is in the next entry.

31 August to 7 September: a modularity equivalent of the Fiedler value

Paper 1 throws the learned graphs away after reading λ_2 off them, so Algorithm 2 was re-run with its exact constants to recover all 243 Laplacians, $\mathbf{A}_t = -\operatorname{offdiag}(\mathbf{L}_t)$ recovered from each, and 22 modularity-derived quantities scored against 6 Laplacian references. Gates first: recomputed λ_2 matches the cached series to 1.2×10^{-10}, and the backtest reproduces the published $S1 = +0.078$ against $S2 = -0.011$.

The question dissolves under normalisation. Writing $\mathbf{u} = \mathbf{D}^{1/2}\mathbf{1}/\sqrt{2m}$, the normalised modularity matrix is $\mathbf{D}^{-1/2}\mathbf{B}\mathbf{D}^{-1/2} = \mathbf{I} - \mathbf{L}_{\text{sym}} - \mathbf{u}\mathbf{u}^\top$, and $\mathbf{u}$ is $\mathbf{L}_{\text{sym}}$'s own null vector. The two normalised matrices are the same operator: eigenvalues paired one-to-one by $\mu = 1 - \lambda$ (full spectrum, all 243 windows, 1.8×10^{-15}) and eigenvectors shared to 1.000000. So no normalised-$\mathbf{B}$ indicator can be new; it is a Laplacian indicator in different notation. Unnormalised, the two do *not* share a basis (overlap 0.871; on some windows the Fiedler vector and $\mathbf{B}$'s leading mode are nearly orthogonal, 0.039), so that is where anything new must live.

The best-scoring candidate was the spurious one. $\bar{d} - \mu_1(\mathbf{B})$ reproduced 91% of λ_2's daily invest/cash calls, and it is exactly the $\bar{d}$ substitution flagged at the meeting. Whole-spectrum error 1.25, i.e. 5.3× λ_2's own sd; residual sd 0.208, **89% of λ_2's sd**; only 40% of the proxy's variance is λ_2; and the residual is *smaller* in-crash (0.308 vs 0.512), so the error term was doing some of the detecting. Degree CV here is 0.305, nowhere near regular. The $\bar{d}$ term also does no work: controlling for it, $\mu_1(\mathbf{B})$ alone gives -0.851 against the proxy's $+0.851$.

Correlation was the wrong screen. It was the working comparison metric between $\mathbf{B}$ and $\mathbf{L}$, and it fails both ways. It promoted the spurious candidate above; and spectral entropy of the normalised $\mathbf{B}$ correlates only -0.264 with λ_2, which looks independent, yet rebuilds from the $\mathbf{L}_{\text{sym}}$ spectrum alone to 8.0×10^{-15}. Novelty has to mean *not reconstructible from* spec($\mathbf{L}$), not *decorrelated from* λ_2.

λ_2 is itself inside the noise band on Paper 1's own setup. Seven stocks, 243 windows, one event, 24 crash days. Raw-level COVID AUC is **0.472**, below chance; only a strictly-prior z-score lifts it to 0.759. It clears a bootstrap over days (CI $[0.634, 0.875]$) but fails the harder null: sliding a 24-day window anywhere in the series puts COVID at the **76th percentile,** $p = 0.243$. A random gate at the same invest rate beats buy-and-hold 39.2% of the time, so ∼11 of 28 candidates were expected to beat $S1$ by luck; 5 did, none with $p < 0.115$. No multiplicity correction exists anywhere in this study.

What survives, and it reframes the project. λ_2 is read off the *unnormalised* $\mathbf{L}$, so it is a blend: $d_{\min}\lambda_2(\mathbf{L}_{\text{sym}}) \le \lambda_2(\mathbf{L}) \le d_{\max}\lambda_2(\mathbf{L}_{\text{sym}})$ holds on 243/243 windows. Splitting the detection: structure alone 0.700, scale alone 0.537 (a coin flip), the two combined 0.759. **Degree information is a modifier, not a signal**: worthless alone, but it improves the signal when mixed in, and it is exactly what normalisation destroys. $\mathbf{L}$ combines it additively on the diagonal; $\mathbf{B}$ combines it through a random-rewiring null. That difference is the only thing $\mathbf{B}$ offers, and it is the question worth taking forward. It also explains rather than merely observes why $\mu_N(\mathbf{B})$ keeps winning: z-AUC 0.775 here against λ_2's 0.759, redundancy only 0.51, and independently the best modularity signal in `reports/probe/` (0.760 vs 0.617 over 17 crises). Not significant on one event (ΔAUC $+0.015$, CI $[-0.115, +0.148]$, and μ_N fails the same sliding null at $p = 0.345$), so it is a lead, not a result.

Written up in full in `reports/2026-09-08_modularity_equivalence.tex` (10 pages); code and the number-checking script in `modularity_indicator/`.

16 to 18 October: the proposal, and what had to be fixed first

Prof. Anguluri's proposal *Modularity Matrix Approach for Graph Learning?* returns to Paper 1 and asks what modularity's null-model view adds to Laplacian graph learning: Part I, the exact relation under diag($\mathbf{L}$) = $\mathbf{1}$; Part II, the market factor as an implicit null; Part III, modularity as a prior replacing the degree constraint. Checking it against the code turned up three issues.

Paper 1's Fig. 2 had not recovered the sectors. It was marked reproduced on rank alone; its three components (77/12/8 stocks) had ARI 0.012 against the sectors. Two causes: a four-year data window where the paper uses three (753 days), and a uniform starting graph whose degenerate spectrum makes Algorithm 1's first $\mathbf{V}$-step, and so its components, depend on round-off. Starting from the Problem (1) estimate recovers the sectors exactly (52/28/17, ARI 1.000), stable across BLAS thread counts. Two false notes were corrected on the way: the Fig. 2 baseline is not SGL, and our solver *can* produce isolated nodes.

Part I(c) applies only to unit-degree graphs. The paper's time-varying Problem (11) has no degree constraint; our FAAMUNG graphs have mean degree 2.13.

The Part III objective is unbounded below. Its modularity term scales linearly with edge weight. On the 753-day data it is bounded for $\eta < 0.186$ and unbounded for $\eta > 0.64$. Fixing the total weight $\mathbf{1}^\top \mathbf{d} = p$ keeps convexity and repairs it.

Written up for the professor in `reports/2026-10-18_proposal_issues.tex`; code, lemma tests and the number-checking script in `modularity_graph_learning/`.

27 October: Part I finished

The lemma, proved. Under $\mathrm{diag}(\mathbf{L}) = \mathbf{1}$, $\mathbf{B} = \mathbf{I} - \mathbf{L} - \mathbf{1}\mathbf{1}^\top/p$, so (a), (b) and (c) follow from one reflection lemma. Part (b) holds in the exact form $\sum_{i \le k} \lambda_i = (k-1) - \sum_{i<k} \mu_i + \max\{0, \lambda_k - 1\}$, so the condition $\lambda_k \le 1$ is exactly what removes the correction; the K_4 and 6-cycle counterexamples show it.

On the repaired Fig. 2 graph the prediction holds to the solver's degree tolerance: degrees 1.0012 to 1.0069, $\mathbf{B}$'s top eigenvalues 1.0052 and 1.0041 (Part I says 1), and k-means on them returns the sectors exactly. Most of the mismatch is the mean degree 1.0047: reflecting about it instead of 1 cuts the spectral gap from 0.0059 to 0.0020.

Loosening the degree penalty (ρ from 100 to 0.3) breaks the algebra (residual 1.7% to 110%, $\mu_1(\mathbf{B})$ up to 1.65) but not the partition, and the normalised $\mathbf{B}_n$ keeps eigenvalue 1 twice at every ρ, as the 8 September identity predicts.

Written up in `reports/2026-10-27_part1_lemma.tex`; (c) stated for unit degree only, pending the professor's reply.

dd Month: next entry here

Pointers. Papers: arXiv:2005.09958 (Cardoso & Palomar), arXiv:1706.03161 (TICC), arXiv:1501.05040 (Silva et al.). Key outside references: Kritzman et al. 2011 (absorption ratio, the baseline to beat); Newman 2006 (modularity matrix); Fasino & Tudisco 2014/2016/2017 (spectrum, group counts, anti-groups); Nadakuditi & Newman 2012 (detectability); MacMahon & Garlaschelli 2015 (the correlation-specific null Paper 3 needed and did not use); Masuda et al., *Physics Reports* 2025 (the "beyond thresholding" review); Kang et al. 2025 (largest Laplacian gap). Two Laplacian-energy papers (Eur. J. Finance 29(9) 2023; Int. Rev. Econ. Finance 2025) have unverified author lists, do not cite as-is. In this repo: `reports/archive/EXPLORATION_EXPLAINED.md` for the ground-up version of all of the above, `reports/probe/` for code and outputs, `modularity_indicator/` for the 31 August to 7 September work and `reports/2026-09-08_modularity_equivalence.tex` for its write-up. Paper 3 needs `python` and `python-igraph`.