

MINISTRY OF AGRICULTURE AND FARMERS WELFARE**(Department of Agriculture, and Farmers Welfare)****ORDER**

New Delhi, the 25th March, 2026

S.O. 1574(E).— In exercise of the powers conferred by section 3 of the Essential Commodities Act, 1955 (10 of 1955), the Central Government hereby makes the following order further to amend the Fertiliser (Inorganic, Organic or Mixed) (Control) Order, 1985, namely: -

1. (1) This Order may be called the Fertiliser (Inorganic, Organic or Mixed) (Control) Fourth Amendment Order, 2026.
(2) It shall come into force on the date of its publication in the Official Gazette.
2. In the Fertiliser (Inorganic, Organic or Mixed) (Control) Order, 1985, in Schedule VI, in Part A, “Specifications of Biostimulants”,-

(a) in sub-heading 1, “Humic and Fulvic acid and their derivatives”, after serial number (31) and the entries relating thereto, the following serial numbers and entries shall be inserted, namely:-

“(32) Humic and fulvic acid 10.48% (Powder)

Composition		
S. No.	Ingredients	Content
(i)	Humic acid (Source: Leonardite) per cent. by weight, minimum	9.96
(ii)	Fulvic acid per cent. by weight, minimum	0.52
(iii)	Dextrin per cent. by weight, maximum	QS
(iv)	Total (per cent.)	100

Specifications		
S. No.	Parameters	Value
(i)	Humic acid per cent. by weight, minimum	9.96
(ii)	Fulvic acid per cent. by weight, minimum	0.52
(iii)	Total organic carbon per cent. by weight, minimum	3.60
(iv)	pH (50% aqueous solution)	5.16 ± 0.5
(v)	Bulk density (gram/cc)	0.572 ± 0.02
(vi)	Solubility per cent. by weight, minimum	98.87

(a) Name of the crop: Tomato

(b) Dose: One soil application at 2.5 litre/ha

(33) Humic acid and Fulvic acid 20.20 % (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Humic acid (Source: Leonardite) per cent. by weight, minimum	18.63
(ii)	Fulvic acid per cent. by weight, minimum	1.57
(iii)	Water per cent. by weight	QS
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Humic acid per cent. by weight, minimum	18.63
(ii)	Fulvic acid per cent. by weight, minimum	1.57
(iii)	Total organic carbon per cent. by weight, minimum	13.29
(iv)	pH (50% aqueous solution)	5.28± 0.5
(v)	Specific gravity	1.186± 0.02
(vi)	Solubility per cent. by weight, minimum	99.24

(a) Name of the crop: Tomato

(b) Dose: Three foliar applications at 2.5 litre/ha

(34) Humic acid and Fulvic acid 51% (Powder/Flake)

Composition		
S. No.	Ingredients	Content
(i)	Humic substances (Source Browncoal/ Leonardite/ weathered coal), per cent. by weight, minimum	80
(ii)	Alkali reagents (Potassium hydroxide, sodium hydroxide, dipotassium hydrogen phosphate, potassium carbonate) per cent. by weight, minimum	10
(iii)	Water per cent. by weight, maximum	QS

(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Humic acid per cent. by weight, minimum	50.0
(ii)	Fulvic acid per cent. by weight, minimum	1.00
(iii)	Total organic carbon per cent. by weight, minimum	29.0
(iv)	pH (50% aqueous solution)	9.0- 12.0
(v)	Bulk density (gram/cc)	0.50- 0.90
(vi)	Solubility per cent. by weight, minimum	70

(a) Name of the crop: Rice

(b) Dose: Three applications (one soil and two foliar) at 1.875 kg/ha”;

(b) in sub-heading 2, “Seaweed Extract”, after serial number (41) and the entries relating thereto, the following serial numbers and entries shall be inserted, namely:-

“(42) Seaweed (*Ascophyllum nodosum*) extract 100% (Powder)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract powder per cent. by weight, minimum	100
(ii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	10.0
(ii)	Fucoidan per cent. by weight, minimum	10.0
(iii)	Mannitol per cent. by weight, minimum	5.0
(iv)	Amino acids per cent. by weight, minimum	2.0
(v)	Total organic carbon per cent. by weight, minimum	26.10
(vi)	pH (1 % aqueous solution)	9.5- 10.5
(vii)	Bulk density (gram/cc)	0.55- 0.65
(viii)	Solubility per cent. by weight, minimum	99.8

(a) Name of the crop: Tomato

(b) Dose: Three foliar applications at 0.75 kg/ha

(43) Seaweed (*Ascophyllum nodosum*) extract 10% (Granule)

Composition		
S. No.	Ingredients	Content

(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract per cent. by weight, minimum	10.00
(ii)	Dolomite granules per cent. by weight, maximum	90.00
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	0.9
(ii)	Total organic carbon per cent. by weight, minimum	5.8
(iii)	pH (1 % aqueous solution)	8.0- 10.0
(iv)	Bulk density (gram/cc)	1.05- 1.09

(a) Name of the crop: Rice

(b) Dose: One soil application at 10 kg/ha

(44) Seaweed (*Ascophyllum nodosum*) extract 99.4% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract liquid per cent. by weight, minimum	99.4
(ii)	Methyl p- hydroxybenzoate per cent. by weight, maximum	0.30
(iii)	Propyl p- hydroxybenzoate per cent. by weight, maximum	0.06
(iv)	2- Propanol per cent. by weight, maximum	0.15
(v)	Sodium hydroxide per cent. by weight, maximum	0.09
(vi)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	1.8
(ii)	Fucoidan per cent. by weight, minimum	2.0
(iii)	Mannitol per cent. by weight, minimum	1.0
(iv)	Total organic carbon per cent. by weight, minimum	5.8
(v)	pH (1 % aqueous solution)	7.5- 8.5
(vi)	Total dissolved solids per cent. by weight, minimum	20
(vii)	Specific gravity	1.10- 1.20
(viii)	Solubility per cent. by weight, minimum	99.8

(a) Name of the crop: Chilli

(b) Dose: Three foliar applications at 2.0 litre/ha

(45) Seaweed (*Ecklonia maxima*) extract 94% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ecklonia maxima</i>) extract per cent. by weight, minimum	94
(ii)	Water per cent. by weight, maximum	6
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	2.0
(ii)	pH (10% aqueous solution)	4.6- 4.9
(iii)	Total organic carbon per cent. by weight, minimum	10
(iv)	Specific gravity	1.10± 0.02
(v)	Total dissolved solids per cent. by weight, minimum	30

(a) Name of the crop: Watermelon

(b) Dose: Three foliar applications at 1.5 liter/ha

(46) Seaweed (*Ascophyllum nodosum*) extract 30% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract per cent. by weight, minimum	30
(ii)	Methylisothiazolinone (preservative) per cent. by weight, minimum	0.1
(iii)	Water per cent. by weight, maximum	QS
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	3.0
(ii)	Mannitol per cent. by weight, minimum	1.0
(iii)	pH (as such)	9.0- 11.0
(iv)	Bulk density (gram/cc)	1.10- 1.25
(v)	Total organic carbon per cent. by weight, minimum	9.50
(vi)	Solubility per cent. by weight, minimum	97
(vii)	Total dissolved solids per cent. by weight, minimum	30.0

(a) Name of the crop: Rice

(b) Dose: Three foliar applications at 1000 ml/ha

(47) Seaweed (*Ascophyllum nodosum*) extract-90 % (Powder/Flake)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract per cent. by weight, minimum	90
(ii)	Methylisothiazolinone (preservative) per cent. by weight, minimum	0.1
(iii)	Water per cent. by weight, maximum	QS
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	8.0
(ii)	Mannitol per cent. by weight, minimum	3.0
(iii)	pH (1% aqueous solution)	9.0- 11.0
(iv)	Bulk density (gram/cc)	0.56- 0.75
(v)	Total organic carbon per cent. by weight, minimum	28.0
(vi)	Solubility per cent. by weight, minimum	96

(a) Name of the crop: Rice

(b) Dose: Three foliar applications at 1000 g/ha

(48) Seaweed (*Ascophyllum nodosum*) extract 20% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract per cent. by weight, minimum	20
(ii)	Water per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	1.02
(ii)	Total organic carbon per cent. by weight, minimum	1.04
(iii)	pH (50% aqueous solution)	7.35 ± 0.5
(iv)	Specific gravity	1.047 ± 0.02
(v)	Total dissolved solids per cent. by weight, minimum	2.4
(vi)	Solubility per cent. by weight, minimum	99.99

(a) Name of the crop: Chilli

(b) Dose: Three foliar applications at 2.5 litre/ha

(49) Seaweed (*Ascophyllum nodosum*) extract 5% (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract per cent. by weight, minimum	5.0
(ii)	Bentonite per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	0.25
(ii)	Total organic carbon per cent. by weight, minimum	2.18
(iii)	pH (50% aqueous solution)	5.28± 0.5
(iv)	Bulk density (gram/cc)	1.055± 0.02

(a) Name of the crop: Chilli

(b) Dose: Three soil applications at 10 kg/ha

(50) Seaweed (*Ascophyllum nodosum*) extract 34.5% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i> extract) per cent. by weight, minimum	34.5
(ii)	Water per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	1.40
(ii)	Mannitol per cent. by weight, minimum	1.1
(iii)	Total organic carbon per cent. by weight, minimum	10
(iv)	pH (1% aqueous solution)	8.0- 10.0
(v)	Specific gravity	1.1- 1.2
(vi)	Total dissolved solids per cent. by weight, minimum	34
(vii)	Solubility per cent. by weight, minimum	99

(a) Name of the crop: Tomato

(b) Dose: Two foliar applications at 950 ml/ha

(51) Seaweed (*Kappaphycus alvarezii*) extract 13.5% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Kappaphycus alvarezii</i>) extract per cent. by weight, minimum	13.5
(ii)	Organic (Acetic acid, Citric acid, Lactic acid) per cent. by weight, minimum	1.5
(iii)	Water per cent. by weight, maximum	85.0
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	D- Galactose-4-O-sulphate per cent. by weight, minimum	2.5
(ii)	Total carbohydrates per cent. by weight, minimum	4.8
(iii)	pH (50% aqueous solution)	3.0- 5.0
(iv)	Total organic carbon per cent. by weight, minimum	3.8
(v)	Total dissolved solids per cent. by weight, minimum	15
(vi)	Specific gravity	1.09- 1.14
(vii)	Solubility per cent. by weight, minimum	80

(a) Name of the crop: Cucumber and Tomato

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Cucumber	Three foliar applications at 2 litre/ha
(ii)	Tomato	Four foliar applications at 1.25 litre/ha

(52) Seaweed (*Sargassum tennerimum*) extract 10% (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Sargassum tennerimum</i>) extract per cent. by weight, minimum	10.0
(ii)	Ferrous sulphate per cent. by weight, minimum	1.29
(iii)	Bentonite per cent. by weight, maximum	88.71
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	1.02
(ii)	Total organic carbon per cent. by weight, minimum	7.0

(iii)	pH (10% aqueous solution)	7.0 – 8.0
(iv)	Bulk density (gram/cc)	1.15- 1.19

(a) Name of the crop: Mustard

(b) Dose: Three soil applications at 10kg/ha

(53) Seaweed (*Gracilaria salicornia*, *Gracilaria edulis*, *Sargassum tenerrimum*, *Sargassum wightii* and *Sargassum polycystum*) extract 3% (Granule)

Composition		
S. No.	Ingredients	Content
(i)	<i>Gracilaria salicornia</i> powder per cent. by weight, minimum	0.75
(ii)	<i>Gracilaria edulis</i> powder per cent. by weight, minimum	0.75
(iii)	<i>Sargassum tennerimum</i> powder extract per cent. by weight, minimum	0.50
(iv)	<i>Sargassum wightii</i> powder per cent. by weight, minimum	0.50
(v)	<i>Sargassum polycystum</i> powder per cent. by weight, minimum	0.50
(vi)	Bentonite per cent. by weight, minimum	20
(vii)	Dolomite per cent. by weight, minimum	77
(viii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginate acid per cent. by weight, minimum	0.02
(ii)	Agar per cent. by weight, minimum	0.03
(iii)	pH (10% aqueous solution)	6.5- 9.5
(iv)	Bulk density (gram/cc)	0.8- 1.3
(v)	Total organic carbon per cent. by weight, minimum	3.8

(a) Name of the crop: Wheat

(b) Dose: One soil application at 25 kg/ha

(54) Seaweed (*Laminaria digitata*) extract 10% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Laminaria digitata</i>) extract per cent. by weight, minimum	10
(ii)	Water per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginate acid per cent. by weight, minimum	1.0

(ii)	Total organic carbon per cent. by weight, minimum	1.8
(iii)	pH (10% aqueous solution)	3.60 ± 0.5
(iv)	Specific gravity	1.0- 1.10
(v)	Total dissolved solids per cent. by weight, minimum	9.0
(vi)	Solubility per cent. by weight, minimum	99.0

(a) Name of the crop: Chilli

(b) Dose: Four foliar applications at 625 ml/ha”;

(c) in sub-heading 3, “Botanical Extract”, after serial number (12) and the entries relating thereto, the following serial numbers and entries shall be inserted, namely:-

“(13) Botanical (*Medicago sativa*, *Camellia sinensis*, *Tinospora cordifolia*) extract 64% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	<i>Medicago sativa</i> extract per cent. by weight, minimum	24
(ii)	<i>Camellia sinensis</i> extract per cent. by weight, minimum	22
(iii)	<i>Tinospora cordifolia</i> extract per cent. by weight, minimum	18
(iv)	Clove oil per cent. by weight, maximum	1
(v)	Water per cent. by weight, maximum	35
(vi)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Polyphenol per cent. by weight, minimum	0.1
(ii)	Total organic carbon per cent. by weight, minimum	25
(iii)	Total dissolved solids per cent. by weight, minimum	55
(iv)	pH (1% aqueous solution)	4.0- 6.0
(v)	Specific gravity	1.0 ± 0.02
(vi)	Solubility per cent. by weight, minimum	95

(a) Name of the crop: Tomato

(b) Dose: Five foliar applications at 1.5 litre/ha

(14) Botanical (*Citrus aurantium* and palm oil) extract 18% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	<i>Citrus aurantium</i> (bitter orange) extract per cent. by weight, maximum	8

(ii)	Palm oil fatty acid blend (Octanoic acid) per cent. by weight, maximum	10
(iii)	Glycerine per cent. by weight, maximum	35
(iv)	Ethanol per cent. by weight, maximum	6
(v)	Water per cent. by weight, maximum	39
(vi)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Total flavonoids per cent. by weight, minimum	5
(ii)	Saponins per cent. by weight, minimum	5
(iii)	Total dissolved solids per cent. by weight, minimum	20.08
(iv)	pH (50% aqueous solution)	2.75 ± 0.5
(v)	Specific gravity	1.09 ± 0.02
(vi)	Solubility per cent. by weight, minimum	99.62
(vii)	Total organic carbon per cent. by weight, minimum	34.53

(a) Name of the crop: Chilli

(b) Dose: Three foliar applications at 0.5 liter/ha

(15) Green algae (*Chlorella vulgaris*) extract 98% (liquid)

Composition		
S. No.	Ingredients	Content
(i)	<i>Chlorella vulgaris</i> aqueous extract per cent. by weight, minimum	98
(ii)	Sodium benzoate per cent. by weight, minimum	1
(iii)	EDTA as stabilizer per cent. by weight, maximum	1
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Free amino acids per cent. by weight, minimum	3.0
(ii)	Flavonoids per cent. by weight, minimum	0.05
(iii)	Total phenolics per cent. by weight, minimum	0.02
(iv)	Total dissolved solids per cent. by weight, minimum	10.0
(v)	Total organic carbon dry weight basis per cent. by weight, minimum	45
(vi)	pH (10% aqueous solution)	7.5- 8.5
(vii)	Specific gravity	1.01- 1.11
(viii)	Solubility per cent. by weight, minimum	90

(a) Name of the crop: Rice

(b) Dose: One foliar application at 2.5 litre/ha”;

(d) in sub-heading 4, “Mixed formulation of Biostimulants”, after serial number (48) and the entries relating thereto, the following serial numbers and entries shall be inserted, namely:-

“(49) Mixture of Botanical extract and Seaweed extract (Liquid)”

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Sargassum tenerrimum</i>) extract per cent. by weight, minimum	3
(ii)	<i>Camellia</i> seed extract per cent. by weight, minimum	20
(iii)	Emulsifier (MEI- 25) per cent. by weight, maximum	20
(iv)	Dimethylformamide per cent. by weight, maximum	57
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, maximum	0.005
(ii)	Total saponin content per cent. by weight, minimum	1.60
(iii)	pH (1% aqueous solution)	7.05± 0.5
(iv)	Specific gravity	1.032± 0.02
(v)	Total dissolved solids per cent. by weight, minimum	36
(vi)	Total organic carbon per cent. by weight, minimum	6.35
(vii)	Solubility per cent. by weight, minimum	99.65

(a) Name of the crop: Brinjal

(b) Dose: Three foliar applications at 750 ml/ha

“(50) Mixture of Seaweed extract and protein hydrolysate (plant source) (Liquid)”

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) fermented extract per cent. by weight, minimum	10
(ii)	Protein hydrolysate derived from maize kernel (Plant source) through enzymatic (<i>Saccharomyces protease</i>)) hydrolysis per cent. by weight, minimum	20
(iii)	Lactic acid per cent. by weight, maximum	0.5
(iv)	Water per cent. by weight, maximum	69.5
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	0.30
(ii)	Total amino acid per cent. by weight, minimum	20

(iii)	Free amino acid per cent. by weight, minimum	20
(iv)	Mannitol per cent. by weight, minimum	0.20
(v)	pH (10% aqueous solution)	7.5- 9.5
(vi)	Specific gravity	1.0- 1.2
(vii)	Total soluble solids per cent. by weight, minimum	22
(viii)	Organic carbon per cent. by weight, minimum	14.53
(ix)	Solubility per cent. by weight, minimum	99.0

(a) Name of the crop: Grapes

(b) Dose: Two foliar applications at 1250 ml/ha

(51) Mixture of Seaweed extract and protein hydrolysate (plant source) (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) fermented extract per cent. by weight, minimum	8.0
(ii)	Protein hydrolysate derived from de oiled cake of soybean (Plant source) through enzymatic (Papain) hydrolysis per cent. by weight, minimum	3.0
(iii)	Polysaccharides (corn starch) per cent. by weight, minimum	37.0
(iv)	China clay per cent. by weight, maximum	52.0
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Total amino acid per cent. by weight, minimum	2.0
(ii)	Free amino acids per cent. by weight, minimum	2.0
(iii)	Alginic acid per cent. by weight, minimum	0.02
(iv)	pH (10% aqueous solution)	8.0 – 10.0
(v)	Bulk density (gram/cc)	0.5- 0.75
(vi)	Total organic carbon per cent. by weight, minimum	29

(a) Name of the crop: Tomato

(b) Dose: Two foliar applications at 1000 g/ha

(52) Mixture of Seaweed extract and protein hydrolysate (plant source) (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) fermented extract per cent. by weight, minimum	2.0

(ii)	Protein hydrolysate derived from de oiled cake of soybean (Plant source) through enzymatic (Papain) hydrolysis per cent. by weight, minimum	0.20
(iii)	Bentonite per cent. by weight, maximum	97.80
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	0.001
(ii)	Total amino acid per cent. by weight, minimum	0.1
(iii)	Free amino acids per cent. by weight, minimum	0.1
(iv)	pH (10% aqueous solution)	7.0- 8.0
(v)	Bulk density (gram/cc)	1.0- 1.1
(vi)	Total organic carbon per cent. by weight, minimum	4.65

(a) Name of the crop: Cotton

(b) Dose: Two soil applications at 15 kg/ha

(53) Mixture of Seaweed extract, protein hydrolysate (plant source) and humic acid (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) fermented extract per cent. by weight, minimum	2.10
(ii)	Protein hydrolysate derived from maize kernel (Plant source) through enzymatic (Saccharomyces protease)) hydrolysis per cent. by weight, minimum	0.25
(iii)	Humic acid (as Potassium humate) (Source: Leonardite) per cent. by weight, minimum	0.10
(iv)	Mixture of Zeolite and Dolomite (45: 55), per cent. by weight, maximum	97.55
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	0.01
(ii)	Total amino acid per cent. by weight, minimum	0.20
(iii)	Free amino acid per cent. by weight, minimum	0.20
(iv)	Humic acid per cent. by weight, minimum	0.03
(v)	pH (10% aqueous solution)	8.0 – 10.0
(vi)	Bulk density (gram/cc)	1.02- 1.13
(vii)	Total organic carbon per cent. by weight, minimum	4.65

(a) Name of the crop: Onion

(b) Dose: One soil application at 15 kg/ha

(54) Mixture of Seaweed extract and amino acid (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Potassium humate per cent. by weight, minimum	0.95
(ii)	Amino acid per cent. by weight, minimum	0.10
(iii)	Water per cent. by weight, maximum	12.0
(iv)	Bentonite per cent. by weight, maximum	QS
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Humic acid per cent. by weight, minimum	0.50
(ii)	Amino acid per cent. by weight, minimum	0.10
(iii)	pH (50% aqueous solution)	6.0- 7.5
(iv)	Bulk density (gram/cc)	0.9- 1.20
(v)	Total organic carbon per cent. by weight, minimum	0.75

(a) Name of the crop: Paddy

(b) Dose: Two soil applications at 12.5 kg/ha

(55) Mixture of fulvic acid, seaweed extract and vitamin (Powder)

Composition		
S. No.	Ingredients	Content
(i)	Fulvic acid per cent. by weight, minimum	18.0
(ii)	Seaweed (<i>Ascophyllum nodosum</i>) Extract per cent. by weight, minimum	8.0
(iii)	Ascorbic acid per cent. by weight, minimum	8.0
(iv)	Anhydrous dextrose per cent. by weight	QS
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Fulvic acid per cent. by weight, minimum	18.0
(ii)	Alginic acid per cent. by weight, minimum	0.60
(iii)	Ascorbic acid per cent. by weight, minimum	8.0
(iv)	pH (1 % aqueous solution)	4.0 ± 1.0
(v)	Bulk density (gram/cc)	0.8 - 1.0
(vi)	Total organic carbon per cent. by weight, minimum	12
(vii)	Solubility per cent. by weight, minimum	96

(a) Name of the crop: Okra

(b) Dose: Two foliar applications at 1 kg/ha

(56) Mixture of vitamin and biochemical (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Vitamin B3 per cent. by weight, minimum	3.75
(ii)	Salicylic Acid	2.00
(iii)	Nonyl phenol ethoxylate (Emulsifier)	10
(iv)	Dimethyl sulfoxide (Solvent)	84.25
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Vitamin B3 per cent. by weight, minimum	3.75
(ii)	Salicylic acid per cent. by weight, minimum	2.00
(iii)	pH (10% aqueous solution)	3.0 - 4.0
(iv)	Specific gravity	1.0-1.15
(v)	Total organic carbon per cent. by weight, minimum	57.70
(vi)	Solubility per cent. by weight, minimum	96.59

(a) Name of the crop: Okra

(b) Dose: Two foliar applications at 12.5 mL/ha

(57) Mixture of Seaweed (*Ascophyllum nodosum*) extract, Vinasse and corn steep liquor- 63.6% (Liquid)*

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Ascophyllum nodosum</i>) extract per cent. by weight, minimum	18.2
(ii)	Vinasse per cent. by weight, minimum	40.0
(iii)	Corn steep liquor per cent. by weight, minimum	5.4
(iv)	Disodium molybdate per cent. by weight, minimum	1.30
(v)	Mn- EDTA per cent. by weight, minimum	15.40
(vi)	Asparagine per cent. by weight, minimum	0.1
(vii)	L- tryptophan per cent. by weight, minimum	0.1
(viii)	Vitamin B1 per cent. by weight, minimum	0.05
(ix)	Vitamin B6 per cent. by weight, minimum	0.05
(x)	Vitamin B3 per cent. by weight, minimum	0.1
(xi)	Vitamin B5 per cent. by weight, minimum	0.05
(xii)	Starch per cent. by weight, minimum	0.50
(xiii)	BIT (Benzisothiazolinone)10% water solution per cent. by weight, minimum	0.475

(xiv)	2- phenoxyethanol per cent. by weight, minimum	0.50
(xv)	Alkylpolyglucosides per cent. by weight, minimum	0.75
(xvi)	Water per cent. by weight	Q.S.
(xvii)	Total (per cent.)	100

Specifications

S. No.	Parameters	Value
(i)	Alginic acid per cent. by weight, minimum	0.50
(ii)	Fucoidan per cent. by weight, minimum	2.00
(iii)	Specific gravity	1.25 ± 0.05
(iv)	Total dissolved solids per cent. by weight, minimum	6.14
(v)	pH (10% aqueous solution)	7.5 ± 0.5
(vi)	Organic carbon per cent. by weight, minimum	14.30
(vii)	Solubility per cent. by weight, minimum	100

* Note: The product label shall include caution: Use only as directed. Avoid direct contact with eyes and skin. Keep out of reach of children. In case of accidental exposure or ingestion, seek medical advice immediately.

(a) Name of the crop: Maize, Rice, Soybean

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Maize	Seed treatment at 200 ml/100kg seed
(ii)	Rice	Seed treatment at 100 ml/100kg seed
(iii)	Soybean	Seed treatment at 200 ml/100kg seed

(58) Mixture of amino acid and biochemical (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	L-Proline per cent. by weight, minimum	1.72
(ii)	Betaine hydrochloride per cent. by weight, minimum	1.21
(iii)	Potassium phosphate tribasic anhydrous per cent. by weight, minimum	2.49
(iv)	Propylene glycol per cent. by weight, minimum	36.84
(v)	Water per cent. by weight	QS
(vi)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	L-Proline per cent. by weight, minimum	1.72

(ii)	Betaine hydrochloride per cent. by weight, minimum	1.21
(iii)	pH (1% aqueous solution)	7.84 ± 0.5
(iv)	Specific gravity	1.06 ± 0.02
(v)	Total organic carbon per cent. by weight, minimum	6.83
(vi)	Solubility per cent. by weight, minimum	100

(a) Name of the crop: Potato and chilli

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Potato	Three foliar applications at 700 ml/ha
(ii)	Chilli	Two foliar applications at 350 ml/ha

(59) Mixture of humic substances and protein hydrolysate (plant source) (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Humic substances (Source: Renewable Agri biomass) per cent. by weight, minimum	20.00
(ii)	Protein hydrolysate derived from soybean meal through enzymatic (protease) hydrolysis, per cent. by weight, minimum	17.00
(iii)	Water per cent. by weight, maximum	QS
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Humic acid per cent. by weight, minimum	1.50
(ii)	Fulvic acid per cent. by weight, minimum	6.00
(iii)	Total amino acid per cent. by weight, minimum	11.90
(iv)	Free amino acid per cent. by weight, minimum	0.50
(v)	Total dissolved solids per cent. by weight, minimum	25.0
(vi)	Total organic carbon per cent. by weight, minimum	10.46
(vii)	pH (1% aqueous solution)	7.5 - 9.5
(viii)	Specific gravity	1.15 - 1.25
(ix)	Solubility per cent. by weight, minimum	100

(a) Name of the crop: Pomegranate

(b) Dose: Two soil applications at 3750 ml/ha

(60) Mixture of organic molasses and protein hydrolysate (plant source) (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Organic molasses, per cent. weight by volume, minimum	20.00
(ii)	Protein hydrolysate derived from defatted soy flour (plant source) through enzymatic (alkaline protease) hydrolysis per cent. weight by volume, minimum	18.00
(iii)	Water per cent. by weight, maximum	62.00
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Total free amino acid per cent. by weight, minimum	1.24
(ii)	Total organic carbon per cent. by weight, minimum	8.0
(iii)	pH (10 % aqueous solution)	4.04 ± 0.5
(iv)	Solubility per cent. by weight, minimum	99.94
(v)	Specific gravity	1.016 ± 0.02
(vi)	Total dissolved solids per cent. by weight, minimum	38

(a) Name of the crop: Chickpea

(b) Dose: Three soil applications at 2.5 litre/ha

(61) Mixture of Seaweed extract and humic acid (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Seaweed (<i>Sargassum tenerrimum</i>) extract per cent. by weight, minimum	10.0
(ii)	Humate compound (Source: Lignite) per cent. by weight, minimum	10.0
(iii)	Octyl phenol ethoxylate (surfactant) per cent. by weight, minimum	0.2
(iv)	Sodium lignosulfonate (dispersing agent) per cent. by weight, minimum	2.2
(v)	Sodium sulfate (stabilizer) per cent. by weight, minimum	2.0
(vi)	Benzisothiazolinone (preservative) per cent. by weight, minimum	0.10
(vii)	Humectant (Glycerol) per cent. by weight, minimum	3.5
(viii)	Water per cent. by weight	QS
(ix)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value

(i)	Alginic acid per cent. by weight, minimum	0.10
(ii)	Humic acid per cent. by weight, minimum	5.0
(iii)	pH (1 % aqueous solution)	9.5± 0.5
(iv)	Specific gravity	1.08 ± 0.02
(v)	Total organic carbon per cent. by weight, minimum	6.0
(vi)	Solubility per cent. by weight, minimum	98
(vii)	Total dissolved solids per cent. weight by volume, minimum	20

(a) Name of the crop: Paddy, sugarcane, chilli, tomato and cotton

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Paddy	Two foliar applications at 500 ml/ha
(ii)	Sugarcane	Two foliar applications at 500 ml/ha
(iii)	Chilli	Two foliar applications at 500 ml/ha
(iv)	Tomato	Two foliar applications at 500 ml/ha
(v)	Cotton	Two foliar applications at 500 ml/ha”;

(e) in sub-heading 5, “Protein Hydrolysates and amino acids”, after serial number (37) and the entries relating thereto, the following serial numbers and entries shall be inserted, namely:-

“(38) Fermentation extract from (*Glycine max* and *Zea mays* leaves) (Plant source) 12% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Fermentation extract from (<i>Glycine max</i> and <i>Zea mays</i> leaves) (Plant origin) per cent. by weight, minimum	12
(ii)	Glutamic acid per cent. by weight, minimum	2
(iii)	Octhilinone per cent. by weight, minimum	0.3
(iv)	Water per cent. by weight, maximum	QS
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Total amino acids per cent. by weight, minimum	12
(ii)	Free amino acids per cent. by weight, minimum	10
(iii)	pH (1% aqueous solution)	4.5- 6.0
(iv)	Total organic carbon per cent. by weight, minimum	22.5
(v)	Specific gravity	1.19± 0.02
(vi)	Solubility per cent. by weight, minimum	100
(vii)	Total dissolved solids per cent. by weight, minimum	6.0

(a) Name of the crop: Chickpea

(b) Dose: Two foliar applications at 1.25 litre/ha

(39) Glycine 0.2% (Liquid)*

S. No.	Ingredients	Content
(i)	Glycine per cent. by weight, minimum	0.2
(ii)	Calcium alkyl benzene sulphonate per cent. by weight, minimum	3
(iii)	Polyoxyethylene ether per cent. by weight, minimum	3
(iv)	Cyclohexanone (solvent) per cent. by weight	QS
(v)	Total (per cent.)	100
S. No.	Parameters	
(i)	Glycine per cent. by weight, minimum	0.2
(ii)	Total organic carbon per cent. by weight, minimum	57.66
(iii)	pH (1% aqueous solution)	5.14 ± 0.5
(iv)	Specific gravity	0.966 ± 0.02
(v)	Solubility per cent. by weight, minimum	100

* Note: The product label shall include caution: Use only as directed. Avoid direct contact with eyes. Keep out of reach of children. In case of accidental exposure or ingestion, seek medical advice immediately.

(a) Name of the crop: Tomato

(b) Dose: Three foliar applications at 375 ml/ha

(40) Tryptophan 0.1% (Liquid)*

Composition		
S. No.	Ingredients	Content
(i)	Tryptophan per cent. by weight, minimum	0.1
(ii)	Calcium alkyl benzene sulphonate per cent. by weight, minimum	3.0
(iii)	Polyoxyethylene ether per cent. by weight, minimum	3.0
(iv)	Cyclohexanone per cent. by weight, minimum	93.9
(v)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Tryptophan per cent. by weight, minimum	0.1
(ii)	Total organic carbon per cent. by weight, minimum	57.69
(iii)	pH (1% aqueous solution)	6.61 ± 0.5
(iv)	Specific gravity	0.951 ± 0.02
(v)	Solubility per cent. by weight, minimum	100

* Note: The product label shall include caution: Use only as directed. Avoid direct contact with eyes. Keep out of reach of children. In case of accidental exposure or ingestion, seek medical advice immediately.

(a) Name of the crop: Tomato

(b) Dose: Three foliar applications at 870 ml/ha

(41) Protein hydrolysate (plant source) 16.94% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Protein hydrolysate derived from maize kernel through acid (hydrochloric acid) hydrolysis per cent. by weight, minimum	16.94
(ii)	Polysorbate 80 per cent. by weight, minimum	2.0
(iii)	Water per cent. by weight, maximum	Q.S
(iv)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Total amino acids per cent. by weight, minimum	14.40
(ii)	Free amino acids per cent. by weight, minimum	3.41
(iii)	Total organic carbon per cent. by weight, minimum	18.10
(iv)	pH (50% aqueous solution)	6.07 ± 0.5
(v)	Specific gravity	1.179 ± 0.02
(vi)	Solubility per cent. by weight, minimum	99.86

(a) Name of the crop: Tomato

(b) Dose: Two foliar applications at 2.5 litre/ha

(42) Protein hydrolysate (plant source) 7.63 % (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Protein hydrolysate derived from maize kernel through acid (hydrochloric acid) hydrolysis per cent. by weight, minimum	7.63
(ii)	Bentonite per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Total amino acids per cent. by weight, minimum	3.41
(ii)	Free amino acids per cent. by weight, minimum	3.41
(iii)	Total organic carbon per cent. by weight, minimum	6.67
(iv)	pH (50% aqueous solution)	6.53 ± 0.5
(v)	Bulk density (gram/cc)	0.907 ± 0.02

(a) Name of the crop: Tomato

(b) Dose: One soil application at 11.25 kg/ha

(43) Protein hydrolysate (plant source) 15.38% (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Protein hydrolysate derived from wheat seeds (plant source) by enzymatic (alkaline protease) hydrolysis per cent. by weight, minimum	15.38
(ii)	Bentonite per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Free amino acids per cent. by weight, minimum	1.47
(ii)	Total amino acids per cent. by weight, minimum	13.39
(iii)	Total organic carbon per cent. by weight, minimum	6.76
(iv)	pH (50% aqueous solution)	6.48 ± 0.5
(v)	Bulk density (gram/cc)	0.901 ± 0.02

(a) Name of the crop: Chilli

(b) Dose: Three soil applications at 12.5 kg/ha

(44) Protein hydrolysate 57% (Plant source) (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Protein hydrolysate derived from Guar Korma through alkaline (KOH) hydrolysis per cent. by weight, minimum	57.0
(ii)	Potassium salt per cent. by weight, maximum	5.5
(iii)	Sodium methyl parabens per cent. by weight, maximum	0.9
(iv)	Sodium propyl parabens per cent. by weight, maximum	0.1
(v)	Water per cent. by weight, maximum	36.5
(vi)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Amino acids per cent. by weight, minimum	30
(ii)	pH (1% aqueous solution)	6.8- 7.4
(iii)	Specific gravity	1.180 ± 0.02
(iv)	Total dissolved solids per cent. by weight, minimum	55
(v)	Total organic carbon per cent. by weight, minimum	22.20
(vi)	Solubility per cent. by weight, minimum	95

(a) Name of the crop: Bottlegourd

(b) Dose: Two foliar applications at 500 ml/ha

(45) Vinasse fermentation metabolites 52% (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Vinasse (sugarcane molasses) fermentation metabolites by yeast (<i>Sacchromyces cerevisiae</i>) per cent. by weight, minimum	52
(ii)	Water per cent. by weight, maximum	48
(iii)	Total (per cent.)	QS
Specifications		
S. No.	Parameters	Value
(i)	Total amino acids per cent. by weight, minimum	2.0
(ii)	Free amino acids per cent. by weight, minimum	0.8
(iii)	pH (1% aqueous solution)	5.7-6.7
(iv)	Total organic carbon per cent. by weight, minimum	12.75
(v)	Bulk density (gram/cc)	1.31± 0.02

(a) Name of the crop: Soybean, blackgram, cotton, sugarcane and wheat

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Soybean	One foliar application at 2500 mL/ha
(ii)	Blackgram	One foliar application at 1500 mL/ha
(iii)	Cotton	Two foliar application at 1500 mL/ha
(iv)	Sugarcane	One foliar application at 1000 mL/ha
(v)	Wheat	One foliar application at 1000 mL/ha

(46) Amino acid (L- Proline) 1.72 % (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	L-Proline per cent. by weight, minimum	1.72
(ii)	Potassium phosphate tribasic anhydrous per cent. by weight, minimum	2.50
(iii)	Phosphoric acid (1 M) per cent. by weight, minimum	3.97
(iv)	Propylene glycol per cent. by weight, minimum	36.84
(v)	Water per cent. by weight	QS
(vi)	Total (per cent.)	100
Specifications		

S. No.	Parameters	Value
(i)	L-Proline per cent. by weight, minimum	1.72
(ii)	pH (1% aqueous solution)	8.63 ± 0.5
(iii)	Specific gravity	1.05 ± 0.02
(iv)	Total organic carbon per cent. by weight, minimum	6.03
(v)	Solubility per cent. by weight, minimum	100

(a) Name of the crop: Maize and cotton

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Maize	Two foliar applications at 500 ml/ha
(ii)	Cotton	One foliar application at 500 ml/ha

(47) Protein hydrolysate 15.68 % (plant source) (Liquid)

Composition		
S. No.	Ingredients	Content
(i)	Protein hydrolysate derived from hydrolysis of wheat (plant source) by enzymatic (alkaline protease) hydrolysis per cent. by weight, minimum	15.68
(ii)	Water per cent. by weight, maximum	QS
(iii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Free amino acids per cent. by weight, minimum	12.98
(ii)	Total organic carbon per cent. by weight, minimum	18.01
(iii)	pH (50% aqueous solution)	6.18 ± 0.5
(iv)	Specific gravity	1.179 ± 0.02
(v)	Solubility per cent. by weight, minimum	99.91

(a) Name of the crop: Chilli

(b) Dose: Three foliar applications at 2.5 litre/ha”;

(f) in sub-heading 8, “Bio-chemicals”, after serial number (5) and the entries relating thereto, the following serial numbers and entries shall be inserted, namely:-

“(6) Biochemical (Stigmasterol and Campesterol) 0.01 % (Liquid)*

Composition		
S. No.	Ingredients	Content
(i)	Stigmasterol and Campesterol (Source: <i>Glycine max</i>) per cent. by weight, minimum	0.01
(ii)	1- (3, 4-dichlorophenyl)-2- (diethylamino) ethanol (adjuvant) per cent. by weight, minimum	0.70
(iii)	Sodium benzoate (stabilizer) per cent. by weight, maximum	0.30
(iv)	Mono ethylene glycol (Humectant) per cent. by weight, maximum	2.00
(v)	Alcohol alkoxylate (emulsifier) per cent. by weight, maximum	2.00
(vi)	Mustard seed oil base organic emulsifier (emulsifier) per cent. by weight, maximum	1.00
(vii)	Isopropyl alcohol (as solvent) per cent. by weight	QS
(viii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Stigmasterol and Campesterol per cent. by weight, minimum	0.01
(ii)	pH (1% aqueous solution)	5.0 – 8.0
(iii)	Specific gravity	0.79 - 0.82
(iv)	Solubility per cent. by weight, minimum	100
(v)	Total organic carbon per cent. by weight, minimum	30

* Note: The product label shall include caution: Use only as directed. Avoid direct contact with eyes. Keep out of reach of children. In case of accidental exposure or ingestion, seek medical advice immediately.

(a) Name of the crop: Tomato, Grapes, Mentha and Chilli

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Tomato	Two foliar applications at 500 ml/ha
(ii)	Grapes	Two foliar applications at 625ml/ha
(iii)	Mentha	Two foliar applications at 500 ml/ha
(iv)	Chilli	Two foliar applications at 1000 ml/ha

(7) Biochemical (β - Sitosterol and Stigmasterol) 0.05 % (Dry flowable)*

Composition		
S. No.	Ingredients	Content
(i)	β - Sitosterol and Stigmasterol (from natural extract of deodorized vegetable oil distillate meal) (Source: <i>Helianthus annuus</i>) per cent. by weight, minimum	0.05

(ii)	α -hydroxy 1 carboxy- 1,3,5- cyclohexatriene per cent. by weight, minimum	10.0
(iii)	Hexanoic, 2 (diethyl amino) ethyl ester citrate per cent. by weight, minimum	12.0
(iv)	Sodium benzoate (stabilizer) per cent. by weight, maximum	1.20
(v)	Polyethylene glycol-6000 (binder) per cent. by weight, maximum	2.00
(vi)	Alcohol alkoxylate (surfactant) per cent. by weight, maximum	2.00
(vii)	Sodium naphthalene formaldehyde sulphonate (surfactant) per cent. by weight, maximum	2.0
(viii)	Mustard seed oil base per cent. by weight, maximum	2.0
(ix)	Lactose, glucose and galactose (non-food grade) per cent. by weight	QS
(x)	Total (per cent.)	100

Specifications

S. No.	Parameters	Value
(i)	β - Sitosterol and Stigmasterol per cent. by weight, minimum	0.05
(ii)	pH (1% aqueous solution)	3.0 – 6.0
(iii)	Bulk density (gram/cc)	0.53-0.65
(iv)	Solubility per cent. by weight, minimum	99.95
(v)	Total organic carbon per cent. by weight, minimum	31

* Note: The product label shall include caution: Use only as directed. Avoid direct contact with eyes. Keep out of reach of children. In case of accidental exposure or ingestion, seek medical advice immediately.

(a) Name of the crop: Green Gram, grapes and chilli

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Green gram	Two foliar applications at 75 g/ha
(ii)	Grapes	Two foliar applications at 100 g/ha
(iii)	Chilli	Three foliar applications at 150g/ha

(8) Biochemical (Stigmasterol and Campesterol) 0.01 % (Granule)

Composition		
S. No.	Ingredients	Content
(i)	Stigmasterol and Campesterol (Source: <i>Glycine max</i>) per cent. by weight, minimum	0.01
(ii)	Potassium humate per cent. by weight, maximum	0.05
(iii)	Sodium benzoate (as stabilizer) per cent. by weight, maximum	0.02

(iv)	Mustard seed oil base (as emulsifier) per cent. by weight, maximum	0.50
(v)	Soybean seed oil base (as emulsifier) per cent. by weight, maximum	0.50
(vi)	Sodium alkyl benzene sulphonate (as emulsifier) per cent. by weight, maximum	1.0
(vii)	Gypsum (as carrier) per cent. by weight, maximum	QS
(viii)	Total (per cent.)	100
Specifications		
S. No.	Parameters	Value
(i)	Stigmasterol and Campesterol per cent. by weight, minimum	0.01
(ii)	pH (1% aqueous solution)	7.0 – 10.0
(iii)	Bulk density (gram/cc)	1.03 - 1.15
(iv)	Total organic carbon per cent. by weight, minimum	0.18

(a) Name of the crop: Paddy, Wheat, Potato, Sugarcane and Onion

(b) Dose: As given in table below:

Table

S. No.	Crop	Dose
(i)	Paddy	One soil application at 7.5 kg/ha
(ii)	Wheat	One soil application at 10 kg/ha
(iii)	Potato	One soil application at 10 kg/ha
(iv)	Sugarcane	Two soil applications at 10 kg/ha
(v)	Onion	One soil application at 10 kg/ha”

[F.No. 3-10/2025-Biostimulants]

FRANKLIN L KHOBUNG, Jt. Secy.

Note: The principal order was published in the Gazette of India, Extraordinary, Part-II, Section 3, Sub-section (i), *vide* G.S.R. number 758 (E), dated the 25th September, 1985 and was last amended, *vide* number S.O. 1335 (E), dated the 13th March, 2026.