

DATA ANALYTICS WITH R

HOMEWORK 3

JASNEEK SINGH CHUGH
JC2433

```
#Loading Required Packages
library(UsingR)
library(ggplot2)
```

#A

```
transactionData <- read.csv(file="/Users/jasneekchugh/Desktop/DS_NJIT/CS636-\ DA\ with\
R/Assignment/Assignment3/transactions.csv")
```

#1

#Numerical data: Data that takes numerical values and for which arithmetic operations make sense

#In transaction data, numerical variables are: payment_plan_days, plan_list_price, actual_amount_paid.


#Categorical data: Qualitative data that is limited to number of distinct categories.

#In transaction data, Categorical variables are: payment_method_id , is_auto_renew(0,1), is_cancel(0,1)

#2


```
hist(transactionData$payment_plan_days)
```

Histogram of transactionData\$payment_plan_days




```
hist(transactionData$plan_list_price)
```

Histogram of transactionData\$plan_list_price



```
hist(transactionData$actual_amount_paid)
```

Histogram of transactionData\$actual_amount_



#3

```
> table(transactionData$payment_method_id)
```

```

 1    2    3    4    5    6    7    8   10
12   52  210   15  474  466 1094  657 1326
11   12   13   14   15   16   17   18   19
2129 3834 6571 13621 1479 11064 7437 16177 32073
20   21   22   23   24   25   26   27   28
28278 22883 20130 42386 16196 13780 4591 62525 95733
29   30   31   32   33   34   35   36   37
113885 160957 252342 146481 411164 731539 541399 855115 1007689
38   39   40   41
1703590 1466655 2225283 11526454
```

```
> table(transactionData$is_auto_renew)
```

```

 0    1
3189796 18357950
```

```
> table(transactionData$is_cancel)
```

```

 0    1
20690895 856851
```

```
> #B
```

```
> #2.4
```

```
> weather1<- table(central.park$WX)
```

```
> weather1
```

```
1 18
```

```
10 11
```

```
> weather2<- table(central.park$WX, exclude = FALSE)
```

```
> weather2
```

```
1 18 <NA>
```

```
10 11 10
```

Table 2 (weather2) is better because it shows the number of NAs as well present in the "WX" column. So it gives more clarity of the data.

> #2.8

```
> attach(npdb)
> head(npdb)
> state<- sort(table(npdb$state))
> state[which.max(state)]
```

CA

1566

```
> detach(npdb)
> #CA has the maximum malpractice awards i.e. 1566
```


#2.9

```
table(ID)
sort(table(ID))
```


#the command table(ID) returns the frequency of each element in column ID. It's interesting because we can quickly see if there are any duplicates values in the column ID. Since there should be only one ID for each doctor.

> #2.10

```
> data(MLBattend)
> attach(MLBattend)
> win = wins[franchise == "NYA"]
> names(win) = c(1969:2000)
> detach(MLBattend)
> barplot(win,xlab = "YEAR", ylab = "Wons", col="red")
```




```
> dotchart(win)
```




```
> #2.16
> #1) It's a proportion so we divide it by the total number of elements in rivers dataset
> length(rivers[rivers < 500])/length(rivers)
[1] 0.5815603
> #2)
> length(rivers[rivers < mean(rivers)])/length(rivers)
[1] 0.6666667
> #3) We find quartile using the summary function. 0.75 is the third quartile
> rivers_summary<-summary(rivers)
> rivers_summary[5]
3rd Qu.
  680
```

```
> #2.23
> mean(npdb$amount)
[1] 166257.2
> median(npdb$amount)
[1] 37500
> quantile(npdb$amount)
  0%   25%   50%   75%  100%
  50  8750 37500 175000 25000000
> # So from output of quantile(npdb$amount) we can say that percentile of mean lies
between 50th and 75th percentile approx. 74.85 percentile.
> #It might be because the data is right skewed and Mean is greater than Median.
```


```
> #2.30
> hist(bumpers,xlab = "Bumpers",col = "green",border = "red")
```



```
> mean(bumpers)
[1] 2122.478
> median(bumpers)
[1] 2129
> sd(bumpers)
[1] 798.4574
> hist(firstchi,xlab = "firstchi",col = "green",border = "red")
```




```
> mean(firstchi)
[1] 23.97701
> median(firstchi)
[1] 23
> sd(firstchi)
[1] 6.254258
> hist(math,xlab = "math",col = "green",border = "red")
```




```
> mean(math)
[1] 54.9
> median(math)
[1] 54
> sd(math)
[1] 9.746264
```

```
> #2.32
> hist(pi2000, prob=TRUE, col = "green",border = "red")
```




```
> #The above distribution looks flat as all digits are equal So we subtract 0.1. So that the
bins for 0 and 1 do not gets combined, by using the argument breaks=0:10-.5.
> hist(pi2000,breaks=c(0:10,.5), prob=TRUE, col = "green",border = "red")
```




```
> #2.34
> DDT
[1] 2.79 2.93 3.22 3.78 3.22 3.38 3.18 3.33 3.34 3.06 3.07 3.56 3.08 4.64 3.34
```

```
> histogram(DDT, col = "grey",border = "red")
```



```
> boxplot(DDT, col = "grey",border = "red") #the small circle point highlights the outlier in the data
```



```
> mean(DDT)
```

```
[1] 3.328
```

```
> sd(DDT)
```

```
[1] 0.4371531
```

```
> #2.35
```

```
> names(state.area) <-state.abb
```

```
> length(state.area)
```

```
[1] 50
```

```
> length(state.area[state.area < state.area['NJ']])/50*100 #percentage of states with area less than NJ
```


```
[1] 8
```

```
> length(state.area[state.area < state.area['NY']])/50*100 #percentage of states with area less than NY
```

```
[1] 40
```




```
> hist(state.area)
```



```
> state.area[state.area > 5e5]#Outlier  
AK  
589757
```

```
> #2.36
```

```
> hist(nym.2002$time, xlab="Time", col = "grey", border = "red")
```



```
> mean(nym.2002$time)
```

```
[1] 268.5707
```

```
> median(nym.2002$time)
```


```
[1] 262.8417
```

#Mean is greater than median therefore, the data is right skewed and the tail of the histogram is towards right. The bulk data is represented in the middle by median.


```
> #2.39
```

```
> #Histograms are very useful to understand the pattern of variability in the data.
```

```
> hist(hall.fame$HR, xlab="HR", main="Histogram of HR", col = "red") #Shape: Right Skewed
```




```
> mean(hall.fame$HR)
[1] 85.1097
> median(hall.fame$HR)
[1] 51
> hist(hall.fame$BA,xlab="BA", main="Histogram of BA",col = "red")#Shape: Symmetric
```



(Mean and Median are almost equal), Smooth curve

```
> mean(hall.fame$BA)
[1] 0.2687739
> median(hall.fame$BA)
[1] 0.267
> hist(hall.fame$OBP,xlab="OBP", main="Histogram of OBP",col = "red") #Shape: Symmetric
```




(Mean and Median are almost equal), Smooth curve


```
> mean(hall.fame$OBP)
```

```
[1] 0.3360642
> median(hall.fame$OBP)
[1] 0.335
```

```
> #2.41
> x=rnorm(1000)
> boxplot(x,range=1.5,main="Box plot with Range 1.5")
> boxplot(x,range=0.5,main="Box plot with Range 0.5")
```



```
> boxplot(x,range=1,main="Box plot with Range 1")
> boxplot(x,range=2,main="Box plot with Range 2")
```




> #Boxplots whiskers are chosen with factor of 1.5 because, the width of the notches is proportional to the IQR and inversely proportion to square root of the size of the sample. The box plot has equal whisker length of $1.5 \times \text{IQR}$ for both whiskers.


> #Upper whisker- Above third quartile: $Q3 + 1.5 \times \text{IQR}$

> #Lower whisker- Below first quartile: $Q1 - 1.5 \times \text{IQR}$


```
> #2.42
> Mode <- function(x) {
+   uq <- unique(x)
+   uq[which.max(tabulate(match(x, uq)))]
+ }
> #Age
> hist(cfb$AGE,xlab="Age", col="blue")
```




```
>
> Mode(cfb$AGE) #We can also see from the histogram that the bar of the histogram is at
the peak.
[1] 40
> mean(cfb$AGE)
[1] 49.635
> median(cfb$AGE)
[1] 48
> #For Age, Mean is greater than the Median. Therefore, the data is skewed to the right.
Tail
> #is also towards the right of the histogram.
>
> #EDUC
> hist(cfb$EDUC,xlab="Educ",col="blue")
```




```
> Mode(cfb$EDUC)
[1] 12
> mean(cfb$EDUC)
[1] 13.075
> median(cfb$EDUC)
[1] 13
> #For EdDUC, Mean and Median are almost the same. However, as per the histogram the
data is skewed left. Tail of the histogram is on the left side.
>
> #NETWORTH
> as.numeric(cfb$NETWORTH)
> hist(cfb$NETWORTH,xlab="NetWorth",col="blue")
```



```
> mean(cfb$NETWORTH)
[1] 376993.4
> median(cfb$NETWORTH)
[1] 89947.5
> Mode(cfb$NETWORTH)
[1] 0
> #we can see that the data is so much skewed to the right. So we can transform the
NETWORTH using the logarithmic function to improve the skewness.
>
> #SAVINGS
> as.numeric(cfb$SAVING)
> hist(cfb$SAVING,col="blue")
```



```
> SAV<-log(cfb$SAVING+1)
> hist(SAV,col="blue")
```



```
> mean(SAV)
[1] 4.24621
> median(SAV)
```

```
[1] 5.303305
```

```
> Mode(SAV)#most of the people have savings of Zero and that is why mode is zero.
```

```
[1] 0
```

```
> #For SAVINGS, median is greater than mean, therefore data is skewed to left.
```

```
> #Tail of the histogram is also towards left.
```

I was able to convince myself for the charts they have for all the 4 variables. However, for "EDUC" variable after looking at the mean and median almost close, I was expecting the histogram to be nearly normally distributed however, the histogram is left skewed.


```
> #2.43
```

```
> summary(brightness)#it returns the useful summary like Min, Max, Median, of each column.
```

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
------	---------	--------	------	---------	------

2.070	7.702	8.500	8.418	9.130	12.430
-------	-------	-------	-------	-------	--------

```
> hist(brightness,col = "red")
```



```
> mean(brightness)
```

```
[1] 8.417743
```

```
> median(brightness)
```

```
[1] 8.5
```

```
> max(brightness)-min(brightness) #Range
```


```
[1] 10.36
```

```
> #2.44
```


```
> par(mfrow=c(1,2))
```

```
> hist(lawsuits,main="Histogram of Lawsuits")
```

```
> hist(log(lawsuits),main="Histogram of log(Lawsuits)")
```




```
> boxplot(lawsuits,ylab="Lawsuits",main="Boxplot of Lawsuits")
> boxplot(log(lawsuits),ylab="log(lawsuits)",main="Boxplot of log(Lawsuits)")
```




It would be bit difficult to guess where the middle 50% of the observations are, without the help of boxplot because the range of lawsuits is quite high. Hence, Boxplot helps us to see where actually the observations are. Also, after the logarithmic transformation it gives more clear picture of the data in the boxplot and show the outlier clearly.

```
> #2.45
> hist(exec.pay,col = "red")#the data is very much skewed to the right side
```



```
> mean(exec.pay)
[1] 59.88945
> median(exec.pay)
[1] 27
> transformed<-log(1+ exec.pay,10)
```

```
> hist(transformed = "red")
```



```
> mean(transformed)
```

```
[1] 1.438955
```

```
> median(transformed)
```

```
[1] 1.447158
```

```
> #after the transformation the data looks symmetric and the mean and the median of
```

```
> #the transformed data are also almost equal.
```

```
> #Histogram of transformed data is better because it looks symmetric and distributed normally.
```

```
> #With transformed histogram we get to see more clear bars for each.
```

```
> #The mean and Median of transformed data are almost equal.
```


Probability Distribution

Jasneek Singh Chugh

JC243

$$\Rightarrow P(U|W) = ?$$

Q	Pr(W R)	R	-R		Pr(U R)	R	R'	
	W	0.7	0.4	1.1	U	0.9	0.2	1.1
	W'	0.3	0.6	0.9	U'	0.1	0.8	0.9
		1.0	1.0	2		1.0	1.0	2.

$$Pr(R) = 0.8 \Rightarrow Pr(R') = 0.2$$

Given:-

$$Pr(UW|R') = Pr(U|R') \cdot Pr(W|R')$$

$$\text{and, } Pr(UW|R) = Pr(U|R) \cdot Pr(W|R)$$

$$\text{Also, } \boxed{Pr(U|W) = \frac{Pr(UW)}{Pr(W)}}$$

$$Pr(UW) = Pr(R) \cdot Pr(UW|R) + Pr(R') \cdot Pr(UW|R')$$

$$\Rightarrow Pr(R) \cdot Pr(U|R) \cdot Pr(W|R) + Pr(R') \cdot Pr(U|R') \cdot Pr(W|R')$$

$$= 0.8 \times 0.9 \times 0.7 + 0.2 \times 0.2 \times 0.4$$

$$\Rightarrow 0.504 + 0.016$$

$$Pr(UW) = 0.52$$

We know,

$$Pr(W) = Pr(R) \cdot Pr(W|R) + Pr(R') \cdot Pr(W|R')$$

$$= 0.8 \times 0.7 + 0.2 \times 0.4$$

$$\boxed{Pr(W) = 0.64}$$

$$\therefore Pr(U|W) = \frac{0.52}{0.64} = 0.8125$$

Ans