

This expanded module equips the chatbot to generate highly personalized recommendations for reversing and managing insulin resistance (IR). It maps clinical protocols to specific dietary preferences (Vegetarian, Non-Vegetarian, Vegan, Jain), age groups, biological sexes, and special conditions like pregnancy.

All recommendations are rooted in the core therapeutic goals: lowering glycemic load, optimizing fiber intake (25-35 g/day), managing visceral adiposity, and improving cellular insulin sensitivity.

PDF+ 1

2. Dietary Protocols by Preference

2.1 Vegetarian Diet (Focus on North Indian Profiles)

A vegetarian diet can effectively reverse IR if refined carbohydrates are replaced with complex, fiber-rich alternatives.

- **Core Carbohydrates:** Replace white rice and refined wheat (maida) with millets (bajra, jowar, ragi) and whole grains like brown rice or quinoa.
- **Protein Sources:** Emphasize high-quality proteins such as paneer (cottage cheese), moong dal, chana (chickpeas), and rajma (kidney beans). Moong dal chilla or besan chilla are excellent low-glycemic breakfast options.
- **Vegetables & Fiber:** Integrate heavy amounts of green leafy vegetables (palak, methi, sarson) and low-glycemic vegetables like bitter melon (karela) and bottle melon (lauki).
- **Fats:** Use healthy fats like ghee in moderation, alongside flaxseeds, almonds, and walnuts.

2.2 Non-Vegetarian Diet

Non-vegetarian diets naturally offer high protein, which helps stabilize blood sugar, but care must be taken to avoid saturated fats and processed meats.

- **Core Carbohydrates:** Pair proteins with high-fiber whole grains (millets, oats) to slow glucose absorption.
- **Protein Sources:** Prioritize lean meats such as skinless chicken breast, turkey, and eggs. Incorporate fatty fish (salmon, mackerel, sardines) 2-3 times a week for Omega-3 fatty acids, which combat cellular inflammation.

- **Foods to Avoid:** Strictly eliminate processed meats (sausages, bacon) and avoid deep-frying meats, which increases harmful trans fats and caloric load.

2.3 Vegan Diet

A vegan diet is highly anti-inflammatory but requires strategic planning to ensure adequate protein and avoid carbohydrate-heavy meals.

- **Core Carbohydrates:** Rely on low-glycemic complex carbs like quinoa, barley, and steel-cut oats.
- **Protein Sources:** Tofu, tempeh, edamame, lentils, and plant-based protein powders (pea or hemp).
- **Fats & Micronutrients:** Avocados, chia seeds, and pumpkin seeds. Vegans must monitor and supplement Vitamin B12 and Iron, as deficiencies can mimic fatigue associated with metabolic syndrome.

2.4 Jain Diet

The Jain diet excludes all root vegetables (onions, garlic, potatoes, carrots) and focuses strictly on non-violent, above-ground plant consumption.

- **Core Carbohydrates:** Use whole grains, millets, and brown rice. Replace potatoes with raw banana or bottle gourd in traditional recipes to drastically lower the glycemic index.
- **Protein Sources:** Moong dal, tuvar dal, chickpeas, and dairy products like paneer and unsweetened yogurt.
- **Vegetables & Fiber:** Maximize the use of cabbage, capsicum, tomatoes, peas, and leafy greens.

3. Demographic-Specific Guidelines

3.1 Age-Specific Protocols

Pediatric & Adolescents (Under 18)

- **Metabolic Focus:** Prevent early-onset type 2 diabetes without restricting essential calories needed for growth and puberty.
- **Diet:** Do not apply aggressive caloric deficits. Focus on eliminating sugary drinks, packaged juices, and refined bakery items. Encourage whole fruits over fruit juices.
- **Exercise:** Minimum 60 minutes of daily active play, sports, or aerobic exercise.

- **Screen Time:** Strictly limit recreational screen time to reduce sedentary behavior and protect sleep hygiene.

Adults (18-60 Years)

- **Metabolic Focus:** Manage occupational stress, weight gain, and lifestyle-induced IR.
- **Diet:** Follow the 0-100 risk scoring index (Low, Moderate, High) to titrate carbohydrate restriction. Target 5-10% weight loss if BMI is in the overweight/obese category.
- **Exercise:** Combine moderate-intensity aerobic exercise (mins/week) with resistance training 2-3 times a week.

Elderly / Old People (60+ Years)

- **Metabolic Focus:** Combat sarcopenia (age-related muscle loss) and declining basal metabolic rates, which naturally increase insulin resistance.
- **Diet:** Increase easily digestible protein intake to preserve muscle mass. Ensure meals are rich in fiber but soft to accommodate dental constraints.
- **Exercise:** Prioritize light resistance training and mobility work (e.g., resistance bands, water aerobics) to maintain the muscle's glucose storage capacity without risking joint injury.

3.2 Biological Sex-Specific Protocols

Men

- **Physiological Risk:** Men are biologically predisposed to accumulate visceral (abdominal) fat, which actively releases inflammatory cytokines that directly block insulin signaling pathways.
- **Clinical Focus:** Aggressive tracking of waist circumference (cm indicates high risk). Cardiovascular health must be prioritized due to the overlapping risks of high blood pressure and hypertriglyceridemia.
- **Intervention:** Focus heavily on High-Intensity Interval Training (HIIT) and heavy resistance training to mobilize stubborn visceral fat depots.

Women

- **Physiological Risk:** Pre-menopausal women are somewhat protected by estrogen, which optimizes insulin sensitivity. However, post-

menopause, the drop in estrogen leads to rapid central fat accumulation and heightened IR risk. Polycystic Ovary Syndrome (PCOS) is also a major driver of severe IR in younger women.

- **Clinical Focus:** Track waist circumference (cm indicates high risk).
- **Intervention:** Emphasize stress management and sleep, as women are highly susceptible to cortisol-induced glucose spikes. Post-menopausal women must prioritize weight-bearing exercises to combat both IR and osteoporosis.

3.3 Special Conditions: Pregnant Women (Gestational Insulin Resistance)

- **Metabolic Focus:** Pregnancy naturally induces a state of mild insulin resistance to ensure adequate glucose reaches the fetus. In some cases, this escalates to Gestational Diabetes Mellitus (GDM).
- **Dietary Protocol: No fasting or aggressive carbohydrate restriction.** The focus is strictly on *stabilizing* blood sugar. Use the "Plate Method": 50% non-starchy vegetables, 25% lean protein, and 25% complex carbohydrates. Eat smaller, frequent meals (3 meals, 2-3 snacks) to avoid sudden glycemic spikes or dips.
- **Exercise:** Unless contraindicated by an obstetrician, a 15-20 minute light walk after meals is highly effective at clearing postprandial glucose. Prenatal yoga and light swimming are also recommended.

PDF

- **Monitoring:** Requires specialized, frequent clinical tracking of fasting and post-meal glucose under the direct supervision of an endocrinologist and obstetrician.

4. Master Quick-Reference Action Matrix

Category	Recommended Diet Focus	Exercise & Lifestyle Priority	Key Clinical Marker to Track
Vegetarian (North Indian)	Millets (bajra/jowar), paneer, moong dal, heavy leafy greens (palak). PDF	150-300 mins aerobic + resistance . PDF	Fasting Insulin, HbA1c. PDF

Categor y	Recommended Diet Focus	Exercise & Lifestyle Priority	Key Clinical Marker to Track
Non- Vegetar ian	Lean chicken, eggs, fatty fish (omega-3s). Avoid processed/fried meats. PDF	Focus on visceral fat reduction PDF	Triglycerid es, HDL Cholesterol PDF
Vegan	Tofu, lentils, quinoa, chia seeds. Supplement B12 & Iron.	Resistanc e training to build muscle. PDF	Fasting Glucose. PDF
Jain	Bottle gourd, raw banana, moong dal. Avoid potatoes/root carbs.	Daily yoga and post-meal walks. PDF	Postprandi al Glucose.
Pediatric / Teens	No restrictive diets. Eliminate liquid sugars and packaged junk. PDF	60+ mins daily active play. Limit screen time.	BMI Percentile.
Adults (Men)	High protein, low glycemic load. Avoid alcohol & refined carbs. PDF	Heavy resistance training + HIIT. PDF	Waist Circumfere nce (cm). PDF
Adults (Wome n)	Anti- inflammatory foods, adequate	Cortisol managem ent	Waist Circumfere nce (cm).

Category	Recommended Diet Focus	Exercise & Lifestyle Priority	Key Clinical Marker to Track
	healthy fats (flaxseeds/almonds).	(yoga, sleep). PDF	PDF
Elderly (60+)	High-digestibility proteins, soft high-fiber foods to prevent spikes.	Light resistance bands, mobility work to prevent sarcopenia. PDF	Fasting Glucose, Muscle Mass.
Pregnant Women	Small, frequent meals. Complex carbs only. NO fasting.	15 min walks post-meals (if cleared by doctor). PDF	Frequent Glucose Monitoring (Doctor Led).