

Selected Solutions to Probability Theory and Examples by Rick Durrett

Student name: *Jason Kenyon*

Course: *Independent Study* – Professor: *Professor David Biddle*
Due date: *November 6, 2022*

Question 1.1.1

Let $\Omega = \mathbb{R}$, \mathcal{F} be the set of all subsets of \mathbb{R} such that it or its complement is countable. Let $P(A) = 0$ if A is countable and $P(A) = 1$ if A^c is countable. Then (Ω, \mathcal{F}, P) is a probability space.

Answer. We begin by showing that \mathcal{F} is a sigma algebra. Suppose that $x \in \mathcal{F}$. Then either x is countable or $\mathbb{R} \setminus x$ is countable. In the first case, $\mathbb{R} \setminus (\mathbb{R} \setminus x) = x \in \mathcal{F}$. In the latter, $\mathbb{R} \in \mathcal{F}$. Now suppose that $x_1, x_2, \dots \in \mathcal{F}$. If at least one of $\mathbb{R} \setminus x_i$ is countable, then so must $\bigcap_i \mathbb{R} \setminus x_i$. Otherwise, it must be that all x_i are uncountable, in which case $\bigcup_i x_i$ is too. Now, to demonstrate that our probability measure is well-defined, it is clear that by definition $P(x) \geq 0$ for any set $x \in \mathcal{F}$. Additionally, supposing $x_1, x_2, \dots \in \mathcal{F}$ are disjoint sets we have either $P(\bigcup_i x_i) = 0$ or $P(\bigcup_i x_i) = 1$. In the first case, we know that the union is countable, which implies that each of x_i must be countable, lest our union be uncountable. This is to say $P(\bigcup_i x_i) = \sum_i 0 = \sum_i P(x_i)$. In the second case, we have that $P(\bigcup_i x_i) = 1$. And notice that if x_i are disjoint then only a single member of that collection can have an uncountable complement, for otherwise they could not be disjoint. This is to say that $P(\bigcup_i x_i) = 1 + 0 + 0 + \dots = \sum_i P(x_i)$.

Question 1.1.2

Let $S_d = \{(a_1, b_1) \times \dots (a_n, b_n) : \}$, then $\sigma(S_d) = \sigma(\mathcal{B}_d)$.

Answer. We may represent an open set (a, b) where $a < b$ as follows:

$$(a, b) = \lim_{n \rightarrow \infty} \left(a, a + (b - a) - \frac{(b - a)}{n} \right]$$

Therefore, $S_d \subseteq \mathcal{B}_d$, whence $\sigma(S_d) = \sigma(\mathcal{B}_d)$.

Question 1.1.3

The Borel sigma algebra is countably generated; that is there exists a subset $A \subseteq \mathcal{B}_d$ such that $\sigma(A) = \sigma(\mathcal{B}_d)$