

Density-dependent selection in evolutionary genetics: a lottery model of Grime's triangle

Manuscript elements:

Keywords: r/K selection, absolute fitness, eco-evo, interference competition, competition-colonization trade-off.

Manuscript type: Article.

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

[Needs to be revised but won't do that now in case it has to become Nature introductory paragraph] Fitness is typically represented in heavily simplified terms in evolutionary genetics, often using constant selection coefficients. This excludes fundamental ecological factors such as dynamic population size or density-dependence from the most genetically-realistic treatments of evolution, a problem that inspired MacArthur's influential but problematic r/K theory. Following in the spirit of r/K -selection as a general-purpose theory of density-dependent selection, but grounding ourselves empirically in "primary strategy" trait classification schemes like Grime's triangle, we develop a new model of density-dependent selection which revolves around territorial contests. To do so, we generalize the classic lottery model of territorial acquisition, which has primarily been used for studying species co-existence questions, to accommodate arbitrary densities. We use this density-dependent lottery model to predict the direction of trait evolution under different environmental conditions and thereby provide a mathematical underpinning for Grime's verbal scheme. We revisit previous concepts of density-dependent selection, including r and K selection, and argue that our model distinguishes between different aspects of fitness in a more natural and intuitive manner.

“...the concept of fitness is probably too complex to allow of a useful mathematical development. Since it enters fundamentally into many population genetics considerations, it is remarkable how little attention has been paid to it.” — Warren J. Ewens, *Mathematical Population Genetics I*, 2004, pp. 276

Evolutionary models differ greatly in their treatment of fitness. In models of genetic evolution, genotypes are typically assigned constant (or occasionally frequency-dependent) selection coefficients describing the change in their relative frequencies over time. This simplified treatment of selection facilitates explicit time-dependent treatment of genotype frequencies, and can be justified over sufficiently short time intervals (Ewens, 2004, p. 276). The emphasis of population genetics is to infer past selection, migration and demographic change given a sample of nucleotide sequences, or to predict how allele frequencies change over time based on their relative fitness effects together with population structure, genetic drift and linkage. The resulting picture of evolution excludes basic elements of the ecological underpinnings of selection, including density dependence, and how selection affects population size. This complicates the inference of past selection, because demographic changes can look genealogically very similar to selective frequency changes (Barton, 1998).

By contrast, models of phenotypic trait evolution use absolute fitness functions to describe how some traits of interest affect survival and reproduction in particular ecological scenarios (Diekmann et al., 2004; Metz et al., 1992). These fitness functions can be quite problem-specific and often only account for a few traits at a time. The emphasis here is on the conditions for invasion from low frequencies and co-existence, rather than frequency or abundance trajectories over time. For instance, adaptive dynamics uses “invasion fitness” to explore the consequences of eco-evolutionary feedbacks (Diekmann et al., 2004).

It is challenging to generalize beyond particular traits or ecological scenarios to model fundamentally different forms of selection. Perhaps this is not surprising given that fitness is such a complex quantity, dependent on all of a phenotype’s functional traits (Violle et al., 2007) and its environment. A detailed, trait-based, predictive model of fitness would be enormously com-

plicated and situation-specific. It is therefore easy to doubt the feasibility of a simplified, general
45 mathematical treatment of fitness (Ewens, 2004, p. 276). For example, MacArthur's famous r/K
scheme (MacArthur, 1962; MacArthur and Wilson, 1967) is now almost exclusively known as a
framework for understanding life-history traits, and judged on its failure in that role (Boyce, 1984;
48 Pianka, 1970; Reznick et al., 2002; Stearns, 1977). However, the r/K scheme's original purpose
was to extend the existing population-genetic treatment of selection to account for population
density (MacArthur, 1962). Few attempts have been made to develop it further along these lines.

51 Empirical trait classification studies have suggested the existence of a few "primary strate-
gies", reflecting broadly distinct responses to selection (Winemiller et al., 2015). Grime famously
considered (Grime, 1974, 1977, 1988; Westoby, 1998) two broad determinants of population den-
54 sity: stress (persistent hardship e.g. due to resource scarcity or unfavorable temperatures) and
disturbance (intermittent destruction of vegetation e.g. due to trampling, herbivory, pathogens,
extreme weather or fire). The extremes of these two factors define three primary strategies de-
57 noted by C/S/R respectively (Fig. 1): competitors "C" excel in low stress, low disturbance envi-
ronments; stress tolerators "S" excel in high stress, low disturbance environments; and ruderals
"R" excel in low stress, high disturbance environments. Population persistence is not possible in
60 high-stress, high-disturbance environments. Grime showed that measures of C, S and R across
a wide range of plant species are anti-correlated, so that strong C-strategists are weak S and R
strategists, and so on, creating a triangular C/S/R ternary plot (Grime, 1974). Similar schemes
63 were proposed for insects (Southwood, 1977), fishes (Winemiller and Rose, 1992), and zooplank-
ton (Allan, 1976). More recently, modern hierarchical clustering techniques have revealed distinct
trait clusters in corals analogous to Grime's primary strategies (Darling et al., 2012). These em-
66 pirical findings suggest that functional traits contribute to fitness predominantly via a few key
factors such as intrinsic reproductive rate or stress-tolerance.

Here we explore the interplay between some "key factors" of fitness in a simplified, terri-
69 torial model of growth, dispersal and competition. This broadly follows the original spirit of
MacArthur's r/K scheme. More specifically, our aim is to begin merging some ecological realism



Figure 1: Schematic of Grime's triangle. The two axes show increasing levels of environmental stress and disturbance, respectively. Population persistence is not possible if the combination of stress and disturbance is too large (dashed line). This creates a triangle, each corner of which corresponds to a "primary strategy".

to population genetics' time-dependent, genetically-focused view of evolution. We revisit the classic lottery model of Chesson and Warner (1981), which has two features that make it well suited for this role, but one critical flaw that we rectify here.

The first feature is that the lottery representation of competition is particularly concise. Mature individuals ("adults") each require their own territory, whereas newborn individuals ("propagules") disperse to, and subsequently compete for, territories made available by the death of adults. Territorial contest among propagules leaves a single victorious adult per territory, the victor chosen at random from the propagules present, with probabilities weighted by a coefficient for each type representing competitive ability, akin to a lottery (Sale, 1977). By comparison, coefficients for the pairwise effects of types on each other (e.g. the α coefficients in the generalized Lotka-Volterra equations and the associated concept of " α -selection"; Case and Gilpin 1974; Gill 1974; Joshi et al. 2001), or explicit resource consumption (Tilman, 1982), are much more complicated. The second feature is the close connection between the lottery model and one of the foundational models of population genetics, the Wright-Fisher model of genetic drift, which we discuss further below.

The critical flaw of the classic lottery model is that it breaks down at low densities (few propagules dispersing to each territory), precluding density-dependent behaviour. Our first task

is to analytically extend the classic lottery model to correctly account for low density behavior (sections “Model” and “Mean field approximation”).

90 Using our extended lottery model, we then revisit Grime’s C/S/R scheme, and evaluate how C/S/R manifests at the level of within-population genotypic evolution (as opposed to phenotypic divergence between species; section “Primary strategies and Grime’s triangle”). This represents a
93 “sanity check” on our density-dependent lottery model. The resulting formulation of the C/S/R scheme is mathematical, in contrast to Grime’s original verbal and descriptive approach, which is a recognized hindrance to the evaluation or broader application of the C/S/R scheme (e.g.
96 Tilman 2007).

We then explore some time-dependent behavior of our extended lottery model. Taking an example inspired by recent studies of rapid, seasonal evolution in *Drosophila* (Bergland et al.,
99 2014), we discuss how environmental fluctuations might stabilize polymorphisms in the presence of cyclical population density.

Model

102 We assume that reproductively mature individuals (“adults”) each require their own territory to survive and reproduce (Fig. 2). All territories are identical, and the total number of territories is T . Time t advances in discrete iterations, each representing the time from birth to reproductive
105 maturity. In iteration t , the number of adults of the i ’th genotype is $n_i(t)$, the total number of adults is $N(t) = \sum_i n_i(t)$, and the number of unoccupied territories is $U(t) = T - N(t)$.

We assume that the n_i and T are large enough that stochastic fluctuations in the n_i (“drift”) can be ignored. We derive deterministic equations for the expected change in the n_i over time,
108 leaving the evaluation of drift for future work. This is an excellent approximation when the n_i are all large. However, we also do not evaluate the initial stochastic behaviour of adaptive mutant
111 lineages while they are at low abundance. When considering new mutations, we therefore restrict our attention to begin with the earliest (lowest n_i) deterministic behavior of mutant lineages (the

transition to deterministic growth occurs at an abundance n_i of order equal to their inverse
 114 expected absolute growth rate; Uecker and Hermisson 2011).

Each iteration, adults produce new offspring (“propagules”), m_i of which disperse to un-
 occupied territories. We assume that adults cannot be ousted from their territories, so that m_i
 117 only includes propagules landing on unoccupied territories. Propagules disperse at random over
 the unoccupied territories, regardless of distance from their parents, and independently of each
 other. There is no interaction between propagules (e.g. avoidance of territories crowded with
 120 propagules). Loss of propagules during dispersal is subsumed into m_i .

In general, m_i will increase with n_i , and will depend on population density N . For example,
 if b_i is the number of successfully dispersing propagules produced per genotype i adult, then
 123 the loss of propagules due to dispersal to occupied territories implies $m_i = b_i(1 - N/T)n_i$, akin
 to Levins’ competition-colonization model (Levins and Culver, 1971; Tilman, 1994). In section
 “Cyclical birth and death rates” we evaluate Eq. (4) numerically using this functional form for
 126 m_i , with b_i assumed to be constant.

In the sections “Invasion of rare genotypes and coexistence” and “Primary strategies and
 Grime’s triangle”, we assume the simpler form $m_i = b_i n_i$, with constant b_i , meaning that all
 129 propagules land on unoccupied territories (a form of directed dispersal). This simplifies the
 mathematics without affecting the results of those sections, which only depend on the low-
 frequency invasion behavior of Eq. (4). Note that due to our assumption of uniform dispersal, the
 132 parameter b_i can be thought of as a measure of “colonization ability”, which combines fecundity
 and dispersal ability (Bolker and Pacala, 1999; Levins and Culver, 1971; Tilman, 1994).

The number of individuals of the i ’th genotype landing in any particular territory is denoted
 135 x_i . We assume that x_i follows a Poisson distribution $p_i(x_i) = l_i^{x_i} e^{-l_i} / x_i!$, where $l_i = m_i / U$ is the
 mean territorial propagule density. This approximation becomes exact when the n_i are large
 enough that drift in n_i can be ignored (Appendix A).

138 When multiple propagules land on the same territory, the victor is determined by lottery
 competition: genotype i wins a territory with probability $c_i x_i / \sum_j c_j x_j$, where c_i is a constant




Figure 2: Each iteration of our lottery model has three main elements. First, propagules are produced by adults which are dispersed at random over the unoccupied territories (only propagules landing on unoccupied territories are shown). Lottery competition then occurs in each unoccupied territory (competition in only one territory is illustrated): each genotype has a probability proportional to $b_i n_i c_i$ of securing the territory. Then occupied territories are freed up by adult mortality. In Eq. (3) and most of the paper, only adults can die (red crosses), but we will also consider the case where juveniles die (blue cross; section “Primary strategies and Grime’s triangle”).

representing relative competitive ability (Fig. 2).

In the classic lottery model (Chesson and Warner, 1981), unoccupied territories are assumed to be saturated with propagules from every genotype $l_i \gg 1$. From the law of large numbers, the composition of propagules in each territory will then not deviate appreciably from the mean composition l_1, l_2, \dots, l_G (G is the number of genotypes present), and so the probability that genotype i wins any particular unoccupied territory is approximately $c_i l_i / \sum_j c_j l_j$. Let $\Delta_+ n_i$ denote the number of territories won by genotype i . Then $\Delta_+ n_1, \Delta_+ n_2, \dots, \Delta_+ n_G$ follow a multinomial distribution with U trials and success probabilities $\frac{c_1 l_1}{\sum_j c_j l_j}, \frac{c_2 l_2}{\sum_j c_j l_j}, \dots, \frac{c_G l_G}{\sum_j c_j l_j}$, respectively. Genotype i is expected to win $c_i l_i / \sum_j c_j l_j$ of the U available territories, and deviations from this expected outcome are small (since T is large by assumption), giving

$$\Delta_+ n_i(t) = \frac{c_i l_i}{\sum_j c_j l_j} U(t) = b_i n_i \frac{1}{L} \frac{c_i}{\bar{c}}, \quad (1)$$

where $\bar{c} = \sum_j c_j m_j / M$ is the mean propagule competitive ability for a randomly selected propagule, $L = M/U$ is the total propagule density and $M = \sum_j m_j$ is the total number of propagules.

There is a close connection between the classic lottery model and the Wright-Fisher model of genetic drift (Svardal et al., 2015). In the Wright-Fisher model, genotype abundances are sampled each generation from a multinomial distribution with success probabilities $w_i n_i / \sum_j w_j n_j$, where w is relative fitness and the n_i are genotype abundances in the preceding generation. Population size N remains constant. This is mathematically equivalent to the classic lottery model with non-overlapping generations ($d_i = 1$ for all i) and $w_i = b_i c_i$. Thus, the classic lottery model allows us to replace the abstract Wright-Fisher relative fitnesses w_i with more ecologically-grounded fecundity, competitive ability and mortality parameters b_i , c_i and d_i , respectively. Since birth and death rates affect absolute abundances, this allows us to evaluate selection at different densities (after appropriate extensions are made), in an otherwise very similar model to the canonical Wright-Fisher. We therefore expect that drift in realized values of n_i in our extended lottery model should be similar to that in the Wright-Fisher model, but we leave this for future work.

In our extension of the classic lottery model, we do not restrict ourselves to high propagule
 165 densities. Eq. (1) is nonsensical if even a single type has low propagule density ($l_i \ll 1$): genotype
 i can win at most m_i territories, yet Eq. (1) demands $c_i l_i / \sum_j c_j l_j$ of the U unoccupied territories,
 for any value of U . Intuitively, the cause of this discrepancy is that individuals are discrete.
 168 Genotypes with few propagules depend on the outcome of contests in territories where they
 have at least one propagule present, not some small fraction of a propagule as would be implied
 by small l_i in the classic lottery model. In other words, deviations from the mean propagule
 171 composition l_1, l_2, \dots, l_G are important at low density.

We expect that a fraction $p_1(x_1) \dots p_G(x_G)$ of the U unoccupied territories will have the
 propagule composition x_1, \dots, x_G . Genotype i is expected to win $c_i x_i / \sum_j c_j x_j$ of these. Ignoring
 174 fluctuations about these two expectations (due to our no-drift, large T , large n_i approximation),
 genotype i 's territorial acquisition is given by

$$\Delta_+ n_i(t) = U(t) \sum_{x_1, \dots, x_G} \frac{c_i x_i}{\sum_j c_j x_j} p_1(x_1) \dots p_G(x_G), \quad (2)$$

in our extended lottery model, where the sum only includes territories with at least one propag-
 177 ule present. Note that unlike the classic lottery model, not all unoccupied territories are claimed
 each iteration, since under Poisson dispersal a fraction e^{-L} remain unoccupied.

For the majority of this manuscript we assume that mortality only occurs in adults (setting
 180 aside the juvenile deaths implicit in territorial contest), and at a constant, genotype-specific per-
 capita rate $0 \leq d_i \leq 1$, so that the overall change in genotype abundances is

$$\Delta n_i(t) = \Delta_+ n_i(t) - d_i n_i(t). \quad (3)$$

This seems reasonable in the absence of disturbances; when we come to consider the effects of dis-
 183 turbances (Section "Primary strategies and Grime's triangle"), we will incorporate disturbance-
 induced mortality in competing juveniles (Fig. 2).

Results

Mean Field Approximation

Eq. (2) involves an expectation over the time-dependent dispersal distributions p_i , and is thus too complicated to give intuition about the dynamics of density-dependent lottery competition.

We now evaluate this expectation using a “mean field” approximation.

Similarly to the high- l_i approximation of classic lottery model, we replace the x_i with appropriate mean values, although we cannot simply replace x_i with l_i . For a genotype with low propagule density $l_i \ll 1$, we have $x_i = 1$ in the territories where its propagules land, and so its growth comes entirely from territories which deviate appreciably from l_i . To account for this, we separate Eq. (2) into $x_i = 1$ and $x_i > 1$ parts. Our more general mean field approximation only requires that there are no large discrepancies in competitive ability (i.e. we do not have $c_i/c_j \gg 1$ for any two genotypes). We obtain (details in Appendix B)

$$\Delta_+ n_i(t) \approx b_i n_i \left[e^{-L} + (R_i + A_i) \frac{c_i}{\bar{c}} \right], \quad (4)$$

where

$$R_i = \frac{\bar{c} e^{-l_i} (1 - e^{-(L-l_i)})}{c_i + \frac{L-1+e^{-L}}{1-(1+l_i)e^{-L}} \frac{\bar{c} L - c_i l_i}{L-l_i}}, \quad (5)$$

and

$$A_i = \frac{\bar{c}(1 - e^{-l_i})}{\frac{1-e^{-l_i}}{1-(1+l_i)e^{-L}} c_i l_i + \frac{1}{L-l_i} \left(L \frac{1-e^{-L}}{1-(1+L)e^{-L}} - l_i \frac{1-e^{-l_i}}{1-(1+l_i)e^{-L}} \right) \sum_{j \neq i} c_j l_j}. \quad (6)$$

To supplement our analytical mean field derivation, we did numerical simulations of our exact density-dependent lottery model, and verified that Eq. (4) is a good approximation (Appendix B). Thus, Eq. (4) describes how type abundances change over time in a lottery model where population density can itself vary with time.

Comparing Eq. (4) to Eq. (1), the classic lottery per-propagule success rate $c_i/\bar{c}L$ has been replaced by three separate terms. The first, e^{-L} , accounts for propagules which land alone on

unoccupied territories; these territories are won without contest. The second, $R_i c_i / \bar{c}$ represents competitive victories when the i genotype is a rare invader in a high density population: from Eq. (5), $R_i \rightarrow 0$ when the i genotype is abundant ($l_i \gg 1$), or other genotypes are collectively rare ($L - l_i \ll 1$). The third term, $A_i c_i / \bar{c}$, represents competitive victories when the i genotype is abundant: $A_i \rightarrow 0$ if $l_i \ll 1$. The relative importance of these three terms varies with both the overall propagule density L and the relative propagule frequencies m_i/M . If $l_i \gg 1$ for all genotypes, we recover the classic lottery model (only the $A_i c_i / \bar{c}$ term remains, and $A_i \rightarrow 1/L$).

Primary strategies and Grime's triangle

Here we describe how Grime's "disturbed", "stressful" and "ideal" environments can be captured mathematically in our model, and evaluate the direction of evolution for each. To proceed, we map these verbally-defined environments to quantitative parameter regimes in our model, summarized in Fig. 3.

Ideal environments are characterized by the near-absence of stress and disturbance, so that survival is easy $d \ll 1$ and growth and reproduction can be rapid $b \gg d$. The propagule density is determined by the magnitude of b/d , and in this case $L \gg 1$ (Appendix C).

Disturbed environments are characterized by short bursts of high extrinsic mortality d caused by physical destruction. We assume that disturbances are equally damaging to adults and juveniles, so that only $(1 - d_i)\Delta_+ n_i$ (rather than $\Delta_+ n_i$) territories are secured by type i each iteration. In the heavily disturbed extreme of Grime's triangle, many such bursts (which could each be spatially localized) will occur each iteration (i.e. over the time required for a propagule to reach reproductive maturity), which we approximate as a high constant mortality rate $d \approx 1$. Population persistence requires $b > d/(1 - d) \gg 1$ (Appendix C), i.e. severe disturbance comes with a high fecundity/dispersal requirement. We expect that it is not possible to greatly exceed this requirement ($b \approx d/(1 - d)$), which implies $L \ll 1$.

Stressful environments are more ambiguous, and have been the subject of an extensive debate in the plant ecology literature (the "Grime-Tilman" debate; Aerts 1999 and references therein).

	Ideal	Disturbance*	Stress (G)	Stress (HD)
Constraints	$d \ll 1$	$d \approx 1$	$b \ll 1$	$b \ll 1$
Other parameters	$b/d \gg 1$	$(1 - d)b \approx d$	$b \approx d$	$b/d \gg 1$
Propagule density L	$\gg 1$	$\ll 1$	$\ll 1$	$\gg 1$

Figure 3: The mathematical representation of Grime’s three environmental extremes in our model, as well as the high-density (HD) variant of the stressful environment. *Mortality affects both adults and juveniles in the disturbed environment, with $\Delta_+ n_i$ replaced by $(1 - d_i)\Delta_+ n_i$.

231 Severe stress inhibits growth and reproduction, so that $b \ll 1$. In Grime’s view, this means that
the rate at which propagules successfully develop to adulthood cannot appreciably exceed the
mortality rate $b/d \approx 1$; coping with the environmental stress is barely possible (Grime, 1974,
234 1977). In our model, this implies $L \ll 1$.

The alternative view is that stressed environments are highly competitive. In spite of inhibited
growth and reproduction, density is actually high relative to carrying capacity of the stressful
237 environment (Taylor et al., 1990). For example, if consumable resources are scarce, we expect
intense resource competition (Davis et al., 1998). Thus, even though $b \ll 1$, we still have $b/d \gg 1$
and $L \ll 1$.

240 The expected direction of evolution is determined by three factors. First, there may be
pleiotropy between b , c , and d , which could be due to trade-offs between them. Second, adaptive
mutations in one trait may occur more frequently than in others (Yampolsky and Stoltzfus, 2001).
243 We do no doubt that these factors operate in nature and contribute towards “primary strategy”
trait patterns. However, modeling how mutations affect traits subject to trade-offs is beyond the
scope of this article, and so we will not evaluate these first two factors: we assume that mutations
246 affect b , c and d independently, and that b , c and d have the same distribution of beneficial fitness
effects.

The third factor is that some traits may have a greater impact on fitness: b , c and d will tend to
249 evolve in the direction of greatest fitness increase because the probability that an adaptive mutant
survives its initial low abundance phase and begins to grow deterministically is proportional to
its expected absolute growth rate (Haldane, 1927; Uecker and Hermisson, 2011). Very few neutral

252 mutations fix (probability of order $1/N$).

Under Eq. (4), mutations which change b or d by the same proportion (i.e. $b_j = (1 + \epsilon)b_i$ or $d_j = (1 - \epsilon)d_i$ where j is the mutant type) have identical fitness effects (Appendix C). When
255 propagule density is high $L \gg 1$, the same is true for proportional c -mutations $c_j = (1 + \epsilon)c_i$.
When propagule density is low $L \ll 1$, c does not contribute to fitness and c -mutations are neutral. Thus, selection under Eq. (4), which applies in ideal and stressed environments, does
258 not intrinsically favor one trait improvement over another, apart from the irrelevance of c at low density.

The situation in disturbed environments, where mortality also affects juveniles, is more com-
261 plicated. The bulk of the high mortality $d \approx 1$ is environmental and unavoidable. Thus, propor-
tional changes in the remaining genetic part of mortality can have a tiny impact on d , suggesting
the primacy of b under disturbance. However, this effect is easily swamped by the fact that
264 disturbance mortality affects both juvenile recruitment and adult mortality.

Following Grime's original argument for a triangular scheme (Grime, 1977), Fig. 4 repre-
sents each environmental extreme schematically as a vertex on a triangular space defined by
267 perpendicular stress and disturbance axes. The ideal environment lies at the origin (no stress or
disturbance), while the stressful and disturbed environments lie at the limits of survival on their
respective axes. The hypotenuse connecting the stress and disturbance endpoints represents the
270 limits of survival in the presence of a combination of stress and disturbance. The direction of
evolutionary change is different at each vertex, leading to the emergence of different trait clusters
or "primary strategies".

273 **Coexistence in constant and cyclical environments**

In the previous section we only considered how b , c and d should respond in Grime's environ-
mental extremes, based on invasion fitness. Here we further explore the low frequency behavior
276 of Eq. (4) to determine which types can coexist in a constant environment, and then consider the
full time-dependent behaviour of Eq. (4) in a cyclical environment.




Figure 4: The realization of Grime’s triangle in our model. Schematic representation of the triangular space bounded by the low/high extremes of stress/disturbance. The high-density (HD) interpretation of stress is also shown. The vertices of the triangles correspond to different environmental extremes. Selection favors different traits at each vertex, leading to different trait clusters.

In a constant environment, stable coexistence is possible in our extended lottery model. A b specialist i and c -specialist j ($b_i > b_j$, $c_j > c_i$) can co-exist because then propagule density L is frequency dependent, and so is the importance of competitive ability (Appendix D). This is a version of the classic competition-colonization trade-off (Levins and Culver, 1971; Tilman, 1994): the competitor (c -specialist) leaves many territories unoccupied (low L) due to its poor colonization ability (low b), which the colonizer (b -specialist) can then exploit. A similar situation holds for coexistence between high- c and low- d specialists; a “competition-longevity” trade-off (Tilman, 1994). These forms of co-existence require density dependence (being mediated by L), and are not present in the classic lottery model. Coexistence is not possible between b - and d -specialists (Appendix D).

Now suppose that birth and death rates vary periodically with amplitude sufficient to cause large changes in population density. This example is inspired by natural *Drosophila* populations, which expand rapidly in the warmer months when fruit is abundant, but largely die off in the colder months. Within this seasonal population density cycle, hundreds of polymorphisms also cycle in frequency (Bergland et al., 2014). Some of these polymorphisms may be adaptive and potentially millions of years old, suggesting stable coexistence (Bergland et al., 2014; Messer

et al., 2016). Selection on allele frequencies thus occurs on the same time scale as population demography, a situation vastly more complicated than classical sweeps in demographically stable populations (Messer et al., 2016).

The classical population genetic treatment of fluctuating selection suggests that environmental fluctuations do not promote coexistence. Allele frequencies are successively multiplied by relative fitness values for each environmental iteration, and so two alleles favored in different environments can only stably coexist if the product of fitnesses for one type exactly equals the product for the other (Dempster, 1955). Thus, stable coexistence still requires frequency-dependent selection or heterozygote advantage (as is required in a constant environment).

This classical argument overlooks two general effects that promote coexistence in fluctuating environments. The first is the storage effect, which occurs when part of the population is protected from selection (due to overlapping generations in the lottery model; Chesson and Warner 1981). The second is the bounded population size effect of Yi and Dean (2013), which occurs when each environmental cycle involves growth from low to high density, with the time spent growing each cycle dependent on the fitness of the types present.

Fig. 5a-c shows the behavior of Eq. (4) for an example where b and d cycle between zero and positive values (“summers” with rapid growth and no mortality, and “winters” with mortality and no growth). Both the storage effect (adults are sheltered from selection during the summer growth phase) and the bounded density effect (expansion to high density occurs every cycle) are operating. Two types are present, a b specialist, which is better at rapidly growing in the summer (higher b), and a d specialist which is better at surviving the winter (lower d). Neither type has an advantage over a full environmental cycle, and they stably coexist. This is due to a combination of the storage and bounded density effects (recall that stable coexistence between b and d specialists was not possible in a constant environment).

The classic lottery model (Eq. 1) gives dramatically different co-existence predictions because expansion to carrying capacity occurs immediately at the start of the summer (Fig. 5d-f). As a result, coexistence requires that the winter survivor’s b must be about 5 times smaller than

required when we properly account for the growth in the abundance of each type using Eq. (4) (keeping the other parameters the same). Previous models of the promotion of genetic variation via the storage effect (Ellner and Hairston Jr, 1994) similarly assume that the total number of offspring per iteration is constant, and would produce a similar error.

Discussion

It is interesting to compare the predictions of the extended lottery model with earlier approaches, such as the r/K scheme, where $r = b - d$ is the maximal, low-density growth rate (Pianka, 1972). Confusingly, the term “ K -selection” sometimes refers generally to selection at high density (Pianka, 1972), encompassing both selection for higher saturation density (MacArthur and Wilson, 1967) and competitive ability (Gill, 1974). Contrary to predictions of an r/K trade-off, empirical studies have shown that maximal growth rate and saturation density (measured by abundance) are positively correlated, both between species/strains (Fitzsimmons et al., 2010; Hendriks et al., 2005; Kuno, 1991; Luckinbill, 1979), and as a result of experimental evolution (Luckinbill, 1978, 1979). From the perspective of our model, this positive correlation is not surprising since the saturation density, which is determined by a balance between births and deaths, increases with b .

There is support for a negative relationship between competitive success at high density and maximal growth rate (Luckinbill, 1979), consistent with a tradeoff between r and the competitive aspect of K . This could be driven by a tradeoff between individual size and reproductive rate. To avoid confusion with other forms of “ K -selection”, selection for competitive ability has been called “ α -selection” after the competition coefficients in the Lotka-Volterra equation (Case and Gilpin, 1974; Gill, 1974; Joshi et al., 2001). However, competitive success as measured by α (i.e. the per-capita effect of one genotype on another genotype’s growth rate) is only partly determined by individual competitive ability — in the presence of age-structured competition and territoriality, it also includes the ability of each genotype to produce contestants i.e. b in our model. Our




Figure 5: Stable coexistence between b and d specialists in a fluctuating environment requires a much greater b advantage in the classic lottery model compared to our density-dependent extension of it when population density is seasonally cyclical. (a) Birth and death rates seasonally alternate being nonzero (white for winter, green for summer). The b specialist (black) has higher b and d ($b = 0.5$, $d = 0.2$) than the d specialist ($b = 0.217$, $d = 0.1$) (blue). (b) Both types grow during the positive b phase, and decline during the positive d phase, but the d specialist does so at a lower rate. Total height (blue+black) is population density N/T . (c) Summer favors the b specialist, winter the d specialist, and they stably coexist. (d-f) Same as (a-c) for the classic lottery model; the types no longer coexist. (g-i) Same as (d-f) where now $b = 0.0421$ for the d specialist and the types coexist. For illustration, the propagule abundances are assumed to have the form $m_i = b_i(1 - N/T)n_i$, reflecting non-directed dispersal.

c is strictly competitive ability only — as such, changes in c do not directly affect population density (the total number of territories occupied per iteration is $\Delta_+ N = U(1 - e^{-L})$, which
 348 does not depend directly on the c_i). The clean separation of a strictly-relative c parameter is particularly useful from an evolutionary genetics perspective, essentially embedding a zero-sum relative fitness trait within a non-zero-sum fitness model. This could have interesting applications
 351 for modeling the impacts of intra-specific competition on species extinction, for example due to clonal interference (Desai and Fisher, 2007; Gerrish and Lenski, 1998) between c -strategists on the one hand, and b - and d - strategists on the other.

354 K -selection in the narrow logistic sense of selection for a greater environmental carrying capacity for given birth and death rates, sometimes referred to as “efficiency” (MacArthur and Wilson, 1967), could be represented in our model by smaller individual territorial requirements.
 357 To a first approximation, two co-occurring genotypes which differ by a small amount in their territorial requirements only should have the same fitness since the costs or benefits of a change in the amount of unoccupied territory is shared equally among genotypes via the propagule density
 360 per territory L . The situation is more complicated when the differences in territorial requirements become large enough that territorial contests can occur on different scales for different genotypes. We leave these complications for future work.

363 Our realization of Grime’s triangle (Fig. 4) differs from approaches which identify primary strategies as trait combinations which can stably co-exist (Bolker and Pacala, 1999), referring instead to the direction and rate of ongoing adaptive trait evolution under different regimes of
 366 stress and disturbance, which is closer in spirit to Grime’s arguments (Grime, 1974, 1977). In addition, we have not assumed any kind of trade-offs or pleiotropy between b , c and d , only constraints imposed by the environment on the order of magnitude of b and d . As an example of
 369 a trade-off, corals which rapidly out-shade neighbors have a tall, branched morphology which is vulnerable to disturbances, and so, all else being equal, ideal environment c -strategists will suffer higher mortality from disturbances (Darling et al., 2012). Fig. 4 gives the same conclusion without
 372 invoking trade-offs; mutations which reduce disturbance vulnerability are essentially neutral

under ideal conditions, leading to no improvements in mortality from disturbances, whereas c will tend to increase over time. Thus, while trade-offs may amplify specialization, and are sometimes invoked to explain primary strategy schemes (Aerts, 1999; MacArthur and Wilson, 1967; Winemiller and Rose, 1992), they are not necessary for it.

One potential limitation of our model as a general-purpose model of density-dependent selection is its restriction to interference competition between juveniles for durable resources (lottery recruitment to adulthood), analogous to the ubiquitous assumption of viability selection in population genetics (Ewens, 2004, p. 45). In some respects this is the complement of consumable resource competition models, which restrict their attention to indirect exploitation competition, typically without age structure (Tilman, 1982). In the particular case that consumable resources are spatially localized (e.g. due to restricted movement through soils), resource competition and territorial acquisition effectively coincide, and in principle resource competition could be represented by a competitive ability c (or conversely, c should be derivable from resource competition). The situation is more complicated if the resources are well-mixed, since, in general, resource levels then need to be explicitly tracked. It seems plausible that explicit resource tracking may not be necessary when the focus is on the evolution of similar genotypes that use identical resources rather than the stable co-existence of widely differing species with different resource preferences (Ram et al., 2016). We are not aware of any attempts to delineate conditions under which explicit resource tracking is unnecessary even if it is assumed that community structure is ultimately determined by competition for consumable resources. More work is needed connecting resource competition models to the density-dependent selection literature, since most of the former has to date been focused on narrower issues of the role of competition at low resource availability (Aerts, 1999; Davis et al., 1998; Tilman, 2007).

While our model can be applied to species rather than genotypes (e.g. ecological invasions), our focus is genotype evolution i.e. the change in allele frequencies over time. Our assumption that there are no large c discrepancies (section “Mean field approximation”) amounts to a restriction on the amount of genetic variation in c in the population. Since beneficial mutation effect

sizes will typically not be much larger than a few percent, large c discrepancies can only arise if the mutation rate is extremely large, and so the assumption will not be violated in most cases.

402 However, this restriction could become important when looking at species interactions rather than genotype evolution.

In the introduction we mentioned the recurring difficulties with confounding selection and
405 demography in population genetic inference. It seems that Eq. (4) or something similar (and hopefully more analytically tractable) is unavoidable for the analysis of time-course genetic data because, fundamentally, selective births and deaths affect both abundances and frequencies, not
408 one or the other in isolation. Moreover, some aspects of allele frequency change are intrinsically density-dependent. In the classic lottery model, which as we have seen is essentially the Wright-Fisher model with overlapping generations, b_i and c_i are equivalent in the sense that the number
411 of territorial victories only depends on the product $b_i c_i$ (see “Model”). This is no longer the case in our extension, where b and c specialists can co-exist. This “colonization-competition trade-off” is well known in the co-existence literature (Tilman, 1994). It and similar forms of
414 “spatial co-existence” in stable environments have previously been modeled either with Levin’s qualitative representation of competition (Levins and Culver, 1971; Tilman, 1994), as opposed to the quantitative c of lottery competition, or with a more sophisticated treatment of space (non-
417 uniform dispersal; Bolker and Pacala 1999; Shmida and Ellner 1984). In cyclical environments, polymorphisms can be stabilized by the bounded density effect which is completely lost if there is an exclusive focus on allele frequencies (Yi and Dean, 2013). We leave the details of how our
420 model might be applied in inference, including the crucial issue of its genetic drift predictions (providing a null model for neutral sites), for future work.

References

423 Aerts, R. 1999. Interspecific competition in natural plant communities: mechanisms, trade-offs and plant-soil feedbacks. *Journal of Experimental Botany* 50:29–37.

Allan, J. D. 1976. Life history patterns in zooplankton. *The American Naturalist* 110:165–180.

426 Arenbaev, N. 1977. Asymptotic behavior of the multinomial distribution. *Theory of Probability
& Its Applications* 21:805–810.

Barton, N. H. 1998. The effect of hitch-hiking on neutral genealogies. *Genetical Research* 72:123–
429 133.

Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt, and D. A. Petrov. 2014. Genomic
evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLOS*
432 *Genetics* 10:1–19.

Bolker, B. M., and S. W. Pacala. 1999. Spatial moment equations for plant competition: Under-
standing spatial strategies and the advantages of short dispersal. *The American Naturalist*
435 153:575–602.

Boyce, M. S. 1984. Restitution of *r*- and *K*-selection as a model of density-dependent natural
selection. *Annual Review of Ecology and Systematics* 15:427–447.

438 Case, T. J., and M. E. Gilpin. 1974. Interference competition and niche theory. *Proceedings of the
National Academy of Sciences* 71:3073–3077.

Chesson, P. L., and R. R. Warner. 1981. Environmental variability promotes coexistence in lottery
441 competitive systems. *American Naturalist* pages 923–943.

Darling, E. S., L. Alvarez-Filip, T. A. Oliver, T. R. McClanahan, and I. M. Côté. 2012. Evaluating
life-history strategies of reef corals from species traits. *Ecology Letters* 15:1378–1386.

444 Davis, M. A., K. J. Wragg, and P. B. Reich. 1998. Competition between tree seedlings and herba-
ceous vegetation: support for a theory of resource supply and demand. *Journal of Ecology*
86:652–661.

447 Dempster, E. R. 1955. Maintenance of genetic heterogeneity. *Cold Spring Harb Symp Quant Biol*
20:25–31.

Desai, M. M., and D. S. Fisher. 2007. Beneficial mutation–selection balance and the effect of
linkage on positive selection. *Genetics* 176:1759–1798.

Diekmann, O., et al. 2004. A beginner’s guide to adaptive dynamics. *Banach Center Publications*
63:47–86.

Ellner, S., and N. G. Hairston Jr. 1994. Role of overlapping generations in maintaining genetic
variation in a fluctuating environment. *The American Naturalist* 143:403–417.

Ewens, W. J. 2004. *Mathematical Population Genetics 1: Theoretical Introduction*. Springer
Science & Business Media.

Fitzsimmons, J. M., S. E. Schoustra, J. T. Kerr, and R. Kassen. 2010. Population consequences of
mutational events: effects of antibiotic resistance on the r/k trade-off. *Evolutionary Ecology*
24:227–236.

Gerrish, P. J., and R. E. Lenski. 1998. The fate of competing beneficial mutations in an asexual
population. *Genetica* 102:127–144.

Gill, D. E. 1974. Intrinsic rate of increase, saturation density, and competitive ability. ii. the
evolution of competitive ability. *American Naturalist* 108:103–116.

Grime, J. P. 1974. Vegetation classification by reference to strategies. *Nature* 250:26–31.

———. 1977. Evidence for the existence of three primary strategies in plants and its relevance to
ecological and evolutionary theory. *American Naturalist* 111:1169–1194.

———. 1988. *Plant Evolutionary Biology*, chap. The C-S-R model of primary plant strategies —
origins, implications and tests, pages 371–393. Springer Netherlands, Dordrecht.

Haldane, J. B. S. 1927. A mathematical theory of natural and artificial selection, part v: selection
and mutation 23:838–844.

- 471 Hendriks, A. J., J. L. Maas-Diepeveen, E. H. Heugens, and N. M. van Straalen. 2005. Meta-analysis
of intrinsic rates of increase and carrying capacity of populations affected by toxic and other
stressors. *Environmental Toxicology and Chemistry* 24:2267–2277.
- 474 Joshi, A., N. Prasad, and M. Shakarad. 2001. K-selection, α -selection, effectiveness, and tolerance
in competition: density-dependent selection revisited. *Journal of Genetics* 80:63–75.
- Kuno, E. 1991. Some strange properties of the logistic equation defined with r and K : Inherent
477 defects or artifacts? *Researches on Population Ecology* 33:33–39.
- Levins, R., and D. Culver. 1971. Regional coexistence of species and competition between rare
species. *Proceedings of the National Academy of Sciences* 68:1246–1248.
- 480 Luckinbill, L. S. 1978. r and k selection in experimental populations of *escherichia coli*. *Science*
(New York, NY) 202:1201–1203.
- . 1979. Selection and the r/k continuum in experimental populations of protozoa. *Ameri-*
483 *can Naturalist* pages 427–437.
- MacArthur, R. H. 1962. Some generalized theorems of natural selection. *Proceedings of the*
National Academy of Sciences 48:1893–1897.
- 486 MacArthur, R. H., and E. O. Wilson. 1967. *Theory of Island Biogeography*. Princeton University
Press.
- Messer, P. W., S. P. Ellner, and N. G. Hairston. 2016. Can population genetics adapt to rapid
489 evolution? *Trends in Genetics* 32:408–418.
- Metz, J. A., R. M. Nisbet, and S. A. Geritz. 1992. How should we define ‘fitness’ for general
ecological scenarios? *Trends in Ecology & Evolution* 7:198–202.
- 492 Pianka, E. R. 1970. On r - and K -Selection. *The American Naturalist* 104:592–597.
- . 1972. r and K selection or b and d selection? *The American Naturalist* 106:581–588.

Ram, Y., E. Dellus-Gur, M. Bibi, U. Obolski, J. Berman, and L. Hadany. 2016. Predicting microbial
relative growth in a mixed culture from growth curve data. *bioRxiv* 10.1101/022640 .

Reznick, D., M. J. Bryant, and F. Bashey. 2002. r- and k-selection revisited: The role of population
regulation in life-history evolution. *Ecology* 83:1509–1520.

Sale, P. F. 1977. Maintenance of high diversity in coral reef fish communities. *The American
Naturalist* 111:337–359.

Shmida, A., and S. Ellner. 1984. Coexistence of plant species with similar niches. *Vegetatio*
58:29–55.

Southwood, T. R. E. 1977. Habitat, the templet for ecological strategies? *Journal of Animal
Ecology* 46:337–365.

Stearns, S. C. 1977. The evolution of life history traits: A critique of the theory and a review of
the data. *Annual Review of Ecology and Systematics* 8:145–171.

Svardal, H., C. Rueffler, and J. Hermisson. 2015. A general condition for adaptive genetic poly-
morphism in temporally and spatially heterogeneous environments. *Theoretical Population
Biology* 99:76 – 97.

Taylor, D. R., L. W. Aarssen, and C. Loehle. 1990. On the relationship between r/k selection and
environmental carrying capacity: a new habitat templet for plant life history strategies. *Oikos*
pages 239–250.

Tilman, D. 1982. Resource competition and community structure. Princeton University Press.

———. 1994. Competition and biodiversity in spatially structured habitats. *Ecology* 75:2–16.

———. 2007. Resource competition and plant traits: a response to craine et al. 2005. *Journal of
Ecology* 95:231–234.

516 Uecker, H., and J. Hermisson. 2011. On the fixation process of a beneficial mutation in a variable environment. *Genetics* 188:915–930.

Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let
519 the concept of trait be functional! *Oikos* 116:882–892.

Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and soil* 199:213–227.

522 Winemiller, K. O., D. B. Fitzgerald, L. M. Bower, and E. R. Pianka. 2015. Functional traits, convergent evolution, and periodic tables of niches. *Ecology Letters* 18:737–751.

Winemiller, K. O., and K. A. Rose. 1992. Patterns of life-history diversification in north american
525 fishes: implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences* 49:2196–2218.

Yampolsky, L. Y., and A. Stoltzfus. 2001. Bias in the introduction of variation as an orienting
528 factor in evolution. *Evolution & Development* 3:73–83.

Yi, X., and A. M. Dean. 2013. Bounded population sizes, fluctuating selection and the tempo and mode of coexistence. *Proceedings of the National Academy of Sciences* 110:16945–16950.

531 **Appendix A: Poisson approximation**

For simplicity of presentation, we have assumed a Poisson distribution for the x_i as our model of dispersal. Strictly speaking, the total number of i propagules $\sum x_i$ (summed over unoccupied
534 territories) is then no longer a constant m_i , but fluctuates between generations for a given mean m_i , which is more biologically realistic. Nevertheless, since we do not consider the random fluctuations in type abundances here, and for ease of comparison with the classic lottery model, we
537 ignore the fluctuations in m_i . Instead we focus, on Poisson fluctuations in propagule composition in each territory.

In the exact model of random dispersal, the counts of a genotype's propagules across un-
 540 noccupied territories follows a multinomial distribution with dimension U , total number of trials
 equal to m_i , and equal probabilities $1/U$ for a propagule to land in a given territory. Thus, the
 x_i in different territories are not independent random variables. However, for sufficiently large
 543 U and m_i , this multinomial distribution for the x_i across territories is closely approximated by
 a product of independent Poisson distributions for each territory, each with rate parameter l_i
 (Arenbaev, 1977, Theorem 1). Since we are ignoring finite population size effects, we effectively
 546 have $T \rightarrow \infty$, in which case U can be only be small enough to violate the Poisson approximation
 if there is vanishing population turnover, and then the dispersal distribution is irrelevant any-
 way. Likewise, in ignoring stochastic finite population size for the n_i , we have effectively already
 549 assumed that m_i is large enough to justify the Poisson approximation (the error scales as $1/\sqrt{m_i}$;
 Arenbaev 1977).

Appendix B: Derivation of growth equation

552 We separate the right hand side of Eq. (2) into three components $\Delta_+ n_i = \Delta_u n_i + \Delta_r n_i + \Delta_a n_i$ which
 vary in relative magnitude depending on the propagule densities l_i . Following the notation in
 the main text, the Poisson distributions for the x_i (or some subset of the x_i) will be denoted p ,
 555 and we use P as a general shorthand for the probability of particular outcomes.

Growth without competition

The first component, $\Delta_u n_i$, accounts for territories where only one focal propagule is present
 558 $x_i = 1$ and $x_j = 0$ for $j \neq i$ (u stands for "uncontested"). The proportion of territories where this
 occurs is $l_i e^{-L}$, and so

$$\Delta_u n_i = U l_i e^{-L} = m_i e^{-L}. \quad (7)$$

Competition when rare

561 The second component, $\Delta_r n_i$, accounts for territories where a single focal propagule is present along with at least one non-focal propagule (r stands for “rare”) i.e. $x_i = 1$ and $X_i \geq 1$ where $X_i = \sum_{j \neq i} x_j$ is the number of nonfocal propagules. The number of territories where this occurs
564 is $Up_i(1)P(X_i \geq 1) = b_i n_i e^{-l_i}(1 - e^{-(L-l_i)})$. Thus

$$\Delta_r n_i = m_i e^{-l_i}(1 - e^{-(L-l_i)}) \left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}}, \quad (8)$$

where $\langle \rangle_{\tilde{p}}$ denotes the expectation with respect to \tilde{p} , and \tilde{p} is the probability distribution of nonfocal propagule abundances x_j *after* dispersal, in those territories where exactly one focal propagule, and at least one non-focal propagule, landed.
567

Our “mean field” approximation is to replace x_j with its mean in the last term in Eq. (8),

$$\left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}} \approx \frac{c_i}{c_i + \sum_{j \neq i} c_j \langle x_j \rangle_{\tilde{p}}}. \quad (9)$$

Below we justify this replacement by arguing that the standard deviation $\sigma_{\tilde{p}}(\sum_{j \neq i} c_j x_j)$ (with
570 respect to \tilde{p}), is much smaller than $\langle \sum_{j \neq i} c_j x_j \rangle_{\tilde{p}}$.

We first calculate $\langle x_j \rangle_{\tilde{p}}$. Let $X = \sum_j x_j$ denote the total number of propagules in a territory and $\mathbf{x}_i = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_G)$ denote the vector of non-focal abundances, so that $p(\mathbf{x}_i) = p_1(x_1) \dots p_{i-1}(x_{i-1}) p_{i+1}(x_{i+1}) \dots p_G(x_G)$. Then, \tilde{p} can be written as

$$\begin{aligned} \tilde{p}(\mathbf{x}_i) &= p(\mathbf{x}_i | X \geq 2, x_i = 1) \\ &= \frac{P(\mathbf{x}_i, X \geq 2 | x_i = 1)}{P(X \geq 2)} \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{X=2}^{\infty} P(X) p(\mathbf{x}_i | X_i = X - 1), \end{aligned} \quad (10)$$

and so

$$\begin{aligned}\langle x_j \rangle_{\tilde{p}} &= \sum_{\mathbf{x}_i} \tilde{p}(\mathbf{x}_i) x_j \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{X=2}^{\infty} P(X) \sum_{\mathbf{x}_i} p(\mathbf{x}_i | X_i = X - 1) x_j.\end{aligned}\quad (11)$$

The inner sum over \mathbf{x}_i is the mean number of propagules of a given nonfocal type j that will be found in a territory which received $X - 1$ nonfocal propagules in total, which is equal to $\frac{l_j}{L - l_i}(X - 1)$. Thus,

$$\begin{aligned}\langle x_j \rangle_{\tilde{p}} &= \frac{l_j}{1 - (1 + L)e^{-L}} \frac{1}{L - l_i} \sum_{k=2}^{\infty} P(X)(X - 1) \\ &= \frac{l_j}{1 - (1 + L)e^{-L}} \frac{L - 1 + e^{-L}}{L - l_i},\end{aligned}\quad (12)$$

where the last line follows from $\sum_{X=2}^{\infty} P(X)(X - 1) = \sum_{X=1}^{\infty} P(X)(X - 1) = \sum_{X=1}^{\infty} P(X)X - \sum_{X=1}^{\infty} P(X)$.

The exact analysis of the fluctuations in $\sum_{j \neq i} c_j x_j$ is complicated because the x_j are not independent with respect to \tilde{p} . These fluctuations are part of the “drift” in type abundances which we leave for future work. Here we use the following approximation to give some insight into the magnitude of these fluctuations and also the nature of the correlations between the x_j . We replace \tilde{p} with \tilde{q} , defined as the \mathbf{x}_i Poisson dispersal probabilities conditional on $X_i \geq 1$ (which are independent). The distinction between \tilde{p} with \tilde{q} will be discussed further below. The \tilde{q} approximation

gives $\langle x_j \rangle_{\tilde{q}} = \langle x_j \rangle_p / C = l_j / C$,

$$\begin{aligned}
\sigma_{\tilde{q}}^2(x_j) &= \langle x_j^2 \rangle_{\tilde{q}} - \langle x_j \rangle_{\tilde{q}}^2 \\
&= \frac{1}{C} \langle x_j^2 \rangle_p - \frac{l_j^2}{C^2} \\
&= \frac{1}{C} (l_j^2 + l_j) - \frac{l_j^2}{C^2} \\
&= \frac{l_j^2}{C} \left(1 - \frac{1}{C} \right) + \frac{l_j}{C},
\end{aligned} \tag{13}$$

and

$$\begin{aligned}
\sigma_{\tilde{q}}(x_j, x_k) &= \langle x_j x_k \rangle_{\tilde{q}} - \langle x_j \rangle_{\tilde{q}} \langle x_k \rangle_{\tilde{q}} \\
&= \frac{1}{C} \langle x_j x_k \rangle_p - \frac{l_j l_k}{C^2} \\
&= \frac{l_j l_k}{C} \left(1 - \frac{1}{C} \right),
\end{aligned} \tag{14}$$

573 where $C = 1 - e^{-(L-l_i)}$ and $j \neq k$.

The exact distribution \tilde{p} assumes that exactly one of the propagules present in a given site after dispersal belongs to the focal type, whereas \tilde{q} assumes that there is a focal propagule present before non-focal dispersal commences. As a result, \tilde{q} predicts that the mean propagule density is greater than L (in sites with only one focal propagule is present) when the focal type is rare and the propagule density is high. This is erroneous, because the mean number of propagules in every site is L by definition. Specifically, if $L - l_i \approx L \gg 1$, then the mean propagule density predicted by \tilde{q} is approximately $L + 1$. The discrepancy causes rare invaders to have an intrinsic rarity disadvantage (territorial contests under \tilde{q} are more intense than they should be). In contrast, Eq. (12) correctly predicts that there are on average $\sum_{j \neq i} \langle x_j \rangle_{\tilde{p}} \approx L - 1$ nonfocal propagules because \tilde{p} accounts for potentially large negative covariances between the x_j “after dispersal”. By neglecting the latter covariences, \tilde{q} overestimates the fluctuations in $\sum_{j \neq i} c_j x_j$; thus \tilde{q} gives an upper bound on the fluctuations. The discrepancy between \tilde{q} and \tilde{p} will be largest

579
582
585

when L is of order 1 or smaller, because then the propagule assumed to already be present under \tilde{q} is comparable to, or greater than, the entire propgaule density.

588 Decomposing the variance in $\sum_{j \neq i} c_j x_j$,

$$\sigma_{\tilde{q}}^2(\sum_{j \neq i} c_j x_j) = \sum_{j \neq i} \left[c_j^2 \sigma_{\tilde{q}}^2(x_j) + 2 \sum_{k > j, k \neq i} c_j c_k \sigma_{\tilde{q}}(x_j, x_k) \right], \quad (15)$$

and using the fact that $\sigma_{\tilde{q}}(x_j, x_k)$ and the first term in Eq. (13) are negative because $C < 1$, we obtain an upper bound on the relative fluctuations in $\sum_{j \neq i} c_j x_j$,

$$\frac{\sigma(\sum_{j \neq i} c_j x_j)}{\langle \sum_{j \neq i} c_j x_j \rangle} = C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j + (1 - 1/C) \left(\sum_{j \neq i} c_j l_j \right)^2 \right)^{1/2}}{\sum_{j \neq i} c_j l_j} < C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j \right)^{1/2}}{\sum_{j \neq i} c_j l_j}. \quad (16)$$

591 Suppose that the c_j are all of similar magnitude (their ratios are of order one). Then Eq. (16) is $\ll 1$ for the case when $L - l_i \ll 1$ (due to the factor of $C^{1/2}$), and also for the case when at least some of the nonfocal propagule densities are large $l_j \gg 1$ (since it is then of order $1/\sqrt{L - l_i}$).
 594 The worst case scenario occurs when $L - l_i$ is of order one. Then Eq. (16) gives a relative error of approximately 50%, which from our earlier discussion we know to be a substantial overestimate when L is of order 1. Our numerical results (Fig. 6) confirm that the relative errors are indeed
 597 small.

However, the relative fluctuations in $\sum_{j \neq i} c_j x_j$ can be large if some of the c_j are much larger than the others. Specifically, in the presence of a rare, extremely strong competitor ($c_j l_j \gg c_{j'} l_{j'}$ for all other nonfocal genotypes j' , and $l_j \ll 1$), then the RHS of Eq. (16) can be large and we
 600 cannot make the replacement Eq. (9).

Substituting Eqs. (9) and (12) into Eq. (8), we obtain

$$\Delta_r n_i \approx m_i R_i \frac{c_i}{\bar{c}}, \quad (17)$$

603 where R_i is defined in Eq. (5).

Competition when abundant

The final contribution, $\Delta_a n_i$, accounts for territories where two or more focal propagules are present (a stands for “abundant”). Similarly to Eq. (8), we have

$$\Delta_a n_i = U(1 - (1 + l_i)e^{l_i}) \left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \quad (18)$$

where \hat{p} is the probability distribution of both focal and nonfocal propagule abundances *after* dispersal in those territories where at least two focal propagules landed.

Again, we argue that the relative fluctuations in $\sum c_j x_j$ are much smaller than 1 (with respect to \hat{p}), so that,

$$\left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \approx \frac{c_i \langle x_i \rangle_{\hat{p}}}{\sum_j c_j \langle x_j \rangle_{\hat{p}}}. \quad (19)$$

Following a similar procedure as for $\Delta_r n_i$, where the vector of propagule abundances is denoted \mathbf{x} , the mean focal genotype abundance is,

$$\begin{aligned} \langle x_i \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_i p(\mathbf{x} | x_i \geq 2) \\ &= \sum_{x_i} x_i p(x_i | x_i \geq 2) \\ &= \frac{1}{1 - (1 + l_i)e^{-l_i}} \sum_{x_i \geq 2} p(x_i) x_i \\ &= l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}}. \end{aligned} \quad (20)$$

For nonfocal genotypes $j \neq i$, we have

$$\begin{aligned}
\langle x_j \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2) \\
&= \sum_X P(X | x_i \geq 2) \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2, X) \\
&= \sum_X P(X | x_i \geq 2) \sum_{x_i} p(x_i | x_i \geq 2, X) \sum_{\mathbf{x}_i} x_j p(\mathbf{x}_i | X_i = X - x_i) \\
&= \sum_X P(X | x_i \geq 2) \sum_{x_i} p(x_i | x_i \geq 2, X) \frac{l_j(X - x_i)}{L - l_i} \\
&= \frac{l_j}{L - l_i} \left[\sum_X P(X | x_i \geq 2) X - \sum_{x_i} p(x_i | x_i \geq 2) x_i \right] \\
&= \frac{l_j}{L - l_i} \left(L \frac{1 - e^{-L}}{1 - (1 + L)e^{-L}} - l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}} \right). \tag{21}
\end{aligned}$$

To calculate the relative fluctuations in $\sum_{j \neq i} c_j x_j$, we use a similar approximation as for $\Delta_r n_i$: \hat{p} is approximated by \hat{q} , defined as the \mathbf{x} dispersal probabilities in a territory conditional on $x_i > 2$ (that is, treating the x_j as independent). All covariances between nonfocal genotypes are now zero, so that $\sigma_{\hat{q}}^2(\sum c_j x_j) = \sum c_j^2 \sigma_{\hat{q}}^2(x_j)$, where $\sigma_{\hat{q}}^2(x_j) = l_j$ for $j \neq i$, and

$$\sigma_{\hat{q}}^2(x_i) = \frac{l_i}{D} \left(l_i + 1 - e^{-l_i} - \frac{l_i}{D} (1 - e^{-l_i})^2 \right), \tag{22}$$

where $D = 1 - (1 + l_i)e^{-l_i}$, and

$$\frac{\sigma_{\hat{q}}(\sum c_j x_j)}{\langle \sum c_j x_j \rangle} = \frac{\left(\sum_{j \neq i} c_j^2 l_j + c_i^2 \sigma_{\hat{q}}^2(x_i) \right)^{1/2}}{\sum_{j \neq i} c_j l_j + c_i l_i (1 - e^{-l_i}) / D}. \tag{23}$$

Similarly to Eq. (16), the RHS of Eq. (23) is $\ll 1$ for the case that $L \ll 1$ (due to a factor of $D^{1/2}$), and also for the case when at least some of the propagule densities (focal or nonfocal) are large — provided that c_i and the c_j are all of similar magnitude. Again, the worst case scenario occurs when l_i and $L - l_i$ are of order 1, in which case Eq. (23) is around 35%, which is again where the \hat{q} approximation produces the biggest overestimate of the fluctuations in \mathbf{x} . Similarly

621 to Eq. (16), the RHS of (23) will not be $\ll 1$ in the presence of a rare, extremely strong competitor.

Combining Eqs. (18) and (19), we obtain

$$\Delta_a n_i = m_i A_i \frac{c_i}{\bar{c}}, \quad (24)$$

where A_i is defined in Eq. (6).

624 **Comparison with simulations**

Fig. 6 shows that Eq. (4) and its components closely approximate our density-dependent lottery model over a wide range of propagule densities (the latter is evaluated by direct simulations of
627 uniform random dispersal and lottery competition). Two genotypes are present, one of which is at low frequency. The growth of the low-frequency genotype relies crucially on the low-density competition term $R_i c_i / \bar{c}$, and also to a lesser extent on the high density competition term $A_i c_i / \bar{c}$
630 if l_1 is large enough (Fig. 6b). On the other hand, $R_i c_i / \bar{c}$ is negligible for the high-frequency genotype, which depends instead on high density territorial victories (Fig. 6d). Fig. 3 also shows the breakdown of the classic lottery model at low propagule densities.

633 **Appendix C: Mutant invasion under different environments**

In this Appendix we evaluate the invasion of novel mutants in a population with a single resident type i (such that $N = n_i$), which is in equilibrium.

636 For a single type i in equilibrium ($\Delta n_i = 0$), we have $R_i = 0$, $\bar{c} = c_i$, $A_i = (1 - (1 + L)e^{-L})/L$, and Eq. (4) becomes

$$b_i(1 - e^{-L})/L - d_i = 0. \quad (25)$$

(Alternatively, Eq. (25) can be deduced directly from Eq. (2)). This implies $L \approx b_i/d_i$ when
639 $b_i/d_i \gg 1$ and $L \ll 1$ when $b_i/d_i \approx 1$. Now suppose that a novel mutant j , which is initially rare, appears in the population. Then $A_j/R_j \ll 0$, $l_j \approx 0$ and $\bar{c} \approx c_i$, and so, from Eq. (4), the mutant




Figure 6: The change in genotype abundances in a density dependent lottery model is closely approximated by Eq. (4). $\Delta_+ n_i / m_i$ from Eq. (4) (and its separate components) are shown, along with direct simulations of random dispersal and lottery competition over one iteration over a range of propagule densities (U is varied between 5×10^3 and 10^6 with $m_1 = 10^4$ and $m_2 = 9 \times 10^4$). Two genotypes are present. (a) and (b) show the low-frequency genotype with c -advantage ($c_1 = 1.5$), (c) and (d) show the high-frequency predominant genotype ($c_2 = 1$). Simulation points are almost invisible in (c) and (d) due to near exact agreement with Eq. (4). Dashed lines in (a) and (c) show the breakdown of the classic lottery model.

lineage's fitness is

$$\Delta n_j/n_j \approx b_j \left(e^{-L} + R_j \frac{c_j}{c_i} \right) - d_j \quad (26)$$

642 where $R_j \approx (1 - e^{-L}) / \left(\frac{c_j}{c_i} + \frac{L-1+e^{-L}}{1-(1+L)e^{-L}} \right)$ since $l_j \ll 1$.

We now consider mutant invasion under the different environments from section "Primary strategies and Grime's triangle". In our representation of Grime's stressful environment, $b \ll 1$,
645 $d \approx b$ and $L \ll 1$, and so Eq. (26) becomes $\Delta n_j/n_j \approx b_j - d_j$. Mutations which improve b or d by the same fraction ϵ , such that $b_j = (1 + \epsilon)b_i$ or $d_j = (1 - \epsilon)d_i$, yield identical fitness benefits.

In our representation of Grime's ideal environment, $d \ll 1$, $b_i/d_i \gg 1$ and $L \gg 1$, and so Eq.
648 (26) becomes

$$\Delta n_j/n_j \approx \frac{b_j}{c_j/c_i + L - 1} \frac{c_j}{c_i} - d_j \approx \frac{b_j}{L} \frac{c_j}{c_i} - d_j, \quad (27)$$

where the last approximation follows from the fact that c_j will be of similar magnitude to c_i (ignoring mutations with very large effect sizes; incidentally, Eq. (1) yields the same expression
651 by assuming $L \rightarrow \infty$ even though this assumption is unrealistic for a rare mutant). Similarly to Grime's stressful environment, improving b , c , or d by a fraction ϵ yields identical fitness benefits, even though they may have very different magnitudes, because the ancestral resident
654 type's density-regulated birth rate $b_i c_i / L$ exactly balances its mortality d_i in equilibrium. Biases in the direction of evolution of b , c or d must again follow from biases in mutation availability or trade-offs/pleiotropy. The same argument applies for high- L interpretation of the stress regime.

657 To represent Grime's disturbed environment, $\Delta_+ n_i$ is replaced with $(1 - d_i)\Delta_+ n_i$, and a single type reaches equilibrium when $L/(1 - e^{-L}) = (1 - d_i)b_i/d_i$ (population persistence now requires $(1 - d_i)b_i > d_i$ rather than $b_i > d_i$). We assumed $d_i \approx 1$ and $(1 - d_i)b_i \approx d_i$, so that $L \ll 1$. A
660 mutant's invasion fitness is then given by

$$\Delta n_j/n_j = (1 - d_j)b_j - d_j. \quad (28)$$

Another distinct feature of disturbed environments is that the bulk of mortality is environmental

and unavoidable. That is, $d_i = d_{ie} + d_{ig}$, where the environmental part $d_{ie} \approx 1$ does not evolve,
 663 and the genetic part $d_{ig} \ll 1$ is affected by mutations in the same way as before $d_{ig} = (1 - \epsilon)d_{ig}$.

The effects of improving b and d are now no longer identical. Increasing b by a fraction ϵ gives
 $\Delta n_j/n_j = \epsilon(1 - d_i)b_i \approx \epsilon$, whereas reducing d by the same fraction gives $\Delta n_j/n_j = \epsilon d_{ig}/(1 - d_i)$.
 666 Thus, even though the genetic component of mortality d_g is small compared to the environmental
 component d_e , if disturbance is intense enough ($1 - d_i$ is small enough), selection for reduced
 mortality is potentially much stronger than that for propagule production.

669 Appendix D: Coexistence in a stable environment

To determine whether coexistence is possible in a constant environment, we check for “mutual
 invasion”, that is, we check that j will invade an i -dominated population, but i will also invade a
 672 j -dominated population.

We consider the case of coexistence between a b specialist i and a c specialist j ($b_i > b_j$, $c_j > c_i$
 and $d_i = d_j$). Suppose that b_i is so large that $L \gg 1$ when i is dominant, and b_j is so small that $L \ll$
 675 1 when j is dominant. Then, when j is dominant, we have $\Delta n_i/n_i = b_i - d_i = b_i - d_j = b_i - b_j > 0$.
 When i is dominant, Eq. (27) applies, where Eq. (25) implies $d_j = d_i = b_i(1 - e^{-L})/L \approx b_i/L$,
 and so

$$\Delta n_j/n_j \approx \frac{b_j c_j}{L c_i} - \frac{b_i}{L}. \quad (29)$$

678 Therefore, coexistence occurs if c_j/c_i is sufficiently large. The analogous argument for d and c
 specialists ($d_i < d_j$ with $L \gg 1$ when i dominates, $L \ll 1$ when j dominates, and $b_i = b_j$) gives
 $\Delta n_j/n_j \approx d_i \frac{c_j}{c_i} - d_j$, which again implies coexistence if c_j/c_i is sufficiently large.

681 For b and d specialists ($c_i = c_j$), we have $\Delta n_j/n_j \approx b_j d_i/b_i - d_j$ when i dominates and $\Delta n_i/n_i \approx$
 $b_i d_j/b_j - d_i$ when j dominates. Thus, either i or j grows when rare, but not both, and stable
 coexistence is not possible in a constant environment.