

A lottery model of density-dependent selection in evolutionary genetics

Manuscript elements:

Keywords: r/K selection, absolute fitness, eco-evo, Grime's triangle, competition-colonization trade-off, fluctuating selection, storage effect.

Manuscript type: Article.

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

Fitness is typically represented in heavily simplified terms in evolutionary genetics, often using constant selection coefficients, to make it easier to infer or predict how type frequencies change over time. This excludes fundamental ecological factors such as dynamic population size or density-dependence from the most genetically-realistic treatments of evolution, a problem that inspired MacArthur's influential but problematic r/K theory. Following the spirit of r/K -selection as a general-purpose theory of density-dependent selection, we develop a new model of density-dependent selection by generalizing the fixed-density classic lottery model of territorial acquisition to accommodate arbitrary population densities. We show that, with density dependence, co-existence is possible in the lottery model in a stable environment. Inspired by natural *Drosophila* populations, we consider co-existence under strong, seasonally-fluctuating selection coupled to large cycles in population density, and show that co-existence (stable polymorphism) is promoted via a combination of the classic storage effect and density-regulated population growth. We also show that the only significant bias introduced by selection at the different environmental extremes of Grime's triangle is the relative importance of competitive ability at different densities, confirming an important role for phenotypic constraints in shaping "primary strategies".

18 “...the concept of fitness is probably too complex to allow of a useful mathematical devel-
opment. Since it enters fundamentally into many population genetics considerations, it is re-
markable how little attention has been paid to it.” — Warren J. Ewens, *Mathematical Population*
21 *Genetics I*, 2004, pp. 276

Introduction

Evolutionary models differ greatly in their treatment of fitness. In models of genetic evolution,
24 genotypes are typically assigned constant (or occasionally frequency-dependent) selection coef-
ficients describing the change in their relative frequencies over time. This simplified treatment
of selection facilitates explicit time-dependent treatment of genotype frequencies, and can be jus-
27 tified over sufficiently short time intervals (Ewens, 2004, p. 276). The emphasis of population
genetics is to infer past selection, migration and demographic change given a sample of nu-
cleotide sequences, or to predict how allele frequencies change over time based on their relative
30 fitness effects together with population structure, genetic drift and linkage. The resulting pic-
ture of evolution excludes basic elements of the ecological underpinnings of selection, including
density dependence, and how selection affects population size. This complicates the inference
33 of past selection, because demographic changes can look genealogically very similar to selective
frequency changes (Barton, 1998).

By contrast, models of phenotypic trait evolution use absolute fitness functions to describe
36 how some traits of interest affect survival and reproduction in particular ecological scenarios
(Diekmann et al., 2004; Metz et al., 1992). These fitness functions can be quite problem-specific
and often only account for a few traits at a time. The emphasis here is on the conditions for
39 invasion from low frequencies and co-existence, rather than frequency or abundance trajectories
over time. For instance, adaptive dynamics uses “invasion fitness” to explore the consequences
of eco-evolutionary feedbacks (Diekmann et al., 2004).

42 It is challenging to generalize beyond particular traits or ecological scenarios to model fun-

damentally different forms of selection. Perhaps this is not surprising given that fitness is such a complex quantity, dependent on all of a phenotype's functional traits (Violle et al., 2007) and its environment. A detailed, trait-based, predictive model of fitness would be enormously complicated and situation-specific. It is therefore easy to doubt the feasibility of a simplified, general mathematical treatment of fitness (Ewens, 2004, p. 276). For example, MacArthur's famous r/K scheme (MacArthur, 1962; MacArthur and Wilson, 1967) is now almost exclusively known as a framework for understanding life-history traits, and judged on its failure in that role (Boyce, 1984; Pianka, 1970; Reznick et al., 2002; Stearns, 1977). However, the r/K scheme's original purpose was to extend the existing population-genetic treatment of selection to account for population density (MacArthur, 1962). Few attempts have been made to develop it further along these lines.

However, empirical trait classification studies have suggested the existence of a few "primary strategies", reflecting broadly distinct responses to selection (Winemiller et al., 2015). Grime famously considered (Grime, 1974, 1977, 1988; Westoby, 1998) two broad determinants of population density: stress (persistent hardship e.g. due to resource scarcity or unfavorable temperatures) and disturbance (intermittent destruction of vegetation e.g. due to trampling, herbivory, pathogens, extreme weather or fire). The extremes of these two factors define three primary strategies denoted by C/S/R respectively (Fig. 1): competitors "C" excel in low stress, low disturbance environments; stress tolerators "S" excel in high stress, low disturbance environments; and ruderals "R" excel in low stress, high disturbance environments. Population persistence is not possible in high-stress, high-disturbance environments. Grime showed that measures of C, S and R across a wide range of plant species are anti-correlated, so that strong C-strategists are weak S and R strategists, and so on, creating a triangular C/S/R ternary plot (Grime, 1974). Similar schemes were proposed for insects (Southwood, 1977), fishes (Winemiller and Rose, 1992), and zooplankton (Allan, 1976). More recently, modern hierarchical clustering techniques have also revealed, with greater statistical rigor, distinct trait clusters in corals analogous to Grime's primary strategies (Darling et al., 2012). These empirical findings suggest that functional traits contribute to fitness predominantly via a few key factors such as intrinsic reproductive rate or




Figure 1: Schematic of Grime’s triangle and the associated direction of selection in our model. The two axes show increasing levels of environmental stress and disturbance. Population persistence is not possible if the combination of stress and disturbance is too large (below dashed line). This creates a triangle, each corner of which corresponds to an environmental extreme and associated “primary strategy”. The high-density (HD) interpretation of stress is shown as a rotation of the triangle with respect to density. The strength of selection on c depends on density L , but b and d experience the same selective force in all three environmental extremes in both versions of the triangle.

stress-tolerance.

Here we explore the interplay between some “key factors” of fitness in a simplified, territorial model of growth, dispersal and competition. This broadly follows the original spirit of MacArthur’s r/K scheme. More specifically, our aim is to begin merging some ecological realism to population genetics’ time-dependent, genetically-focused view of evolution. We revisit the classic lottery model of Chesson and Warner (1981), which has two features that make it well suited for this role, but one critical flaw that we rectify here.

The first feature is that the lottery representation of competition is particularly concise. Mature individuals (“adults”) each require their own territory, whereas newborn individuals (“propagules”) disperse to, and subsequently compete for, territories made available by the death of adults. Territorial contest among propagules leaves a single victorious adult per territory, the victor chosen at random from the propagules present, with probabilities weighted by a coefficient for each type representing competitive ability, akin to a lottery (Sale, 1977). By comparison, coefficients for the pairwise effects of types on each other (e.g. the α coefficients in the gen-

eralized Lotka-Volterra equations and the associated concept of “ α -selection”; Case and Gilpin 1974; Gill 1974; Joshi et al. 2001), or explicit resource consumption (Tilman, 1982), are much more complicated. The second feature is the close connection between the lottery model and one of the foundational models of population genetics, the Wright-Fisher model of genetic drift, which we discuss further below.

The critical flaw of the classic lottery model is that it breaks down at low densities (few propagules dispersing to each territory), precluding density-dependent behaviour. Our first task is to analytically extend the classic lottery model to correctly account for low density behavior (sections “Model” and “Mean field approximation”).

Using our extended lottery model, we then revisit Grime’s C/S/R scheme, and evaluate the extent to which selection favors fecundity, competitive ability or lower mortality under Grime’s environments (section “Primary strategies and Grime’s triangle”). This represents a simple “sanity check” on the invasion behavior of our extended lottery model under different extremes.

We then explore some time-dependent behavior of our extended lottery model. Taking an example inspired by recent studies of rapid, seasonal evolution in *Drosophila* (Bergland et al., 2014), we discuss how environmental fluctuations might stabilize polymorphisms in the presence of cyclical population density.

Model

We assume that reproductively mature individuals (“adults”) each require their own territory to survive and reproduce (Fig. 2). All territories are identical, and the total number of territories is T . Time t advances in discrete iterations, each representing the time from birth to reproductive maturity. In iteration t , the number of adults of the i ’th genotype is $n_i(t)$, the total number of adults is $N(t) = \sum_i n_i(t)$, and the number of unoccupied territories is $U(t) = T - N(t)$.

We assume that the n_i and T are large enough that stochastic fluctuations in the n_i (“drift”) can be ignored. We derive deterministic equations for the expected change in the n_i over time,

leaving the evaluation of drift for future work. This is an excellent approximation when the n_i are all large. However, we also do not evaluate the initial stochastic behaviour of adaptive mutant lineages while they are at low abundance. When considering new mutations, we therefore restrict our attention to begin with the earliest (lowest n_i) deterministic behavior of mutant lineages (the transition to deterministic growth occurs at an abundance n_i of order equal to their inverse expected absolute growth rate; Uecker and Hermisson 2011).

Each iteration, adults produce new offspring (“propagules”), m_i of which disperse to unoccupied territories. We assume that adults cannot be ousted from their territories, so that m_i only includes propagules landing on unoccupied territories. Propagules disperse at random over the unoccupied territories, regardless of distance from their parents, and independently of each other. There is no interaction between propagules (e.g. avoidance of territories crowded with propagules). Loss of propagules during dispersal is subsumed into m_i .

In general, m_i will increase with n_i , and will depend on population density N . For example, if b_i is the number of successfully dispersing propagules produced per genotype i adult, then the loss of propagules due to dispersal to occupied territories implies $m_i = b_i(1 - N/T)n_i$, akin to Levins’ competition-colonization model (Levins and Culver, 1971; Tilman, 1994). In section “Cyclical birth and death rates” we evaluate Eq. (4) numerically using this functional form for m_i , with b_i assumed to be constant.

In the sections “Invasion of rare genotypes and coexistence” and “Primary strategies and Grime’s triangle”, we assume the simpler form $m_i = b_i n_i$, with constant b_i , meaning that all propagules land on unoccupied territories (a form of directed dispersal). This simplifies the mathematics without affecting the results of those sections, which only depend on the low-frequency invasion behavior of Eq. (4). Note that due to our assumption of uniform dispersal, the parameter b_i can be thought of as a measure of “colonization ability”, which combines fecundity and dispersal ability (Bolker and Pacala, 1999; Levins and Culver, 1971; Tilman, 1994).

The number of individuals of the i ’th genotype landing in any particular territory is denoted x_i . We assume that x_i follows a Poisson distribution $p_i(x_i) = l_i^{x_i} e^{-l_i} / x_i!$, where $l_i = m_i / U$ is the




Figure 2: Each iteration of our lottery model has three main elements. First, propagules are produced by adults which are dispersed at random over the unoccupied territories (only propagules landing on unoccupied territories are shown). Lottery competition then occurs in each unoccupied territory (competition in only one territory is illustrated): each genotype has a probability proportional to $b_i n_i c_i$ of securing the territory. Then occupied territories are freed up by adult mortality. In Eq. (3) and most of the paper, only adults can die (red crosses), but we will also consider the case where juveniles die (blue cross; section “Primary strategies and Grime’s triangle”).

mean territorial propagule density. This approximation becomes exact when the n_i are large enough that drift in n_i can be ignored (Appendix A).

When multiple propagules land on the same territory, the victor is determined by lottery competition: genotype i wins a territory with probability $c_i x_i / \sum_j c_j x_j$, where c_i is a constant representing relative competitive ability (Fig. 2).

In the classic lottery model (Chesson and Warner, 1981), unoccupied territories are assumed to be saturated with propagules from every genotype $l_i \gg 1$. From the law of large numbers, the composition of propagules in each territory will then not deviate appreciably from the mean composition l_1, l_2, \dots, l_G (G is the number of genotypes present), and so the probability that genotype i wins any particular unoccupied territory is approximately $c_i l_i / \sum_j c_j l_j$. Let $\Delta_+ n_i$ denote the number of territories won by genotype i . Then $\Delta_+ n_1, \Delta_+ n_2, \dots, \Delta_+ n_G$ follow a multinomial distribution with U trials and success probabilities $\frac{c_1 l_1}{\sum_j c_j l_j}, \frac{c_2 l_2}{\sum_j c_j l_j}, \dots, \frac{c_G l_G}{\sum_j c_j l_j}$, respectively. Genotype i is expected to win $c_i l_i / \sum_j c_j l_j$ of the U available territories, and deviations from this expected outcome are small (since T is large by assumption), giving

$$\Delta_+ n_i(t) = \frac{c_i l_i}{\sum_j c_j l_j} U(t) = b_i n_i \frac{1}{L} \frac{c_i}{\bar{c}}, \quad (1)$$

where $\bar{c} = \sum_j c_j m_j / M$ is the mean propagule competitive ability for a randomly selected propagule, $L = M/U$ is the total propagule density and $M = \sum_j m_j$ is the total number of propagules.

There is a close connection between the classic lottery model and the Wright-Fisher model of genetic drift (Svardal et al., 2015). In the Wright-Fisher model, genotype abundances are sampled each generation from a multinomial distribution with success probabilities $w_i n_i / \sum_j w_j n_j$, where w is relative fitness and the n_i are genotype abundances in the preceding generation. Population size N remains constant. This is mathematically equivalent to the classic lottery model with non-overlapping generations ($d_i = 1$ for all i) and $w_i = b_i c_i$. Thus, the classic lottery model allows us to replace the abstract Wright-Fisher relative fitnesses w_i with more ecologically-grounded fecundity, competitive ability and mortality parameters b_i , c_i and d_i , respectively. Since birth and

death rates affect absolute abundances, this allows us to evaluate selection at different densities (after appropriate extensions are made), in an otherwise very similar model to the canonical Wright-Fisher. We therefore expect that drift in realized values of n_i in our extended lottery model should be similar to that in the Wright-Fisher model, but we leave this for future work.

In our extension of the classic lottery model, we do not restrict ourselves to high propagule densities. Eq. (1) is nonsensical if even a single type has low propagule density ($l_i \ll 1$): genotype i can win at most m_i territories, yet Eq. (1) demands $c_i l_i / \sum_j c_j l_j$ of the U unoccupied territories, for any value of U . Intuitively, the cause of this discrepancy is that individuals are discrete. Genotypes with few propagules depend on the outcome of contests in territories where they have at least one propagule present, not some small fraction of a propagule as would be implied by small l_i in the classic lottery model. In other words, deviations from the mean propagule composition l_1, l_2, \dots, l_G are important at low density.

We expect that a fraction $p_1(x_1) \dots p_G(x_G)$ of the U unoccupied territories will have the propagule composition x_1, \dots, x_G . Genotype i is expected to win $c_i x_i / \sum_j c_j x_j$ of these. Ignoring fluctuations about these two expectations (due to our no-drift, large T , large n_i approximation), genotype i 's territorial acquisition is given by

$$\Delta_+ n_i(t) = U(t) \sum_{x_1, \dots, x_G} \frac{c_i x_i}{\sum_j c_j x_j} p_1(x_1) \dots p_G(x_G), \quad (2)$$

in our extended lottery model, where the sum only includes territories with at least one propagule present. Note that unlike the classic lottery model, not all unoccupied territories are claimed each iteration, since under Poisson dispersal a fraction e^{-L} remain unoccupied.

For the majority of this manuscript we assume that mortality only occurs in adults (setting aside the juvenile deaths implicit in territorial contest), and at a constant, genotype-specific per-capita rate $0 \leq d_i \leq 1$, so that the overall change in genotype abundances is

$$\Delta n_i(t) = \Delta_+ n_i(t) - d_i n_i(t). \quad (3)$$

This seems reasonable in the absence of disturbances; when we come to consider the effects of disturbances (Section “Primary strategies and Grime’s triangle”), we will incorporate disturbance-induced mortality in competing juveniles (Fig. 2).

Results

Mean Field Approximation

Eq. (2) involves an expectation over the time-dependent dispersal distributions p_i , and is thus too complicated to give intuition about the dynamics of density-dependent lottery competition.

We now evaluate this expectation using a “mean field” approximation.

Similarly to the high- l_i approximation of classic lottery model, we replace the x_i with appropriate mean values, although we cannot simply replace x_i with l_i . For a genotype with low propagule density $l_i \ll 1$, we have $x_i = 1$ in the territories where its propagules land, and so its growth comes entirely from territories which deviate appreciably from l_i . To account for this, we separate Eq. (2) into $x_i = 1$ and $x_i > 1$ parts. Our more general mean field approximation only requires that there are no large discrepancies in competitive ability (i.e. we do not have $c_i/c_j \gg 1$ for any two genotypes). We obtain (details in Appendix B)

$$\Delta_+ n_i(t) \approx b_i n_i \left[e^{-L} + (R_i + A_i) \frac{c_i}{\bar{c}} \right], \quad (4)$$

where

$$R_i = \frac{\bar{c} e^{-l_i} (1 - e^{-(L-l_i)})}{c_i + \frac{L-1+e^{-L}}{1-(1+L)e^{-L}} \frac{\bar{c} L - c_i l_i}{L-l_i}}, \quad (5)$$

and

$$A_i = \frac{\bar{c} (1 - e^{-l_i})}{\frac{1-e^{-l_i}}{1-(1+l_i)e^{-l_i}} c_i l_i + \frac{1}{L-l_i} \left(L \frac{1-e^{-L}}{1-(1+L)e^{-L}} - l_i \frac{1-e^{-l_i}}{1-(1+l_i)e^{-l_i}} \right) \sum_{j \neq i} c_j l_j}. \quad (6)$$

To supplement our analytical mean field derivation, we did numerical simulations of our exact density-dependent lottery model, and verified that Eq. (4) is a good approximation (Ap-

pendix B). Thus, Eq. (4) describes how type abundances change over time in a lottery model where population density can itself vary with time.

Comparing Eq. (4) to Eq. (1), the classic lottery per-propagule success rate $c_i/\bar{c}L$ has been replaced by three separate terms. The first, e^{-L} , accounts for propagules which land alone on unoccupied territories; these territories are won without contest. The second, $R_i c_i/\bar{c}$ represents competitive victories when the i genotype is a rare invader in a high density population: from Eq. (5), $R_i \rightarrow 0$ when the i genotype is abundant ($l_i \gg 1$), or other genotypes are collectively rare ($L - l_i \ll 1$). The third term, $A_i c_i/\bar{c}$, represents competitive victories when the i genotype is abundant: $A_i \rightarrow 0$ if $l_i \ll 1$. The relative importance of these three terms varies with both the overall propagule density L and the relative propagule frequencies m_i/M . If $l_i \gg 1$ for all genotypes, we recover the classic lottery model (only the $A_i c_i/\bar{c}$ term remains, and $A_i \rightarrow 1/L$).

Primary strategies and Grime's triangle

Here we describe how Grime's "disturbed", "stressful" and "ideal" environments can be captured mathematically in our model, and evaluate role of selection in shaping the corresponding "primary strategies". To proceed, we map these verbally-defined environments to quantitative parameter regimes in our model, summarized in Fig. 3.

Ideal environments are characterized by the near-absence of stress and disturbance, so that survival is easy $d \ll 1$ and growth and reproduction can be rapid $b \gg d$. The propagule density is determined by the magnitude of b/d , and in this case $L \gg 1$ (Appendix C).

Disturbed environments are characterized by short bursts of high extrinsic mortality d caused by physical destruction. We assume that disturbances are equally damaging to adults and juveniles, so that only $(1 - d_i)\Delta_+ n_i$ (rather than $\Delta_+ n_i$) territories are secured by type i each iteration. In the heavily disturbed extreme of Grime's triangle, many such bursts (which could each be spatially localized) will occur each iteration (i.e. over the time required for a propagule to reach reproductive maturity), which we approximate as a high constant mortality rate $d \approx 1$. Population persistence requires $b > d/(1 - d)$, which implies $b \gg 1$ (severe disturbance comes with

	Ideal	Disturbance*	Stress (G)	Stress (HD)
Constraints	$d \ll 1$	$d \approx 1$	$b \ll 1$	$b \ll 1$
Other parameters	$b/d \gg 1$	$(1-d)b \approx d$	$b \approx d$	$b/d \gg 1$
Propagule density L	$\gg 1$	$\ll 1$	$\ll 1$	$\gg 1$

Figure 3: The mathematical representation of Grime’s three environmental extremes in our model, as well as the high-density (HD) variant of the stressful environment. *Mortality affects both adults and juveniles in the disturbed environment, with $\Delta_+ n_i$ replaced by $(1 - d_i)\Delta_+ n_i$.

a high fecundity/dispersal requirement). We assume this requirement can just be met, so that

228 $b \approx d/(1 - d)$ and $L \ll 1$ (Appendix C).

Stressful environments are more ambiguous, and have been the subject of an extensive debate in the plant ecology literature (the “Grime-Tilman” debate; Aerts 1999 and references therein).

231 Severe stress inhibits growth and reproduction, so that $b \ll 1$. In Grime’s view, this means that the rate at which propagules successfully develop to adulthood cannot appreciably exceed the mortality rate $b/d \approx 1$; it is just possible to cope with the environmental stress (Grime, 1974, 234 1977). In our model, this implies $L \ll 1$.

The alternative view is that stressed environments are highly competitive. In spite of inhibited growth and reproduction, density is actually high relative to carrying capacity of the stressful environment (Taylor et al., 1990). For example, if consumable resources are scarce, we expect 237 intense resource competition (Davis et al., 1998). Thus, even though $b \ll 1$, we still have $b/d \gg 1$ and $L \ll 1$.

240 The expected direction of evolution is determined by three factors. First, there may be pleiotropy between b , c , and d , which could be due to trade-offs between them. Second, adaptive mutations in one trait may occur more frequently than in others (Yampolsky and Stoltzfus, 243 2001). Third, selection favors some trait changes over others: b , c and d will tend to evolve in the direction of greatest fitness increase because the probability that an adaptive mutant survives its initial low abundance phase and begins to grow deterministically is proportional to its expected 246 absolute growth rate (Haldane, 1927; Uecker and Hermisson, 2011). Only a tiny proportion of neutral and deleterious mutations fix (probability of order $1/N$ for neutrals).

Accounting for the first two factors would require an evaluation of the effects of mutations
 249 on traits subject to trade-offs, which is beyond the scope of this article. We only consider how
 selection shapes the direction of evolution for b , c and d , assuming that mutations affect them in-
 dependently and with the same distribution of beneficial fitness effects. Specifically, we consider
 252 the effect of mutations that change b , c , or d by a fixed proportion $b_j = (1 + \epsilon)b_i$, $c_j = (1 + \epsilon)c_i$
 or $d_j = (1 - \epsilon)d_i$, where j is a novel mutant (Appendix C). Mutations with magnitude ϵ are
 assumed to occur with the same frequency for all traits .

255 Not surprisingly, mutations which increase c are effectively neutral when $L \ll 1$ and adaptive
 when $L \gg 1$. However, beyond this, selection in our density-dependent lottery model does
 not favor one trait improvement over another (Fig. 3). This applies regardless of the initial
 258 magnitude of b or d . More specifically, in ideal and stressed environments, where Eq. (4) applies,
 b and d mutations of size ϵ have identical fitness effects (Appendix C), and the same is true for
 proportional c -mutations $c_j = (1 + \epsilon)c_i$ when $L \gg 1$.

261 In disturbed environments, $\Delta_+ n_i$ is replaced by $(1 - d_i)\Delta_+ n_i$. The bulk of the high mortality
 $d \approx 1$ is environmental and unavoidable. This implies that ϵ mutations in the remaining genetic
 part of mortality have a reduced impact on d , suggesting the primacy of b under disturbance.
 264 However, this effect is easily swamped by the fact that $d_i \approx 1$ and that disturbance mortality
 affects both juvenile recruitment and adult mortality. In Appendix C, we argue that b and d
 mutations should have comparable effects on fitness (assuming that d can evolve at all) because
 267 the fitness benefit of improving d declines as d improves and so d cannot run ahead of b for long
 (diminishing returns).

Coexistence in constant and cyclical environments

270 In the previous section we only considered how b , c and d should respond in Grime's environ-
 mental extremes, based on invasion fitness. Here we further explore the low frequency behavior
 of Eq. (4) to determine which types can coexist in a constant environment, and then consider the
 273 full time-dependent behaviour of Eq. (4) in a cyclical environment.

In a constant environment, stable coexistence is possible in our extended lottery model. A b specialist i and c -specialist j ($b_i > b_j$, $c_j > c_i$) can co-exist because then propagule density L is frequency dependent, and so is the importance of competitive ability (Appendix D). This is a version of the classic competition-colonization trade-off (Levins and Culver, 1971; Tilman, 1994): the competitor (c -specialist) leaves many territories unoccupied (low L) due to its poor colonization ability (low b), which the colonizer (b -specialist) can then exploit. A similar situation holds for coexistence between high- c and low- d specialists; a “competition-longevity” trade-off (Tilman, 1994). These forms of co-existence require density dependence (being mediated by L), and are not present in the classic lottery model. Coexistence is not possible between b - and d -specialists (Appendix D).

Now suppose that birth and death rates vary periodically with amplitude sufficient to cause large changes in population density. This example is inspired by natural *Drosophila* populations, which expand rapidly in the warmer months when fruit is abundant, but largely die off in the colder months. Within this seasonal population density cycle, hundreds of polymorphisms also cycle in frequency (Bergland et al., 2014). Some of these polymorphisms may be adaptive and potentially millions of years old, suggesting stable coexistence (Bergland et al., 2014; Messer et al., 2016). Selection on allele frequencies thus occurs on the same time scale as population demography, a situation vastly more complicated than classical sweeps in demographically stable populations (Messer et al., 2016).

The classical population genetic treatment of fluctuating selection suggests that environmental fluctuations do not promote coexistence. Allele frequencies are successively multiplied by relative fitness values for each environmental iteration, and so two alleles favored in different environments can only stably coexist if the product of fitnesses for one type exactly equals the product for the other (Dempster, 1955). Thus, stable coexistence still requires frequency-dependent selection or heterozygote advantage (as is required in a constant environment).

This classical argument overlooks two general effects that promote coexistence in fluctuating environments. The first is the storage effect, which occurs when part of the population is pro-

tected from selection (due to overlapping generations in the lottery model; Chesson and Warner 1981). The second is the bounded population size effect of Yi and Dean (2013), which occurs
 303 when each environmental cycle involves growth from low to high density, with the time spent growing each cycle dependent on the fitness of the types present.

Fig. 4a-c shows the behavior of Eq. (4) for an example where b and d cycle between zero and
 306 positive values (“summers” with rapid growth and no mortality, and “winters” with mortality and no growth). Both the storage effect (adults are sheltered from selection during the summer growth phase) and the bounded density effect (expansion to high density occurs every cycle)
 309 are operating. Two types are present, a b specialist, which is better at rapidly growing in the summer (higher b), and a d specialist which is better at surviving the winter (lower d). Neither type has an advantage over a full environmental cycle, and they stably coexist. This is due to a
 312 combination of the storage and bounded density effects (recall that stable coexistence between b and d specialists was not possible in a constant environment).

The classic lottery model (Eq. 1) gives dramatically different co-existence predictions because
 315 expansion to carrying capacity occurs immediately at the start of the summer (Fig. 4d-f). As a result, coexistence requires that the winter survivor’s b must be about 5 times smaller than required when we properly account for the growth in the abundance of each type using Eq. (4)
 318 (keeping the other parameters the same). Previous models of the promotion of genetic variation via the storage effect (Ellner and Hairston Jr, 1994) similarly assume that the total number of offspring per iteration is constant, and would produce a similar error.

321 Discussion

It is interesting to compare the predictions of the extended lottery model with earlier approaches, such as the r/K scheme, where $r = b - d$ is the maximal, low-density growth rate (Pianka, 1972).
 324 Confusingly, the term “ K -selection” sometimes refers generally to selection at high density (Pianka, 1972), encompassing both selection for higher saturation density (MacArthur and Wilson,




Figure 4: Stable coexistence between b and d specialists in a fluctuating environment requires a much greater b advantage in the classic lottery model compared to our density-dependent extension of it when population density is seasonally cyclical. (a) Birth and death rates seasonally alternate being nonzero (white for winter, green for summer). The b specialist (black) has higher b and d ($b = 0.5$, $d = 0.2$) than the d specialist ($b = 0.217$, $d = 0.1$) (blue). (b) Both types grow during the positive b phase, and decline during the positive d phase, but the d specialist does so at a lower rate. Total height (blue+black) is population density N/T . (c) Summer favors the b specialist, winter the d specialist, and they stably coexist. (d-f) Same as (a-c) for the classic lottery model; the types no longer coexist. (g-i) Same as (d-f) where now $b = 0.0421$ for the d specialist and the types coexist. For illustration, the propagule abundances are assumed to have the form $m_i = b_i(1 - N/T)n_i$, reflecting non-directed dispersal.

1967) and competitive ability (Gill, 1974). Contrary to predictions of an r/K trade-off, empirical studies have shown that maximal growth rate and saturation density (measured by abundance) are positively correlated, both between species/strains (Fitzsimmons et al., 2010; Hendriks et al., 2005; Kuno, 1991; Luckinbill, 1979), and as a result of experimental evolution (Luckinbill, 1978, 1979). From the perspective of our model, this positive correlation is not surprising since the saturation density, which is determined by a balance between births and deaths, increases with b .

There is support for a negative relationship between competitive success at high density and maximal growth rate (Luckinbill, 1979), consistent with a tradeoff between r and the competitive aspect of K . This could be driven by a tradeoff between individual size and reproductive rate. To avoid confusion with other forms of “ K -selection”, selection for competitive ability has been called “ α -selection” after the competition coefficients in the Lotka-Volterra equation (Case and Gilpin, 1974; Gill, 1974; Joshi et al., 2001). However, competitive success as measured by α (i.e. the per-capita effect of one genotype on another genotype’s growth rate) is only partly determined by individual competitive ability — in the presence of age-structured competition and territoriality, it also includes the ability of each genotype to produce contestants i.e. b in our model. Our c is strictly competitive ability only — as such, changes in c do not directly affect population density (the total number of territories occupied per iteration is $\Delta_+ N = U(1 - e^{-L})$, which does not depend directly on the c_i). The clean separation of a strictly-relative c parameter is particularly useful from an evolutionary genetics perspective, essentially embedding a zero-sum relative fitness trait within a non-zero-sum fitness model. This could have interesting applications for modeling the impacts of intra-specific competition on species extinction, for example due to clonal interference (Desai and Fisher, 2007; Gerrish and Lenski, 1998) between c -strategists on the one hand, and b - and d - strategists on the other.

K -selection in the narrow logistic sense of selection for a greater environmental carrying capacity for given birth and death rates, sometimes referred to as “efficiency” (MacArthur and Wilson, 1967), could be represented in our model by smaller individual territorial requirements.

To a first approximation, two co-occurring genotypes which differ by a small amount in their territorial requirements only should have the same fitness since the costs or benefits of a change in the amount of unoccupied territory is shared equally among genotypes via the propagule density per territory L . The situation is more complicated when the differences in territorial requirements become large enough that territorial contests can occur on different scales for different genotypes. We leave these complications for future work.

Our realization of Grime's triangle (Fig. 1) differs from approaches which identify primary strategies as trait combinations which can stably co-exist (Bolker and Pacala, 1999), referring instead to the direction and rate of ongoing adaptive trait evolution under different regimes of stress and disturbance, which is closer in spirit to Grime's arguments (Grime, 1974, 1977). Moreover, our formulation is mathematical, in contrast to Grime's original verbal and descriptive approach, which is a recognized hindrance to the evaluation or broader application of the C/S/R scheme (e.g. Tilman 2007). However, section "Primary strategies and Grime's triangle" suggests that, apart from the obvious irrelevance of territorial contests at low density, selection alone has no preference for fecundity, competitive ability or longevity. This could mean that one or both of the factors we set aside — pleiotropy/trade-offs and mutation bias — are needed to get true "primary strategy" trait differentiation. A separate model of trade-offs and mutation effects would be needed to explore this possibility.

Nevertheless, it is interesting to note that ruderals, which are typically thought of as high fecundity dispersers (b specialists), may also be strongly d -selected, consistent with our findings. An effective way to reduce d in the face of unavoidable physical destruction is to shorten the time to reproductive maturity — short life cycles are a characteristically ruderal trait. Moreover, a recent hierarchical cluster analysis of coral traits did find a distinct "ruderal" cluster, but high fecundity was not its distinguishing feature. Rather, ruderals used brood- (as opposed to broadcast-) spawning, which could plausibly be a mechanism for improving propagule survivorship in disturbed environments (Darling et al., 2012).

One potential limitation of our model as a general-purpose model of density-dependent selec-

tion is its restriction to interference competition between juveniles for durable resources (lottery
 381 recruitment to adulthood), analogous to the ubiquitous assumption of viability selection in pop-
 ulation genetics (Ewens, 2004, p. 45). In some respects this is the complement of consumable
 resource competition models, which restrict their attention to indirect exploitation competition,
 384 typically without age structure (Tilman, 1982). In the particular case that consumable resources
 are spatially localized (e.g. due to restricted movement through soils), resource competition and
 territorial acquisition effectively coincide, and in principle resource competition could be repre-
 387 sented by a competitive ability c (or conversely, c should be derivable from resource competition).
 The situation is more complicated if the resources are well-mixed, since, in general, resource lev-
 els then need to be explicitly tracked. It seems plausible that explicit resource tracking may not
 390 be necessary when the focus is on the evolution of similar genotypes that use identical resources
 rather than the stable co-existence of widely differing species with different resource preferences
 (Ram et al., 2016). We are not aware of any attempts to delineate conditions under which explicit
 393 resource tracking is unnecessary even if it is assumed that community structure is ultimately
 determined by competition for consumable resources. More work is needed connecting resource
 competition models to the density-dependent selection literature, since most of the former has
 396 to date been focused on narrower issues of the role of competition at low resource availability
 (Aerts, 1999; Davis et al., 1998; Tilman, 2007).

While our model can be applied to species rather than genotypes (e.g. ecological invasions),
 399 our focus is genotype evolution i.e. the change in allele frequencies over time. Our assumption
 that there are no large c discrepancies (section “Mean field approximation”) amounts to a restric-
 tion on the amount of genetic variation in c in the population. Since beneficial mutation effect
 402 sizes will typically not be much larger than a few percent, large c discrepancies can only arise if
 the mutation rate is extremely large, and so the assumption will not be violated in most cases.
 However, this restriction could become important when looking at species interactions rather
 405 than genotype evolution.

In the introduction we mentioned the recurring difficulties with confounding selection and

demography in population genetic inference. It seems that Eq. (4) or something similar (and
 408 hopefully more analytically tractable) is unavoidable for the analysis of time-course genetic data
 because, fundamentally, selective births and deaths affect both abundances and frequencies, not
 one or the other in isolation. Moreover, some aspects of allele frequency change are intrinsically
 411 density-dependent. In the classic lottery model, which as we have seen is essentially the Wright-
 Fisher model with overlapping generations, b_i and c_i are equivalent in the sense that the number
 of territorial victories only depends on the product $b_i c_i$ (see “Model”). This is no longer the
 414 case in our extension, where b and c specialists can co-exist. This “colonization-competition
 trade-off” is well known in the co-existence literature (Tilman, 1994). It and similar forms of
 “spatial co-existence” in stable environments have previously been modeled either with Levin’s
 417 qualitative representation of competition (Levins and Culver, 1971; Tilman, 1994), as opposed to
 the quantitative c of lottery competition, or with a more sophisticated treatment of space (non-
 uniform dispersal; Bolker and Pacala 1999; Shmida and Ellner 1984). In cyclical environments,
 420 polymorphisms can be stabilized by the bounded density effect which is completely lost if there
 is an exclusive focus on allele frequencies (Yi and Dean, 2013). We leave the details of how our
 model might be applied in inference, including the crucial issue of its genetic drift predictions
 423 (providing a null model for neutral sites), for future work.

References

- Aerts, R. 1999. Interspecific competition in natural plant communities: mechanisms, trade-offs
 426 and plant-soil feedbacks. *Journal of Experimental Botany* 50:29–37.
- Allan, J. D. 1976. Life history patterns in zooplankton. *The American Naturalist* 110:165–180.
- Arenbaev, N. 1977. Asymptotic behavior of the multinomial distribution. *Theory of Probability*
 429 & Its Applications 21:805–810.

Barton, N. H. 1998. The effect of hitch-hiking on neutral genealogies. *Genetical Research* 72:123–133.

432 Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt, and D. A. Petrov. 2014. Genomic evidence of rapid and stable adaptive oscillations over seasonal time scales in *Drosophila*. *PLOS Genetics* 10:1–19.

435 Bolker, B. M., and S. W. Pacala. 1999. Spatial moment equations for plant competition: Understanding spatial strategies and the advantages of short dispersal. *The American Naturalist* 153:575–602.

438 Boyce, M. S. 1984. Reconstitution of *r*- and *K*-selection as a model of density-dependent natural selection. *Annual Review of Ecology and Systematics* 15:427–447.

Case, T. J., and M. E. Gilpin. 1974. Interference competition and niche theory. *Proceedings of the National Academy of Sciences* 71:3073–3077.

Chesson, P. L., and R. R. Warner. 1981. Environmental variability promotes coexistence in lottery competitive systems. *American Naturalist* pages 923–943.

444 Darling, E. S., L. Alvarez-Filip, T. A. Oliver, T. R. McClanahan, and I. M. Côté. 2012. Evaluating life-history strategies of reef corals from species traits. *Ecology Letters* 15:1378–1386.

Davis, M. A., K. J. Wragg, and P. B. Reich. 1998. Competition between tree seedlings and herbaceous vegetation: support for a theory of resource supply and demand. *Journal of Ecology* 86:652–661.

Dempster, E. R. 1955. Maintenance of genetic heterogeneity. *Cold Spring Harb Symp Quant Biol* 20:25–31.

Desai, M. M., and D. S. Fisher. 2007. Beneficial mutation–selection balance and the effect of linkage on positive selection. *Genetics* 176:1759–1798.

- 453 Diekmann, O., et al. 2004. A beginner's guide to adaptive dynamics. Banach Center Publications
63:47–86.
- Ellner, S., and N. G. Hairston Jr. 1994. Role of overlapping generations in maintaining genetic
456 variation in a fluctuating environment. *The American Naturalist* 143:403–417.
- Ewens, W. J. 2004. *Mathematical Population Genetics 1: Theoretical Introduction*. Springer
Science & Business Media.
- 459 Fitzsimmons, J. M., S. E. Schoustra, J. T. Kerr, and R. Kassen. 2010. Population consequences of
mutational events: effects of antibiotic resistance on the r/k trade-off. *Evolutionary Ecology*
24:227–236.
- 462 Gerrish, P. J., and R. E. Lenski. 1998. The fate of competing beneficial mutations in an asexual
population. *Genetica* 102:127–144.
- Gill, D. E. 1974. Intrinsic rate of increase, saturation density, and competitive ability. ii. the
465 evolution of competitive ability. *American Naturalist* 108:103–116.
- Grime, J. P. 1974. Vegetation classification by reference to strategies. *Nature* 250:26–31.
- . 1977. Evidence for the existence of three primary strategies in plants and its relevance to
468 ecological and evolutionary theory. *American Naturalist* 111:1169–1194.
- . 1988. *Plant Evolutionary Biology*, chap. The C-S-R model of primary plant strategies —
origins, implications and tests, pages 371–393. Springer Netherlands, Dordrecht.
- 471 Haldane, J. B. S. 1927. A mathematical theory of natural and artificial selection, part v: selection
and mutation 23:838–844.
- Hendriks, A. J., J. L. Maas-Diepeveen, E. H. Heugens, and N. M. van Straalen. 2005. Meta-analysis
474 of intrinsic rates of increase and carrying capacity of populations affected by toxic and other
stressors. *Environmental Toxicology and Chemistry* 24:2267–2277.

Joshi, A., N. Prasad, and M. Shakarad. 2001. K-selection, α -selection, effectiveness, and tolerance
477 in competition: density-dependent selection revisited. *Journal of Genetics* 80:63–75.

Kuno, E. 1991. Some strange properties of the logistic equation defined with r and K : Inherent
defects or artifacts? *Researches on Population Ecology* 33:33–39.

480 Levins, R., and D. Culver. 1971. Regional coexistence of species and competition between rare
species. *Proceedings of the National Academy of Sciences* 68:1246–1248.

Luckinbill, L. S. 1978. r and k selection in experimental populations of *Escherichia coli*. *Science*
483 (New York, NY) 202:1201–1203.

———. 1979. Selection and the r/k continuum in experimental populations of protozoa. *American
Naturalist* pages 427–437.

486 MacArthur, R. H. 1962. Some generalized theorems of natural selection. *Proceedings of the
National Academy of Sciences* 48:1893–1897.

MacArthur, R. H., and E. O. Wilson. 1967. *Theory of Island Biogeography*. Princeton University
489 Press.

Messer, P. W., S. P. Ellner, and N. G. Hairston. 2016. Can population genetics adapt to rapid
evolution? *Trends in Genetics* 32:408–418.

492 Metz, J. A., R. M. Nisbet, and S. A. Geritz. 1992. How should we define ‘fitness’ for general
ecological scenarios? *Trends in Ecology & Evolution* 7:198–202.

Pianka, E. R. 1970. On r - and K -Selection. *The American Naturalist* 104:592–597.

495 ———. 1972. r and K selection or b and d selection? *The American Naturalist* 106:581–588.

Ram, Y., E. Dellus-Gur, M. Bibi, U. Obolski, J. Berman, and L. Hadany. 2016. Predicting microbial
relative growth in a mixed culture from growth curve data. *bioRxiv* 10.1101/022640 .

- 498 Reznick, D., M. J. Bryant, and F. Bashey. 2002. r- and k-selection revisited: The role of population
regulation in life-history evolution. *Ecology* 83:1509–1520.
- Sale, P. F. 1977. Maintenance of high diversity in coral reef fish communities. *The American*
501 *Naturalist* 111:337–359.
- Shmida, A., and S. Ellner. 1984. Coexistence of plant species with similar niches. *Vegetatio*
58:29–55.
- 504 Southwood, T. R. E. 1977. Habitat, the templet for ecological strategies? *Journal of Animal*
Ecology 46:337–365.
- Stearns, S. C. 1977. The evolution of life history traits: A critique of the theory and a review of
507 the data. *Annual Review of Ecology and Systematics* 8:145–171.
- Svardal, H., C. Rueffler, and J. Hermisson. 2015. A general condition for adaptive genetic poly-
morphism in temporally and spatially heterogeneous environments. *Theoretical Population*
510 *Biology* 99:76 – 97.
- Taylor, D. R., L. W. Aarssen, and C. Loehle. 1990. On the relationship between r/k selection and
environmental carrying capacity: a new habitat templet for plant life history strategies. *Oikos*
513 pages 239–250.
- Tilman, D. 1982. Resource competition and community structure. Princeton University Press.
- . 1994. Competition and biodiversity in spatially structured habitats. *Ecology* 75:2–16.
- 516 ———. 2007. Resource competition and plant traits: a response to craine et al. 2005. *Journal of*
Ecology 95:231–234.
- Uecker, H., and J. Hermisson. 2011. On the fixation process of a beneficial mutation in a variable
519 environment. *Genetics* 188:915–930.

Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let the concept of trait be functional! *Oikos* 116:882–892.

522 Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and soil* 199:213–227.

Winemiller, K. O., D. B. Fitzgerald, L. M. Bower, and E. R. Pianka. 2015. Functional traits, convergent evolution, and periodic tables of niches. *Ecology Letters* 18:737–751.

Winemiller, K. O., and K. A. Rose. 1992. Patterns of life-history diversification in north american fishes: implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences* 49:2196–2218.

Yampolsky, L. Y., and A. Stoltzfus. 2001. Bias in the introduction of variation as an orienting factor in evolution. *Evolution & Development* 3:73–83.

531 Yi, X., and A. M. Dean. 2013. Bounded population sizes, fluctuating selection and the tempo and mode of coexistence. *Proceedings of the National Academy of Sciences* 110:16945–16950.

Appendix A: Poisson approximation

534 For simplicity of presentation, we have assumed a Poisson distribution for the x_i as our model of dispersal. Strictly speaking, the total number of i propagules $\sum x_i$ (summed over unoccupied territories) is then no longer a constant m_i , but fluctuates between generations for a given mean
537 m_i , which is more biologically realistic. Nevertheless, since we do not consider the random fluctuations in type abundances here, and for ease of comparison with the classic lottery model, we ignore the fluctuations in m_i . Instead we focus, on Poisson fluctuations in propagule composition
540 in each territory.

In the exact model of random dispersal, the counts of a genotype’s propagules across unoccupied territories follows a multinomial distribution with dimension U , total number of trials

543 equal to m_i , and equal probabilities $1/U$ for a propagule to land in a given territory. Thus, the
 x_i in different territories are not independent random variables. However, for sufficiently large
 U and m_i , this multinomial distribution for the x_i across territories is closely approximated by
546 a product of independent Poisson distributions for each territory, each with rate parameter l_i
(Arenbaev, 1977, Theorem 1). Since we are ignoring finite population size effects, we effectively
have $T \rightarrow \infty$, in which case U can be only be small enough to violate the Poisson approximation
549 if there is vanishing population turnover, and then the dispersal distribution is irrelevant any-
way. Likewise, in ignoring stochastic finite population size for the n_i , we have effectively already
assumed that m_i is large enough to justify the Poisson approximation (the error scales as $1/\sqrt{m_i}$;
552 Arenbaev 1977).

Appendix B: Derivation of growth equation

We separate the right hand side of Eq. (2) into three components $\Delta_+ n_i = \Delta_u n_i + \Delta_r n_i + \Delta_a n_i$ which
555 vary in relative magnitude depending on the propagule densities l_i . Following the notation in
the main text, the Poisson distributions for the x_i (or some subset of the x_i) will be denoted p ,
and we use P as a general shorthand for the probability of particular outcomes.

558 Growth without competition

The first component, $\Delta_u n_i$, accounts for territories where only one focal propagule is present
 $x_i = 1$ and $x_j = 0$ for $j \neq i$ (u stands for “uncontested”). The proportion of territories where this
561 occurs is $l_i e^{-L}$, and so

$$\Delta_u n_i = U l_i e^{-L} = m_i e^{-L}. \quad (7)$$

Competition when rare

The second component, $\Delta_r n_i$, accounts for territories where a single focal propagule is present
564 along with at least one non-focal propagule (r stands for “rare”) i.e. $x_i = 1$ and $X_i \geq 1$ where

$X_i = \sum_{j \neq i} x_j$ is the number of nonfocal propagules. The number of territories where this occurs is $Up_i(1)P(X_i \geq 1) = b_i n_i e^{-l_i}(1 - e^{-(L-l_i)})$. Thus

$$\Delta_r n_i = m_i e^{-l_i}(1 - e^{-(L-l_i)}) \left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}}, \quad (8)$$

567 where $\langle \rangle_{\tilde{p}}$ denotes the expectation with respect to \tilde{p} , and \tilde{p} is the probability distribution of nonfocal propagule abundances x_j *after* dispersal, in those territories where exactly one focal propagule, and at least one non-focal propagule, landed.

570 Our “mean field” approximation is to replace x_j with its mean in the last term in Eq. (8),

$$\left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}} \approx \frac{c_i}{c_i + \sum_{j \neq i} c_j \langle x_j \rangle_{\tilde{p}}}. \quad (9)$$

Below we justify this replacement by arguing that the standard deviation $\sigma_{\tilde{p}}(\sum_{j \neq i} c_j x_j)$ (with respect to \tilde{p}), is much smaller than $\langle \sum_{j \neq i} c_j x_j \rangle_{\tilde{p}}$.

We first calculate $\langle x_j \rangle_{\tilde{p}}$. Let $X = \sum_j x_j$ denote the total number of propagules in a territory and $\mathbf{x}_i = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_G)$ denote the vector of non-focal abundances, so that $p(\mathbf{x}_i) = p_1(x_1) \dots p_{i-1}(x_{i-1}) p_{i+1}(x_{i+1}) \dots p_G(x_G)$. Then, \tilde{p} can be written as

$$\begin{aligned} \tilde{p}(\mathbf{x}_i) &= p(\mathbf{x}_i | X \geq 2, x_i = 1) \\ &= \frac{P(\mathbf{x}_i, X \geq 2 | x_i = 1)}{P(X \geq 2)} \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{X=2}^{\infty} P(X) p(\mathbf{x}_i | X_i = X - 1), \end{aligned} \quad (10)$$

and so

$$\begin{aligned} \langle x_j \rangle_{\tilde{p}} &= \sum_{\mathbf{x}_i} \tilde{p}(\mathbf{x}_i) x_j \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{X=2}^{\infty} P(X) \sum_{\mathbf{x}_i} p(\mathbf{x}_i | X_i = X - 1) x_j. \end{aligned} \quad (11)$$

The inner sum over \mathbf{x}_i is the mean number of propagules of a given nonfocal type j that will be found in a territory which received $X - 1$ nonfocal propagules in total, which is equal to $\frac{l_j}{L-l_i}(X-1)$. Thus,

$$\begin{aligned}\langle x_j \rangle_{\tilde{p}} &= \frac{l_j}{1 - (1+L)e^{-L}} \frac{1}{L-l_i} \sum_{k=2}^{\infty} P(X)(X-1) \\ &= \frac{l_j}{1 - (1+L)e^{-L}} \frac{L-1+e^{-L}}{L-l_i},\end{aligned}\tag{12}$$

573 where the last line follows from $\sum_{X=2}^{\infty} P(X)(X-1) = \sum_{X=1}^{\infty} P(X)(X-1) = \sum_{X=1}^{\infty} P(X)X - \sum_{X=1}^{\infty} P(X)$.

The exact analysis of the fluctuations in $\sum_{j \neq i} c_j x_j$ is complicated because the x_j are not independent with respect to \tilde{p} . These fluctuations are part of the “drift” in type abundances which we leave for future work. Here we use the following approximation to give some insight into the magnitude of these fluctuations and also the nature of the correlations between the x_j . We replace \tilde{p} with \tilde{q} , defined as the \mathbf{x}_i Poisson dispersal probabilities conditional on $X_i \geq 1$ (which are independent). The distinction between \tilde{p} with \tilde{q} will be discussed further below. The \tilde{q} approximation gives $\langle x_j \rangle_{\tilde{q}} = \langle x_j \rangle_p / C = l_j / C$,

$$\begin{aligned}\sigma_{\tilde{q}}^2(x_j) &= \langle x_j^2 \rangle_{\tilde{q}} - \langle x_j \rangle_{\tilde{q}}^2 \\ &= \frac{1}{C} \langle x_j^2 \rangle_p - \frac{l_j^2}{C^2} \\ &= \frac{1}{C} (l_j^2 + l_j) - \frac{l_j^2}{C^2} \\ &= \frac{l_j^2}{C} \left(1 - \frac{1}{C}\right) + \frac{l_j}{C},\end{aligned}\tag{13}$$

and

$$\begin{aligned}
\sigma_{\tilde{q}}(x_j, x_k) &= \langle x_j x_k \rangle_{\tilde{q}} - \langle x_j \rangle_{\tilde{q}} \langle x_k \rangle_{\tilde{q}} \\
&= \frac{1}{C} \langle x_j x_k \rangle_p - \frac{l_j l_k}{C^2} \\
&= \frac{l_j l_k}{C} \left(1 - \frac{1}{C} \right),
\end{aligned} \tag{14}$$

where $C = 1 - e^{-(L-l_i)}$ and $j \neq k$.

576 The exact distribution \tilde{p} assumes that exactly one of the propagules present in a given site after dispersal belongs to the focal type, whereas \tilde{q} assumes that there is a focal propagule present before non-focal dispersal commences. As a result, \tilde{q} predicts that the mean propagule
579 density is greater than L (in sites with only one focal propagule is present) when the focal type is rare and the propagule density is high. This is erroneous, because the mean number of propagules in every site is L by definition. Specifically, if $L - l_i \approx L \gg 1$, then the mean
582 propagule density predicted by \tilde{q} is approximately $L + 1$. The discrepancy causes rare invaders to have an intrinsic rarity disadvantage (territorial contests under \tilde{q} are more intense than they should be). In contrast, Eq. (12) correctly predicts that there are on average $\sum_{j \neq i} \langle x_j \rangle_{\tilde{p}} \approx L - 1$
585 nonfocal propagules because \tilde{p} accounts for potentially large negative covariances between the x_j “after dispersal”. By neglecting the latter covariances, \tilde{q} overestimates the fluctuations in $\sum_{j \neq i} c_j x_j$; thus \tilde{q} gives an upper bound on the fluctuations. The discrepancy between \tilde{q} and \tilde{p} will be largest
588 when L is of order 1 or smaller, because then the propagule assumed to already be present under \tilde{q} is comparable to, or greater than, the entire propagaule density.

Decomposing the variance in $\sum_{j \neq i} c_j x_j$,

$$\sigma_{\tilde{q}}^2(\sum_{j \neq i} c_j x_j) = \sum_{j \neq i} \left[c_j^2 \sigma_{\tilde{q}}^2(x_j) + 2 \sum_{k > j, k \neq i} c_j c_k \sigma_{\tilde{q}}(x_j, x_k) \right], \tag{15}$$

591 and using the fact that $\sigma_{\tilde{q}}(x_j, x_k)$ and the first term in Eq. (13) are negative because $C < 1$, we

obtain an upper bound on the relative fluctuations in $\sum_{j \neq i} c_j x_j$,

$$\frac{\sigma(\sum_{j \neq i} c_j x_j)}{\langle \sum_{j \neq i} c_j x_j \rangle} = C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j + (1 - 1/C) \left(\sum_{j \neq i} c_j l_j \right)^2 \right)^{1/2}}{\sum_{j \neq i} c_j l_j} < C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j \right)^{1/2}}{\sum_{j \neq i} c_j l_j}. \quad (16)$$

Suppose that the c_j are all of similar magnitude (their ratios are of order one). Then Eq. (16) is $\ll 1$ for the case when $L - l_i \ll 1$ (due to the factor of $C^{1/2}$), and also for the case when at least some of the nonfocal propagule densities are large $l_j \gg 1$ (since it is then of order $1/\sqrt{L - l_i}$). The worst case scenario occurs when $L - l_i$ is of order one. Then Eq. (16) gives a relative error of approximately 50%, which from our earlier discussion we know to be a substantial overestimate when L is of order 1. Our numerical results (Fig. 5) confirm that the relative errors are indeed small.

However, the relative fluctuations in $\sum_{j \neq i} c_j x_j$ can be large if some of the c_j are much larger than the others. Specifically, in the presence of a rare, extremely strong competitor ($c_j l_j \gg c_{j'} l_{j'}$ for all other nonfocal genotypes j' , and $l_j \ll 1$), then the RHS of Eq. (16) can be large and we cannot make the replacement Eq. (9).

Substituting Eqs. (9) and (12) into Eq. (8), we obtain

$$\Delta_r n_i \approx m_i R_i \frac{c_i}{C}, \quad (17)$$

where R_i is defined in Eq. (5).

606 Competition when abundant

The final contribution, $\Delta_a n_i$, accounts for territories where two or more focal propagules are present (a stands for “abundant”). Similarly to Eq. (8), we have

$$\Delta_a n_i = U(1 - (1 + l_i)e^{l_i}) \left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \quad (18)$$

609 where \hat{p} is the probability distribution of both focal and nonfocal propagaule abundances *after* dispersal in those territories where at least two focal propagules landed.

Again, we argue that the relative fluctuations in $\sum c_j x_j$ are much smaller than 1 (with respect
612 to \hat{p}), so that,

$$\left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \approx \frac{c_i \langle x_i \rangle_{\hat{p}}}{\sum_j c_j \langle x_j \rangle_{\hat{p}}}. \quad (19)$$

Following a similar procedure as for $\Delta_r n_i$, where the vector of propagule abundances is denoted \mathbf{x} , the mean focal genotype abundance is,

$$\begin{aligned} \langle x_i \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_i p(\mathbf{x} | x_i \geq 2) \\ &= \sum_{x_i} x_i p(x_i | x_i \geq 2) \\ &= \frac{1}{1 - (1 + l_i)e^{-l_i}} \sum_{x_i \geq 2} p(x_i) x_i \\ &= l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}}. \end{aligned} \quad (20)$$

For nonfocal genotypes $j \neq i$, we have

$$\begin{aligned} \langle x_j \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2) \\ &= \sum_X P(X | x_i \geq 2) \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2, X) \\ &= \sum_X P(X | x_i \geq 2) \sum_{x_i} p(x_i | x_i \geq 2, X) \sum_{\mathbf{x}_i} x_j p(\mathbf{x}_i | X_i = X - x_i) \\ &= \sum_X P(X | x_i \geq 2) \sum_{x_i} p(x_i | x_i \geq 2, X) \frac{l_j (X - x_i)}{L - l_i} \\ &= \frac{l_j}{L - l_i} \left[\sum_X P(X | x_i \geq 2) X - \sum_{x_i} p(x_i | x_i \geq 2) x_i \right] \\ &= \frac{l_j}{L - l_i} \left(L \frac{1 - e^{-L}}{1 - (1 + L)e^{-L}} - l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}} \right). \end{aligned} \quad (21)$$

To calculate the relative fluctuations in $\sum_{j \neq i} c_j x_j$, we use a similar approximation as for $\Delta_r n_i$: \hat{p}

is approximated by \hat{q} , defined as the \mathbf{x} dispersal probabilities in a territory conditional on $x_i > 2$
 615 (that is, treating the x_j as independent). All covariances between nonfocal genotypes are now zero,
 so that $\sigma_{\hat{q}}^2(\sum c_j x_j) = \sum c_j^2 \sigma_{\hat{q}}^2(x_j)$, where $\sigma_{\hat{q}}^2(x_j) = l_j$ for $j \neq i$, and

$$\sigma_{\hat{q}}^2(x_i) = \frac{l_i}{D} \left(l_i + 1 - e^{-l_i} - \frac{l_i}{D} (1 - e^{-l_i})^2 \right), \quad (22)$$

where $D = 1 - (1 + l_i)e^{-l_i}$, and

$$\frac{\sigma_{\hat{q}}(\sum c_j x_j)}{\langle \sum c_j x_j \rangle} = \frac{\left(\sum_{j \neq i} c_j^2 l_j + c_i^2 \sigma_{\hat{q}}^2(x_i) \right)^{1/2}}{\sum_{j \neq i} c_j l_j + c_i l_i (1 - e^{-l_i}) / D}. \quad (23)$$

618 Similarly to Eq. (16), the RHS of Eq. (23) is $\ll 1$ for the case that $L \ll 1$ (due to a factor of
 $D^{1/2}$), and also for the case when at least some of the propagule densities (focal or nonfocal) are
 large — provided that c_i and the c_j are all of similar magnitude. Again, the worst case scenario
 621 occurs when l_i and $L - l_i$ are of order 1, in which case Eq. (23) is around 35%, which is again
 where the \hat{q} approximation produces the biggest overestimate of the fluctuations in \mathbf{x} . Similarly
 to Eq. (16), the RHS of (23) will not be $\ll 1$ in the presence of a rare, extremely strong competitor.

624 Combining Eqs. (18) and (19), we obtain

$$\Delta_a n_i = m_i A_i \frac{c_i}{\bar{c}}, \quad (24)$$

where A_i is defined in Eq. (6).

Comparison with simulations

627 Fig. 5 shows that Eq. (4) and its components closely approximate our density-dependent lottery
 model over a wide range of propagule densities (the latter is evaluated by direct simulations of
 uniform random dispersal and lottery competition). Two genotypes are present, one of which is
 630 at low frequency. The growth of the low-frequency genotype relies crucially on the low-density
 competition term $R_i c_i / \bar{c}$, and also to a lesser extent on the high density competition term $A_i c_i / \bar{c}$




Figure 5: The change in genotype abundances in a density dependent lottery model is closely approximated by Eq. (4). $\Delta_+ n_i/m_i$ from Eq. (4) (and its separate components) are shown, along with direct simulations of random dispersal and lottery competition over one iteration over a range of propagule densities (U is varied between 5×10^3 and 10^6 with $m_1 = 10^4$ and $m_2 = 9 \times 10^4$). Two genotypes are present. (a) and (b) show the low-frequency genotype with c -advantage ($c_1 = 1.5$), (c) and (d) show the high-frequency predominant genotype ($c_2 = 1$). Simulation points are almost invisible in (c) and (d) due to near exact agreement with Eq. (4). Dashed lines in (a) and (c) show the breakdown of the classic lottery model.

if l_1 is large enough (Fig. 5b). On the other hand, $R_i c_i/\bar{c}$ is negligible for the high-frequency
 633 genotype, which depends instead on high density territorial victories (Fig. 5d). Fig. 3 also shows
 the breakdown of the classic lottery model at low propagule densities.

Appendix C: Mutant invasion under different environments

636 In this Appendix we evaluate the invasion of novel mutants in a population with a single resident
 type i (such that $N = n_i$), which is in equilibrium.

For a single type i in equilibrium ($\Delta n_i = 0$), we have $R_i = 0$, $\bar{c} = c_i$, $A_i = (1 - (1 + L)e^{-L})/L$,

639 and Eq. (4) becomes

$$b_i(1 - e^{-L})/L - d_i = 0. \quad (25)$$

(Alternatively, Eq. (25) can be deduced directly from Eq. (2)). This implies $L \approx b_i/d_i$ when $b_i/d_i \gg 1$ and $L \ll 1$ when $b_i/d_i \approx 1$. Now suppose that a novel mutant j , which is initially rare,

642 appears in the population. Then $A_j/R_j \ll 0$, $l_j \approx 0$ and $\bar{c} \approx c_i$, and so, from Eq. (4), the mutant lineage's fitness is

$$\Delta n_j/n_j \approx b_j \left(e^{-L} + R_j \frac{c_j}{c_i} \right) - d_j \quad (26)$$

where $R_j \approx (1 - e^{-L}) / \left(\frac{c_j}{c_i} + \frac{L-1+e^{-L}}{1-(1+L)e^{-L}} \right)$ since $l_j \ll 1$.

645 We now consider mutant invasion under the different environments from section "Primary strategies and Grime's triangle". In our representation of Grime's stressful environment, $b \ll 1$, $d \approx b$ and $L \ll 1$, and so Eq. (26) becomes $\Delta n_j/n_j \approx b_j - d_j$. Mutations which improve b or d by
648 the same fraction ϵ , such that $b_j = (1 + \epsilon)b_i$ or $d_j = (1 - \epsilon)d_i$, yield identical fitness benefits.

In our representation of Grime's ideal environment, $d \ll 1$, $b_i/d_i \gg 1$ and $L \gg 1$, and so Eq. (26) becomes

$$\Delta n_j/n_j \approx \frac{b_j}{c_j/c_i + L - 1} \frac{c_j}{c_i} - d_j \approx \frac{b_j}{L} \frac{c_j}{c_i} - d_j, \quad (27)$$

651 where the last approximation follows from the fact that c_j will be of similar magnitude to c_i (ignoring mutations with very large effect sizes; incidentally, Eq. (1) yields the same expression by assuming $L \rightarrow \infty$ even though this assumption is unrealistic for a rare mutant). Similarly
654 to Grime's stressful environment, improving b , c , or d by a fraction ϵ yields identical fitness benefits, even though they may have very different magnitudes, because the ancestral resident type's density-regulated birth rate $b_i c_i / L$ exactly balances its mortality d_i in equilibrium. Biases
657 in the direction of evolution of b , c or d must again follow from biases in mutation availability or trade-offs/pleiotropy. The same argument applies for the high- L interpretation of the stress regime.

660 To represent Grime's disturbed environment, $\Delta_+ n_i$ is replaced with $(1 - d_i)\Delta_+ n_i$, and a single type reaches equilibrium when $L/(1 - e^{-L}) = (1 - d_i)b_i/d_i$ (population persistence now requires $(1 - d_i)b_i > d_i$ rather than $b_i > d_i$). We assumed $d_i \approx 1$ and $(1 - d_i)b_i \approx d_i$, so that $L \ll 1$. A
663 mutant's invasion fitness is then given by

$$\Delta n_j/n_j \approx (1 - d_j)b_j - d_j. \quad (28)$$

Another distinct feature of disturbed environments is that the bulk of mortality is environmental and unavoidable. That is, $d_i = d_{ie} + d_{ig}$, where the environmental part d_{ie} does not evolve, and
666 the genetic part d_{ig} is affected proportionally by mutations $d_{ig} = (1 - \epsilon)d_{ig}$.

The effects of improving b and d are now no longer identical for given values of d_{ie} and d_{ig} . A b -mutation $b_j = (1 + \epsilon)b_i$ gives $\Delta n_j/n_j = \epsilon(1 - d_i)b_i \approx \epsilon d_i \approx \epsilon$, whereas a d -mutation
669 $d_{ig} = (1 - \epsilon)d_{ig}$ gives $\Delta n_j/n_j = \epsilon d_{ig}/(1 - d_i)$. Since overall mortality is high $d_i \approx 1$, d_g can be small compared to d_e and yet still make a greater contribution to invasion fitness than b . The case where d_{ig} is so small that $d_{ig} \ll 1 - d_i$ (i.e. d_{ig} is an order of magnitude smaller than $1 - d_i$ which
672 itself an order of magnitude smaller than d_{ie}) effectively represents a hard constraint that d cannot evolve. We do not deny this possibility, but this bring us back to the issue of mutations affect traits (the first two factors controlling the direction of evolution in section "Primary strategies and Grime's triangle"), which we do not address in this manuscript.
675

Thus, assuming that d can evolve in this sense, d_{ig} is either comparable to or greater than $1 - d_i$. Yet d_{ig} cannot be appreciably greater than $1 - d_i$ for long, because selection will then
678 strongly favor reductions in d_g over increases in b , thereby bringing d_{ig} closer to $1 - d_i$. Thus, we expect selection on b and d to be of comparable strength in disturbed environments as well.

Appendix D: Coexistence in a stable environment

681 To determine whether coexistence is possible in a constant environment, we check for "mutual invasion", that is, we check that j will invade an i -dominated population, but i will also invade a

j -dominated population.

684 We consider the case of coexistence between a b specialist i and a c specialist j ($b_i > b_j$, $c_j > c_i$
and $d_i = d_j$). Suppose that b_i is so large that $L \gg 1$ when i is dominant, and b_j is so small that $L \ll$
1 when j is dominant. Then, when j is dominant, we have $\Delta n_i/n_i = b_i - d_i = b_i - d_j = b_i - b_j > 0$.
687 When i is dominant, Eq. (27) applies, where Eq. (25) implies $d_j = d_i = b_i(1 - e^{-L})/L \approx b_i/L$,
and so

$$\Delta n_j/n_j \approx \frac{b_j}{L} \frac{c_j}{c_i} - \frac{b_i}{L}. \quad (29)$$

Therefore, coexistence occurs if c_j/c_i is sufficiently large. The analogous argument for d and c
690 specialists ($d_i < d_j$ with $L \gg 1$ when i dominates, $L \ll 1$ when j dominates, and $b_i = b_j$) gives
 $\Delta n_j/n_j \approx d_i \frac{c_j}{c_i} - d_j$, which again implies coexistence if c_j/c_i is sufficiently large.

For b and d specialists ($c_i = c_j$), we have $\Delta n_j/n_j \approx b_j d_i / b_i - d_j$ when i dominates and $\Delta n_i/n_i \approx$
693 $b_i d_j / b_j - d_i$ when j dominates. Thus, either i or j grows when rare, but not both, and stable
coexistence is not possible in a constant environment.