

A preliminary rK derivation

J Bertram and J Masel

October 24, 2016

“...the concept of fitness is probably too complex to allow of a useful mathematical development. Since it enters fundamentally into many population genetics considerations, it is remarkable how little attention has been paid to it.” — Warren J. Ewens, *Mathematical Population Genetics I*, 2004

1 Introduction

Evolutionary models differ greatly in their treatment of fitness. In models of genetic evolution, genotypes are typically assigned constant (or frequency-dependent) selection coefficients describing the change in their relative frequencies over time due to differences in viability. This considerably simplifies the mathematics of selection, facilitating greater genetic realism, and can be justified over sufficiently short time intervals [1, p. 276]. However, selection can have very different effects when operating on different types of traits, and evolutionary changes in one population can lead to complicated ecological responses.

By contrast, models of phenotypic trait evolution represent the change in phenotypic abundances over time using absolute fitness functions which describe how those traits affect survival and reproduction in particular ecological scenarios. This approach is powerful enough to model eco-evolutionary feedbacks between co-evolving traits, but is generally problem-specific and restricted to only a few traits at a time.

Far less work has been done to model fitness in more general terms than particular traits or ecological scenarios, while still capturing key distinctions between different forms of selection. Perhaps this is not surprising given that fitness is such a complex quantity, dependent on all of a phenotype's

functional traits [2] as well as its biotic and abiotic environment. In most cases, a detailed, trait-based, predictive model of fitness would be enormously complicated and have narrow applicability. It is therefore easy to doubt the feasibility of a simplified, general mathematical treatment of fitness [1, p. 276]. Even MacArthur’s famous r/K selection scheme is now almost exclusively known as a framework for understanding life-history traits, and judged on its failure in that role [3, 4, 5, 6]. In spite of the r/K scheme’s original purpose as an extension of the existing population-genetic treatment of selection to account for population density [7], comparatively few attempts have been made to develop it further as a mathematical analysis of the major different forms of selection.

Nevertheless, there are strong indications there are broader principles governing the operation of selection. In many groups of organisms (including corals [8], insects [9], fishes [10] and plants [11]), species can be partitioned into a small number of distinct trait clusters corresponding to fundamentally distinct “primary strategies” [12]. The most famous example is Grime’s plant trait classification scheme [13, 14, 11]. Grime considered two broad determinants of population density: stress (persistent hardship e.g. due to resource scarcity, unfavorable temperatures or toxins) and disturbance (intermittent destruction of vegetation e.g. due to trampling, herbivory, pathogens, extreme weather or fire). The extremes of these two factors define three primary strategies denoted by C/S/R respectively: competitors “C” excel in low stress, low disturbance environments; stress tolerators “S” excel in high stress, low disturbance environments; and ruderals “R” excel in low stress, high disturbance environments. Survival is not possible in high-stress, high-disturbance environments. Grime showed that measures of C, S and R across a wide range of plant species are anti-correlated, so that strong C-strategists are weak S and R strategists, and so on. Thus, plant species can be classified on a triangular C/S/R ternary plot [13]. Trait classification schemes for other organisms closely parallel Grime’s scheme [12].

Trait classification schemes show empirically that, beneath the complicated details of trait variation, even among closely-related species, fitness is predominantly determined by a few key factors such as intrinsic reproductive rate or stress-tolerance. However, while trait classification schemes are firmly grounded in trait data, they are verbal and descriptive rather than mathematical, a recognized hinderance to their broader applicability (e.g. [15]).

The aim of this paper is explore the interplay between some major di-

mensions of fitness in a simplified, spatially-homogeneous model of genotype growth, dispersal and competition. Building on the earlier r/K and C/S/R schemes, a central question is how fitness depends on the interaction between population density, intrinsic birth/death rates and competitive ability.

We broadly follow the spirit of MacArthur’s r/K selection scheme in that our model is intended to account for fundamentally different forms of selection without getting entangled in the intricacies of particular ecological scenarios. However, rather than building directly on MacArthur’s formalism and its later extensions [?] (e.g. using Lotka-Volterra equations), our model is devised primarily with Grime’s C/S/R scheme in mind, and represents a quantitative formalization of how C/S/R manifests at the level of genotype evolution (as opposed to divergence between species). This choice is motivated in part by the substantial empirical support for the C/S/R scheme, and in part by the failings of the r/K low/high density dichotomy — many growth ability traits will confer advantages at both low and high densities, in which case r- and K- selection will effectively coincide (more details in the Discussion).

In section

2 Model

We assume that each individual in a population requires its own territory to survive and reproduce (a site-occupancy model). All territories are identical, and the total number of territories is K . Time t advances in discrete steps, each step representing one generation. In generation t , the number of mature individuals (henceforth called “adults”) of the i ’th genotype is $n_i(t)$, the total number of adults is $N(t) = \sum_i n_i(t)$, and the number of unoccupied territories is $U(t) = K - N(t)$.

Each generation, adults produce m_i new offspring (henceforth called “propagules”) which disperse at random over the U unoccupied territories (there is no dispersal limitation). We ignore propagules that land on occupied territories — these are assumed to have no chance of survival and are not included in m_i . For simplicity, we assume $m_i = b_i n_i$, where b_i is a constant, genotype-specific birth rate.

The number of individuals of the i ’th genotype landing in any particular territory is denoted x_i . Random dispersal implies that in the limit $K \rightarrow \infty$, with n_i/K held fixed, x_i is Poisson distributed with mean territorial

propagule density $l_i = m_i/U$. Although K is finite in our model, we assume that K and the n_i are large enough that x_i is Poisson-distributed to a good approximation (details in Appendix A). This dispersal Poisson distribution is denoted $p_i(x_i)$. Note that the large n_i , large K approximation places no restrictions on our densities n_i/K , but it does preclude consideration of demographic stochasticity when n_i itself is very small (this will be discussed further in Section 3.2).

When multiple propagules land on the same territory, they compete to secure the territory as they develop. This territorial contest is modeled as a weighted lottery: the probability that genotype i wins a given territory by the next generation is $c_i x_i / \sum_j c_j x_j$ where c_i is a constant representing relative competitive ability.

The increase in n_i over one generation due to territorial acquisition, $\Delta_+ n_i$, is the sum of genotype i 's victories over all U unoccupied territories. Since $p_1(x_1) \dots p_G(x_G)$ is equal to the proportion of unoccupied territories with x_1, \dots, x_G of the respective propagules (again, we assume that K is large enough that fluctuations in this proportion are negligible), this sum can be replaced by an expectation over the p_i . This gives

$$\Delta_+ n_i(t) = U(t) \sum_{x_1, \dots, x_G} \frac{c_i x_i}{\sum_j c_j x_j} p_1(x_1) \dots p_G(x_G). \quad (1)$$

In addition to propagule birth and competition, occupied territories become unoccupied due to mortality. For the vast majority of this manuscript we assume that mortality only occurs in adults, and at a constant, genotype-specific per-capita rate d_i , so that the overall change in genotype abundances is

$$\Delta n_i(t) = \Delta_+ n_i(t) - d_i n_i(t). \quad (2)$$

We will introduce a different mortality model when we consider the effects of disturbances (Section 3.3), which will also affect competing juveniles.

Note that the competitive ability coefficients c_i represent a strictly relative aspect of fitness in the sense that they only influence population size N indirectly by changing genotype frequencies; that may in turn change the population mean birth and death rates. This can be seen by summing Eq. (2) over genotypes to get the change in population size N ,

$$\Delta N = U(1 - e^{-L}) - \sum_i d_i n_i, \quad (3)$$

which is independent of c_i (here $L = \sum_j l_j$ is the overall propagule density).

3 Results

3.1 Mean Field Approximation

Eq. (2) gives little intuition about the dynamics of density-dependent lottery competition, since (1) involves an expectation over the random dispersal distributions p_i , which depend on how the n_i change over time. We now evaluate this expectation using a “mean field” approximation; the intuition behind this approximation is as follows.

If the unoccupied territories are saturated with propagules from every genotype ($l_i \gg 1$ for all genotypes), the fluctuations in the x_i are small compared to their means l_i (since the x_i are Poisson distributed), and so the composition of propagules in a territory will only rarely differ appreciably from the mean composition l_1, l_2, \dots, l_G . Consequently, we can replace x_i with l_i in Eq. (1). This gives the classic lottery model [21],

$$\Delta_+ n_i(t) = U(t) \frac{c_i m_i}{\sum_j c_j m_j} = b_i n_i \frac{1}{L} \frac{c_i}{\bar{c}}, \quad (4)$$

where $\bar{c} = \sum_j c_j m_j / M$ is the mean propagule competitive ability for a randomly selected propagule ($M = \sum_j m_j$ is the total number of propagules).

However, in general the l_i are not all large, and the x_i cannot simply be replaced by their means in Eq. (1). Indeed, Eq. (4) is nonsensical if l_i is sufficiently small: genotype i can win at most m_i territories, yet Eq. (4) demands a fraction $c_i m_i / \sum_j c_j m_j$ of the unoccupied territories U , no matter how large U is. The source of this pathological behavior when $l_i \ll 1$ is that $x_i = 1$ in the few territories where i propagules do land, and so i ’s growth comes entirely from territories which deviate appreciably from the mean.

Our mean field approximation is similar to the high- l_i approximation leading to Eq. (4) in that we replace the x_i with appropriate mean values. The key distinction is that territories with a single propagule from the focal genotype, which are critical at low densities, are handled separately. In place of the requirement of $l_i \gg 1$ for all i , our approximation only requires that there are no large discrepancies in competitive ability (discussed further below). We obtain (details in Appendix B)

$$\Delta_+ n_i(t) \approx b_i n_i \left[e^{-L} + (R_i + A_i) \frac{c_i}{\bar{c}} \right], \quad (5)$$

where

$$R_i = \frac{\bar{c}e^{-l_i}(1 - e^{-(L-l_i)})}{c_i + \frac{L-1+e^{-L}}{1-(1+L)e^{-L}} \frac{\bar{c}L-c_i l_i}{L-l_i}}, \quad (6)$$

and

$$A_i = \frac{\bar{c}(1 - e^{-l_i})}{c_i l_i \frac{1-e^{-l_i}}{1-(1+l_i)e^{-l_i}} + \sum_{j \neq i} \frac{c_j l_j}{L-l_j} \left(L \frac{1-e^{-L}}{1-(1+L)e^{-L}} - l_j \frac{1-e^{-l_j}}{1-(1+l_j)e^{-l_j}} \right)}. \quad (7)$$

Comparing Eq. (5) to Eq. (4), the classic lottery per-propagule success rate $c_i/\bar{c}L$ has been replaced by three separate terms. The first, e^{-L} , accounts for propagules which land alone on unoccupied territories; these territories are won without contest. The second term, $R_i c_i/\bar{c}$ represents competitive victories when the i genotype is a rare invader in a high density population: from Eq. (6), $R_i \rightarrow 0$ when the i genotype is abundant ($l_i \gg 1$), or other genotypes are collectively rare ($L-l_i \ll 1$). The third term, $A_i c_i/\bar{c}$, represents competitive victories when the i genotype is abundant: $A_i \rightarrow 0$ if $l_i \ll 1$. The relative importance of these three terms varies with both the overall propagule density L and the relative propagule frequencies l_i/L . If $l_i \gg 1$ for all genotypes, we recover the classic lottery model (only the $A_i c_i/\bar{c}$ term remains, and $A_i \rightarrow 1/L$). Thus, Eq. (5) generalizes the classic lottery model to account for arbitrary propagule densities for each genotype.

Fig. 1 shows that Eq. (5) (and its components) closely approximate direct simulations of random dispersal and lottery competition over a wide range of propagule densities (obtained by varying U). Two genotypes are present, one of which has a c -advantage and is at low frequency. The growth of the low-frequency genotype relies crucially on the low-density competition term $R_i c_i/\bar{c}$, and also to a lesser extent on the high density competition term $A_i c_i/\bar{c}$ if l_1 is large enough (Fig. 1b). On the other hand, $R_i c_i/\bar{c}$ is negligible for the high-frequency genotype, which depends instead on high density territorial victories (Fig. 1d).

3.2 Invasion of rare genotypes and coexistence

To determine how b , c and d will evolve in a population where those traits are being modified by mutations, we need to know whether mutant lineages will grow (or decline) starting from low densities. In this section we discuss the behavior of rare genotypes predicted by Eq. (5).




Figure 1: The change in genotype abundances in a density dependent lottery model is closely approximated by Eq. (5). $\Delta+n_i/m_i$ from Eq. (5) (and its separate components) are shown, along with direct simulations of random dispersal and lottery competition over one generation over a range of propagule densities (varied by changing U with the m_i fixed). Two genotypes are present. (a) and (b) show low-frequency genotype with c -advantage ($m_1/M = 0.1$, $c_1 = 1.5$), (c) and (d) show the high-frequency predominant genotype ($m_2/M = 0.9$, $c_2 = 1$).

Suppose that a population with a single genotype i is in equilibrium. Then $R_i = 0$, $\bar{c} = c_i$ and $\Delta n_i = 0$, and so Eq. (5) gives

$$b_i (e^{-L} + A_i) - d_i = 0. \quad (8)$$

Now suppose that a new genotype j , which is initially rare, appears in the population. Then $A_j \ll 1$, $l_j \ll L$ and $\bar{c} \approx c_i$, and so, from Eq. (5), n_j will increase if

$$b_j \left(e^{-L} + R_j \frac{c_j}{c_i} \right) - d_j > 0. \quad (9)$$

Combining Eqs. (8) and (9), it is easily verified that if j is superior in one trait, but otherwise identical to i , it will invade. Moreover, j will eventually exclude i , since it is strictly superior. However, stable coexistence is possible between genotypes that are superior in different traits. To illustrate, suppose that j is better at securing territories ($c_j > c_i$), that i is better at producing propagules ($b_i > b_j$), and that $d_i = d_j$. Coexistence occurs if j will invade an i -dominated population, but i will also invade a j -dominated population (“mutual invasion”). It is not hard to show that this is possible, since if b_i is so large that $L \gg 1$ when i is dominant, and b_j is so small that $L \ll 1$ when j is dominant, then, combining Eqs. (8) and (9), we find that i invades j because $b_i > b_j$, while j invades i provided that

$$b_j c_j R_j - b_i c_i A_i > 0. \quad (10)$$

Thus, coexistence occurs if c_j is large enough. Intuitively, the mechanism for coexistence is that territorial contests are important in an i -dominated population (high L), ensuring that the c -specialist j is not excluded, yet territorial contests are irrelevant in a j -dominated population (low L), ensuring that the b -specialist i is not excluded. Fig. 2 shows an example of this coexistence between b and c specialists.

A similar argument applies for coexistence between high- c and low- d specialists; again coexistence occurs because the importance of territorial contests declines along with propagule density L as the c -specialist increases in frequency. Coexistence is technically possible between b - and d -specialists which exactly satisfy $b_i/d_i = b_j/d_j$ (this follows from the fact that all propagules have the same probability of success when $c_i = c_j$ i.e. $A_i + R_i = A_j + R_j$). However, this coexistence scenario is not biologically relevant, since the tiniest deviation from $b_i/d_i = b_j/d_j$ will lead to the eventual exclusion of the genotype with greater b_i/d_i .




Figure 2: Coexistence between b ($c_i = 1$, $b_i = 1$) and c ($c_j = 2$, $b_j = 0.7$) specialists, where $d_i = d_j = 0.3$. Vertical axis shows frequency of the c -specialist predicted by Eq. (5).

If the rare genotype j arises due to mutation, then its initial low-density behavior is more complicated than the above invasion analysis suggests. The mutant lineage starts with one individual $n_j = 1$, and remains at low abundance for many generations after its initial appearance. During this period, the mutant abundance n_j will behave stochastically, and the deterministic equations (1) and (5) do not apply (Section 2). However, if n_j becomes large enough, its behavior will become effectively deterministic, and governed by Eq. (5). For mutants with fitness greater than the population mean fitness, this process is known as “establishment”, and occurs when n_j is of order $1/s$, where s is the mutant’s fitness advantage relative to the mean [22]. Here we do not consider the initial stochastic behavior of novel mutants, and have restricted our attention to the earliest deterministic behavior of rare genotypes. In particular, for beneficial mutations we have only considered the case where s is large enough that deterministic behavior starts when $n_j \ll N$.

3.3 Environmental archetypes and Grime’s triangle

We now discuss which changes in the traits b, c and d will be most favored under different environmental conditions. Of particular interest are Grime’s “disturbance”, “stress” and “ideal” environmental archetypes. To proceed, we need to map these verbal archetypes to quantitative parameter regimes in our model.

The ideal environmental archetype is characterized by the near-absence of stress and disturbance. Consequently, $d_i \ll 1$, whereas b_i can easily be of order 1 or larger. From Eq. (3), the equilibrium value of L only depends on the ratio of birth and death rates. For one genotype, $L/(1 - e^{-L}) = b_i/d_i$, and so the propagule density is high $L \approx b_i/d_i \gg 1$. Moreover, since $L = b_i N/(N - K)$ by definition, population density is also high $N/K \approx 1$. Thus, territorial contests are decisively important.

The disturbance archetype is characterized by unavoidably high extrinsic mortality caused by physical destruction. Disturbances do not only affect adults as in Eq. (2), but also juveniles in the process of territorial contest. These juvenile deaths can be represented as a fractional reduction in the number of territories secured. To illustrate, we assume that the disturbance is equally damaging to adults and juveniles, so that only $(1 - d_i)\Delta_+ n_i$ rather than $\Delta_+ n_i$ territories are secured by genotype i each generation. Then, the disturbance archetype is characterized by d_i being close to 1 (almost all adults and juveniles are killed each generation), which can only be tolerated by those organisms with large enough b_i to ensure population persistence. The single genotype equilibrium then gives $L \approx 2(1 - d_i/[(1 - d_i)b_i])$, and we have $L \ll 1$ and $N/K \ll 1$. The terms proportional to c_i/\bar{c} in Eq. (5) are then negligible, and $\Delta_+ n_i$ depends primarily on b_i .

The stress archetype is more ambiguous, and has been the subject of an extensive debate in the plant ecology literature (the ‘‘Grime-Tilman’’ debate [23]). In Grime’s view, the archetypal stressful environmental imposes such severe challenges that success depends primarily on the ability to tolerate the stressors. Population density N/K is suppressed to such low levels that competition between individuals is not important. In our model, this corresponds to $b_i \ll 1$ and $b_i/d_i \approx 1$. Thus $L \ll 1$ and $N/K \ll 1$ similar to the disturbance archetype, except that stress now constrains b_i to be small.

The alternative view is that the stress archetype should rather be interpreted as a large reduction in the maximum number of individuals that can be supported by the environment [24]. For example, in the case that the stress is induced by a scarcity of consumable resources, competition for resources would likely be intense, and the stressed population should actually be regarded as having a high population density. In our model, this would imply a large reduction in K (greater per-individual territorial requirement). That is, N under stress is much lower than under ideal conditions, but it is not much lower than K for the stressful environment. Since our model accommodates both of these alternatives, we include them both here.

	Ideal	Disturbance*	Stress (G)	Stress (K)
Parameter-	$d_i \ll 1$	$d_i \approx 1$	$b_i \ll 1$	$b_i \ll 1$
regime	$b_i/d_i \gg 1$	$b_i/d_i \gg 1$	$b_i \approx d_i$	$b_i > d_i$
Density N/K	High	Low	Low	High
$\Delta_+ n_i \propto$	$b_i c_i$	b_i	b_i	$b_i c_i$
Evolution for	$\uparrow b, \uparrow c$	$\uparrow b$	$\downarrow d$	$\uparrow c$

Figure 3: The realization of Grime’s environmental archetypes in our model, as well as the low- K variant of the stress archetype. Shown are the mapping to our parameters of each archetype, the approximate dependence of $\Delta_+ n_i$ on b_i and c_i , as well as the corresponding expected evolutionary changes in b_i , c_i and d_i . *Mortality affects both adults and juveniles in the disturbance archetype, with $\Delta_+ n_i$ replaced by $(1 - d_i)\Delta_+ n_i$ in Eq. (2).

The mapping of environmental archetypes to our model parameters is summarized in the first two rows of Fig. 3. Also shown is the approximate dependence of $\Delta_+ n_i$ on b_i and c_i for each archetype (third row), which can be used infer the expected direction of evolution for the traits b , c and d (fourth row).

The latter is obtained as follows. As noted in the previous section, if beneficial mutations can survive the low-abundance stochastic regime, their behavior is governed deterministically by Eq. (5). They will then proceed to grow deterministically (establishment). The probability of establishment increases with the mutant fitness advantage, and is therefore typically on the order of a few percent, whereas the fixation of neutral mutations is exceedingly unlikely (probability of order $1/N$). Consequently, the direction of evolutionary change is determined by which trait changes confer an appreciable benefit, subject to the constraints imposed by the environment.

For example, in Grime’s version of the stress archetype, population density is low, so competition is not important, and so only mutants with greater b or lower d will have an appreciably greater Δn_i . Mutations in c are effectively neutral, and will rarely fix. However, by definition of the stress archetype, b is constrained to be very small. Thus, while some rare mutations may produce small improvements in b , it is much more likely that mutations will arise that lower d , making this the expected direction of evolutionary change for Grime’s stress archetype.

Following Grime’s original argument for a triangular scheme [14], Fig. 4 represents each environmental archetype schematically as a vertex on a




Figure 4: The realization of Grime’s triangle in our model. Schematic representation of the triangular space bounded by the low/high extremes of stress/disturbance. The low- K interpretation of stress is also shown. The vertices of the triangles correspond to environmental archetypes. Selection favors different traits at each vertex, leading to different trait clusters.

triangular space defined by perpendicular stress and disturbance axes. The ideal archetype lies at the origin (no stress or disturbance), while the stress and disturbance archetypes lie at the limits of survival on their respective axes. The hypotenuse connecting the stress and disturbance endpoints represents the limits of survival in the presence of a combination of stress and disturbance. The direction of evolutionary change is different at each vertex, leading to the emergence of different trait clusters or “primary strategies”.

4 Discussion

Grime’s triangle without constraints

Our model differs from both Grime’s C/S/R and MacArthur’s r/K schemes in the role of the propagule production rate b , a measure of intrinsic fecundity closely related to the growth rate at low densities $r = b - d$. In both of those schemes, the essential feature of life at high densities is competition. This is less in the r/K scheme, which does not explicitly it is better to have a

contributes just as much to

In the Introduction, we noted that the r - K dichotomy is not consistent with “ K -selection” (i.e. selection at high densities) for growth ability traits.

Specifically, positive correlations between measures of r and K are common, both between species and strains [16, 17, 18, 19], and as a result of experimental evolution [16, 20]). From the perspective of our model, this correlation is not at all surprising;

r-K correlation, meaning of K selection

Actual K selection

Significance of stage structure

caveats: large c discrepancy

References

- [1] Warren J Ewens. *Mathematical Population Genetics 1: Theoretical Introduction*, volume 27. Springer Science & Business Media, 2012.
- [2] Cyrille Violle, Marie-Laure Navas, Denis Vile, Elena Kazakou, Claire Fortunel, Irne Hummel, and Eric Garnier. Let the concept of trait be functional! *Oikos*, 116(5):882–892, 2007.
- [3] Eric R. Pianka. On r- and K-Selection. *The American Naturalist*, 104(940):592–597, 1970.
- [4] Stephen C. Stearns. The evolution of life history traits: A critique of the theory and a review of the data. *Annual Review of Ecology and Systematics*, 8:145–171, 1977.
- [5] Mark S Boyce. Restitution of r-and k-selection as a model of density-dependent natural selection. *Annual Review of Ecology and Systematics*, 15:427–447, 1984.
- [6] David Reznick, Michael J. Bryant, and Farrah Bashey. r- and k-selection revisited: The role of population regulation in life-history evolution. *Ecology*, 83(6):1509–1520, 2002.
- [7] Robert H MacArthur. Some generalized theorems of natural selection. *Proceedings of the National Academy of Sciences*, 48(11):1893–1897, 1962.
- [8] Emily S. Darling, Lorenzo Alvarez-Filip, Thomas A. Oliver, Timothy R. McClanahan, and Isabelle M. Ct. Evaluating life-history strategies of reef corals from species traits. *Ecology Letters*, 15(12):1378–1386, 2012.

- [9] T. R. E. Southwood. Habitat, the templet for ecological strategies? *Journal of Animal Ecology*, 46(2):337–365, 1977.
- [10] Kirk O Winemiller and Kenneth A Rose. Patterns of life-history diversification in north american fishes: implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences*, 49(10):2196–2218, 1992.
- [11] J. Philip Grime. *Plant Evolutionary Biology*, chapter The C-S-R model of primary plant strategies — origins, implications and tests, pages 371–393. Springer Netherlands, Dordrecht, 1988.
- [12] Kirk O. Winemiller, Daniel B. Fitzgerald, Luke M. Bower, and Eric R. Pianka. Functional traits, convergent evolution, and periodic tables of niches. *Ecology Letters*, 18(8):737–751, 2015.
- [13] J P Grime. Vegetation classification by reference to strategies. *Nature*, 250:26–31, 1974.
- [14] J P Grime. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *American naturalist*, pages 1169–1194, 1977.
- [15] D Tilman. Resource competition and plant traits: a response to craine et al. 2005. *Journal of Ecology*, 95(2):231–234, 2007.
- [16] Leo S Luckinbill. Selection and the r/k continuum in experimental populations of protozoa. *American Naturalist*, pages 427–437, 1979.
- [17] Eizi Kuno. Some strange properties of the logistic equation defined withr andk: Inherent defects or artifacts? *Researches on population ecology*, 33(1):33–39, 1991.
- [18] A Jan Hendriks, Johanna LM Maas-Diepeveen, Evelyn HW Heugens, and Nico M van Straalen. Meta-analysis of intrinsic rates of increase and carrying capacity of populations affected by toxic and other stressors. *Environmental toxicology and chemistry*, 24(9):2267–2277, 2005.
- [19] Jay M Fitzsimmons, Sijmen E Schoustra, Jeremy T Kerr, and Rees Kassen. Population consequences of mutational events: effects of antibiotic resistance on the r/k trade-off. *Evolutionary ecology*, 24(1):227–236, 2010.

- [20] Leo S Luckinbill. r and k selection in experimental populations of *Escherichia coli*. *Science (New York, NY)*, 202(4373):1201–1203, 1978.
- [21] Peter L Chesson and Robert R Warner. Environmental variability promotes coexistence in lottery competitive systems. *American Naturalist*, pages 923–943, 1981.
- [22] Michael M. Desai and Daniel S. Fisher. Beneficial mutation-selection balance and the effect of linkage on positive selection. *Genetics*, 176(3):1759–1798, 2007.
- [23] Rien Aerts. Interspecific competition in natural plant communities: mechanisms, trade-offs and plant-soil feedbacks. *Journal of Experimental Botany*, 50(330):29–37, 1999.
- [24] Douglas R Taylor, Lonnie W Aarssen, and Craig Loehle. On the relationship between r/k selection and environmental carrying capacity: a new habitat template for plant life history strategies. *Oikos*, pages 239–250, 1990.

Appendix A: Poisson approximation

The propagule numbers x_i in different territories are not independent random variables. To determine the dispersal outcomes in all unoccupied territories exactly, we would need to proceed territory-by-territory as follows. In the first territory we evaluate, x_i drawn from a binomial distribution with m_i trials and success probability $1/U$. In the second, x_i is drawn from a binomial distribution with $m_i - x$ trials and success probability $1/(U - 1)$, where x is the number of propagules that landed in the first territory. And so on.

For sufficiently large K , holding n_i/K fixed, the Poisson limit theorem implies that the binomial distributions for x_i at each successive stage of this procedure are all closely approximated by a Poisson distribution with mean l_i , where we have used the fact that large K implies large U except in the biologically uninteresting case that there is vanishing population turnover $d_i \sim 1/K$.

Under the Poisson approximation, the total number of genotype i propagules $\sum x_i$ (summed over unoccupied territories) will deviate about its mean value m_i . Since the coefficient of variation of $\sum x_i$ is proportional to $1/\sqrt{m_i}$,

these deviations are negligible unless m_i is very small (say of order 100 or less).

Appendix B: Derivation of growth equation

We separate the right hand side of Eq. (1) into three components $\Delta_+ n_i = \Delta_u n_i + \Delta_r n_i + \Delta_a n_i$ which vary in relative magnitude depending on the propagule densities l_i . Following the notation in the main text, the Poisson distributions for the x_i (or some subset of the x_i) will be denoted p ; for instance $p(x_1, \dots, x_G) = p_1(x_1) \dots p_G(x_G)$ and $p(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_G) = p_1(x_1) \dots p_{i-1}(x_{i-1}) p_{i+1}(x_{i+1}) \dots p_G(x_G)$. We use P as a general shorthand for the probability of particular outcomes.

Growth without competition

The first component, $\Delta_u n_i$, accounts for territories where only one focal propagule is present $x_i = 1$ and $x_j = 0$ for $j \neq i$ (u stands for “uncontested”). The proportion of territories where this occurs is $l_i e^{-L}$, and so

$$\Delta_u n_i = U l_i e^{-L} = m_i e^{-L}. \quad (11)$$

Competition when rare

The second component, $\Delta_r n_i$, accounts for territories where a single focal genotype propagule is present along with at least one non-focal propagule (r stands for “rare”) i.e. $x_i = 1$ and $\sum_{j \neq i} x_j \geq 1$. The number of territories where this occurs is $U p_i(1) P(\sum_{j \neq i} x_j \geq 1) = b_i n_i e^{-l_i} (1 - e^{-(L-l_i)})$. Thus

$$\Delta_r n_i = m_i e^{-l_i} P \left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}}, \quad (12)$$

where $\langle \rangle_{\tilde{p}}$ denotes the expectation with respect to \tilde{p} , and \tilde{p} is the probability distribution of nonfocal propagaule abundances x_j *after* dispersal, in those territories where exactly one focal propagule, and at least one non-focal propagule, landed.

We now show that, with respect to \tilde{p} , the standard deviation in $\sum_{j \neq i} c_j x_j$, $\sigma(\sum_{j \neq i} c_j x_j)$, is much smaller than its mean $\langle \sum_{j \neq i} c_j x_j \rangle_{\tilde{p}}$. Then x_j can be

replaced by its mean in the last term in Eq. (12),

$$\left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}} \approx \frac{c_i}{c_i + \sum_{j \neq i} c_j \langle x_j \rangle_{\tilde{p}}}, \quad (13)$$

which will give us Eq. (6).

The exact expression for $\langle x_j \rangle_{\tilde{p}}$ is somewhat complicated. Letting k denote the total number of propagules in a territory, and $\mathbf{x}_i = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_G)$ denote the vector of non-focal abundances, \tilde{p} can be written as

$$\begin{aligned} \tilde{p}(\mathbf{x}_i) &= p(\mathbf{x}_i | k \geq 2, x_i = 1), \\ &= \frac{P(k \geq 2 | \mathbf{x}_i, x_i = 1) p(\mathbf{x}_i | x_i = 1)}{P(k \geq 2)}, \\ &= \frac{p(\mathbf{x}_i | x_i = 1)}{1 - (1 + L)e^{-L}}, \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{k=2}^{\infty} P(k) p(\mathbf{x}_i | \sum_{j \neq i} x_j = k - 1), \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{k=2}^{\infty} \frac{P(k) \delta_k^{\sum_{j \neq i} x_j}}{P(\sum_{j \neq i} x_j = k - 1)} p(\mathbf{x}_i), \\ &= \frac{Le^{-l_i}}{1 - (1 + L)e^{-L}} \sum_{k=1}^{\infty} \left(\frac{L}{L - l_i} \right)^k \frac{\delta_k^{\sum_{j \neq i} x_j}}{k + 1} p(\mathbf{x}_i), \end{aligned} \quad (14)$$

where $\delta_k^{\sum_{j \neq i} x_j} = 1$ if $\sum_{j \neq i} x_j = k$, and equals zero otherwise. Then, since

$$\begin{aligned} \sum_{\mathbf{x}_i} \delta_k^{\sum_{j \neq i} x_j} p(\mathbf{x}_i) x_j &= \frac{l_j}{L - L_j} k P(\sum_{j \neq i} x_j = k) \\ &= l_j P(\sum_{j \neq i} x_j = k - 1), \end{aligned} \quad (15)$$

after some algebra we obtain,

$$\langle x_j \rangle_{\tilde{p}} = \frac{l_j}{1 - (1 + L)e^{-L}} \frac{L - 1 + e^{-L}}{L - l_i}. \quad (16)$$

To calculate the relative fluctuations in $\sum_{j \neq i} c_j x_j$, we use the following approximation, which gives considerably simpler expressions for the means, variances and covariances of the x_j compared with the exact expressions using \tilde{p} . Rather than evaluating the situation in each territory after dispersal as above, we let \tilde{p} instead be the \mathbf{x}_i dispersal probabilities in a territory where one focal propagule is assumed to be present, conditional on $\sum_{j \neq i} x_j > 1$. This gives $\langle x_j \rangle_{\tilde{p}} = l_j/C$,

$$\sigma^2(x_j) = \frac{l_j^2}{C} \left(1 - \frac{1}{C}\right) + \frac{l_j}{C}, \quad (17)$$

and

$$\sigma(x_j, x_k) = \frac{l_j l_k}{C} \left(1 - \frac{1}{C}\right), \quad (18)$$

where $C = 1 - e^{-(L-l_i)}$ (note the difference from Eq. (16) for $\langle x_j \rangle_{\tilde{p}}$). Then, since

$$\sigma^2\left(\sum_{j \neq i} c_j x_j\right) = \sum_{j \neq i} \left[c_j^2 \sigma^2(x_j) + 2 \sum_{k > j} c_j c_k \sigma(x_j, x_k) \right], \quad (19)$$

and $1/C > 1$, we have

$$\frac{\sigma(\sum_{j \neq i} c_j x_j)}{\langle \sum_{j \neq i} c_j x_j \rangle} < C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j\right)^{1/2}}{\sum_{j \neq i} c_j l_j}. \quad (20)$$

Without loss of generality, we restrict attention to the case that the total nonfocal density $L - l_i$ is of order 1 or larger (otherwise $\Delta_r n_i$ does not contribute significantly to $\Delta_+ n_i$ because $\Delta_r n_i$ is proportional to $C = 1 - e^{-(L-l_i)}$).

When at least some of the nonfocal propagule densities are large $l_j \gg 1$, then the RHS of Eq. (20) is $\ll 1$, as desired. This is also the case if none of the nonfocal genotype densities are large and the c_j are all of similar magnitude (their ratios are of order one); the worst case scenario occurs when $(L - l_i) \sim O(1)$, in which case the negative covariances (Eq. (18)) which were neglected in the RHS of Eq. (20) significantly reduce the overall variance $\sigma^2(\sum_{j \neq i} c_j x_j)$.

However, the relative fluctuations in $\sum_{j \neq i} c_j x_j$ can be large if some of the c_j are much larger than the others. Specifically, if $c_j l_j \gg c_k l_k$ ($j, k \neq i$, $j \neq k$) and $l_j \ll 1$ (i.e. in the presence of a rare, extremely strong competitor), then we cannot make the replacement Eq. (13).

Substituting Eqs. (13) and (16) into Eq. (12), we obtain

$$\Delta_r n_i \approx m_i R_i \frac{c_i}{\bar{c}}, \quad (21)$$

where R_i is defined in Eq. (6).

Competition when abundant

The final contribution, $\Delta_a n_i$, accounts for territories where two or more focal propagules are present (a stands for “abundant”). Similarly to Eq. (12), we have

$$\Delta_a n_i = U(1 - (1 + l_i)e^{l_i}) \left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \quad (22)$$

where \hat{p} is the probability distribution of both focal and nonfocal propagule abundances *after* dispersal in those territories where at least two focal propagules landed.

Again, we wish to show that the relative fluctuations in $\sum c_j x_j$ are much smaller than 1 (with respect to \hat{p}), so that we have

$$\left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \approx \frac{c_i \langle x_i \rangle_{\hat{p}}}{\sum_j c_j \langle x_j \rangle_{\hat{p}}}. \quad (23)$$

Following a similar procedure as for $\Delta_r n_i$, where the vector of propagule abundances is denoted \mathbf{x} , we have

$$\begin{aligned} \langle x_j \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2) \\ &= \sum_k P(k | x_i \geq 2) \sum_{x_i} \sum_{\mathbf{x}_i} x_j p(\mathbf{x}_i | \sum_{j \neq i} x_j = k - x_i) p(x_i | x_i \geq 2, k) \\ &= \sum_k P(k | x_i \geq 2) \sum_{x_i} \frac{l_j (k - x_i)}{L - l_j} p(x_i | x_i \geq 2, k) \\ &= \frac{l_j}{L - l_j} \left(L \frac{1 - e^{-L}}{1 - (1 + L)e^{-L}} - l_j \frac{1 - e^{-l_j}}{1 - (1 + l_j)e^{-l_j}} \right) \end{aligned} \quad (24)$$

for $j \neq i$, and

$$\langle x_i \rangle_{\hat{p}} = l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}}. \quad (25)$$

To calculate the relative fluctuations in $\sum_{j \neq i} c_j x_j$, we use a similar approximation as for $\Delta_r n_i$: \tilde{p} is approximated by the \mathbf{x} dispersal probabilities in a territory where at least two focal propagule is assumed to be present. All covariances are now zero, so that $\sigma^2(\sum c_j x_j) = \sum c_j^2 \sigma^2(x_j)$, where $\sigma^2(x_j) = l_j$ for $j \neq i$. The expression for $\sigma^2(x_i)$ is more complicated. We assume $p(x_i = 0) \approx 0$ without loss of generality (since otherwise $D \gg 1$ and Δn_a is negligible). Then

$$\sigma^2(x_i) = \frac{l_i^2}{D} \left(1 - \frac{1}{D}\right) + \frac{l_i}{D}, \quad (26)$$

where $D = 1 - (1 + l_i)e^{-l_i}$, analogous to Eq. (17), and

$$\frac{\sigma(\sum c_j x_j)}{\langle \sum c_j x_j \rangle} \approx \frac{\left(\sum_{j \neq i} c_j^2 l_j + c_i^2 \sigma^2(x_i)\right)^{1/2}}{\sum_{j \neq i} c_j l_j + c_i l_i / D}. \quad (27)$$

Similarly to Eq. (20), the RHS of (27) will not be $\ll 1$ if there is a nonfocal genotype j with $l_j \ll 1$ and $c_j l_j \gg c_k l_k$ for $j, k \neq i$, $j \neq k$. When this is not the case, then since l_i must be of order 1 or larger for $\Delta_a n$ to make an appreciable contribution to $\Delta_+ n_i$, the RHS of Eq. (27) is $\ll 1$ as desired.

Combining Eqs. (22) and (23), we obtain

$$\Delta_a n_i = m_i A_i \frac{c_i}{\bar{c}}, \quad (28)$$

where A_i is defined in Eq. (7).