

Density-dependent selection in evolutionary genetics: a lottery model of Grime's triangle

Manuscript elements:

Keywords: r/K selection, absolute fitness, eco-evo, interference competition, competition-colonization trade-off.

Manuscript type: Article.

Prepared using the suggested L^AT_EX template for *Am. Nat.*

Abstract

Fitness is typically represented in heavily simplified terms in evolutionary genetics, often using
3 constant selection coefficients. This excludes fundamental ecological factors such as dynamic
population size or density-dependence from the most genetically-realistic treatments of evolu-
tion, a problem that inspired MacArthur’s influential but problematic r/K theory. Following
6 in the spirit of r/K -selection as a general-purpose theory of density-dependent selection, but
grounding ourselves empirically in “primary strategy” trait classification schemes like Grime’s
triangle, we develop a new model of density-dependent selection which revolves around terri-
9 torial contests. To do so, we generalize the classic lottery model of territorial acquisition, which
has primarily been used for studying species co-existence questions, to accommodate arbitrary
densities. We use this density-dependent lottery model to predict the direction of trait evolution
12 under different environmental conditions and thereby provide a mathematical underpinning for
Grime’s verbal scheme. We revisit previous concepts of density-dependent selection, including r
and K selection, and argue that our model distinguishes between different aspects of fitness in a
15 more natural and intuitive manner.

“...the concept of fitness is probably too complex to allow of a useful mathematical development. Since it enters fundamentally into many population genetics considerations, it is remarkable how little attention has been paid to it.” — Warren J. Ewens, *Mathematical Population Genetics I*, 2004

Evolutionary models differ greatly in their treatment of fitness. In models of genetic evolution, genotypes are typically assigned constant (or frequency-dependent) selection coefficients describing the change in their relative frequencies over time. This simplified treatment of selection facilitates genetic realism, and can be justified over sufficiently short time intervals (Ewens, 2004, p. 276). The emphasis here is to infer past selection, migration and demographic change given a sample of nucleotide sequences, or to predict how allele frequencies change over time based on their fitness effects, population structure, genetic drift and linkage. The resulting picture of evolution excludes basic elements of the ecological underpinnings of selection, including density dependence, and how selection affects population size. This complicates the inference of past selection, because demographic changes can look genealogically very similar to selective frequency changes (Barton, 1998).

By contrast, models of phenotypic trait evolution use absolute fitness functions to describe how some traits of interest affect survival and reproduction in particular ecological scenarios (Diekmann et al., 2004; Metz et al., 1992). These fitness functions can be quite problem-specific and often only account for a few traits at a time. The emphasis here is on the conditions for invasion from low frequencies and co-existence, rather than frequency or abundance trajectories over time. For instance, adaptive dynamics uses “invasion fitness” to explore the consequences of eco-evolutionary feedbacks (Diekmann et al., 2004).

It is challenging to generalize beyond particular traits or ecological scenarios to model fundamentally different forms of selection. Perhaps this is not surprising given that fitness is such a complex quantity, dependent on all of a phenotype’s functional traits (Violle et al., 2007) and its environment. A detailed, trait-based, predictive model of fitness would be enormously complicated and situation-specific. It is therefore easy to doubt the feasibility of a simplified, general

mathematical treatment of fitness (Ewens, 2004, p. 276). For example, MacArthur's famous r/K scheme (MacArthur, 1962; MacArthur and Wilson, 1967) is now almost exclusively known as a framework for understanding life-history traits, and judged on its failure in that role (Boyce, 1984; Pianka, 1970; Reznick et al., 2002; Stearns, 1977). However, the r/K scheme's original purpose was to extend the existing population-genetic treatment of selection to account for population density (MacArthur, 1962). Few attempts have been made to develop it further along these lines.

Empirical trait classification studies have suggested the existence of a few "primary strategies", reflecting broadly distinct responses to selection (Winemiller et al., 2015). Grime famously considered (Grime, 1974, 1977, 1988; Westoby, 1998) two broad determinants of population density: stress (persistent hardship e.g. due to resource scarcity or unfavorable temperatures) and disturbance (intermittent destruction of vegetation e.g. due to trampling, herbivory, pathogens, extreme weather or fire). The extremes of these two factors define three primary strategies denoted by C/S/R respectively (Fig. 1): competitors "C" excel in low stress, low disturbance environments; stress tolerators "S" excel in high stress, low disturbance environments; and ruderals "R" excel in low stress, high disturbance environments. Population persistence is not possible in high-stress, high-disturbance environments. Grime showed that measures of C, S and R across a wide range of plant species are anti-correlated, so that strong C-strategists are weak S and R strategists, and so on, creating a triangular C/S/R ternary plot (Grime, 1974). Similar schemes were proposed for insects (Southwood, 1977), fishes (Winemiller and Rose, 1992), and zooplankton (Allan, 1976). More recently, modern hierarchical clustering techniques have revealed distinct trait clusters in corals analogous to Grime's primary strategies (Darling et al., 2012). These empirical findings suggest that functional traits contribute to fitness predominantly via a few key factors such as intrinsic reproductive rate or stress-tolerance.

Here we explore the interplay between some "key factors" of fitness in a simplified, territorial model of growth, dispersal and competition. This broadly follows the original spirit of MacArthur's r/K scheme. More specifically, our aim is to begin adding some ecological realism to population genetics' time-dependedent, genetically-focused view of evolution. We revisit the



Figure 1: Schematic of Grime's triangle. The two axes show increasing levels of environmental stress and disturbance, respectively. Population persistence is not possible if the combination of stress and disturbance is too large (dashed line). This creates a triangle, each corner of which corresponds to a "primary strategy".

classic lottery model of Chesson and Warner (1981), which has two features that make it well suited for this role, but one critical flaw that we rectify here.

72 The first feature is that the lottery representation of competition is particularly concise. Mature individuals ("adults") each require their own territory, whereas newborn individuals ("propagules") disperse to, and subsequently compete for, territories made available by the death
75 of adults. Territorial contest among propagules leaves a single victorious adult per territory, the victor chosen at random from the propagules present, with probabilities weighted by a coefficient for each type representing competitive ability, akin to a lottery (Sale, 1977). By comparison,
78 coefficients for the pairwise effects of types on each other (e.g. the α coefficients in the generalized Lotka-Volterra equations and the associated concept of " α -selection"; Case and Gilpin 1974; Gill 1974; Joshi et al. 2001), or explicit resource consumption (Tilman, 1982), are much more
81 complicated. The second feature is the close connection between the lottery model and one of the foundational models of population genetics, the Wright-Fisher model of genetic drift, which we discuss further below.

84 The critical flaw of the classic lottery model is that it breaks down at low densities (few propagules dispersing to each territory), precluding density-dependent behaviour. Our first task is to analytically extend the classic lottery model to correctly account for low density behavior

87 (sections “Model” and “Mean field approximation”).

Using our extended lottery model, we then revisit Grime’s C/S/R scheme, and evaluate how C/S/R manifests at the level of within-population genotypic evolution (as opposed to phenotypic
90 divergence between species; sections “Invasion of rare genotypes and coexistence” and “Primary strategies and Grime’s triangle”). This represents a “sanity check” on our density-dependent lottery model. The resulting formulation of the C/S/R scheme is mathematical, in contrast
93 to Grime’s original verbal and descriptive approach, which is a recognized hindrance to the evaluation or broader application of the C/S/R scheme (e.g. Tilman 2007).

We then explore some time-dependent behavior of our extended lottery model. Taking an
96 example inspired by recent studies of rapid, seasonal evolution in *Drosophila* (Bergland et al., 2014), we discuss how environmental fluctuations might stabilize polymorphisms in the presence of cyclical population density.

99 **Model**

We assume that reproductively mature individuals (“adults”) each require their own territory to survive and reproduce (Fig. 2). All territories are identical, and the total number of territories is
102 T . Time t advances in discrete iterations, each representing the time from birth to reproductive maturity. In iteration t , the number of adults of the i ’th genotype is $n_i(t)$, the total number of adults is $N(t) = \sum_i n_i(t)$, and the number of unoccupied territories is $U(t) = T - N(t)$. We
105 assume that the n_i and T are large enough that stochastic fluctuations in the n_i (“drift”) can be ignored. We derive deterministic equations for the expected change in the n_i over time, leaving the evaluation of drift for future work.

108 Each iteration, adults produce new offspring (“propagules”), m_i of which disperse to unoccupied territories. We assume that adults cannot be ousted from their territories, so that m_i only includes propagules landing on unoccupied territories. Propagules disperse at random over
111 the unoccupied territories, regardless of distance from their parents, and independently of each




Figure 2: Each iteration of our lottery model has three main elements. First, propagules are produced by adults which are dispersed at random over the unoccupied territories (only propagules landing on unoccupied territories are shown). Lottery competition then occurs in each unoccupied territory (competition in only one territory is illustrated): each genotype has a probability proportional to $b_i n_i c_i$ of securing the territory. Then occupied territories are freed up by adult mortality. In Eq. (3) and most of the paper, only adults can die (red crosses), but we will also consider the case where juveniles die (blue cross; section “Primary strategies and Grime’s triangle”).

other. There is no interaction between propagules (e.g. avoidance of territories crowded with propagules). Loss of propagules during dispersal is subsumed into m_i .

In general, m_i will increase with n_i , and will depend on population density N . For example, if b_i is the number of successfully dispersing propagules produced per genotype i adult, then the loss of propagules due to dispersal to occupied territories implies $m_i = b_i(1 - N/T)n_i$, akin to Levins' competition-colonization model (Levins and Culver, 1971; Tilman, 1994). In section "Cyclical birth and death rates" we evaluate Eq. (4) numerically using this functional form for m_i , with b_i assumed to be constant.

In the sections "Invasion of rare genotypes and coexistence" and "Primary strategies and Grime's triangle", we assume the simpler form $m_i = b_i n_i$, with constant b_i , meaning that all propagules land on unoccupied territories (a form of directed dispersal). This simplifies the mathematics without affecting the results of those sections, which only depend on the low-frequency invasion behavior of Eq. (4). Note that due to our assumption of uniform dispersal, the parameter b_i can be thought of as a measure of "colonization ability", which combines fecundity and dispersal ability (Bolker and Pacala, 1999; Levins and Culver, 1971; Tilman, 1994).

The number of individuals of the i 'th genotype landing in any particular territory is denoted x_i . We assume that x_i follows a Poisson distribution $p_i(x_i) = l_i^{x_i} e^{-l_i} / x_i!$, where $l_i = m_i / U$ is the mean territorial propagule density. This only approximates uniform random dispersal, but is essentially exact provided that the n_i are large enough that drift can be ignored (Appendix A).

When multiple propagules land on the same territory, the victor is determined by lottery competition: genotype i wins a territory with probability $c_i x_i / \sum_j c_j x_j$, where c_i is a constant representing relative competitive ability (Fig. 2).

In the classic lottery model (Chesson and Warner, 1981), unoccupied territories are assumed to be saturated with propagules from every genotype $l_i \gg 1$. From the law of large numbers, the composition of propagules in each territory will then not deviate appreciably from the mean composition l_1, l_2, \dots, l_G (G is the number of genotypes present), and so the probability that genotype i wins any particular unoccupied territory is approximately $c_i l_i / \sum_j c_j l_j$. Let $\Delta_+ n_i$ denote the

number of territories won by genotype i . Then $\Delta_+n_1, \Delta_+n_2, \dots, \Delta_+n_G$ follow a multinomial distribution with U trials and success probabilities $\frac{c_1 l_1}{\sum_j c_j l_j}, \frac{c_2 l_2}{\sum_j c_j l_j}, \dots, \frac{c_G l_G}{\sum_j c_j l_j}$, respectively. Genotype i

141 is expected to win $c_i l_i / \sum_j c_j l_j$ of the U available territories, and deviations from this expected outcome are small (since T is large by assumption), giving

$$\Delta_+n_i(t) = \frac{c_i l_i}{\sum_j c_j l_j} U(t) = b_i n_i \frac{1}{L} \frac{c_i}{\bar{c}}, \quad (1)$$

where $\bar{c} = \sum_j c_j m_j / M$ is the mean propagule competitive ability for a randomly selected propagule, $L = M/U$ is the total propagule density and $M = \sum_j m_j$ is the total number of propagules.

There is a close connection between the classic lottery model and the Wright-Fisher model of genetic drift (Svardal et al., 2015). In the Wright-Fisher model, genotype abundances are sampled each generation from a multinomial distribution with success probabilities $w_i n_i / \sum_j w_j n_j$, where w is relative fitness and the n_i are genotype abundances in the preceding generation. Population size N remains constant. This is mathematically equivalent to the classic lottery model with non-overlapping generations ($d_i = 1$ for all i) and $w_i = b_i c_i$. Thus, the classic lottery model allows us to replace the abstract Wright-Fisher relative fitnesses w_i with more ecologically-grounded fecundity, competitive ability and mortality parameters b_i , c_i and d_i , respectively. Since birth and death rates affect absolute abundances, this allows us to evaluate selection at different densities (after appropriate extensions are made), in an otherwise very similar model to the canonical Wright-Fisher. We therefore expect that drift in our extended lottery model should be similar to that in the Wright-Fisher model, but we leave this for future work.

In our extension of the classic lottery model, we do not restrict ourselves to high propagule densities. Eq. (1) is nonsensical at low densities ($l_i \ll 1$): genotype i can win at most m_i territories, yet Eq. (1) demands $c_i l_i / \sum_j c_j l_j$ of the U unoccupied territories, for any value of U . Intuitively, the cause of this discrepancy is that individuals are discrete. Genotypes with few propagules depend on the outcome of contests in territories where they have at least one propagule present, not some small fraction of a propagule as would be implied by small l_i in the classic lottery model.

In other words, deviations from the mean propagule composition l_1, l_2, \dots, l_G are important at low density.

165 We expect that a fraction $p_1(x_1) \dots p_G(x_G)$ of the U unoccupied territories will have the propagule composition x_1, \dots, x_G . Genotype i is expected to win $c_i x_i / \sum_j c_j x_j$ of these. Ignoring fluctuations about these two expectations (due to our no-drift, large T , large n_i approximation),
168 genotype i 's territorial acquisition is given by

$$\Delta_+ n_i(t) = U(t) \sum_{x_1, \dots, x_G} \frac{c_i x_i}{\sum_j c_j x_j} p_1(x_1) \dots p_G(x_G), \quad (2)$$

in our extended lottery model, where the sum only includes territories with at least one propagule present. Note that unlike the classic lottery model, not all unoccupied territories are claimed
171 each iteration, since under Poisson dispersal a fraction e^{-L} remain unoccupied.

For the majority of this manuscript we assume that mortality only occurs in adults (setting aside the juvenile deaths implicit in territorial contest), and at a constant, genotype-specific per-
174 capita rate d_i , so that the overall change in genotype abundances is

$$\Delta n_i(t) = \Delta_+ n_i(t) - d_i n_i(t). \quad (3)$$

This is reasonable approximation in the absence of disturbances; when we come to consider the effects of disturbances (Section “Primary strategies and Grime’s triangle”), we will incorporate
177 disturbance-induced mortality in competing juveniles (Fig. 2).

Results

Mean Field Approximation

180 Eq. (2) involves an expectation over the time-dependent dispersal distributions p_i , and is thus too complicated to give intuition about the dynamics of density-dependent lottery competition. We now evaluate this expectation using a “mean field” approximation.

Similarly to the high- l_i approximation of classic lottery model, we replace the x_i with appropriate mean values, although we cannot simply replace x_i with l_i . For a genotype with low propagule density $l_i \ll 1$, we have $x_i = 1$ in the territories where its propagules land, and so its growth comes entirely from territories which deviate appreciably from l_i . To account for this, we separate Eq. (2) into $x_i = 1$ and $x_i > 1$ parts. Our more general mean field approximation only requires that there are no large discrepancies in competitive ability (i.e. we do not have $c_i/c_j \gg 1$ for any two genotypes). We obtain (details in Appendix B)

$$\Delta_+ n_i(t) \approx b_i n_i \left[e^{-L} + (R_i + A_i) \frac{c_i}{\bar{c}} \right], \quad (4)$$

where

$$R_i = \frac{\bar{c} e^{-l_i} (1 - e^{-(L-l_i)})}{c_i + \frac{L-1+e^{-L}}{1-(1+L)e^{-L}} \frac{\bar{c} L - c_i l_i}{L-l_i}}, \quad (5)$$

and

$$A_i = \frac{\bar{c} (1 - e^{-l_i})}{\frac{1-e^{-l_i}}{1-(1+l_i)e^{-l_i}} c_i l_i + \frac{1}{L-l_i} \left(L \frac{1-e^{-L}}{1-(1+L)e^{-L}} - l_i \frac{1-e^{-l_i}}{1-(1+l_i)e^{-l_i}} \right) \sum_{j \neq i} c_j l_j}. \quad (6)$$

Comparing Eq. (4) to Eq. (1), the classic lottery per-propagule success rate $c_i/\bar{c}L$ has been replaced by three separate terms. The first, e^{-L} , accounts for propagules which land alone on unoccupied territories; these territories are won without contest. The second, $R_i c_i/\bar{c}$ represents competitive victories when the i genotype is a rare invader in a high density population: from Eq. (5), $R_i \rightarrow 0$ when the i genotype is abundant ($l_i \gg 1$), or other genotypes are collectively rare ($L - l_i \ll 1$). The third term, $A_i c_i/\bar{c}$, represents competitive victories when the i genotype is abundant: $A_i \rightarrow 0$ if $l_i \ll 1$. The relative importance of these three terms varies with both the overall propagule density L and the relative propagule frequencies m_i/M . If $l_i \gg 1$ for all genotypes, we recover the classic lottery model (only the $A_i c_i/\bar{c}$ term remains, and $A_i \rightarrow 1/L$).

Fig. 3 shows that Eq. (4) (and its components) closely approximate direct simulations of random dispersal and lottery competition over a wide range of propagule densities. Two genotypes are present, one of which is at low frequency. The growth of the low-frequency genotype




Figure 3: The change in genotype abundances in a density dependent lottery model is closely approximated by Eq. (4). $\Delta+n_i/m_i$ from Eq. (4) (and its separate components) are shown, along with direct simulations of random dispersal and lottery competition over one iteration over a range of propagule densities (U is varied between 5×10^3 and 10^6 with $m_1 = 10^4$ and $m_2 = 9 \times 10^4$). Two genotypes are present. (a) and (b) show the low-frequency genotype with c -advantage ($c_1 = 1.5$), (c) and (d) show the high-frequency predominant genotype ($c_2 = 1$). Simulation points are almost invisible in (c) and (d) due to near exact agreement with Eq. (4). Dashed lines in (a) and (c) show the breakdown of the classic lottery model.

204 relies crucially on the low-density competition term $R_i c_i / \bar{c}$, and also to a lesser extent on the
high density competition term $A_i c_i / \bar{c}$ if l_1 is large enough (Fig. 3b). On the other hand, $R_i c_i / \bar{c}$
is negligible for the high-frequency genotype, which depends instead on high density territorial
207 victories (Fig. 3d). Fig. 3 also shows the breakdown of the classic lottery model at low propagule
densities.

Primary strategies and Grime's triangle

Here we describe how Grime's "disturbed", "stressful" and "ideal" environments can be captured mathematically in our scheme. To proceed, we need to map these verbally-defined environments to quantitative parameter regimes in our model.

The ideal environment is characterized by the near-absence of stress and disturbance: $d_i \ll 1$, whereas b_i is potentially much larger than 1.

Disturbed environments are characterized by short bursts of high elevated extrinsic mortality caused by physical destruction

high extrinsic mortality, or, more accurately,

In the disturbance extreme of Grime's triangle we can

, we instead represent disturbance by high constant mortality rates d_i , and extend this mortality to juveniles in the process of territorial contest (since disturbances do not only affect adults as in Eq. (3)).

We assume that disturbances are equally damaging to adults and juveniles, so that only $(1 - d_i)\Delta_+ n_i$ rather than $\Delta_+ n_i$ territories are secured by genotype i each iteration. Disturbed environments then correspond to d_i being close to 1 for all genotypes (almost all adults and juveniles are killed each iteration).

Stressful environments are more ambiguous, and have been the subject of an extensive debate in the plant ecology literature (the "Grime-Tilman" debate; Aerts 1999 and references therein).

Severe stress inhibits growth and reproduction, so that $b \ll 1$ (Grime, 1974, 1977). Mutations which appreciably improve b will be either non-existent or extremely unlikely, so b is constrained to remain low. In Grime's view, under these conditions the rate at which propagules successfully develop to adulthood cannot appreciably exceed the mortality rate. This implies $b/d \approx 1$ in our model, and so the propagule density L is suppressed to such low levels that there are essentially no territorial contests occurring.

The alternative view is that, while stressful environments imply lower b and support a lower number of individuals per unit area compared what is attainable in ideal environments, stressed

populations are actually at high densities relative to the environmental carrying capacity, and are
 237 highly competitive (Taylor et al., 1990). In the particular case that stress is caused by scarcity of
 consumable resources, we might expect intense resource competition (for empirical support, see
 Davis et al. 1998). Thus, b may actually appreciably exceed d under stressful conditions, even
 240 though the absolute value of b is small.

The mapping of different environments to our model parameters is summarized in the first
 two rows of Fig. 4. Also shown is the approximate dependence of Δn_i on b_i and c_i for each
 243 environment (fourth row). These can be used infer the expected direction of evolution for the
 traits b , c and d (fifth row) using a standard invasion analysis.

When a new mutant genotype j appears, it starts with one individual $n_j = 1$. While the mu-
 246 tant lineage remains at low-abundance, n_j will behave stochastically, but if its expected growth
 rate is positive and n_j becomes large enough, n_j will effectively grow deterministically according
 to Eq. (4). This transition occurs at an abundance of order $1/r$ (Uecker and Hermisson, 2011),
 249 where $r = \Delta n_i / n_i$ is the mutant lineage's growth rate. Since here we do not evaluate drift in
 the extended lottery model, we do not attempt to calculate the probability that mutants escape
 the initial stochastic phase, and restrict our attention to the earliest deterministic behavior of rare
 252 genotypes while they are still at negligibly low frequency. We simply invoke the well known
 result this probability is proportional to r , with a proportionality factor typically of order one
 (Haldane's formula; Uecker and Hermisson 2011). The fixation of neutral mutations is exceed-
 255 ingly unlikely (probability of order $1/N$). Consequently, the direction of evolutionary change is
 determined by the mutational trait changes which are available and also confer an appreciable
 fitness benefit, where availability is subject to constraints imposed by the environment.

For example, in Grime's interpretation of stressful environments, L is low, so competition is
 258 not important, and only mutants with greater b or lower d will have an appreciably greater Δn_i .
 Mutations in c are effectively neutral, and will rarely fix. However, b is constrained to be small.
 261 Thus, while some rare mutations may produce small improvements in b , it is much more likely
 that mutations lower d , making this the expected direction of evolutionary change.

	Ideal	Disturbance*	Stress (G)	Stress (HD)
Constraints	$d \ll 1$	$d \approx 1$	$b \ll 1$	$b \ll 1$
Other parameters	$b \gg d$	$b \gg d$	$b \approx d$	$b > d$
Density N/T	High	Low	Low	High
$\Delta_+ n_i \propto$	$b_i c_i$	b_i	b_i	$b_i c_i$
Evolution for	$\uparrow b, \uparrow c$	$\uparrow b, \downarrow d$	$\downarrow d$	$\uparrow c, \downarrow d$

Figure 4: The realization of Grime’s three environmental extremes in our model, as well as the high-density variant of the stressful environment. Shown are the mapping of each environment to our parameters, the approximate dependence of $\Delta_+ n_i$ on b_i and c_i , as well as the corresponding expected evolutionary changes in b_i , c_i and d_i . *Mortality affects both adults and juveniles in the disturbed environment, with $\Delta_+ n_i$ replaced by $(1 - d_i)\Delta_+ n_i$ in Eq. (3).




Figure 5: The realization of Grime’s triangle in our model. Schematic representation of the triangular space bounded by the low/high extremes of stress/disturbance. The low- T interpretation of stress is also shown. The vertices of the triangles correspond to different environmental extremes. Selection favors different traits at each vertex, leading to different trait clusters.

Following Grime’s original argument for a triangular scheme (Grime, 1977), Fig. 5 represents each environmental extreme schematically as a vertex on a triangular space defined by perpendicular stress and disturbance axes. The ideal environment lies at the origin (no stress or disturbance), while the stressful and disturbed environments lie at the limits of survival on their respective axes. The hypotenuse connecting the stress and disturbance endpoints represents the limits of survival in the presence of a combination of stress and disturbance. The direction of evolutionary change is different at each vertex, leading to the emergence of different trait clusters or “primary strategies”.

How does Fig. 4 compare to empirical analyses of Grime’s C/S/R strategies? In our compar-

ison we will stick to fishes, corals and plants, for which three-way primary strategy schemes are well developed (Darling et al., 2012; Grime, 1977; Winemiller and Rose, 1992). The connection of our model to fish strategies is necessarily more tentative, given that fishes are motile and not all territorial.

In disturbed environments, we predict evolution for higher b and lower d , but not higher c . Higher b means higher fecundity and/or dispersal ability (see “Model”). This is consistent with a ruderal strategy. Plant ruderals devote a large proportion of their productivity to seed production (Grime, 1977), whereas the analogous “opportunistic” fishes have large intrinsic growth rates (Winemiller and Rose, 1992). In corals, ruderals are distinguished by brood spawning (rather than broadcast spawning; Darling et al. 2012). This corresponds to higher parental investment and lower overall propagule production — counter-intuitively, a stereotypical “K-selected”, high-density trait (Pianka, 1970). However, since broadcast spawners are vulnerable to an Allee effect at the egg fertilization stage (Knowlton, 2001), brood spawning could actually be a way to ensure high b at low densities (Darling et al., 2012). Lower d could be achieved by improved individual resistance to physical destruction, but it is hard to reduce mortality in the face of severe disturbances. Alternatively, shortening the time to reproductive maturity (the iteration time in our model) is an effective way of reducing the chance of death per iteration, d , for a given frequency of disturbance. An exceptionally short life cycle is probably the most defining characteristic of ruderals (Darling et al., 2012; Grime, 1977; Winemiller and Rose, 1992). Note that if evolution manages to appreciably reduce d for a given disturbance intensity, then the population no longer lies at the extreme disturbance vertex of Grime’s triangle. Thus, ruderals are characterized by both high b and high d , but there is a constant pressure for a shorter life cycle, extending the limits of disturbance that can be tolerated.

In stressful environments, we predict evolution for lower d , and also for higher c in the high-density interpretation of stressful environments. Low d is essential when $b \ll 1$, and stress tolerant plants and corals have long life spans, allowing for long intervals between successful recruitments (and episodic broadcast spawning in corals). For fishes, the “equilibrium” strategy

is the analogue of Grime's stress tolerator. This strategy is associated with consumable resource
 300 limitation, and is also characterized by long life span, as well as high parental investment in tiny
 broods. This may reflect a high- c strategy in the face of intense competition for severely limited
 resources (the high-density interpretation).

303 In ideal environments, we predict evolution for higher b and c , but not lower d . In plants
 and corals, a key mechanism for winning territorial contests (higher c) is rapidly outgrowing
 and "shading out" competitors; empirically, rapid individual growth is a defining feature of the
 306 competitor trait cluster (Darling et al., 2012; Grime, 1977). The situation for b is more ambiguous.
 Competitor strategies in plants and corals span a range of b . For fishes, the analogous "periodic"
 strategy is characterized by enormous spawn sizes as well as rapid development (Winemiller
 309 et al., 2015; Winemiller and Rose, 1992). The role of b in ideal environments will be discussed
 further below.

Coexistence in constant and cyclical environments

312 In the previous section we only considered the how b , c and d should respond in Grime's envi-
 ronmental extremes. Here we further explore the low frequency behavior of Eq. (4) to determine
 which types can coexist in a constant environment, and then consider the full time-dependent
 315 behaviour of Eq. (4) in a cyclical environment.

In a population with a single genotype i which is in equilibrium, $R_i = 0$, $\bar{c} = c_i$ and $\Delta n_i = 0$,
 and Eq. (4) gives

$$b_i \left(e^{-L} + A_i \right) - d_i = 0, \quad (7)$$

318 where $A_i = (1 - (1 + L)e^{-L})/L$. Now suppose that a new genotype j , which is initially rare,
 appears in the population. Then $A_j \ll R_j$, $l_j \approx 0$ and $\bar{c} \approx c_i$, and so, from Eq. (4), n_j will increase
 if

$$b_j \left(e^{-L} + R_j \frac{c_j}{c_i} \right) - d_j > 0, \quad (8)$$

321 where $R_j \approx (1 - e^{-L}) / \left(\frac{c_j}{c_i} + \frac{L-1-e^{-L}}{1-(1+L)e^{-L}} \right)$.

Stable coexistence is possible between genotypes that are superior in different traits. Suppose that j is better at securing territories ($c_j > c_i$), that i is better at producing propagules ($b_i > b_j$),
 324 and that $d_i = d_j$. Coexistence occurs if j will invade an i -dominated population, but i will also
 invade a j -dominated population (“mutual invasion”). If b_i is so large that $L \gg 1$ when i is
 dominant, and b_j is so small that $L \ll 1$ when j is dominant, then, combining Eqs. (7) and (8),
 327 we find that i invades j because $b_i > b_j$, while j invades i provided that

$$b_j c_j R_j - b_i c_i A_i > 0. \quad (9)$$

Thus, coexistence occurs if c_j/c_i is large enough. This is a version of the classic competition-
 colonization trade-off (Levins and Culver, 1971; Tilman, 1994): the competitor (c -specialist) leaves
 330 many territories unoccupied (low L) due to its poor colonization ability (low b), which the colo-
 nizer (b -specialist) can then exploit. A similar argument applies for coexistence between high- c
 and low- d specialists; a “competition-longevity” trade-off (Tilman, 1994). Mutual invasibility is
 333 not possible between b - and d -specialists.

Now suppose that birth and death rates vary periodically with amplitude sufficient to cause
 large changes in population density. This example is inspired by natural *Drosophila* populations,
 336 which expand rapidly in the warmer months when fruit is abundant, but largely die off in the
 colder months. Within this seasonal population density cycle, hundreds of polymorphisms also
 cycle in frequency (Bergland et al., 2014). Some of these polymorphisms may be adaptive and
 339 potentially millions of years old, suggesting stable coexistence (Bergland et al., 2014; Messer
 et al., 2016). Selection on allele frequencies thus occurs on the same time scale as population
 demography, a situation vastly more complicated than classical sweeps in demographically stable
 342 populations (Messer et al., 2016).

The classical population genetic treatment of fluctuating selection suggests that environmen-
 tal fluctuations do not promote coexistence. Allele frequencies are successively multiplied by
 345 relative fitness values for each environmental iteration, and so two alleles favored in different

environments can only stably coexist if the product of fitnesses for one type exactly equals the product for the other (Dempster, 1955). Thus, stable coexistence still requires frequency dependent selection or heterozygote advantage (as is required in a constant environment).

This classical argument overlooks two general mechanisms by which fluctuating selection promotes coexistence. The first is the “storage effect”, which introduces a form of frequency dependent selection that promotes coexistence in the presence of environmental fluctuations but not in a constant environment. The storage effect occurs when some individuals are protected from selection; in the lottery model a fraction $(1 - d_i)n_i$ of each type’s adults do not experience selection in a given iteration. Protection from selection promotes coexistence in fluctuating environments because abundant types cannot fully exploit environmental periods that favor them (since only a fraction of the rare type can be displaced), whereas rare types gain the full benefits of their favorable periods (far more adults from the abundant type die than can possibly be replaced by the rare types) (Chesson and Warner, 1981).

The second mechanism we will call the “bounded density effect”, since it is a consequence of the inhibition of reproduction at high population densities (Dempster’s (1955) argument ignores density-dependent effects). If there is growth from low to high density in each environmental cycle, then types that are abundant determine the time available for growth every cycle, with less time spent growing in cycles where they are favored, and more in cycles where they are not. This promotes coexistence even in the absence of frequency dependent selection (Yi and Dean, 2013).

Figure 6 shows the behavior of Eq. (4) for an example where b and d cycle between zero and positive values (“summers” with rapid growth and no mortality, and “winters” with mortality and no growth). Both the storage effect (adults are sheltered from selection during the summer growth phase) and the bounded density effect (expansion to high density occurs every cycle) are operating. Two types are present, a b specialist, which is better at rapidly growing in the summer (higher b), and a d specialist which is better at surviving the winter (lower d).

Neither type has an advantage over a full environmental cycle, and they stably coexist. This is due to some combination of the storage and bounded population size effects (stable coexistence

between b and d specialists was not possible in a constant environment). It is clear that the classic lottery model, which has a storage effect but no bounded population size effect, will give very different coexistence predictions from our extension of it, because population size will immediately return to capacity $N = T$ in the first summer iteration, after which type frequencies remain constant until the winter. The d specialist thus effectively has infinitely many propagules to secure its winter frequency gains, an enormous advantage compared to the finite propagule density dynamics in Fig. 6. Similar difficulties arise in previous models of how the storage effect promotes genetic variation (Ellner and Hairston Jr, 1994), which assume that the total number of offspring per iteration is constant. Beyond this observation, disentangling the storage and bounded population size effects is not straightforward, and requires a more detailed discussion of each effect than we have space for here. Our model is well suited for such a disentangling since it extends the canonical example of the storage effect to allow for density dependent effects.

Discussion

In the introduction we mentioned the recurring difficulties with confounding selection and demography in population genetic inference. While we have not directly attempted to perform inference with our model, and Eq. (4) may not be appropriate for particular inference problems, it seems that something similar (and hopefully more analytically tractable) to our extension of the lottery model is unavoidable because, fundamentally, selective births and deaths affect both abundances and frequencies, not one or the other in isolation. Moreover, some aspects of allele frequency change are intrinsically density dependent. In the classic lottery model, which as we have seen is essentially the Wright-Fisher model with overlapping generations, b_i and c_i are equivalent in the sense that the number of territorial victories only depends on the product $b_i c_i$ (see “Model”). This is no longer the case in our extension, where b and c specialists can co-exist. This “colonization-competition trade-off” is well known in the co-existence literature (Tilman, 1994). It and similar forms of “spatial co-existence” in stable environments have previously been




Figure 6: Stable coexistence between b and d specialists in a fluctuating environment. (a) Birth and death rates seasonally alternate being nonzero (white for winter, green for summer). The b specialist (black) has higher b and d ($b = 0.5$, $d = 0.2$) than the d specialist ($b = 0.2$, $d = 0.1$) (blue). (b) Both types grow during the positive b phase, and decline during the positive d phase, but the d specialist does so at a lower rate. Total height (blue+black) is population density N/T . (c) Summer favors the b specialist, winter the d specialist, and they stably coexist. For illustration, the propagule abundances are assumed to have the form $m_i = b_i(1 - N/T)n_i$, reflecting non-directed dispersal.

modeled either with Levin's qualitative representation of competition (Levins and Culver, 1971;
399 Tilman, 1994), as opposed to the quantitative c of lottery competition, or with a more sophisticated treatment of space (non-uniform dispersal; Bolker and Pacala 1999; Shmida and Ellner 1984). In cyclical environments, polymorphisms can be stabilized by the bounded density effect
402 which is completely lost if there is an exclusive focus on allele frequencies (Yi and Dean, 2013). We leave the details of how our model might be applied in inference, including the crucial issue of its genetic drift predictions, for future work.

405 It is interesting to compare the predictions of the extended lottery model with earlier approaches, such as the r/K scheme. While adaptive evolution in the direction predicted by our model does produce traits broadly consistent with Grime's scheme, trait data on selection for
408 higher b at high density was ambiguous. This prediction is also counter to the expectations of MacArthur's r/K dichotomy (MacArthur and Wilson, 1967), since b is closely related to the maximal, low-density growth rate $r = b - d$ (Pianka, 1972), yet in the r/K scheme, high density
411 populations should be subject to K , not r , selection. Yet it is not surprising that b can matter at high densities. In our model (or any lottery model of competition), b matters at high densities because territorial contests among juveniles are intrinsically unpredictable. This is a realistic
414 feature of the model. Even if one genotype is guaranteed to win a territory in a "fair" contest (e.g. it is the most efficient exploiter of a limiting consumable resource; Tilman 1982), inferior competitors can win by chance. For example, an inferior competitor's propagules may happen
417 to arrive first, gaining a decisive developmental advantage. First arrivals are more likely to occur for genotypes with a fecundity and/or dispersal advantage, as represented by higher b in lottery models. The analogous intuition in the Wright-Fisher model is that fecundity confers a relative
420 fitness advantage, even though population size is not changing. The logistic model for which r and K are named, does not capture this intuition.

Confusingly, the term " K -selection" sometimes refers generally to selection at high density
423 (Pianka, 1972), encompassing both selection for higher saturation density (MacArthur and Wilson, 1967) and competitive ability (Gill, 1974). Contrary to an r/K dichotomy, empirical studies

have shown that maximal growth rate and saturation density (measured by abundance) are positively correlated, both between species/strains (Fitzsimmons et al., 2010; Hendriks et al., 2005; Kuno, 1991; Luckinbill, 1979), and as a result of experimental evolution (Luckinbill, 1978, 1979). From the perspective of our model, this correlation is not surprising since the saturation density, which is determined by a balance between births and deaths, increases with b .

There is support for a negative relationship between competitive success at high density and maximal growth rate (Luckinbill, 1979), consistent with an r/K dichotomy. This could be driven by a tradeoff between individual size and reproductive rate. To avoid confusion with other forms of “ K -selection”, selection for competitive ability has been called “ α -selection” after the competition coefficients in the Lotka-Volterra equation (Case and Gilpin, 1974; Gill, 1974; Joshi et al., 2001). However, competitive success as measured by α (i.e. the per-capita effect of one genotype on another genotype’s growth rate) is only partly determined by individual competitive ability — in the presence of age-structured competition and territoriality, it also includes the ability of each genotype to produce contestants i.e. b in our model. Our c is strictly competitive ability only — as such, changes in c do not directly affect population density (the total number of territories occupied in an iteration is $\Delta_+ N = U(1 - e^{-L})$, which does not depend directly on the c_i). The clean separation of a strictly-relative c parameter is particularly useful from an evolutionary genetics perspective, essentially embedding a zero-sum relative fitness trait within a non-zero-sum fitness model. This could have interesting applications for modeling the impacts of intra-specific competition on species extinction, for example due to clonal interference (Desai and Fisher, 2007; Gerrish and Lenski, 1998) between c -strategists on the one hand, and b - and d - strategists on the other.

K -selection in the narrow logistic sense of selection for a greater environmental carrying capacity for given birth and death rates, sometimes referred to as “efficiency” (MacArthur and Wilson, 1967), could be represented in our model by smaller individual territorial requirements. To a first approximation, two co-occurring genotypes which differ by a small amount in their territorial requirements only should have the same fitness since the costs or benefits of a change in

the amount of unoccupied territory is shared equally among genotypes via the propagule density
453 per territory L . The situation is more complicated when the differences in territorial requirements
become large enough that territorial contests can occur on different scales for different genotypes.
We leave these complications for future work.

456 Our realization of Grime's triangle (Fig. 4) differs from approaches which identify primary
strategies as trait combinations which can co-exist (Bolker and Pacala, 1999), referring instead to
the direction of adaptive trait evolution under different regimes of stress and disturbance, which
459 is closer in spirit to Grime's arguments (Grime, 1974, 1977). In addition, we have not assumed any
kind of trade-offs or pleiotropy between b , c and d , only constraints imposed by the environment
on the order of magnitude of b and d . As an example of a trade-off, corals which rapidly out-
462 shade neighbors have a tall, branched morphology which is vulnerable to disturbances, and so,
all else being equal, ideal environment c -strategists will suffer higher mortality from disturbances
(Darling et al., 2012). Fig. 5 gives the same conclusion without invoking trade-offs; mutations
465 which reduce disturbance vulnerability are essentially neutral under ideal conditions, leading
to no improvements in mortality from disturbances, whereas c will tend to increase over time.
Thus, while trade-offs may amplify specialization, and are sometimes invoked to explain primary
468 strategy schemes (Aerts, 1999; MacArthur and Wilson, 1967; Winemiller and Rose, 1992), they are
not necessary for it.

One limitation of our model as a general-purpose model of density-dependent selection is
471 the restriction of competition to interference competition between juveniles for durable resources
(lottery recruitment to adulthood), analogous to the ubiquitous assumption of viability selection
in population genetics (Ewens, 2004, p. 45). In some respects this is the complement of resource
474 competition models, which restrict their attention to exploitation competition, typically without
age structure (Tilman, 1982). In the particular case that resources are spatially localized (e.g. due
to restricted movement through soils), resource competition and territorial acquisition effectively
477 coincide, and in principle resource competition could be represented by a competitive ability c (or
conversely, c should be derivable from resource competition). The situation is more complicated

if the resources are well-mixed, since, in general, resource levels then need to be explicitly tracked.

480 It seems plausible that explicit resource tracking may not be necessary when the focus is on the evolution of similar genotypes rather than the stable co-existence of widely differing species (Ram et al., 2016). We are not aware of any attempts to delineate conditions under which explicit
483 resource tracking is unnecessary even if it is assumed that community structure is ultimately determined by competition for consumable resources. More work is needed connecting resource competition models to the density-dependent selection literature, since most of the former has
486 to date been focused on narrower issues of the role of competition at low resource availability (Aerts, 1999; Davis et al., 1998; Tilman, 2007).

While our model can be applied to species rather than genotypes (e.g. ecological invasions),
489 our focus is genotype evolution. Our assumption that there are no large c discrepancies (section “Mean field approximation”) amounts to a restriction on the amount of genetic variation in c in the population. Since beneficial mutation effect sizes will typically not be much larger than a few
492 percent, large c discrepancies can only arise if the mutation rate is extremely large, and so the assumption will not be violated in most cases. However, this restriction could become important when looking at species interactions rather than genotype evolution.

495 **References**

Aerts, R. 1999. Interspecific competition in natural plant communities: mechanisms, trade-offs and plant-soil feedbacks. *Journal of Experimental Botany* 50:29–37.

498 Allan, J. D. 1976. Life history patterns in zooplankton. *The American Naturalist* 110:165–180.

Arenbaev, N. 1977. Asymptotic behavior of the multinomial distribution. *Theory of Probability & Its Applications* 21:805–810.

501 Barton, N. H. 1998. The effect of hitch-hiking on neutral genealogies. *Genetical Research* 72:123–133.

Bergland, A. O., E. L. Behrman, K. R. O'Brien, P. S. Schmidt, and D. A. Petrov. 2014. Genomic
504 evidence of rapid and stable adaptive oscillations over seasonal time scales in drosophila. PLOS
Genetics 10:1–19.

Bolker, B. M., and S. W. Pacala. 1999. Spatial moment equations for plant competition: Under-
507 standing spatial strategies and the advantages of short dispersal. The American Naturalist
153:575–602.

Boyce, M. S. 1984. Restitution of r-and k-selection as a model of density-dependent natural
510 selection. Annual Review of Ecology and Systematics 15:427–447.

Case, T. J., and M. E. Gilpin. 1974. Interference competition and niche theory. Proceedings of the
National Academy of Sciences 71:3073–3077.

513 Chesson, P. L., and R. R. Warner. 1981. Environmental variability promotes coexistence in lottery
competitive systems. American Naturalist pages 923–943.

Darling, E. S., L. Alvarez-Filip, T. A. Oliver, T. R. McClanahan, and I. M. Côté. 2012. Evaluating
516 life-history strategies of reef corals from species traits. Ecology Letters 15:1378–1386.

Davis, M. A., K. J. Wragge, and P. B. Reich. 1998. Competition between tree seedlings and herba-
ceous vegetation: support for a theory of resource supply and demand. Journal of Ecology
519 86:652–661.

Dempster, E. R. 1955. Maintenance of genetic heterogeneity. Cold Spring Harb Symp Quant Biol
20:25–31.

522 Desai, M. M., and D. S. Fisher. 2007. Beneficial mutation–selection balance and the effect of
linkage on positive selection. Genetics 176:1759–1798.

Diekmann, O., et al. 2004. A beginner's guide to adaptive dynamics. Banach Center Publications
525 63:47–86.

Ellner, S., and N. G. Hairston Jr. 1994. Role of overlapping generations in maintaining genetic variation in a fluctuating environment. *The American Naturalist* 143:403–417.

528 Ewens, W. J. 2004. *Mathematical Population Genetics 1: Theoretical Introduction*, vol. 27. Springer Science & Business Media.

Fitzsimmons, J. M., S. E. Schoustra, J. T. Kerr, and R. Kassen. 2010. Population consequences of
531 mutational events: effects of antibiotic resistance on the r/k trade-off. *Evolutionary Ecology* 24:227–236.

Gerrish, P. J., and R. E. Lenski. 1998. The fate of competing beneficial mutations in an asexual
534 population. *Genetica* 102:127–144.

Gill, D. E. 1974. Intrinsic rate of increase, saturation density, and competitive ability. ii. the evolution of competitive ability. *American Naturalist* 108:103–116.

537 Grime, J. P. 1974. Vegetation classification by reference to strategies. *Nature* 250:26–31.

———. 1977. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *American Naturalist* 111:1169–1194.

540 ———. 1988. *Plant Evolutionary Biology*, chap. The C-S-R model of primary plant strategies — origins, implications and tests, pages 371–393. Springer Netherlands, Dordrecht.

Hendriks, A. J., J. L. Maas-Diepeveen, E. H. Heugens, and N. M. van Straalen. 2005. Meta-analysis
543 of intrinsic rates of increase and carrying capacity of populations affected by toxic and other stressors. *Environmental Toxicology and Chemistry* 24:2267–2277.

Joshi, A., N. Prasad, and M. Shakarad. 2001. K-selection, α -selection, effectiveness, and tolerance
546 in competition: density-dependent selection revisited. *Journal of Genetics* 80:63–75.

Knowlton, N. 2001. The future of coral reefs. *Proceedings of the National Academy of Sciences of the United States of America* 98:5419–5425.

- 549 Kuno, E. 1991. Some strange properties of the logistic equation defined with r and K : Inherent defects or artifacts? *Researches on Population Ecology* 33:33–39.
- Levins, R., and D. Culver. 1971. Regional coexistence of species and competition between rare
552 species. *Proceedings of the National Academy of Sciences* 68:1246–1248.
- Luckinbill, L. S. 1978. r and k selection in experimental populations of *Escherichia coli*. *Science* (New York, NY) 202:1201–1203.
- 555 ———. 1979. Selection and the r/k continuum in experimental populations of protozoa. *American Naturalist* pages 427–437.
- MacArthur, R. H. 1962. Some generalized theorems of natural selection. *Proceedings of the*
558 *National Academy of Sciences* 48:1893–1897.
- MacArthur, R. H., and E. O. Wilson. 1967. *Theory of Island Biogeography*. Princeton University Press.
- 561 Messer, P. W., S. P. Ellner, and N. G. Hairston. 2016. Can population genetics adapt to rapid evolution? *Trends in Genetics* 32:408–418.
- Metz, J. A., R. M. Nisbet, and S. A. Geritz. 1992. How should we define ‘fitness’ for general
564 ecological scenarios? *Trends in Ecology & Evolution* 7:198–202.
- Pianka, E. R. 1970. On r - and K -Selection. *The American Naturalist* 104:592–597.
- . 1972. r and K selection or b and d selection? *The American Naturalist* 106:581–588.
- 567 Ram, Y., E. Dellus-Gur, M. Bibi, U. Obolski, J. Berman, and L. Hadany. 2016. Predicting microbial relative growth in a mixed culture from growth curve data. *bioRxiv* .
- Reznick, D., M. J. Bryant, and F. Bashey. 2002. r - and k -selection revisited: The role of population
570 regulation in life-history evolution. *Ecology* 83:1509–1520.

Sale, P. F. 1977. Maintenance of high diversity in coral reef fish communities. *The American Naturalist* 111:337–359.

573 Shmida, A., and S. Ellner. 1984. Coexistence of plant species with similar niches. *Vegetatio* 58:29–55.

Southwood, T. R. E. 1977. Habitat, the templet for ecological strategies? *Journal of Animal Ecology* 46:337–365.

Stearns, S. C. 1977. The evolution of life history traits: A critique of the theory and a review of the data. *Annual Review of Ecology and Systematics* 8:145–171.

579 Svoldal, H., C. Rueffler, and J. Hermisson. 2015. A general condition for adaptive genetic polymorphism in temporally and spatially heterogeneous environments. *Theoretical Population Biology* 99:76 – 97.

582 Taylor, D. R., L. W. Aarssen, and C. Loehle. 1990. On the relationship between r/k selection and environmental carrying capacity: a new habitat templet for plant life history strategies. *Oikos* pages 239–250.

585 Tilman, D. 1982. Resource competition and community structure. 17. Princeton University Press.

———. 1994. Competition and biodiversity in spatially structured habitats. *Ecology* 75:2–16.

———. 2007. Resource competition and plant traits: a response to craine et al. 2005. *Journal of Ecology* 95:231–234.

Uecker, H., and J. Hermisson. 2011. On the fixation process of a beneficial mutation in a variable environment. *Genetics* 188:915–930.

591 Violle, C., M.-L. Navas, D. Vile, E. Kazakou, C. Fortunel, I. Hummel, and E. Garnier. 2007. Let the concept of trait be functional! *Oikos* 116:882–892.

Westoby, M. 1998. A leaf-height-seed (LHS) plant ecology strategy scheme. *Plant and soil*
199:213–227.

Winemiller, K. O., D. B. Fitzgerald, L. M. Bower, and E. R. Pianka. 2015. Functional traits, convergent evolution, and periodic tables of niches. *Ecology Letters* 18:737–751.

Winemiller, K. O., and K. A. Rose. 1992. Patterns of life-history diversification in north american fishes: implications for population regulation. *Canadian Journal of Fisheries and Aquatic Sciences* 49:2196–2218.

Yi, X., and A. M. Dean. 2013. Bounded population sizes, fluctuating selection and the tempo and mode of coexistence. *Proceedings of the National Academy of Sciences* 110:16945–16950.

Appendix A: Poisson approximation

For simplicity of presentation, we have assumed a Poisson distribution for the x_i as our model of dispersal. Strictly speaking, the total number of i propagules $\sum x_i$ (summed over unoccupied territories) is then no longer a constant m_i , but fluctuates between generations for a given mean m_i , which is more biologically realistic. Nevertheless, since we do not consider the random fluctuations in type abundances here, and for ease of comparison with the classic lottery model, we ignore the fluctuations in m_i . Instead we focus, on Poisson fluctuations in propagule composition in each territory.

In the exact model of random dispersal, the counts of a genotype’s propagules across unoccupied territories follows a multinomial distribution with dimension U , total number of trials equal to m_i , and equal probabilities $1/U$ for a propagule to land in a given territory. Thus, the x_i in different territories are not independent random variables. However, for sufficiently large U and m_i , this multinomial distribution for the x_i across territories is closely approximated by a product of independent Poisson distributions for each territory, each with rate parameter l_i (Arenbaev, 1977, Theorem 1). Since we are ignoring finite population size effects, we effectively

have $T \rightarrow \infty$, in which case U can be only be small enough to violate the Poisson approximation
 618 if there is vanishing population turnover, and then the dispersal distribution is irrelevant any-
 way. Likewise, in ignoring stochastic finite population size for the n_i , we have effectively already
 assumed that m_i is large enough to justify the Poisson approximation (the error scales as $1/\sqrt{m_i}$;
 621 Arenbaev 1977).

Appendix B: Derivation of growth equation

We separate the right hand side of Eq. (2) into three components $\Delta_+ n_i = \Delta_u n_i + \Delta_r n_i + \Delta_a n_i$ which
 624 vary in relative magnitude depending on the propagule densities l_i . Following the notation in
 the main text, the Poisson distributions for the x_i (or some subset of the x_i) will be denoted p ,
 and we use P as a general shorthand for the probability of particular outcomes.

627 Growth without competition

The first component, $\Delta_u n_i$, accounts for territories where only one focal propagule is present
 $x_i = 1$ and $x_j = 0$ for $j \neq i$ (u stands for “uncontested”). The proportion of territories where this
 630 occurs is $l_i e^{-L}$, and so

$$\Delta_u n_i = U l_i e^{-L} = m_i e^{-L}. \quad (10)$$

Competition when rare

The second component, $\Delta_r n_i$, accounts for territories where a single focal propagule is present
 633 along with at least one non-focal propagule (r stands for “rare”) i.e. $x_i = 1$ and $X_i \geq 1$ where
 $X_i = \sum_{j \neq i} x_j$ is the number of nonfocal propagules. The number of territories where this occurs
 is $U p_i(1) P(X_i \geq 1) = b_i n_i e^{-l_i} (1 - e^{-(L-l_i)})$. Thus

$$\Delta_r n_i = m_i e^{-l_i} (1 - e^{-(L-l_i)}) \left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\bar{p}}, \quad (11)$$

636 where $\langle \rangle_{\tilde{p}}$ denotes the expectation with respect to \tilde{p} , and \tilde{p} is the probability distribution of
 nonfocal propagule abundances x_j *after* dispersal, in those territories where exactly one focal
 propagule, and at least one non-focal propagule, landed.

639 Our “mean field” approximation is to replace x_j with its mean in the last term in Eq. (11),

$$\left\langle \frac{c_i}{c_i + \sum_{j \neq i} c_j x_j} \right\rangle_{\tilde{p}} \approx \frac{c_i}{c_i + \sum_{j \neq i} c_j \langle x_j \rangle_{\tilde{p}}}. \quad (12)$$

Below we justify this replacement by arguing that the standard deviation $\sigma_{\tilde{p}}(\sum_{j \neq i} c_j x_j)$ (with respect to \tilde{p}), is much smaller than $\langle \sum_{j \neq i} c_j x_j \rangle_{\tilde{p}}$.

We first calculate $\langle x_j \rangle_{\tilde{p}}$. Let $X = \sum_j x_j$ denote the total number of propagules in a territory and $\mathbf{x}_i = (x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_G)$ denote the vector of non-focal abundances, so that $p(\mathbf{x}_i) = p_1(x_1) \dots p_{i-1}(x_{i-1}) p_{i+1}(x_{i+1}) \dots p_G(x_G)$. Then, \tilde{p} can be written as

$$\begin{aligned} \tilde{p}(\mathbf{x}_i) &= p(\mathbf{x}_i | X \geq 2, x_i = 1) \\ &= \frac{P(\mathbf{x}_i, X \geq 2 | x_i = 1)}{P(X \geq 2)} \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{X=2}^{\infty} P(X) p(\mathbf{x}_i | X_i = X - 1), \end{aligned} \quad (13)$$

and so

$$\begin{aligned} \langle x_j \rangle_{\tilde{p}} &= \sum_{\mathbf{x}_i} \tilde{p}(\mathbf{x}_i) x_j \\ &= \frac{1}{1 - (1 + L)e^{-L}} \sum_{X=2}^{\infty} P(X) \sum_{\mathbf{x}_i} p(\mathbf{x}_i | X_i = X - 1) x_j. \end{aligned} \quad (14)$$

The inner sum over \mathbf{x}_i is the mean number of propagules of a given nonfocal type j that will be found in a territory which received $X - 1$ nonfocal propagules in total, which is equal to

$\frac{l_j}{L-l_i}(X-1)$. Thus,

$$\begin{aligned}\langle x_j \rangle_{\tilde{p}} &= \frac{l_j}{1 - (1+L)e^{-L}} \frac{1}{L-l_i} \sum_{k=2}^{\infty} P(X)(X-1) \\ &= \frac{l_j}{1 - (1+L)e^{-L}} \frac{L-1+e^{-L}}{L-l_i},\end{aligned}\tag{15}$$

642 where the last line follows from $\sum_{X=2}^{\infty} P(X)(X-1) = \sum_{X=1}^{\infty} P(X)(X-1) = \sum_{X=1}^{\infty} P(X)X - \sum_{X=1}^{\infty} P(X)$.

The exact analysis of the fluctuations in $\sum_{j \neq i} c_j x_j$ is complicated because the x_j are not independent with respect to \tilde{p} . These fluctuations are part of the “drift” in type abundances which we leave for future work. Here we use the following approximation to give some insight into the magnitude of these fluctuations and also the nature of the correlations between the x_j . We replace \tilde{p} with \tilde{q} , defined as the \mathbf{x}_i Poisson dispersal probabilities conditional on $X_i \geq 1$ (which are independent). The distinction between \tilde{p} with \tilde{q} will be discussed further below. The \tilde{q} approximation gives $\langle x_j \rangle_{\tilde{q}} = \langle x_j \rangle_p / C = l_j / C$,

$$\begin{aligned}\sigma_{\tilde{q}}^2(x_j) &= \langle x_j^2 \rangle_{\tilde{q}} - \langle x_j \rangle_{\tilde{q}}^2 \\ &= \frac{1}{C} \langle x_j^2 \rangle_p - \frac{l_j^2}{C^2} \\ &= \frac{1}{C} (l_j^2 + l_j) - \frac{l_j^2}{C^2} \\ &= \frac{l_j^2}{C} \left(1 - \frac{1}{C}\right) + \frac{l_j}{C},\end{aligned}\tag{16}$$

and

$$\begin{aligned}\sigma_{\tilde{q}}(x_j, x_k) &= \langle x_j x_k \rangle_{\tilde{q}} - \langle x_j \rangle_{\tilde{q}} \langle x_k \rangle_{\tilde{q}} \\ &= \frac{1}{C} \langle x_j x_k \rangle_p - \frac{l_j l_k}{C^2} \\ &= \frac{l_j l_k}{C} \left(1 - \frac{1}{C}\right),\end{aligned}\tag{17}$$

where $C = 1 - e^{-(L-l_i)}$ and $j \neq k$.

645 The exact distribution \tilde{p} assumes that exactly one of the propagules present in a given site after dispersal belongs to the focal type, whereas \tilde{q} assumes that there is a focal propagule present before non-focal dispersal commences. As a result, \tilde{q} predicts that the mean propagule
648 density is greater than L (in sites with only one focal propagule is present) when the focal type is rare and the propagule density is high. This is erroneous, because the mean number of propagules in every site is L by definition. Specifically, if $L - l_i \approx L \gg 1$, then the mean
651 propagule density predicted by \tilde{q} is approximately $L + 1$. The discrepancy causes rare invaders to have an intrinsic rarity disadvantage (territorial contests under \tilde{q} are more intense than they should be). In contrast, Eq. (15) correctly predicts that there are on average $\sum_{j \neq i} \langle x_j \rangle_{\tilde{p}} \approx L - 1$
654 nonfocal propagules because \tilde{p} accounts for potentially large negative covariances between the x_j “after dispersal”. By neglecting the latter covariences, \tilde{q} overestimates the fluctuations in $\sum_{j \neq i} c_j x_j$; thus \tilde{q} gives an upper bound on the fluctuations. The discrepancy between \tilde{q} and \tilde{p} will be largest
657 when L is of order 1 or smaller, because then the propagule assumed to already be present under \tilde{q} is comparable to, or greater than, the entire propgaule density.

Decomposing the variance in $\sum_{j \neq i} c_j x_j$,

$$\sigma_{\tilde{q}}^2(\sum_{j \neq i} c_j x_j) = \sum_{j \neq i} \left[c_j^2 \sigma_{\tilde{q}}^2(x_j) + 2 \sum_{k > j, k \neq i} c_j c_k \sigma_{\tilde{q}}(x_j, x_k) \right], \quad (18)$$

660 and using the fact that $\sigma_{\tilde{q}}(x_j, x_k)$ and the first term in Eq. (16) are negative because $C < 1$, we obtain an upper bound on the relative fluctuations in $\sum_{j \neq i} c_j x_j$,

$$\frac{\sigma(\sum_{j \neq i} c_j x_j)}{\langle \sum_{j \neq i} c_j x_j \rangle} = C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j + (1 - 1/C) \left(\sum_{j \neq i} c_j l_j \right)^2 \right)^{1/2}}{\sum_{j \neq i} c_j l_j} < C^{1/2} \frac{\left(\sum_{j \neq i} c_j^2 l_j \right)^{1/2}}{\sum_{j \neq i} c_j l_j}. \quad (19)$$

Suppose that the c_j are all of similar magnitude (their ratios are of order one). Then Eq. (19)
663 is $\ll 1$ for the case when $L - l_i \ll 1$ (due to the factor of $C^{1/2}$), and also for the case when at least some of the nonfocal propagule densities are large $l_j \gg 1$ (since it is then of order $1/\sqrt{L - l_i}$).

The worst case scenario occurs when $L - l_i$ is of order one. Then Eq. (19) gives a relative error of
 666 approximately 50%, which from our earlier discussion we know to be a substantial overestimate
 when L is of order 1. Our numerical results (Fig. 3) confirm that the relative errors are indeed
 small.

669 However, the relative fluctuations in $\sum_{j \neq i} c_j x_j$ can be large if some of the c_j are much larger
 than the others. Specifically, in the presence of a rare, extremely strong competitor ($c_j l_j \gg c_{j'} l_{j'}$
 for all other nonfocal genotypes j' , and $l_j \ll 1$), then the RHS of Eq. (19) can be large and we
 672 cannot make the replacement Eq. (12).

Substituting Eqs. (12) and (15) into Eq. (11), we obtain

$$\Delta_r n_i \approx m_i R_i \frac{c_i}{\bar{c}}, \quad (20)$$

where R_i is defined in Eq. (5).

675 **Competition when abundant**

The final contribution, $\Delta_a n_i$, accounts for territories where two or more focal propagules are
 present (a stands for “abundant”). Similarly to Eq. (11), we have

$$\Delta_a n_i = U(1 - (1 + l_i)e^{l_i}) \left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \quad (21)$$

678 where \hat{p} is the probability distribution of both focal and nonfocal propagaule abundances *after*
 dispersal in those territories where at least two focal propagules landed.

Again, we argue that the relative fluctuations in $\sum c_j x_j$ are much smaller than 1 (with respect
 681 to \hat{p}), so that,

$$\left\langle \frac{c_i x_i}{\sum_j c_j x_j} \right\rangle_{\hat{p}} \approx \frac{c_i \langle x_i \rangle_{\hat{p}}}{\sum_j c_j \langle x_j \rangle_{\hat{p}}}. \quad (22)$$

Following a similar procedure as for $\Delta_r n_i$, where the vector of propagule abundances is denoted

\mathbf{x} , the mean focal genotype abundance is,

$$\begin{aligned}
\langle x_i \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_i p(\mathbf{x} | x_i \geq 2) \\
&= \sum_{x_i} x_i p(x_i | x_i \geq 2) \\
&= \frac{1}{1 - (1 + l_i)e^{-l_i}} \sum_{x_i \geq 2} p(x_i) x_i \\
&= l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}}.
\end{aligned} \tag{23}$$

For nonfocal genotypes $j \neq i$, we have

$$\begin{aligned}
\langle x_j \rangle_{\hat{p}} &= \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2) \\
&= \sum_X P(X | x_i \geq 2) \sum_{\mathbf{x}} x_j p(\mathbf{x} | x_i \geq 2, X) \\
&= \sum_X P(X | x_i \geq 2) \sum_{x_i} p(x_i | x_i \geq 2, X) \sum_{\mathbf{x}_i} x_j p(\mathbf{x}_i | X_i = X - x_i) \\
&= \sum_X P(X | x_i \geq 2) \sum_{x_i} p(x_i | x_i \geq 2, X) \frac{l_j(X - x_i)}{L - l_i} \\
&= \frac{l_j}{L - l_i} \left[\sum_X P(X | x_i \geq 2) X - \sum_{x_i} p(x_i | x_i \geq 2) x_i \right] \\
&= \frac{l_j}{L - l_i} \left(L \frac{1 - e^{-L}}{1 - (1 + L)e^{-L}} - l_i \frac{1 - e^{-l_i}}{1 - (1 + l_i)e^{-l_i}} \right).
\end{aligned} \tag{24}$$

To calculate the relative fluctuations in $\sum_{j \neq i} c_j x_j$, we use a similar approximation as for $\Delta_r n_i$: \hat{p} is approximated by \hat{q} , defined as the \mathbf{x} dispersal probabilities in a territory conditional on $x_i > 2$ (that is, treating the x_j as independent). All covariances between nonfocal genotypes are now zero, so that $\sigma_{\hat{q}}^2(\sum c_j x_j) = \sum c_j^2 \sigma_{\hat{q}}^2(x_j)$, where $\sigma_{\hat{q}}^2(x_j) = l_j$ for $j \neq i$, and

$$\sigma_{\hat{q}}^2(x_i) = \frac{l_i}{D} \left(l_i + 1 - e^{-l_i} - \frac{l_i}{D} (1 - e^{-l_i})^2 \right), \tag{25}$$

where $D = 1 - (1 + l_i)e^{-l_i}$, and

$$\frac{\sigma_{\hat{q}}(\sum c_j x_j)}{\langle \sum c_j x_j \rangle} = \frac{\left(\sum_{j \neq i} c_j^2 l_j + c_i^2 \sigma_{\hat{q}}^2(x_i) \right)^{1/2}}{\sum_{j \neq i} c_j l_j + c_i l_i (1 - e^{-l_i}) / D}. \quad (26)$$

687 Similarly to Eq. (19), the RHS of Eq. (26) is $\ll 1$ for the case that $L \ll 1$ (due to a factor of $D^{1/2}$), and also for the case when at least some of the propagule densities (focal or nonfocal) are large — provided that c_i and the c_j are all of similar magnitude. Again, the worst case scenario
690 occurs when l_i and $L - l_i$ are of order 1, in which case Eq. (26) is around 35%, which is again where the \hat{q} approximation produces the biggest overestimate of the fluctuations in \mathbf{x} . Similarly to Eq. (19), the RHS of (26) will not be $\ll 1$ in the presence of a rare, extremely strong competitor.

693 Combining Eqs. (21) and (22), we obtain

$$\Delta_a n_i = m_i A_i \frac{c_i}{\bar{c}}, \quad (27)$$

where A_i is defined in Eq. (6).