Estimating Same-Source Probabilities from Value Collisions with Unknown Latent Emission Graphs

Jason Bono
Lucor (https://lucor.ai)
jasonbono@lucor.ai
ORCID: 0000-0002-3018-714X
February 2026

Abstract

When two observations share a value, how likely is it that they came from the same latent source? We give a closed-form estimator for this *collision-conditioned same-source probability* that requires only the observed value frequencies and a single scalar summary of source heterogeneity—the across-source degree variance—with no knowledge of the full source–value emission structure.

The setting is a bipartite multigraph whose hidden bottom vertices are latent entities and whose top vertices are observable value labels; a measurement samples an edge uniformly and returns the value endpoint. Under a maximum-entropy (configuration-model) ensemble over graphs consistent with the observed value marginals, we prove that the ensemble mean of the exact same-source posterior depends on the unknown entity side only through the entity-degree second moment, equivalently the degree variance. We also derive an analytic expression for the ensemble variance (uncertainty) of the posterior, which depends on one additional degree-shape statistic. We validate both formulas by exact enumeration in small cases and by end-to-end simulations that sample ground-truth graphs, estimate value marginals from finite data, and compare the true posterior to the estimator's mean and uncertainty bands.

Keywords: same-source probability, value collision, bipartite configuration model, maximum-entropy ensemble, record linkage, entity resolution

1 Introduction

Many systems observe *values* but not the *entities* that produced them. Examples include repeated measurements of latent sources, anonymized logs, records produced by hidden actors, or sensors that emit coarse labels. Even when a value repeats (a "collision"), it may reflect either repeated emission from the same latent entity or emissions from different entities that share common values. Quantifying this ambiguity is useful for debugging pipelines, designing downstream inference, and understanding privacy and identifiability properties.

This paper considers a stylized but broadly applicable model: latent entities emit value labels with different intensities. The (unobserved) emission spectrum is encoded as a bipartite multigraph with integer edge multiplicities. A measurement draw samples an edge uniformly and returns the value label. We focus on the simplest nontrivial question: given two draws (with replacement), what is the posterior probability that they came from the same latent entity, especially conditioned on observing a value collision?

When the full emission graph is known, the posterior is computable exactly. The challenging regime is when the full graph is not known. We assume the observer can estimate the value-label marginal distribution (equivalently, value-side degrees up to scale) and that domain knowledge

provides a summary of entity-side heterogeneity—in this paper, the across-entity degree variance. We then:

- introduce a maximum-entropy bipartite graph ensemble (configuration model) consistent with the observed value marginals,
- prove that the ensemble mean of the exact posterior given a collision is determined solely by the entity-degree second moment (variance),
- derive an analytic expression for the ensemble variance (uncertainty) across graphs,
- validate by exact enumeration in small toy examples, and
- demonstrate end-to-end performance: sampling ground-truth graphs, simulating measurements, estimating value marginals from finite data, and comparing the true posterior to the estimator's mean and uncertainty bands.

2 Related work

Record linkage and the Fellegi–Sunter model. Fellegi and Sunter [14] cast record linkage as a likelihood-ratio test between a *match* hypothesis (same source) and a *non-match* hypothesis (different source), leading to the familiar m- and u-probabilities for field agreement: $m = P(\text{agree} \mid \text{match})$ and $u = P(\text{agree} \mid \text{non-match})$. In our setting with a single equality comparator, the "agree" event is exactly the collision event $\mathcal{C} = \{V_1 = V_2\}$. The Fellegi–Sunter posterior for an agreement,

$$P(\text{match} \mid \text{agree}) = \frac{m\,\pi}{m\,\pi + u\,(1-\pi)},$$

maps directly to our target $\theta(C) = P(S \mid \mathcal{C})$, with prior $\pi = P(S)$. A key difference is that classical linkage frameworks typically treat (m, u, π) as free parameters and estimate them from labeled data or via EM [15, 16]; later statistical models refine the probabilistic foundations [17]. Here we instead start from a generative emission graph model, and derive $\theta(C)$ exactly when C is known (Proposition 1), and in closed form *in expectation* when C is unknown under a maximum-entropy (configuration-model) ensemble (Theorem 1).

Entity resolution and duplicate detection. Entity resolution and duplicate detection aim to infer the full partition or matching structure among records, often leveraging multiple noisy fields and transitive/collective constraints. Surveys and textbooks in the database and data-mining communities emphasize the end-to-end ER pipeline, including blocking/indexing, similarity computation, and clustering/classification [23, 24, 25, 26, 27]. Collective ER models exploit relational structure and dependencies across decisions [28]. Recent Bayesian approaches place priors directly on the latent partition or bipartite matching [18, 19]. In contrast, our goal is a single scalar summary—the collision-conditioned same-source probability—and we derive it analytically under a principled random-graph null (configuration model) consistent with observed value marginals. This scalar can be interpreted as the *precision* of an equality-based blocking key: among colliding values, what fraction of collisions are truly same-source?

Collision probabilities, concentration, and the "birthday" phenomenon. The probability that two draws from a discrete distribution collide is the classical birthday problem and is closely related to concentration indices such as Simpson's concentration (and the Herfindahl index) [1, 20]. Our setting introduces a second layer: values are generated via an unobserved bipartite emission structure, so the same-source probability *given* a collision depends on both value-side and entity-side heterogeneity. Under the configuration-model ensemble, we show that the ensemble mean of this conditional probability depends on the entity side only through the degree second moment (Theorem 1).

Privacy and disclosure risk. Collisions of quasi-identifier values are central to disclosure risk: if a released key value is rare or unique, it can enable re-identification [21], and uniqueness of partial identifiers has been quantified in several domains. De-anonymization attacks show that sparse/high-dimensional keys can be re-linked to individuals even without explicit identifiers [22]. Our $\theta(C) = P(S \mid \mathcal{C})$ addresses a complementary question to group-size metrics such as k-anonymity: *given that a released key value repeats, how likely is it that the repetition is due to the same individual/source rather than different individuals sharing the key?* This provides an interpretable "collision informativeness" statistic that can be used for debugging linkage pipelines or for characterizing match risk under repeated values.

3 Model and exact posteriors when the graph is known

3.1 Latent emission spectrum as a bipartite multigraph

Let $e_1, \ldots, e_n$ denote latent entities and $v_1, \ldots, v_m$ denote value labels. The emission spectrum is represented by a nonnegative integer incidence matrix

$$C = (c_{ia}) \in \mathbb{Z}_{\geq 0}^{n \times m},$$

where c_{ia} is the number of edges between entity e_i and value v_a.

Define entity degrees and value degrees:

$$\ell_i := \sum_{a=1}^{m} c_{ia}, \qquad o_a := \sum_{i=1}^{n} c_{ia}, \qquad T := \sum_{a=1}^{m} o_a = \sum_{i=1}^{n} \ell_i.$$

We interpret T as the total number of edge "stubs" (total emission mass).

3.2 Measurement draws

A single measurement draw samples one edge uniformly at random from the multigraph and returns the value label endpoint. Equivalently: sample an entity-value edge uniformly, observe only the value endpoint. Consider two draws with replacement. Let $E_1, E_2 \in \{e_1, \ldots, e_n\}$ be the latent emitting entities and $V_1, V_2 \in \{v_1, \ldots, v_m\}$ the observed values. Define events:

$$S := \{E_1 = E_2\} \quad \text{(same latent source)}, \qquad \mathcal{C} := \{V_1 = V_2\} \quad \text{(value collision)}.$$



Figure 1: Emission-spectrum bipartite multigraph. Bottom nodes are latent entities (e_1, e_2, e_3); top nodes are value labels (v_1, v_2, v_3). Each edge represents one unit of emission mass; parallel edges encode higher emission strength (e.g., e_1 emits v_1 with multiplicity 3). A measurement draw samples one edge uniformly at random and returns the value endpoint.

3.3 Exact posteriors for a known incidence matrix

The following identities are elementary but form the foundation for the unknown-graph analysis.

Proposition 1 (Known-graph posteriors)**.** *Assume C is known and draws are with replacement. Then for any (a, b) with $o_a, o_b > 0$,*

$$P(S \mid V_1 = v_a, V_2 = v_b) = \frac{\sum_{i=1}^{n} c_{ia} c_{ib}}{o_a o_b}.$$

In particular,

$$P(S \mid V_1 = v_a, V_2 = v_a) = \frac{\sum_{i=1}^{n} c_{ia}^2}{o_a^2}.$$

Conditioned only on the event of a collision,

$$\boxed{P(S \mid \mathcal{C}) = \frac{\sum_{a=1}^{m} \sum_{i=1}^{n} c_{ia}^2}{\sum_{a=1}^{m} o_a^2}.}$$

Proof. For a single draw, $P(E = e_i, V = v_a) = c_{ia}/T$, hence $P(V = v_a) = o_a/T$ and

$$P(E = e_i \mid V = v_a) = \frac{P(E = e_i, V = v_a)}{P(V = v_a)} = \frac{c_{ia}/T}{o_a/T} = \frac{c_{ia}}{o_a}.$$

Because the two draws are independent (with replacement),

$$P(S \mid V_1 = v_a, V_2 = v_b) = \sum_{i=1}^{n} P(E_1 = e_i \mid V_1 = v_a)\, P(E_2 = e_i \mid V_2 = v_b) = \sum_{i=1}^{n} \frac{c_{ia}}{o_a} \frac{c_{ib}}{o_b},$$

which gives the first identity. The collision-at-v_a case is $a = b$.

For the collision-only posterior, write $P(S \mid \mathcal{C}) = P(S \cap \mathcal{C})/P(\mathcal{C})$. Under with-replacement edge sampling,

$$P(S \cap \mathcal{C}) = \sum_{i=1}^{n} \sum_{a=1}^{m} P(E_1 = e_i, V_1 = v_a)\, P(E_2 = e_i, V_2 = v_a) = \sum_{i,a} \left(\frac{c_{ia}}{T}\right)^2 = \frac{\sum_{i,a} c_{ia}^2}{T^2}.$$

Similarly,

$$P(\mathcal{C}) = \sum_{a=1}^{m} P(V_1 = v_a)P(V_2 = v_a) = \sum_a \left(\frac{o_a}{T}\right)^2 = \frac{\sum_a o_a^2}{T^2}.$$

Taking the ratio yields $P(S \mid \mathcal{C}) = \left(\sum_{i,a} c_{ia}^2\right) / \left(\sum_a o_a^2\right)$. □

We will write

$$\theta(C) := P(S \mid \mathcal{C}) = \frac{N(C)}{D}, \qquad N(C) := \sum_{i,a} c_{ia}^2, \qquad D := \sum_a o_a^2.$$

When C is unknown, $\theta(C)$ becomes a random variable over plausible graphs.

4 Unknown-graph regime and maximum-entropy graph ensemble

4.1 Observer information

In the unknown-graph regime, the observer does not know C (and typically does not know ℓ). We assume the observer knows:

- the number of *active emitters* n (entities that exist and emit),
- the value-side degrees $o = (o_1, \ldots, o_m)$ (or equivalently the marginal distribution $P(V = v_a) = o_a/T$),
- and a domain-knowledge parameter describing entity heterogeneity: the across-entity degree variance σ^2.

For active emitters, we assume $\ell_i \geq 1$ for all i, hence $T \geq n$.

Let

$$\mu := \frac{1}{n}\sum_{i=1}^{n} \ell_i = \frac{T}{n}, \qquad \sigma^2 := \frac{1}{n}\sum_{i=1}^{n} (\ell_i - \mu)^2.$$

Then the second moment is fixed:

$$\sum_{i=1}^{n} \ell_i^2 = n(\mu^2 + \sigma^2) = \frac{T^2}{n} + n\sigma^2.$$

Define the normalized concentration statistic (Simpson concentration of the entity weights ℓ_i/T) [1]:

$$\alpha := \sum_{i=1}^{n} \left(\frac{\ell_i}{T}\right)^2 = \frac{\sum_i \ell_i^2}{T^2} = \boxed{\frac{1}{n} + \frac{n\sigma^2}{T^2}.}$$

Sanity check: α is the unconditional same-source probability. Before conditioning on collisions, α already has a direct probabilistic meaning.

Proposition 2 (Sanity check: unconditional same-source probability)**.** *Under the measurement model with replacement,*

$$\boxed{P(S) = \sum_{i=1}^{n} \left(\frac{\ell_i}{T}\right)^2 = \alpha.}$$

Proof. Each draw selects entity i with probability ℓ_i/T, independently across draws, hence $P(S) = \sum_i (\ell_i/T)^2 = \alpha$. □

4.2 Maximum-entropy conditional model: bipartite configuration model

To obtain a distribution over unknown graphs, we adopt a conditional model commonly called the bipartite configuration model.

Definition 1 (Bipartite configuration model)**.** Fix degree sequences $\ell = (\ell_1, \ldots, \ell_n)$ and $o = (o_1, \ldots, o_m)$ with $\sum_i \ell_i = \sum_a o_a = T$. Create ℓ_i stubs on entity i and o_a stubs on value a. Match all stubs uniformly at random among all perfect matchings. The induced random contingency table $C = (c_{ia})$ is a bipartite multigraph with those degrees.

Maximum-entropy clarification (microcanonical on matchings). When we say "maximum entropy" in this paper, we mean the following precise statement: *conditional on the hard constraints imposed by the degree sequences* (i.e., the specified multisets of stubs on each side), we place the *uniform* distribution on the finite set of perfect stub matchings (microstates). This is the maximum Shannon entropy distribution over matchings subject to those hard constraints, and it induces (via many-to-one mapping) a *non-uniform* distribution over contingency tables C. This is the standard configuration-model ensemble for graphs with fixed degrees; see, e.g., [2, 3, 4, 6, 5].

It implies a probability mass function on tables:

$$P(C \mid \ell, o) \propto \frac{1}{\prod_{i,a} c_{ia}!},$$

because many stub matchings correspond to the same matrix C (see, e.g., [6, 7]).

What "mean over all graphs" means here. All expectations in this paper are taken with respect to the configuration-model distribution (uniform stub matchings), not a uniform distribution over adjacency matrices and not an adversarial/worst-case model.

5 Main result: a variance-only point estimator equals the ensemble mean

The exact posterior given a collision is $\theta(C) = N(C)/D$, with $D = \sum_a o_a^2$ fixed given o. So to compute $\mathbb{E}[\theta(C) \mid \ell, o]$ it suffices to compute $\mathbb{E}[N(C) \mid \ell, o]$.

5.1 Collision probabilities without replacement

A key quantity under uniform stub matching is the probability that *two distinct value stubs* share a label. Define the falling factorial $(x)_k := x(x-1)\cdots(x-k+1)$. Then (for $T \geq 2$)

$$p_2 := P(A_1 = A_2) = \sum_{a=1}^{m} \frac{o_a}{T} \frac{o_a - 1}{T - 1} = \frac{\sum_a (o_a)_2}{(T)_2} = \frac{D - T}{T(T-1)}.$$

(We emphasize: p_2 is *without replacement* on stubs. It differs from the with-replacement collision probability $p_{\text{coll}} = \sum_a (o_a/T)^2 = D/T^2$.)

Define the entity-side ordered-pair count:

$$M_2 := \sum_{i=1}^{n} (\ell_i)_2 = \sum_i \ell_i(\ell_i - 1).$$

Note $M_2 = \sum_i \ell_i^2 - T = T^2\alpha - T$.

Theorem 1 (Ensemble mean depends only on α (degree variance))**.** *Under the bipartite configuration model,*

$$\boxed{\mathbb{E}[\theta(C) \mid \ell, o] = \frac{T + M_2\, p_2}{D}.}$$

Equivalently, using $\alpha = \sum_i (\ell_i/T)^2$*,*

$$\boxed{\mathbb{E}[\theta(C) \mid o, \alpha] = \frac{T + (T^2\alpha - T)\, p_2}{D}, \qquad p_2 = \frac{D - T}{T(T-1)}.}$$

If the observer specifies the degree variance σ^2 *(hence* $\alpha = \frac{1}{n} + \frac{n\sigma^2}{T^2}$*), this yields a closed-form point estimator using only* (o, n, σ^2)*.*

Interpretation (model-based estimator). For fixed value marginals o and fixed entity-degree second moment (equivalently fixed σ^2), the configuration-model ensemble mean of the *exact* posterior $\theta(C) = P(S \mid \mathcal{C})$ is uniquely determined. The variance-based estimator is therefore not an ad hoc approximation but exactly the ensemble mean under the uniform-stub-matching (maximum-entropy-over-matchings) graph model. When real emission graphs deviate from the configuration model (e.g., entity–value affinities or block structure), the estimator should be viewed as a principled degree-constrained baseline.

Computation. Computing the estimator requires only $T = \sum_a o_a$, $D = \sum_a o_a^2$, and α. This is $O(m)$ time.

Proof sketch. Write $N(C) = \sum_{i,a} c_{ia}^2 = T + \sum_{i,a} c_{ia}(c_{ia} - 1)$ and interpret the second term as the number of ordered within-entity stub pairs that collide on value label; each such pair collides with probability p_2 under uniform matching. A complete proof is given in Appendix A.

6 Uncertainty across graphs: analytic variance under the configuration model

The estimator in Theorem 1 is an ensemble mean. In many applications we also want uncertainty: how much can the exact $\theta(C)$ fluctuate across graphs consistent with the same marginals?

Because D is fixed given o,

$$\mathrm{Var}(\theta(C) \mid \ell, o) = \frac{\mathrm{Var}(N(C) \mid \ell, o)}{D^2}.$$

We derive $\mathrm{Var}(N(C) \mid \ell, o)$ by expanding second moments of within-entity collision indicators.

6.1 Additional value-side probabilities

For three distinct value stubs sampled without replacement (for $T \geq 3$),

$$p_3 := P(A_1 = A_2 = A_3) = \frac{\sum_a (o_a)_3}{(T)_3}.$$

For four distinct stubs (for $T \geq 4$), we need the probability that two disjoint pairs each collide:

$$p_{22} := P(A_1 = A_2 \text{ and } A_3 = A_4).$$

Let

$$F_2 := \sum_a (o_a)_2\,, \quad F_4 := \sum_a (o_a)_4\,, \quad F_{22} := \sum_a ((o_a)_2)^2\,.$$

Then

$$\boxed{p_{22} = \frac{F_4 + (F_2^2 - F_{22})}{(T)_4}.}$$

6.2 An extra entity-degree moment appears

Define the entity-side ordered-triple count:

$$M_3 := \sum_{i=1}^{n} (\ell_i)_3 = \sum_i \ell_i(\ell_i - 1)(\ell_i - 2).$$

Unlike M_2, M_3 is *not* determined by σ^2 alone (it depends on a third moment / skewness of the degree sequence). So: the mean of $\theta(C)$ is variance-only; the variance of $\theta(C)$ requires one additional degree-shape statistic (or a model for ℓ).

Theorem 2 (Ensemble variance conditional on (ℓ, o))**.** Under the bipartite configuration model,

$$\boxed{\mathrm{Var}(\theta(C) \mid \ell, o) = \frac{\Big(2M_2p_2 + M_2(M_2-2)p_{22} + 4M_3(p_3 - p_{22})\Big) - (M_2p_2)^2}{D^2}.}$$

How to use Theorem 2 in the variance-only regime. If only σ^2 is specified, then M_2 is fixed but M_3 is not. In practice, there are two options:

- **Monte Carlo uncertainty bands (recommended):** Choose a simple degree-shape model for ℓ consistent with (T, n, σ^2), sample ℓ, then sample C via stub matching, compute $\theta(C)$, and report empirical quantiles.
- **Add one extra knob:** Specify M_3 (or an equivalent skewness parameter) as an additional domain-knowledge input. Then Theorem 2 gives analytic error bars in $O(m)$ time.

A proof is given in Appendix B.

7 Implementation recipes

7.1 Point estimator from (o, n, σ^2)

Algorithm 1 computes the variance-only point estimate (ensemble mean).

7.2 Uncertainty bands by Monte Carlo

Algorithm 2 describes a practical uncertainty procedure. It requires choosing a degree-shape model for ℓ consistent with (T, n, σ^2). For the toy experiments we simply fix ℓ exactly; for larger demos we recommend either (i) a simple two-level construction or (ii) a parametric prior (e.g., a Dirichlet–multinomial / compound-multinomial model [12, 13]) that can be tuned to match σ^2.

Algorithm 1 Variance-only point estimate of $P(S \mid \mathcal{C})$ (ensemble mean)

Require: value degrees $o_1, \ldots, o_m$; number of entities n; assumed degree variance σ^2
Ensure: $\widehat{\theta} \approx \mathbb{E}[\theta(C) \mid o, \sigma^2]$

$T \leftarrow \sum_{a=1}^{m} o_a$
$D \leftarrow \sum_{a=1}^{m} o_a^2$
$\alpha \leftarrow \frac{1}{n} + \frac{n\sigma^2}{T^2}$
$p_2 \leftarrow \frac{D-T}{T(T-1)}$ ▷ without-replacement stub collision prob
$M_2 \leftarrow T^2\alpha - T$
$\widehat{\theta} \leftarrow \frac{T+M_2 p_2}{D}$
return $\widehat{\theta}$

Algorithm 2 Monte Carlo distribution of $\theta(C)$ for uncertainty bands

Require: value degrees o; number of entities n; assumed σ^2; degree model $\mathcal{M}$; trials B
Ensure: samples $\{\theta^{(b)}\}_{b=1}^{B}$ approximating the distribution of $\theta(C)$

for $b = 1$ to B **do**
  Sample degrees $\ell^{(b)} \sim \mathcal{M}(T, n, \sigma^2)$ with $\ell_i^{(b)} \geq 1$ and $\sum_i \ell_i^{(b)} = T$
  Sample $C^{(b)} \sim \mathrm{ConfigModel}(\ell^{(b)}, o)$ via uniform stub matching
  Compute $\theta^{(b)} \leftarrow \left(\sum_{i,a} (c_{ia}^{(b)})^2\right) / \left(\sum_a o_a^2\right)$
end for
return $\{\theta^{(b)}\}_{b=1}^{B}$

8 Baselines and connection to Fellegi–Sunter

To understand what the variance-informed estimator (Theorem 1) provides beyond simpler alternatives, we define several baselines and make the Fellegi–Sunter correspondence explicit.

Notation. Let $\alpha = P(S)$ denote the unconditional same-source probability (Proposition 2), and let $p_{\mathrm{coll}} := P(\mathcal{C}) = D/T^2$ be the with-replacement collision probability.

Baseline 0 (prior only). Ignoring collisions entirely gives

$$\widehat{\theta}_{\mathrm{prior}} := P(S) = \alpha.$$

In Fellegi–Sunter terms, this corresponds to $m = u$ (collision rates equal under match and non-match), so the posterior equals the prior. This baseline sets the floor: any estimator that conditions on collisions should exceed α.

Baseline 1 (uniform-entity degrees). Setting $\sigma^2 = \sigma^2_{\min}$ (degrees as equal as possible) and applying Algorithm 1 yields $\widehat{\theta}_{\mathrm{unif}}$. This quantifies what the value marginals alone imply when entity activity is assumed homogeneous. It corresponds to the leftmost point of the Theorem 1 sensitivity curve (e.g., Figures 5 and 8).

Deterministic-emission upper bound. At the opposite extreme, suppose each entity emits exactly one value label (deterministic emission: $P(\mathcal{C} \mid S) = 1$). Because $\sum_a c_{ia}^2 \leq \left(\sum_a c_{ia}\right)^2 = \ell_i^2$ for

any entity i, with equality iff entity i maps to a single value, we obtain $N(C) = \sum_{i,a} c_{ia}^2 \le \sum_i \ell_i^2 = T^2\alpha$, hence

$$\theta(C) = \frac{N(C)}{D} \;\le\; \frac{T^2\alpha}{D} = \frac{\alpha}{p_{\text{coll}}} =: \widehat{\theta}_{\text{det}}.$$

This is a genuine upper bound on $\theta(C)$ for all graphs with entity-degree concentration α. It is informative when $\alpha < p_{\text{coll}}$ (entities less concentrated than values); when $\alpha \ge p_{\text{coll}}$ the bound exceeds 1 and is vacuous.

Fellegi–Sunter parameterization. In the one-field case, the Fellegi–Sunter posterior is $P(S \mid \mathcal{C}) = m\alpha/(m\alpha + u(1-\alpha))$, where $m = P(\mathcal{C} \mid S)$ and $u = P(\mathcal{C} \mid \bar{S})$. When C is known, these can be written explicitly as

$$m(C) = \frac{N(C)}{T^2\alpha}, \qquad u(C) = \frac{D - N(C)}{T^2(1-\alpha)},$$

and the resulting posterior simplifies to $\theta(C) = N(C)/D$, recovering Proposition 1. Our estimator supplies an analytic, model-based prediction of this posterior when C is unknown.

Visual comparison. Figures 5 and 8 overlay the prior $\alpha(\sigma^2)$ curve alongside the Theorem 1 estimate. The vertical gap between the two curves is the "collision lift"—the additional information gained by conditioning on a value collision at each heterogeneity level.

9 Toy experiments: exact enumeration validation

This section validates Theorems 1–2 by exact enumeration of all feasible incidence matrices C in small cases. For fixed margins (ℓ, o), each table C receives weight proportional to $1/\prod_{i,a} c_{ia}!$ (the configuration-model probability induced by uniform stub matchings). For general background on sampling/enumerating contingency tables with fixed margins, see, e.g., [7, 11, 8, 9, 10].

9.1 Toy set A: $n = 3$, $m = 3$, $o = (4, 4, 4)$

Here $T = 12$ and $D = \sum_a o_a^2 = 48$.

Table 1 summarizes the means and standard deviations. Tables 2–4 list the exact discrete distributions of $\theta(C)$.

Table 1: Toy summary: configuration-model ensemble mean and sd of $\theta(C) = P(S \mid \mathcal{C})$. All examples use $n = 3$ entities, $m = 3$ values.

Example	o	ℓ	σ^2	α	$\mathbb{E}[\theta]$	sd(θ)
A1	$(4,4,4)$	$(4,4,4)$	0	1/3	$5/11 \approx 0.4545$	$\sqrt{2/363} \approx 0.0742$
A2	$(4,4,4)$	$(6,3,3)$	2	3/8	$43/88 \approx 0.4886$	$\sqrt{71/14520} \approx 0.0699$
A3	$(4,4,4)$	$(10,1,1)$	18	17/24	$67/88 \approx 0.7614$	$\sqrt{1/2904} \approx 0.0186$
B	$(8,2,2)$	$(4,4,4)$	0	1/3	$13/33 \approx 0.3939$	$\sqrt{62/49005} \approx 0.0356$

9.2 Toy set B: skewed value marginals $o = (8, 2, 2)$ with $\ell = (4, 4, 4)$

Here $T = 12$ and $D = 72$. Table 5 lists the exact distribution.

Table 2: Exact distribution of $\theta(C)$ for Example A1: $o = (4, 4, 4)$, $\ell = (4, 4, 4)$. (Weighted by configuration-model probability $P(C \mid \ell, o) \propto 1/\prod c_{ia}!$.)

θ	Probability	θ	Probability
0.3750	0.29922	0.6250	0.02216
0.4167	0.22442	0.6667	0.00935
0.5000	0.33662	0.7500	0.00831
0.5417	0.09974	1.0000	0.00017

Table 3: Exact distribution of $\theta(C)$ for Example A2: $o = (4, 4, 4)$, $\ell = (6, 3, 3)$.

θ	Probability	θ	Probability
0.3750	0.09351	0.6250	0.04416
0.4583	0.51429	0.7083	0.01558
0.5000	0.18701	0.7500	0.00519
0.5833	0.14026		

Table 4: Exact distribution of $\theta(C)$ for Example A3: $o = (4, 4, 4)$, $\ell = (10, 1, 1)$.

θ	Probability	Notes
0.7500	0.72727	most mass
0.7917	0.27273	remainder

Table 5: Exact distribution of $\theta(C)$ for Example B: $o = (8, 2, 2)$, $\ell = (4, 4, 4)$.

θ	Probability	θ	Probability
0.3611	0.38788	0.4444	0.26667
0.3889	0.33939	0.5556	0.00606



Figure 2: Toy Set A ($n = 3$, $m = 3$, $o = (4, 4, 4)$): exact weighted distribution of $\theta(C)$ across all feasible contingency tables C with margins (ℓ, o). The dashed line marks the ensemble mean; the dotted line marks the σ^2-only point estimate (Theorem 1)—they coincide exactly. The shaded band shows ± 1 sd (Theorem 2).



Figure 3: Toy Set B ($n = 3$, $m = 3$, $o = (8, 2, 2)$, $\ell = (4, 4, 4)$): exact weighted distribution of $\theta(C)$. Dashed line: ensemble mean; dotted line: σ^2-only estimate (Theorem 1); shaded band: ± 1 sd (Theorem 2).

10 Choosing the entity-degree variance in practice

The estimator requires σ^2 (equivalently α), which encodes heterogeneity of emission volume across entities.

Feasible range (active emitters). Because $\ell_i \geq 1$ and $\sum_i \ell_i = T$, not all σ^2 are possible. Let $T = qn + r$ with $0 \leq r < n$. Then:

$$\sigma^2_{\min} = \frac{r(n-r)}{n^2} \quad \text{(degrees as equal as possible, by a standard exchange argument)},$$

$$\sigma^2_{\max} = \frac{(n-1)(T-n)^2}{n^2} \quad \text{(one dominant emitter, all others at 1)}.$$

How to set σ^2 in practice. The primary recommendation is to set σ^2 from domain knowledge about the emission process, then verify robustness with a sensitivity sweep.

Domain elicitation (recommended). In many entity-resolution settings the analyst already knows, at least qualitatively, how heterogeneous emission volumes are. For example, consider linking records on first names, where each person (entity) may appear under one or several given names (values). Census and administrative data consistently show that most individuals use a single first name while a smaller fraction use two or more (due to cultural naming conventions, nicknames, or legal name changes). This skewed-but-bounded pattern implies a moderate σ^2: well above the uniform baseline $\sigma^2 \approx 0$, but far below the "one dominant emitter" extreme. Translating even this rough qualitative knowledge into a plausible range for σ^2 is typically sufficient, because $\widehat{\theta}$ varies smoothly and monotonically in σ^2.

Variance-only sufficiency. Theorem 1 implies that $\widehat{\theta}$ depends on the entity-degree sequence ℓ only through σ^2: two degree sequences with different shapes but the same variance yield the same point estimate. Figure 4 verifies this empirically. Three hand-constructed ℓ vectors—spread, moderate, and peaked—all have $\sigma^2 = 36$ but very different higher moments (M_3 ranges from 16 920 to 22 680). The MC distributions of $\theta(C)$ all centre on the same Theorem 1 estimate (dashed line).

The spread of each histogram differs (consistent with the M_3-dependence in Theorem 2), but the mean is invariant. This means the analyst need only specify a single scalar, not the full shape of ℓ.



Figure 4: Three ℓ vectors with the same $\sigma^2 = 36$ but different shapes ($n = 10$, $T = 100$, uniform o). All three MC distributions centre on the Theorem 1 estimate (dashed), confirming that the mean depends on ℓ only through σ^2.

Sensitivity analysis (robustness check). Given variance-only sufficiency, a sweep of $\widehat{\theta}$ over the feasible σ^2 interval fully characterises sensitivity. The mapping $\sigma^2 \mapsto \widehat{\theta}$ is monotone increasing, so a single plot immediately shows whether the conclusion depends on the heterogeneity assumption.

Figure 5 demonstrates this for $n = 20$, $m = 15$, $T = 200$ with two value marginals (uniform and Zipf-skewed). The Theorem 1 curve (solid) sits inside the MC 5th–95th percentile band across the entire feasible range, confirming the formula at scale. In both panels $\widehat{\theta}$ spans roughly 0.1–0.9 over the full interval. If domain knowledge places σ^2 in a narrow subinterval, the corresponding vertical slice gives tight bounds—the conclusion is robust. If the plausible range maps to a wide band of $\widehat{\theta}$, the heterogeneity assumption matters and must be justified.

Finite-sample estimation of value marginals. The preceding experiments assume that the value-degree vector o is known exactly. In practice, o (equivalently, the marginal distribution $P(V{=}v_a) = o_a/T$) must be estimated from observed measurements. To evaluate the impact of this estimation step, we run an end-to-end pipeline:

1. Sample a ground-truth graph C^* from the configuration model with specified ℓ and o.
2. Simulate N_{obs} single measurement draws from C^* (each draw samples one edge uniformly and returns the value endpoint; the entity endpoint is latent and unobserved).
3. From the N_{obs} observed values, estimate the value-degree vector $\hat{o}$ by scaling empirical frequencies to the known total T.
4. Feed $(\hat{o}, n, \sigma^2)$ into the Theorem 1 estimator to obtain $\widehat{\theta}(\hat{o})$.
5. Compare $\widehat{\theta}(\hat{o})$ to both the oracle estimate $\widehat{\theta}(o)$ (which uses the true marginals) and the exact posterior $\theta(C^*)$ computed from the known graph.



Figure 5: Sensitivity of $\widehat{\theta}$ to σ^2 for $n = 20$, $m = 15$, $T = 200$. Left: uniform o; right: Zipf-skewed o. Solid line: Theorem 1; shaded band: MC 5th–95th percentile ($B = 500$); dashed lines: ± 1 analytic sd (Theorem 2); grey dotted: prior $\alpha = P(S)$ (Section 8).

Figure 6 shows the result for three heterogeneity levels ($\sigma^2 = 0$, moderate, and high) using $n = 20$, $m = 15$, $T = 200$ with uniform value marginals. In each panel the blue curve and band show the median and 10th–90th percentile of $\widehat{\theta}(\hat{o})$ over $R = 300$ independent measurement simulations at each sample size N_{obs}. As N_{obs} grows, this band collapses onto the green dashed oracle line (the Theorem 1 estimate with true o), confirming that the finite-sample estimation error vanishes. Convergence is effectively complete by $N_{\text{obs}} \approx 500$–$1\,000$.

The red horizontal line marks the exact $\theta(C^*)$ for one specific ground-truth graph; the thin red lines show θ for eight additional sampled graphs. All of these fall within the grey ± 1 sd ensemble band (Theorem 2), illustrating that the analytic uncertainty captures the graph-to-graph variation. The gap between the oracle (green dashed) and any individual $\theta(C^*)$ (red) is the irreducible ensemble uncertainty: the estimator targets the ensemble mean, not the realization-specific posterior.

Direct measurement-model validation. As a complementary check, we validate the measurement model end-to-end by simulating draw *pairs* from a known graph C^* and computing the empirical same-source rate among collisions. For each pair, we draw two edges independently and uniformly from C^*, observe both the value endpoints (to detect a collision) and the entity endpoints (to check same-source status—information that would be latent in practice but is available in simulation). The empirical estimator is

$$\hat{\theta}_{\text{pairs}} = \frac{\#\{\text{pairs with collision and same source}\}}{\#\{\text{pairs with collision}\}}.$$

Figure 7 shows $\hat{\theta}_{\text{pairs}}$ as a running average over 200 000 simulated pairs for the moderate-σ^2 scenario. The curve starts noisy (few collisions at small sample sizes, since the collision rate is $\approx 6.7\%$) and converges to the exact $\theta(C^*)$ (red line), confirming that the algebraic formula $\theta(C) = \sum_{i,a} c_{ia}^2 / \sum_a o_a^2$ correctly computes the same-source probability from the incidence matrix.

11 Worked example: quasi-identifier collisions in event logs

We illustrate the estimator on a synthetic-but-realistic scenario where entity identifiers are unavailable but domain knowledge about activity heterogeneity is available.



Figure 6: Convergence of the Theorem 1 estimator when value marginals are estimated from finite data ($n = 20$, $m = 15$, $T = 200$, uniform o). Each panel fixes one ground-truth graph C^* at a different σ^2. Blue curve/band: Theorem 1 estimate using $\hat{o}$ from N_{obs} simulated draws (median and 10th–90th percentile over 300 repetitions). Green dashed: oracle estimate using true o. Red solid: exact $\theta(C^*)$; thin red lines: θ for eight other sampled graphs. Grey band: ± 1 sd from Theorem 2.



Figure 7: Direct measurement-model validation: empirical same-source rate among collisions from simulated draw pairs converges to the exact $\theta(C^*)$. Red solid: exact $\theta(C^*)$; green dashed: Theorem 1 oracle estimate. Blue curve: running empirical $\hat{\theta}_{\text{pairs}}$ over 200 000 pairs ($\sigma^2 \approx 204$, $n = 20$, $T = 200$).

Setting. Consider a de-identified event log with $T = 5{,}000$ rows, each generated by one of $n = 100$ latent entities (users, devices, or patients). Each row carries a quasi-identifier value (e.g., a demographic key such as {age bracket, sex} or a finer {birth year, sex, ZIP3}). The analyst observes the value counts $o_1, \ldots, o_m$ directly from the log and wants to estimate

$$\theta = P(\text{same entity} \mid \text{same quasi-identifier value}).$$

Operationally, θ is the precision of using quasi-identifier equality as a blocking key: among all pairs of rows that agree on the key, what fraction correspond to the same underlying entity?

Inputs. The estimator (Algorithm 1) requires: (i) the value counts o (computed directly from the log), (ii) the number of active entities n, and (iii) a heterogeneity parameter σ^2 describing the across-entity variance in activity volume. In practice, n and σ^2 come from domain knowledge, a privacy-safe aggregate report, or historical data from periods when stable identifiers were available.

Synthetic ground truth. To evaluate the estimator against a known answer, we generate entity activity volumes from a discretised lognormal distribution (log-space standard deviation 1.5), producing heavy-tailed heterogeneity (median degree 17, maximum 1,191; resulting $\sigma^2 \approx 16{,}841$). We construct two quasi-identifier keys of different granularity:

- **Coarse key** ($m = 30$ distinct values, Zipf-distributed): collision rate $P(\mathcal{C}) \approx 6.9\%$;
- **Fine key** ($m = 300$ distinct values, mildly skewed): collision rate $P(\mathcal{C}) \approx 0.6\%$.

For each key, a ground-truth graph C^* is sampled from the configuration model with the generated degrees, giving exact posteriors $\theta(C^*) = 0.083$ (coarse) and $\theta(C^*) = 0.112$ (fine).

Results. Figure 8 shows the Theorem 1 estimate $\widehat{\theta}$ as a function of the assumed σ^2 for both keys. At the true σ^2, the estimator yields $\widehat{\theta} = 0.080$ (coarse) and $\widehat{\theta} = 0.110$ (fine), closely matching the ground truth. Thin horizontal lines show θ for eight additional configuration-model graphs at the same degrees, confirming tight ensemble concentration.

Three practical conclusions emerge:

1. **Key granularity matters.** The fine key has a ten-fold lower collision rate but collisions are more informative ($\theta \approx 0.11$ vs. 0.08): when a fine-grained quasi-identifier repeats, it is more likely to be the same entity.
2. **Collision lift can be small for coarse keys.** The dotted prior curve $\alpha(\sigma^2)$ (Section 8) nearly coincides with the estimator for the coarse key, confirming that collisions on broad quasi-identifiers are barely more informative than the unconditional same-source rate. For the fine key, the gap between the prior and the estimator is substantially larger. This contrast is itself a useful diagnostic: the estimator quantifies *how much* a collision adds beyond the prior, guiding the analyst toward keys where collisions carry real evidential weight.
3. **Heterogeneity matters.** Under the uniform-entity baseline ($\sigma^2 \approx 0$), both estimates drop sharply ($\widehat{\theta} = 0.013$ and 0.045). The monotone $\sigma^2 \mapsto \widehat{\theta}$ curve allows the analyst to read off how sensitive the conclusion is to the heterogeneity assumption.



Figure 8: Worked example ($n = 100$, $T = 5{,}000$): sensitivity of $\widehat{\theta}$ to assumed entity-degree variance σ^2. Left: coarse key ($m = 30$); right: fine key ($m = 300$). Blue curve: Theorem 1 estimate. Grey dotted: prior $\alpha = P(S)$ (Section 8); the gap is the collision lift. Red line: exact $\theta(C^*)$; thin red lines: θ for eight additional sampled graphs. Dotted vertical: true σ^2. Blue dot: $\widehat{\theta}$ at true σ^2.

12 Discussion and conclusion

12.1 Interpretation and limitations

Why the mean is "variance-only". Theorem 1 shows that the ensemble mean depends on the entity side only through $M_2 = \sum_i \ell_i(\ell_i - 1)$, i.e., the degree second moment. Thus a single heterogeneity parameter controls the mean estimate.

Sanity check connection. Proposition 2 shows that α equals the *unconditional* same-source probability $P(S)$. The contribution of this paper is to characterize how conditioning on a collision $\mathcal{C}$ transforms this baseline into $\mathbb{E}[\theta(C) \mid o, \alpha]$ under the configuration-model ensemble.

Why uncertainty needs more than variance. Theorem 2 depends on $M_3 = \sum_i \ell_i(\ell_i - 1)(\ell_i - 2)$, a skewness-sensitive statistic. Therefore, a single variance parameter does not determine full uncertainty. In practice, Monte Carlo bands under a chosen degree model are the most straightforward solution.

Modeling assumptions. The uniform-stub-matching (configuration-model) assumption is a principled default given degree information, but real systems may exhibit structure beyond degrees (e.g., specialization where certain entities preferentially emit certain values). Such structure can shift $\theta(C)$ away from the configuration-model mean. The toy distributions in Section 9 illustrate the spread even under the configuration model; structured deviations can be larger.

Finite-sample estimation of value marginals. The theoretical results assume that the value marginals o (or $P(V{=}v_a)$) are known exactly. In practice they are estimated from finite measurements. Section 10 demonstrates that the plug-in estimator $\widehat{\theta}(\hat{o})$ converges to the oracle $\widehat{\theta}(o)$ with $\approx$500–1 000 draws for the problem sizes tested. A natural extension is to propagate sampling error analytically

(e.g., via bootstrap over observed values or a Bayesian Dirichlet posterior over value probabilities) to quantify the additional uncertainty from marginal estimation.

12.2 Future directions

Value-conditioned collisions. A natural extension is to derive a closed-form expression for the configuration-model ensemble mean of

$$P(S \mid V_1 = V_2 = v_a),$$

i.e., the same-source posterior conditioned on a collision at a specific value label. Such a result would provide a per-value "collision informativeness" statistic that complements the global collision event $\mathcal{C}$.

Multi-field extensions. Record-linkage applications typically compare records on several fields simultaneously (e.g., name, date of birth, postal code), each governed by its own emission spectrum. If fields are conditionally independent given the latent entity, the joint same-source posterior factorises into per-field contributions, and the present framework can be applied field by field. When fields are correlated (e.g., name and ethnicity), a joint bipartite multigraph over field tuples is needed, and the configuration-model ensemble must respect cross-field dependence structure. Characterising the ensemble mean of the joint posterior—and whether a tractable "variance-only" parameterisation survives—is an open question. A related challenge is aggregating value-conditioned posteriors across multiple observed collisions to form a stronger overall same-source estimate, e.g., by treating per-collision posteriors as evidence factors in a probabilistic linkage objective.

Inverting collisions to estimate σ^2. This paper treats the entity-degree variance σ^2 as a domain-knowledge input. A complementary direction is to *estimate* σ^2 from observed data: if many pairs of draws are available, the empirical collision rate constrains α (and hence σ^2) via the relationship $P(\mathcal{C}) = \sum_a (o_a/T)^2 \cdot \mathbb{E}[\theta(C)]$. More generally, higher-order collision statistics (e.g., triple collisions) may identify additional degree-shape parameters such as M_3. Formalising this inverse problem—including identifiability conditions and finite-sample rates—would make the framework fully data-driven.

Robustness beyond the configuration model. The configuration model is a maximum-entropy baseline, but real emission graphs may exhibit structure such as entity–value affinity (block structure) or degree–degree correlations (assortativity). Quantifying how much such departures can shift $\theta(C)$ away from the configuration-model mean—e.g., via worst-case bounds over graphs with given marginals and bounded block structure—would provide robustness guarantees for practitioners.

12.3 Concluding remarks

We introduced a method for estimating the collision-conditioned same-source probability $P(S \mid \mathcal{C})$ in a latent-entity emission model when the full emission graph is unknown. Under a maximum-entropy-over-matchings bipartite configuration-model ensemble consistent with observed value marginals, we proved that the ensemble mean of the exact posterior depends on the unknown entity side only through the entity-degree variance (second moment). We also derived an analytic expression for ensemble variance across graphs, highlighting an additional dependence on degree skewness. Small toy examples exactly enumerate the distribution of $\theta(C)$, validating the mean and variance formulas.

This yields a compact, interpretable framework: specify (or vary) a single heterogeneity parameter, compute a principled point estimate, and optionally quantify uncertainty via Monte Carlo or by specifying an additional skewness statistic.

A Proof of Theorem 1

Proof. Fix ℓ and o. For each entity i, consider its ℓ_i stubs. After stub matching, each stub is assigned a value label; let $A(u)$ denote the label assigned to stub u. Then the count of label a in entity i is c_{ia}, and

$$\sum_{a=1}^{m} c_{ia}^2 = \sum_{u,v\in i} \mathbf{1}\{A(u) = A(v)\} = \ell_i + \sum_{\substack{u,v\in i\\ u\neq v}} \mathbf{1}\{A(u) = A(v)\}.$$

Summing over i yields

$$N(C) = \sum_{i,a} c_{ia}^2 = T + X, \quad X := \sum_{\text{ordered within-entity pairs } (u,v),\, u\neq v} \mathbf{1}\{A(u) = A(v)\}.$$

The number of ordered within-entity pairs is

$$M_2 = \sum_i \ell_i(\ell_i - 1).$$

Under uniform stub matching, for any ordered pair (u, v) of distinct stubs, the probability that $A(u) = A(v)$ equals the probability that two distinct value stubs drawn uniformly without replacement share the same label:

$$p_2 = \sum_a \frac{o_a}{T}\frac{o_a - 1}{T - 1} = \frac{\sum_a (o_a)_2}{(T)_2} = \frac{D - T}{T(T - 1)}.$$

Therefore $\mathbb{E}[X \mid \ell, o] = M_2 p_2$ and

$$\mathbb{E}[N(C) \mid \ell, o] = T + M_2 p_2, \qquad \mathbb{E}[\theta(C) \mid \ell, o] = \frac{\mathbb{E}[N(C) \mid \ell, o]}{D} = \frac{T + M_2 p_2}{D}.$$

Finally, $M_2 = \sum_i \ell_i(\ell_i - 1) = \sum_i \ell_i^2 - T = T^2\alpha - T$, which yields the stated dependence on α. □

B Proof of Theorem 2

Proof. Write $X = \sum_{p\in\mathcal{P}} I_p$ where $\mathcal{P}$ is the set of ordered within-entity pairs (u, v) with $u \neq v$, $|\mathcal{P}| = M_2$, and $I_p = \mathbf{1}\{A(u) = A(v)\}$. Then $\theta(C) = (T + X)/D$, so $\mathrm{Var}(\theta) = \mathrm{Var}(X)/D^2$.

We compute $\mathrm{Var}(X)$ from second moments:

$$\mathbb{E}[X^2] = \sum_p \mathbb{E}[I_p] + 2\sum_{p<q} \mathbb{E}[I_p I_q].$$

For a fixed unordered pair $\{p, q\}$, there are three possible overlap types:

- **Reverse pair:** q is (v, u) when p is (u, v). Joint probability is p_2. There are exactly $M_2/2$ such unordered pairs.

- **Share one stub:** (u, v) and (u, w), or (v, u) and (w, u), etc., involving three distinct stubs in the same entity. Joint probability is p_3. For entity i, the number of unordered pairs of ordered pairs sharing one stub equals $2\,(\ell_i)_3$; summing gives $2M_3$.

- **Disjoint:** four distinct stubs. Joint probability is p_{22}. The remaining unordered pairs count is $\binom{M_2}{2} - \frac{M_2}{2} - 2M_3$.

Using $\mathbb{E}[I_p] = p_2$, we obtain:

$$\mathbb{E}[X^2] = M_2 p_2 + 2\Big[\frac{M_2}{2} p_2 + (2M_3)p_3 + \Big(\binom{M_2}{2} - \frac{M_2}{2} - 2M_3\Big)p_{22}\Big].$$

Algebra simplifies this to:

$$\mathbb{E}[X^2] = 2M_2 p_2 + M_2(M_2 - 2)p_{22} + 4M_3(p_3 - p_{22}).$$

Since $\mathbb{E}[X] = M_2 p_2$, we have

$$\mathrm{Var}(X) = \mathbb{E}[X^2] - (\mathbb{E}[X])^2 = \Big(2M_2 p_2 + M_2(M_2 - 2)p_{22} + 4M_3(p_3 - p_{22})\Big) - (M_2 p_2)^2.$$

Dividing by D^2 gives Theorem 2. □

References

[1] E. H. Simpson. Measurement of diversity. *Nature*, 163:688, 1949.

[2] E. T. Jaynes. Information theory and statistical mechanics. *Physical Review*, 106(4):620–630, 1957.

[3] J. Park and M. E. J. Newman. Statistical mechanics of networks. *Physical Review E*, 70(6):066117, 2004.

[4] D. Garlaschelli and M. I. Loffredo. Maximum likelihood: extracting unbiased information from complex networks. *Physical Review E*, 78(1):015101, 2008.

[5] M. E. J. Newman. *Networks: An Introduction*. Oxford University Press, 2010.

[6] D. S. Fosdick, D. B. Larremore, J. Nishimura, and J. Ugander. Configuring the configuration model. *SIAM Review*, 60(2):315–355, 2018.

[7] P. Diaconis and A. Gangolli. Rectangular arrays with fixed margins. In *Discrete Probability and Algorithms* (Minneapolis, MN, 1993), Vol. 72 of *IMA Volumes in Mathematics and its Applications*, pages 15–41. Springer, 1995.

[8] M. Dyer, R. Kannan, and J. Mount. Sampling contingency tables. *Random Structures & Algorithms*, 10:487–506, 1997.

[9] Y. Chen, P. Diaconis, S. Holmes, and J. S. Liu. Sequential Monte Carlo methods for statistical analysis of tables. *Journal of the American Statistical Association*, 100(469):109–120, 2005.

[10] W. M. Patefield. Algorithm AS 159: An efficient method of generating random $R \times C$ tables with given row and column totals. *Applied Statistics*, 30(1):91–97, 1981.

[11] P. Diaconis and B. Sturmfels. Algebraic algorithms for sampling from conditional distributions. *The Annals of Statistics*, 26(1):363–397, 1998.

[12] J. E. Mosimann. On the compound multinomial distribution, the multivariate β-distribution, and correlations among proportions. *Biometrika*, 49(1–2):65–82, 1962.

[13] T. P. Minka. Estimating a Dirichlet distribution. Technical report, MIT, 2000.

[14] I. P. Fellegi and A. B. Sunter. A theory for record linkage. *Journal of the American Statistical Association*, 64(328):1183–1210, 1969.

[15] M. A. Jaro. Advances in record-linkage methodology as applied to matching the 1985 Census of Tampa, Florida. *Journal of the American Statistical Association*, 84(406):414–420, 1989.

[16] W. E. Winkler. Overview of record linkage and current research directions. Technical Report RR2006/02, U.S. Census Bureau, 2006.

[17] J. B. Copas and F. J. Hilton. Record linkage: Statistical models for matching computer records. *Journal of the Royal Statistical Society, Series A*, 153(3):287–312, 1990.

[18] R. C. Steorts, R. Hall, and S. E. Fienberg. A Bayesian approach to graphical record linkage and deduplication. *Journal of the American Statistical Association*, 111(516):1660–1672, 2016.

[19] M. Sadinle. Bayesian estimation of bipartite matchings for record linkage. *Journal of the American Statistical Association*, 112(518):600–612, 2017.

[20] P. Flajolet, D. Gardy, and L. Thimonier. Birthday paradox, coupon collectors, caching algorithms, and self-organizing search. *Discrete Applied Mathematics*, 39(3):207–229, 1992.

[21] L. Sweeney. k-anonymity: A model for protecting privacy. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 10(5):557–570, 2002.

[22] A. Narayanan and V. Shmatikov. Robust de-anonymization of large sparse datasets. In *IEEE Symposium on Security and Privacy*, pages 111–125, 2008.

[23] A. K. Elmagarmid, P. G. Ipeirotis, and V. S. Verykios. Duplicate record detection: A survey. *IEEE Transactions on Knowledge and Data Engineering*, 19:1–16, 2007.

[24] M. A. Hernández and S. J. Stolfo. The merge/purge problem for large databases. In *Proceedings of ACM SIGMOD*, pages 127–138, 1995.

[25] P. Christen. *Data Matching: Concepts and Techniques for Record Linkage, Entity Resolution, and Duplicate Detection*. Springer, 2012.

[26] P. Christen. A survey of indexing techniques for scalable record linkage and deduplication. *IEEE Transactions on Knowledge and Data Engineering*, 24(9):1537–1555, 2012.

[27] L. Getoor and A. Machanavajjhala. Entity resolution: theory, practice & open challenges. *Proceedings of the VLDB Endowment*, 5(12):2018–2019, 2012.

[28] I. Bhattacharya and L. Getoor. Collective entity resolution in relational data. *ACM Transactions on Knowledge Discovery from Data*, 1(1):Article 5, 2007.