Short remainder-only certificates for the greatest common divisor

October 16, 2026

Abstract

We give a remainder-only straight-line derivation of the greatest common divisor of two positive integers using $O(\log a/\log\log a)$ remainder calls, uniformly for $a > b \geq 1$. Previously generated values may be retained and reused, but numerical constants and other arithmetic operations are not available as instructions. The construction combines a stored Euclidean prefix with three-call contractions and two smaller recursive derivations. It yields the same upper bound for intrinsic remainder-call complexity, and therefore a negative answer to the uniform-process strengthening discussed by Moschovakis. It does not establish a deterministic remainder-only algorithm with this cost. The mathematical proof below and the exact scope of the accompanying Lean verification are distinguished in the final section.

Statement and relation to the book

Throughout, logarithms are to base 2. A derivation starts with its two inputs. An instruction adds $\operatorname{rem}(x,y)$, where x,y are already available and $y > 0$. Earlier values remain available. No constants are supplied. Let $G(a,b)$ be the minimum number of instructions needed to make $\gcd(a,b)$ available, and let $D(a,b) = G(a,b)$ for coprime inputs. Thus $G(a,0) = 0$ and $D(a,1) = 0$. These are costs of generating values.

Theorem 1. *Uniformly for integers $a > b \geq 1$,*

$$G(a,b) = O\left(\frac{\log a}{\log\log a}\right).$$

In particular this bound holds for $D(a,b)$ when $\gcd(a,b) = 1$.

As usual, the asymptotic assertion concerns sufficiently large a; a bound valid at every positive input is $O(1 + \log(2+a)/\log\log(4+a))$.

Moschovakis's Main Conjecture concerns logarithmic lower bounds for gcd algorithms from rem, eq_0 [1, p. 2]. The uniform-process strengthening is discussed after Corollary 6C.7 [1, p. 188]; see also (4F-5) on p. 151. Our conclusion addresses that strengthening. The definitions of generated certificates and intrinsic norms are in [1, secs. 1D, 4C–4E]. Theorem 4F.1 [1, pp. 149–150] transfers pointwise intrinsic bounds to a single uniform process. Deterministic uniform processes are treated separately in [1, sec. 4H].

Normalization

Write $g = \gcd(a,b) > 0$. For positive y,

$$\operatorname{rem}(gx, gy) = g \operatorname{rem}(x,y).$$

Every generated value is divisible by g. Consequently scaling a derivation, or dividing all its values by g, preserves its length and the validity of every instruction. Hence

$$G(a,b) = D(a/g, b/g).$$

The normalized pair $(1, 0)$ is understood to have cost zero. Define

$$T(n) = \max_{0 \le b \le a \le 2^n,\ a > 0} G(a, b).$$

Euclid's algorithm makes this a finite maximum. Normalization reduces an upper bound for T to coprime inputs, with equality, zero inputs, and inputs equal to one handled without calls.

Three-call contractions

Lemma 2. *If $m > 0$ is available and a finite collection S of available values occupies $t \ge 2$ residue classes modulo m, at most three remainder calls generate w with*

$$0 < w < \frac{m}{t - 1}.$$

Proof. Sort the distinct residues. Some adjacent pair $u < v$ has gap $\delta = v - u < m/(t - 1)$, because the sum of the $t - 1$ adjacent gaps is less than m. Each selected residue can be generated with one call from a witness in S. If $u = 0$, use v. If $0 < u \le \delta$, use u. Otherwise $0 < \delta < u$ and $v = u + \delta$, so $\operatorname{rem}(v, u) = \delta$. This uses at most three calls. Subtraction and sorting describe the proof and the selection of instructions; they are not instructions of the derivation. □

Lemma 3 (Euclidean-prefix lemma). *Let $r_0 = a > r_1 = b > \cdots > r_h > 0$ be a Euclidean prefix of a coprime pair, with $r_{i-1} = q_i r_i + r_{i+1}$ and $q_i \ge 1$. For every integer $m > 2a/r_h$, the h pairs*

$$(r_i \bmod m, r_{i+1} \bmod m), \qquad 0 \le i < h,$$

are distinct. The prefix therefore occupies t residue classes with $t^2 \ge h$.

Proof. Suppose the pairs at $i < j$ coincide. Set

$$M = \prod_{\ell=i+1}^{j} \begin{pmatrix} q_\ell & 1 \\ 1 & 0 \end{pmatrix} = \begin{pmatrix} A & B \\ C & E \end{pmatrix}, \qquad \begin{pmatrix} r_i \\ r_{i+1} \end{pmatrix} = M \begin{pmatrix} r_j \\ r_{j+1} \end{pmatrix}.$$

All entries are nonnegative, $E \le A$, and $\det M = (-1)^{j-i}$. Also $A r_j \le r_i \le a$, whence $A \le a/r_h$. Write

$$\Delta = \det(M - I) = 1 + \det M - A - E.$$

For an odd number of factors, $\Delta = -A - E < 0$. A product of at least two factors has $A \ge 2$ and $E \ge 1$; thus for an even positive number of factors, $\Delta = 2 - A - E < 0$. In either case,

$$0 < |\Delta| \le A + E \le 2A \le 2a/r_h < m.$$

The presumed congruence gives $(M - I)(r_j, r_{j+1})^{\mathsf{T}} \equiv 0 \pmod m$. Multiplication by the adjugate shows that m divides both Δr_j and Δr_{j+1}. Consecutive remainders are coprime, so Bézout's identity implies $m \mid \Delta$, a contradiction. Finally, t individual residue classes give at most t^2 ordered pairs. □

Thus a stored prefix can be reused for every available modulus $m > 2a/r_h$. When $\sqrt{h} > 1$, three calls suffice to generate

$$0 < w < \frac{m}{\sqrt{h} - 1}.$$

The universal quantifier over m is essential: the modulus generated by a contraction is not predetermined.

Construction and recurrence

Fix a sufficiently large integer n and put $N = 2^n$ and $h = \lceil \sqrt{n} \rceil$. After normalization we may assume $a > b > 1$ and $a \le N$. Generate the Euclidean prefix until its last positive value is at most $N^{3/4}$ or until r_h is available. Check the input b as part of this rule. The preliminary work costs at most h calls. Euclid cannot encounter zero first without having encountered the positive remainder 1.

If a positive remainder $c \le N^{3/4}$ is encountered, generate the next remainder $s < c$ with one more call. The pair (c, s) is coprime. If $s = 0$, then $c = 1$ is already available; otherwise both inputs to the remaining derivation are at most $2^{\lceil 3n/4 \rceil}$. This gives the bound

$$h + 1 + T(\lceil 3n/4 \rceil).$$

Otherwise $r_h > N^{3/4}$, so $2a/r_h < 2N^{1/4}$. For $n > 12$, $N^{1/3} > 2N^{1/4}$, and Lemma 3 applies to every available $m > N^{1/3}$. Starting at $m = r_h$, repeatedly use the contraction inequality until $0 < m \le N^{1/3}$. Set

$$\kappa_n = \sqrt{h} - 1 > 1, \qquad J_n = \lceil \frac{2n/3}{\log \kappa_n} \rceil.$$

There are at most J_n contractions: if J_n have occurred then $m < N/\kappa_n^{J_n} \le N^{1/3}$. Since $\log \kappa_n = \frac{1}{4} \log n + O(1)$, their cost is $3J_n = O(n/\log n)$.

The small m need not be coprime to either input. Generate $u = \text{ rem } (a, m)$ and $v = \text{ rem } (b, m)$, then concatenate derivations of $d = \text{ gcd } (m, u)$ and $e = \text{ gcd } (d, v)$. These derivations exist by the definition of T and use only inputs bounded by m. They are not gcd-oracle instructions. The gcd identities give

$$e = \text{ gcd } (\text{ gcd } (m, a \mod m), b \mod m) = \text{ gcd } (m, \text{ gcd } (a, b)) = 1.$$

The cost in this case is at most

$$h + 3J_n + 2 + 2T(\lceil n/3 \rceil).$$

Retaining values makes concatenation legitimate. In particular, the second recursive derivation has both d and v available even if the first one uses neither original input.

The two case estimates and normalization imply

$$T(n) \le \max \{T(\lceil 3n/4 \rceil) + O(\sqrt{n}),\ 2T(\lceil n/3 \rceil) + O(n/\log n)\}.$$

All implied constants are independent of the inputs.

Proof of Theorem 1. For $n \ge 2$ let $f(n) = n/\log n$. The ratios

$$\frac{f(\lceil 3n/4 \rceil)}{f(n)} \to \frac{3}{4}, \qquad \frac{2f(\lceil n/3 \rceil)}{f(n)} \to \frac{2}{3}$$

show that, beyond a fixed threshold, the two recursive terms have potentials at most $\frac{7}{8}f(n)$ and $\frac{5}{6}f(n)$, respectively. Both overheads are at most $Af(n)$ for a fixed A. Choose $C \ge 8A$ and increase C to cover the finitely many initial values with indices at least 2. Choose the induction threshold large enough that both recursive indices are at least 2 and strictly less than n. Strong induction applied to this recurrence then gives $T(n) \le Cf(n)$: the two bounds are $(7C/8 + A)f(n)$ and $(5C/6 + A)f(n)$, both at most $Cf(n)$. Finally take $n = \lceil \log a \rceil$. This also gives $G(a, b) = o(\log a)$ uniformly as $a \to \infty$. □

Certificates and uniform processes

Use the book's partial remainder operation: its second argument must be nonzero. Write $\mathcal{N}_0 = (\mathbb{N}, \text{ rem }, \text{ eq}_0)$ and $\mathcal{N}_\varepsilon = (\mathbb{N}, \text{ rem }, \text{ eq}_0, \text{ eq}_1)$; compare [1, pp. (1C–10), pp. 24, 142–143]. A derivation determines a finite generated partial substructure by recording its remainder equations. Repeated equations can be discarded.

For the relation $\text{ gcd } (a, b) = 1$ on all natural inputs, a derivation of 1 together with $\text{eq}_1\ (1) = \text{tt}$ certifies the relation: every common divisor of the images of the inputs divides every transported generated value, hence divides 1. Conversely, without a positive eq_1 fact the map $x \mapsto 2x$ preserves every recorded remainder and zero-test fact, as well as every negative one-test fact. It cannot preserve coprimeness. A certificate therefore reaches 1. Enumerating a finite generated diagram in generation order gives a derivation using no more calls than the number of remainder facts. Consequently, on coprime inputs,

$$\mathrm{calls}_{\mathrm{rem}}\,(\mathcal{N}_{\varepsilon},\ \gcd\ = 1, a, b) = D(a,b), \qquad \mathrm{calls}\,(\mathcal{N}_{\varepsilon},\ \gcd\ = 1, a, b) = D(a,b) + 1.$$

If the relation is instead defined as $a > 0 \wedge b > 0 \wedge\ \gcd\,(a,b) = 1$, add the two input-positivity facts. The remainder count is unchanged, but one should state the full count as $D(a,b) + O(1)$ rather than assert the exact $D + 1$ formula without addressing positivity. For example, the single fact $\mathrm{eq}_1\ (1) = \mathrm{tt}$ at $(2, 1)$ allows the image $(0, 1)$.

For gcd as a function, the certificate must reach its output g, giving a lower bound of $G(a,b)$. For an upper bound, append

$$\mathrm{rem}\,(a,g) = z, \qquad \mathrm{rem}\,(b,g) = z, \qquad \mathrm{eq}_0\ (z) = \mathrm{tt},$$

where the generated z is zero. Every homomorphism π has $\pi(z) = 0$ and $\pi(g) > 0$ (by definedness of either appended remainder). Thus $\pi(g)$ divides both input images. Conversely every common divisor of those images divides $\pi(g)$ by induction over the derivation. Therefore

$$G(a,b) \le\ \mathrm{calls}_{\mathrm{rem}}\,(\mathcal{N}_0,\ \gcd\,, a, b) \le G(a,b) + 2.$$

Positivity facts may be appended if the function is restricted to positive inputs. For a noncoprime input, add $\mathrm{eq}_1\ (g) = \mathrm{ff}$ to this gcd certificate to certify noncoprimeness. Thus the sublogarithmic remainder-call bound holds for both truth values of coprimeness as well as for gcd. Theorem 4F.1 supplies a single uniform process realizing each intrinsic norm pointwise [1, Theorem 4F.1].

Scope and verification

Selecting a close pair of residues is uncharged existential choice in a straight-line derivation. A deterministic remainder-only algorithm cannot assume that choice is available at constant cost. Computing all prefix residues would take about h calls per contraction. The construction therefore does not settle the deterministic conjecture.

The supplied offline Python planner uses unrestricted arithmetic to select instructions; its independent replay checks operand availability, nonzero divisors, every remainder equation, and the final gcd. Running its default seed reproduces:

Input bits	**Derivation calls**	**Euclid calls to reach** 1
512	196	326
2048	669	1204
8192	2059	4754

These are checks of finite instances, not an asymptotic proof. A newly specified reproducible suite also passes 310 cases, including 175 noncoprime pairs, with 3,543 contractions across 335 reservoir phases. Four deliberately invalid traces are rejected. This suite is recorded in the repository; it is not claimed to reconstruct the unspecified original 310-case suite.

Machine-checked statements

The Lean 4.35.0 development, with mathlib pinned to version 4.35.0, proves the uniform real-logarithmic bound, its worst-case Big-O formulation, the Euclidean-prefix lemma, the complete three-call contraction lemma, normalization, and the certificate upper bounds. It contains no proof placeholders or added axioms. An axiom audit of the main results reports only the standard Lean foundations `propext`, `Classical.choice`, and `Quot.sound`. The theorem `ShortGCD.uniform_real_bound` states

$$\exists C > 0\ \forall a, b \in \mathbb{N}, \quad 4 \le a\ \wedge\ b \le a\ \Rightarrow\ G(a,b) \le C \frac{\log_2 a}{\log_2\,(\log_2 a)}.$$

The proof starts from a finite-store instruction semantics in which the only instruction is remainder with a positive, available divisor. Costs are indices of derivations, rather than an assumed recurrence. A formal conversion to finite generated diagrams proves the gcd certificate bound with two extra remainder facts and the bound for positive and negative coprimeness instances. The equality with the existing D definition is also checked.

The intrinsic complexities themselves are also defined in Lean, directly from certificates: `callsCoprimeness` for the coprimeness relation in N_ε and `callsGcd` for the gcd without eq_1 facts. On coprime pairs the former agrees with the existing reduction to D (`callsCoprimeness_eq_callsRem`). The theorems `ShortGCD.not_conjecture_3_4` and `ShortGCD.not_main_conjecture_4F5` state that no $r > 0$ satisfies $\mathrm{calls(rem)}(N_\varepsilon, \bot\ \bot, a, b) \geq r \log_2 a$ for infinitely many $a \geq b \geq 1$, and that no $K > 0$ satisfies $2 \log_2 \max(x, y) \leq K\,\mathrm{calls(rem)}(\gcd, x, y)$ for infinitely many pairs. The general theory of uniform processes and Theorem 4F.1 are cited from the book, not re-formalized here. They are needed only to interpret these intrinsic bounds as statements about a single best uniform process, not for the refutations themselves.

Integer parameters used in Lean

To avoid real rounding in the construction itself, the formal proof uses, for $j = \lfloor \log_2 n \rfloor \geq 32$,

$$p = \lfloor j/4 \rfloor, \quad k = 2^p, \quad h = 2^{2p} + 1, \quad J = \lfloor n/p \rfloor + 1, \quad \ell = \lceil 3n/4 \rceil, \quad s = \lceil n/3 \rceil.$$

The finite construction proves, for arbitrary natural parameters satisfying $k \geq 1$, $k^2 < h$, $L \geq 1$, $2A \leq ML$, and $A \leq Mk^J$,

$$S(A) \leq \max\{h + 1 + S(L),\ h + 3J + 2 + 2S(M)\}, \qquad S(A) = \max_{a,b \leq A} G(a, b).$$

Taking $A = 2^n, L = 2^\ell, M = 2^s$ gives the required hypotheses. Lean checks

$$(h + 1)j \leq 64n, \quad (h + 3J + 2)j \leq 64n, \quad j - 2 \leq \log_2^{\mathrm{nat}} \ell,\ \log_2^{\mathrm{nat}} s, \quad 8\ell \leq 7n, \quad 16s \leq 7n.$$

The last estimates and $8j \leq 9(j - 2)$ give a factor 63/64 for either recursive contribution after clearing denominators. Taking $C \geq 4096$ and large enough for the finite initial segment proves $S(2^n) \lfloor \log_2 n \rfloor \leq Cn$ for all n. Comparing integer floors and ceilings with real base-two logarithms yields the stated real bound. These parameter choices change only absolute constants, not the proof mechanism or its asymptotic order.

References

[1] Moschovakis, Y. N. *Abstract recursion and intrinsic complexity*. Lecture notes in logic, vol. 48. Cambridge University Press; Association for Symbolic Logic, 2019. Author's prepublication PDF, compiled November 30, 2018. Citations use printed page numbers. https://www.math.ucla.edu/~ynm/lectures/aricplace/c-aric.pdf.