A sharp threshold for three-colour targets from four-colour cubes

Jason Davies

November 1, 2026

Abstract

Any collection of 29 cubes, each displaying all four colours of a fixed palette, contains eight that form a $2 \times 2 \times 2$ cube with monochromatic faces using exactly three colours. For this fixed palette, exactly two collections of 28 cubes fail, and reflection exchanges them. This answers the threshold question in Problem 2 of Berkove, Eickemeyer and Schulman (2018). The proof uses their corner-containment reduction and elementary counting. The threshold and classification are formalised in Lean.

1 Introduction

Berkove, Eickemeyer and Schulman [1, Problem 2] ask for the least number of four-colour cubes that guarantees a choice of eight cubes forming a three-colour corner solution: a $2 \times 2 \times 2$ cube whose exterior faces are monochromatic and use exactly three colours. Here every source cube uses all four colours of a fixed palette. A *variety* is a class of face colourings identified under proper rotations. Collections are multisets of varieties, so repeated copies are allowed.

We prove that the least number is 29 and classify all collections of 28 cubes that fail. There are exactly two such collections, exchanged by reflection and by any odd colour permutation, so there is one obstruction up to colour permutation. Each consists of seven copies of each of four varieties; explicit face tuples are given in Theorem 3. The proof first classifies the possible 28-cube obstructions by corner containment and counting, then deduces the upper bound and verifies that both obstructions fail.

2 Model and geometric facts

Reflections are not allowed as moves. Only exterior faces must match the target; internal faces are unrestricted.

A corner is a cyclically ordered triple. Read its face colours counterclockwise as viewed from outside along the corner's body diagonal; equivalently, order the three outward face normals to form a positively oriented basis. BES [1, Section 2] reads corners clockwise; our convention reverses every triple, preserving containment and matching. Thus abc and acb are distinct when the three colours differ. A source cube fits a target corner by a proper rotation if and only if it has the same cyclic corner type: the cyclic alignments of two corner frames are exactly those realised by proper rotations of the cube. A target is represented by its multiset of eight corner types: a collection builds it exactly when these eight occurrences can be matched to distinct cubes having the required types.

Write $Q(a; b, c)$ for a target whose b and c faces are opposite and whose other four faces have colour a. Its corner multiset is $4aab + 4aac$. Write $R(a; b, c)$ for the analogous target with b and c adjacent; its corners are

$$2aaa + 2aab + 2aac + abc + acb.$$

Finally, write $D(a, b, c)$ for the target with opposite pairs aa, bb, cc; it has four corners of each orientation on $\{a, b, c\}$. We need only these three families of exactly-three-colour targets for the sufficient constructions.

If every corner type of a variety v occurs on a variety w, replacing a copy of w by v preserves unsolvability. This is the reduction of [1, Section 2]. For clarity, Table 1 gives the entire reduction needed here. A face tuple is read in the order $+x, -x, +y, -y, +z, -z$. The rows represent the seven orbits under colour permutations, with rotations already identified. These seven orbits of all 68 varieties differ from the seven minimal representatives in [1, Figure 2]; four of our orbits are retained. The three non-retained representatives all contain the corner types of 001123. Colour permutations transport these containments to every member of their orbits.

Representative	Orbit size	Retained type	Replacement
001123	6	checkerboard C	—
010203	8	triple-corner T	—
010213	12	pair type P	—
010223	12	pair type P	—
000123	12	—	001123
001213	12	—	001123
010123	6	—	001123

Table 1: The 68 varieties reduce to 38.

The retained types are precisely the minimal varieties in [1, Table 8]. A reduced variety belongs to bucket a if it has at least two of the three corner types aab with $b \neq a$.

- The eight T varieties have corners aaa and aab for every $b \neq a$, and belong only to bucket a.
- The 24 P varieties belong to two buckets, one for each repeated colour.
- The six checkerboards belong to no bucket. Their repeated colours occupy opposite face pairs; with repeated colours a, b, a checkerboard supports both orientations on $\{a, b, c\}$ and $\{a, b, d\}$.

The classes and checkerboard terminology come from [1, Section 2]. The 32 ordinary varieties (T and P) split into two classes of 16, corresponding to BES's K_o and K_m (up to interchanging the names). Every member of one class has the same four trichromatic corner types, one orientation for each three-element subset of the palette; every member of the other class has the four reversed orientations. For our corner convention the four trichromatic types of one class are

$$\Omega = \{012, 031, 023, 132\}.$$

These are one set of BES's *universal corners*, in our convention. All three ordinary representatives in Table 2 belong to the class of Ω. Even colour permutations preserve Ω and odd ones reverse it, with triples read cyclically. Applying colour permutations to Table 2 gives the stated incidences and orientation classes. The eight T varieties split evenly between the classes, and even permutations act transitively on the four colours, so each class contains exactly one T variety for each repeated colour. A separate geometric checker verifies both tables and all the stated incidences using physical rotations.

Lemma 1 (Bucket bound)**.** *Eight reduced cubes in bucket a can build a three-colour target. Consequently, a reduced collection with no such solution has at most seven cubes in each bucket.*

Representative	Corner-type set
001123	$\{012, 013, 021, 031\}$
010203	$\{000, 001, 002, 003, 012, 023, 031, 132\}$
010213	$\{001, 003, 011, 012, 023, 031, 112, 132\}$
010223	$\{002, 003, 012, 022, 023, 031, 122, 132\}$

Table 2: Corner-type sets of the retained representatives, with cyclic triples written in lexicographically least form.

Proof. Each of the eight cubes misses at most one of the three types aab. Thus at most one type is missed by five or more cubes. Choose two types, aab and aac, each missed by at most four cubes. Assign cubes missing aac to aab positions, and cubes missing aab to aac positions. These groups are disjoint and each has size at most four. Every remaining cube supports both types, so they fill the remaining positions of $Q(a; b, c)$. □

3 The extremal collections

Theorem 1. *A collection of 28 four-colour cubes without a three-colour corner solution consists of seven copies of each of the four T varieties in one ordinary orientation class.*

Proof. Suppose the collection has no solution. Reduce it as above; the reduced collection has no solution either, since reduction preserves unsolvability. Let $A \geq B$ be the sizes of the two ordinary orientation classes, and let C count checkerboards, so $A + B + C = 28$. Call the class of size A the majority class (either one if equal). Call the remaining $M = B + C$ cubes *minority cubes*. A minority cube *supports* a three-element colour set E if it has the orientation on E opposite to the majority class. Smaller-class cubes support every E; a checkerboard supports two.

For $E = \{a, b, c\}$, let z_E count checkerboards supporting E. If $z_E \geq 8$, those eight checkerboards build the target $D(a, b, c)$. Thus $z_E \leq 7$ for each E, and, since each checkerboard supports two triples, $2C \leq 28$. It follows that $A + B \geq 14$, so $A \geq 7$. There are $m_E = B + z_E$ minority cubes supporting E. If $m_E \geq 4$, four such cubes and four majority cubes build $D(a, b, c)$. Hence $m_E \leq 3$ for every E. Summing gives

$$4B + 2C \leq 12, \qquad M \leq 6, \qquad A = 28 - M \geq 22. \tag{1}$$

From now on, let t_a and p_a count only majority-class T and P cubes in bucket a, and let n_T, n_P be their respective totals, so $A = n_T + n_P$. By Lemma 1, $t_a + p_a \leq 7$. Each T cube belongs to one bucket and each P cube to two, so summing gives

$$n_T + 2n_P \leq 28, \qquad n_P \leq M. \tag{2}$$

We will repeatedly use the following construction. If six majority cubes supply $2aaa + 2aab + 2aac$ and a minority cube supports $\{a, b, c\}$, then the minority cube supplies the opposite trichromatic orientation and an unused majority cube supplies the other. Such a cube is available since $A \geq 22$. This builds $R(a; b, c)$.

Suppose $M > 0$. Every minority cube supports some triple containing any given colour a. The construction above therefore rules out $t_a \geq 6$. Thus $t_a \leq 5$ for every a, and $n_T \leq 20$. Since $n_T + n_P + M = 28$ and $n_P \leq M$, we obtain $M \geq 4$.

For any two distinct colours a, b, some minority cube supports a triple containing both. Otherwise every minority cube would have to be a checkerboard with the complementary pair of repeated colours. One of the corresponding m_E would then equal $M \geq 4$, contradicting $m_E \leq 3$.

We claim that $t_a \geq 4$ forces $t_a + p_a \leq 5$. Otherwise, since $t_a \leq 5$, choose six majority cubes in bucket a, with four or five T cubes and the rest P cubes. These one or two P cubes share a type aab, since each has two of the three possibilities. Choose a minority cube supporting $\{a, b, c\}$. The P cubes take aab positions and the T cubes supply the rest of $2aaa + 2aab + 2aac$, so the construction above gives a solution, proving the claim.

If $t_a \leq 3$, the bucket bound gives $2t_a + p_a \leq t_a + 7 \leq 10$. If $4 \leq t_a \leq 5$, the claim gives $2t_a + p_a \leq t_a + 5 \leq 10$. Thus in all cases $2t_a + p_a \leq 10$, and summing gives

$$2A = 2n_T + 2n_P = \sum_a (2t_a + p_a) \leq 40,$$

contradicting $A \geq 22$ from (1). Therefore $M = 0$, and (2) forces $n_P = 0$. All 28 cubes are majority-class T cubes, and the four bucket bounds force each $t_a = 7$.

Finally, this conclusion holds for the original collection, not just its reduction. Every non-retained variety in Table 1 is replaced by a checkerboard; since the reduced collection contains none, no such replacement could have occurred. □

Theorem 2. *Every collection of 29 four-colour cubes contains a three-colour corner solution.*

Proof. Otherwise every 28-cube subcollection would have the form in Theorem 1. Take one such subcollection S and let v be the remaining cube. Remove from S a cube of a variety different from v, and insert v. The multiplicity of v's variety is now 1 or 8, whereas every variety in a collection described by Theorem 1 has multiplicity 0 or 7, a contradiction. □

4 Sharpness

Lemma 2. *In a cube using exactly three colours, the two orientations of a trichromatic corner occur equally often. If no trichromatic corner occurs, the cube is $Q(a; b, c)$ for some distinct a, b, c.*

Proof. For any cube coloured from a palette $\{a, b, c, d\}$, let $n(xyz)$ count corners of cyclic type xyz. Read the three directed transitions around each corner in the chosen cyclic order. Each edge between an a face and a b face contributes ab at one endpoint and ba at the other, so the total counts of these transitions agree. Corners with repeated colours contribute equally to both counts. The remaining contributions give

$$n(abc) - n(acb) + n(abd) - n(adb) = 0. \tag{3}$$

If only a, b, c occur, the last two terms vanish, proving the balance.

For the second assertion, if a colour c occurs on only one face, its four neighbours must all have the same colour a: a change from a to b around this face would produce a trichromatic corner. The opposite face is then the sole b face. Otherwise each colour occurs twice. If the two c faces are opposite, the four neighbours of either contain both a and b, forcing a change. If they are adjacent, the other three neighbours form a path containing both a and b, again forcing a change. Both cases produce a trichromatic corner. □

Theorem 3. *The threshold 29 in Theorem 2 is best possible. For a fixed palette, exactly two 28-cube collections fail, and they are mirror images. Up to colour permutation, there is a single obstruction.*

Proof. Take seven copies of each of the following four face tuples:

$$010203, \qquad 011213, \qquad 021223, \qquad 031323.$$

These are T cubes with repeated colours $0, 1, 2, 3$, respectively, all in the same ordinary orientation class. A target with any trichromatic corner requires both orientations by Lemma 2,

whereas the collection supplies only one. A target without one is $Q(a;b,c)$; every one of its eight corners has colour a twice, so only the seven T cubes of repeated colour a are usable. No three-colour target is possible. Reflection gives the analogous example in the other orientation class. Theorem 1 excludes all other collections. Every odd colour permutation exchanges the two orientation classes and hence these two collections. □

Remark. For the palette $\{0,1,2,3\}$, write $\delta(xyz) = n(xyz) - n(xzy)$. Applying (3) to the different colour pairs gives

$$\delta(012) = \delta(031) = \delta(023) = \delta(132).$$

This common integer is the cube's *degree*, relative to Ω. Reflection negates the degree, as does each transposition of colour labels; thus even colour permutations preserve it and odd ones negate it. Each ordinary representative in Table 2 has eight distinct corner types, so each occurs once; the two ordinary classes therefore have degrees $+1$ and -1. The checkerboard representative has neither 023 nor 032, so its degree, and that of every checkerboard, is zero.

5 Formal verification and computation

The threshold and classification are formalised in Lean 4.35.0 [3] with mathlib v4.35.0 [4]. Inventories assign natural-number multiplicities to the 68 source varieties; solutions select eight distinct copies and proper rotations matching the exposed faces. Kernel checks establish source and target catalogue coverage, source uniqueness up to rotation, and correctness of stored corner codes. A palette-independent lemma, allowing repeated colours, equates cyclic corner matching with physical face matching by a proper rotation. Transport theorems extend the results to arbitrary indexed face colourings: each source uses all four colours, and the existentially chosen target uses exactly three. The sharp threshold also has this table-free statement.

The Lean proof uses a related counting argument and finite Hall certificates for sufficient constructions. It treats checkerboards and the smaller ordinary class separately, whereas the written proof combines them as minority cubes. The lower bound checks Hall deficiencies for all 120 targets and both extremal collections; reflection exchanges the two. No proof enumerates all collections. The axiom audit reports only Lean's standard `propext`, `Classical.choice`, and `Quot.sound`. The Lean proof uses neither proof holes nor `native_decide`. The repository[1] contains sources and reproduction commands, with the proper-rotation and outward-face geometry pinned from the earlier MacMahon formalisation. No classification result from that project is used.

A computational search found the 28-cube example and returned results consistent with the threshold and classification. A separate script checks the finite geometric facts and lower obstructions; neither the written proof nor the Lean proof depends on the search or this script.

Literature and priority

A literature search on November 1, 2026 found no earlier answer; this does not establish priority. Rosenfeld's October 2026 article [2] *MacMahon Cubes and Moshe's Insanity, Puzzle 1* remains unread, and its relevance is undetermined.

Declaration of AI-generated research and review

GPT-6-Astra (OpenAI), used through Codex, developed the original research argument, computational search, Lean formalisation and manuscript under the author's direction. Claude Opus

[1] https://github.com/jasondavies/macmahon-four-to-three

5.5 (Anthropic), used through Claude Code in this repository, contributed the minority-cube argument, edge-count balance proof, degree remark, and shorter bucket and threshold proofs. Subsequent Codex revisions incorporated these contributions. A separate AI reviewer checked an earlier version of the argument and physical model. The human author initiated and directed the research and accepts responsibility for the content of this paper and the accuracy of this disclosure. The paper has not received independent human peer review.

References

[1] E. Berkove, A. Eickemeyer and M. Schulman, *The 4-Color Cubes Puzzle*, Integers **18** (2018), A68. Journal PDF.

[2] M. Rosenfeld, *MacMahon Cubes and Moshe's Insanity, Puzzle 1*, Geombinatorics Quarterly **36**(2) (October 2026), starts at p. 78. Journal listing; full text not examined.

[3] L. de Moura and S. Ullrich, *The Lean 4 Theorem Prover and Programming Language*, in Automated Deduction—CADE 28, LNCS **12699**, Springer (2021), 625–635. doi:10.1007/978-3-030-79876-5_37.

[4] The mathlib Community, *The Lean Mathematical Library*, in Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), ACM (2020), 367–381. doi:10.1145/3372885.3373824. arXiv:1910.09336.