Three-colour solutions from four-colour cubes

Jason Davies

November 1, 2026

Abstract

For every $n \geq 2$, the sharp number of cubes using all four colours of a fixed palette guaranteeing an $n \times n \times n$ frame with monochromatic exterior faces using exactly three colours is $16n-3$. For a fixed palette, exactly two collections of $16n-4$ cubes fail: each has $4n-1$ copies of each of four monochromatic-corner varieties in one orientation class, and reflection exchanges the collections. The threshold for a complete cube is $\max\{n^3, 16n-3\}$; in particular, any n^3 source cubes suffice when $n \geq 4$. The case $n = 2$ answers the threshold question in Problem 2 of Berkove, Eickemeyer and Schulman (2018). The proof combines their corner-containment reduction with orientation constraints and bucket counts.

1 Introduction

Berkove, Eickemeyer and Schulman (BES) [1, Problem 2] ask for the least number of four-colour cubes that guarantees a choice of eight cubes forming a three-colour corner solution: a $2 \times 2 \times 2$ cube whose exterior faces are monochromatic and use exactly three colours. This follows BES's convention that a k-colour cube uses all k colours [1, Section 1, opening definition]. Here every source cube uses all four colours of a fixed palette. A *variety* is a face colouring up to proper rotation. Collections are multisets of varieties, so repeated copies are allowed.

We prove a sharp threshold for every side length. An n-frame consists of the corners and edges of an $n \times n \times n$ cube, using $12n-16$ pieces. The least number of source cubes guaranteeing a three-colour n-frame is $16n-3$ (Theorem 4). Exactly two collections of $16n-4$ cubes fail: $4n-1$ copies of each of four varieties, one with a monochromatic corner of each colour, all in the same orientation class. Reflection or any odd colour permutation exchanges these collections. Figure 1 shows the first case, $n = 2$.

BES [1, Theorem 5] prove that four-colour targets need only the $12n-16$ pieces in the frame when $n \geq 3$. Requiring exactly three exterior colours raises the sharp guarantee to $16n-3$.

The classification explains the form $16n-4 = 4(4n-1)$. Mixing orientation classes allows new corner matches, while repeated-colour demands limit the capacity of each colour bucket. Together these restrictions force an extremal collection into one orientation class and then force the four multiplicities to be equal. We first prove the corner case, including its restricted-target corollary, and then extend the argument to frames. Filling the face interiors gives the complete-cube threshold $\max\{n^3, 16n-3\}$ (Corollary 2).

2 Model and geometric facts

Reflections are not allowed as moves. Only exterior faces must match the target; internal faces are unrestricted.

A corner type is a cyclically ordered triple of face colours, read counterclockwise when viewed from outside along the corner's body diagonal. Equivalently, order the three outward face normals to form a positively oriented basis. A trichromatic corner has three distinct colours and two orientations, *abc* and *acb*. BES [1, Section 2] use clockwise order; our convention

reverses every triple, preserving containment and matching. A source cube fits a target corner by a proper rotation if and only if it has the same cyclic corner type: the cyclic alignments of two corner frames are exactly those realised by proper rotations of the cube. For the corner problem ($n = 2$), a target is represented by its multiset of eight corner types. A collection builds a corner solution modelled on that target exactly when these eight occurrences can be matched to distinct cubes having the required types.

For distinct colours a, b, c, write $Q(a; b, c)$ for a target whose b and c faces are opposite and whose other four faces have colour a. Its corner multiset is $4aab + 4aac$. Write $R(a; b, c)$ for the analogous target with b and c adjacent; its corners are

$$2aaa + 2aab + 2aac + abc + acb.$$

Finally, write $D(a, b, c)$ for the target with opposite pairs aa, bb, cc; it has four corners of each orientation on $\{a, b, c\}$. These three families suffice for the corner argument. Section 5 introduces one further target for larger frames.

If every corner type of a variety v occurs on a variety w, replacing a copy of w by v cannot create a solution to any target. In particular, it preserves unsolvability. Table 1 records the reduction. Face tuples use the order $+x, -x, +y, -y, +z, -z$. The rows represent the seven orbits under colour permutations, with rotations already identified. They cover all 68 varieties; four orbits are retained.

Representative	Orbit size	Retained type	Replacement
001123	6	checkerboard C	—
010203	8	triple-corner T	—
010213	12	pair type P	—
010223	12	pair type P	—
000123	12	—	001123
001213	12	—	001123
010123	6	—	001123

Table 1: The 68 varieties reduce to 38.

A reduced variety belongs to bucket a if it has at least two of the three corner types aab with $b \neq a$. The reduction, 38 minimal varieties, orientation classes, checkerboards and buckets below are from BES [1, Section 2, Table 8 and Definition 3]; we count each bucket's seven mirror pairs as fourteen individual varieties.

Lemma 1 (Finite geometry)**.** *Every non-retained variety in Table 1 contains the corner types of a checkerboard, so every variety reduces to one of the 38 retained varieties. Their bucket incidences are:*

- *There are eight T varieties. Each has a repeated colour a, corners aaa and aab for every $b \neq a$, and belongs only to bucket a.*
- *The 24 P varieties belong to two buckets, one for each repeated colour.*
- *The six checkerboards belong to no bucket. Their repeated colours occupy opposite face pairs; with repeated colours a, b, a checkerboard has both orientations on $\{a, b, c\}$ and $\{a, b, d\}$.*

The 32 ordinary varieties (T and P) split into two classes of 16, corresponding to BES's K_o and K_m (up to interchanging the names). Every member of a class has exactly the same four

trichromatic corner types, one for each three-element colour set. The two classes have opposite orientations. In our convention one class has the trichromatic corner set

$$\Omega = \{012, 031, 023, 132\},$$

one set of BES's universal corners. *Each class contains exactly one T variety for each repeated colour. Even colour permutations preserve the classes; odd colour permutations and reflection exchange them.*

Proof. The face-colour multiplicities are $3+1+1+1$ or $2+2+1+1$. In the first case, the three equal faces either meet at a vertex or contain an opposite pair, giving two orbits. In the second case, both repeated pairs can be opposite, exactly one can be opposite, or both can be adjacent. In the last case, the two monochromatic edges either have perpendicular directions, giving two mirror configurations represented by 010213 and 010223, or are parallel, represented by 010123. These cases cover all varieties. The orbit sizes in Table 1 sum to 68, the total number of varieties [1], so the seven orbits are distinct. The three non-retained representatives contain the corner types of 001123; colour permutations transport these containments to their whole orbits. Table 2 gives the corner sets of the retained representatives. All three ordinary representatives in Table 2 belong to the class of Ω. Even colour permutations preserve Ω and odd ones reverse it, with triples read cyclically. Applying colour permutations to Table 2 gives the stated incidences and orientation classes. The eight T varieties split evenly between the classes, and even permutations act transitively on the four colours, so each class contains exactly one T variety for each repeated colour. Reflection reverses every trichromatic corner. □

Representative	Corner-type set
001123	$\{012, 013, 021, 031\}$
010203	$\{000, 001, 002, 003, 012, 023, 031, 132\}$
010213	$\{001, 003, 011, 012, 023, 031, 112, 132\}$
010223	$\{002, 003, 012, 022, 023, 031, 122, 132\}$

Table 2: Corner-type sets of the retained representatives, with cyclic triples written in lexicographically least form.

Lemma 2 (Bucket bound)**.** *Eight reduced cubes in bucket a can build a target $Q(a; b, c)$ for some b, c. Consequently, a reduced collection with no Q target has at most seven cubes in each bucket.*

Proof. Each of the eight cubes misses at most one of the three types *aab*. Thus at most one type is missed by five or more cubes. Choose two types, *aab* and *aac*, each missed by at most four cubes. Assign cubes missing *aac* to *aab* positions, and cubes missing *aab* to *aac* positions. These groups are disjoint and each has size at most four. Every remaining cube has both types, so they fill the remaining positions of $Q(a; b, c)$. □

3 The extremal collections

Theorem 1. *A collection of 28 four-colour cubes that builds no target of type Q, R or D consists of seven copies of each of the four T varieties in one ordinary orientation class.*

Proof. Replace each non-retained cube by a checkerboard using Lemma 1, leaving retained cubes unchanged. This preserves avoidance of all three target families. Let $A \geq B$ be the sizes of the two ordinary orientation classes, and let C count checkerboards, so $A + B + C = 28$. Call the class of size A the majority class (either one if equal). Call the remaining $M = B + C$ cubes

minority cubes. A minority cube *supports* a three-element colour set E if it has the orientation on E opposite to the majority class. Smaller-class cubes support every E; a checkerboard supports two.

(i) At most six minority cubes: $M \le 6$. For $E = \{a, b, c\}$, let z_E count checkerboards supporting E. If $z_E \ge 8$, those eight checkerboards build the target $D(a, b, c)$. Thus $z_E \le 7$ for each E, and, since each checkerboard supports two triples, $2C \le 28$. It follows that $A + B \ge 14$, so $A \ge 7$. There are $m_E = B + z_E$ minority cubes supporting E. If $m_E \ge 4$, four such cubes and four majority cubes build $D(a, b, c)$. Hence $m_E \le 3$ for every E. Summing gives

$$4B + 2C \le 12, \qquad M \le 6, \qquad A = 28 - M \ge 22. \tag{1}$$

(ii) No minority cubes: $M = 0$. Let t_a and p_a count only majority-class T and P cubes in bucket a, and let n_T, n_P be their respective totals, so $A = n_T + n_P$. By Lemma 2, $t_a + p_a \le 7$. Each T cube belongs to one bucket and each P cube to two. This is BES's double-count [1, equation (1)]; summing the bucket bounds and using $n_T + n_P = 28 - M$ gives

$$n_T + 2n_P \le 28, \qquad n_P \le M. \tag{2}$$

We will repeatedly use the following construction. If six majority cubes supply $2aaa + 2aab + 2aac$ and a minority cube supports $\{a, b, c\}$, then the minority cube supplies the opposite trichromatic orientation and an unused majority cube supplies the other. Such a cube is available since $A \ge 22$. This builds $R(a; b, c)$.

Suppose $M > 0$. Every minority cube supports some triple containing any given colour a. The construction above therefore rules out $t_a \ge 6$. Thus $t_a \le 5$ for every a, and $n_T \le 20$. Since $n_T + n_P + M = 28$ and $n_P \le M$, we obtain $M \ge 4$.

For any two distinct colours a, b, some minority cube supports a triple containing both. If not, every minority cube would be a checkerboard. A checkerboard whose repeated pair meets $\{a, b\}$ supports such a triple, so all would have the complementary repeated pair $\{c, d\}$, supporting exactly $\{a, c, d\}$ and $\{b, c, d\}$. Thus $m_{\{a,c,d\}} = M \ge 4$, contradicting $m_E \le 3$.

We claim that $t_a \ge 4$ forces $t_a + p_a \le 5$. Otherwise, since $t_a \le 5$, choose six majority cubes in bucket a, with four or five T cubes and the rest P cubes. These one or two P cubes share a type aab, since each has two of the three possibilities. By the preceding pair-support argument, choose a minority cube supporting $\{a, b, c\}$ for some $c \notin \{a, b\}$. With four T cubes and two P cubes, the P cubes supply $2aab$ and the T cubes supply $2aaa + 2aac$. With five T cubes and one P cube, the P cube supplies aab and the T cubes supply $2aaa + aab + 2aac$. In either case the six cubes supply $2aaa + 2aab + 2aac$, so the construction above gives a solution, proving the claim.

If $t_a \le 3$, the bucket bound gives $2t_a + p_a \le t_a + 7 \le 10$. If $4 \le t_a \le 5$, the claim gives $2t_a + p_a \le t_a + 5 \le 10$. Thus $2t_a + p_a \le 10$ in all cases. Each majority T cube contributes two through its single bucket, and each majority P cube contributes one through each of its two buckets. Summing gives

$$2A = 2n_T + 2n_P = \sum_a (2t_a + p_a) \le 40,$$

contradicting $A \ge 22$ from (1). Therefore $M = 0$.

(iii) Equal multiplicities: $t_a = 7$. With $M = 0$, (2) forces $n_P = 0$. All 28 cubes are majority-class T cubes, and the four bucket bounds force each $t_a = 7$.

Since the reduced collection contains no checkerboards, no replacement occurred: the original collection was already reduced. □

Theorem 2. *Every collection of 29 four-colour cubes contains a three-colour corner solution.*

Proof. Otherwise every 28-cube subcollection would have the form in Theorem 1. Take one such subcollection S and let v be the remaining cube. Remove from S a cube of a variety different from v, and insert v. The multiplicity of v's variety is now 1 or 8, whereas every variety in a collection described by Theorem 1 has multiplicity 0 or 7, a contradiction. □

4 Sharpness

Lemma 3. *In a cube using exactly three colours, the two orientations of a trichromatic corner occur equally often. If no trichromatic corner occurs, the cube is $Q(a;b,c)$ for some distinct a,b,c.*

Proof. For any cube coloured from a palette $\{a,b,c,d\}$, let $n(xyz)$ count corners of cyclic type xyz. Read the three directed transitions around each corner in the chosen cyclic order. Each edge between an a face and a b face contributes ab at one endpoint and ba at the other, so the total counts of these transitions agree. Corners with repeated colours contribute equally to both counts. The remaining contributions give

$$n(abc) - n(acb) + n(abd) - n(adb) = 0. \tag{3}$$

If only a,b,c occur, the last two terms vanish, proving the balance.

For the second assertion, if a colour c occurs on only one face, its four neighbours must all have the same colour a: a change from a to b around this face would produce a trichromatic corner. The opposite face is then the sole b face. Otherwise each colour occurs twice. If the two c faces are opposite, the four neighbours of either contain both a and b, forcing a change. If they are adjacent, the other three neighbours form a path containing both a and b, since each occurs only twice. Again a change is forced. Both cases produce a trichromatic corner. □

Theorem 3. *The threshold 29 in Theorem 2 is best possible. For a fixed palette, exactly two 28-cube collections fail, and they are mirror images. Up to colour permutation, there is a single obstruction.*

Proof. Take seven copies of each of the following four face tuples (Figure 1):

$$010203, \qquad 011213, \qquad 021223, \qquad 031323.$$

These are T cubes with repeated colours $0,1,2,3$, respectively, all in the same ordinary orientation class. A target with any trichromatic corner requires both orientations by Lemma 3, whereas the collection supplies only one. A target without one is $Q(a;b,c)$; every one of its eight corners has colour a twice, so only the seven T cubes of repeated colour a are usable. No three-colour target is possible. Reflection gives the analogous example in the other orientation class. Theorem 1 excludes all other collections. Every odd colour permutation exchanges the two orientation classes and hence these two collections. □

Remark (Allowing fewer colours). If targets may use at most three colours, 28 cubes suffice. In either exceptional collection, four T cubes of repeated colour a and four of repeated colour b supply $aaa + 3aab + bbb + 3bba$. These are the corners of a two-colour cube whose three a faces meet at one vertex and whose three b faces meet at the opposite vertex. Every other 28-cube collection builds a three-colour target by Theorem 3. This gives an upper bound, not an exact threshold, for that variant.




Figure 1: Why one exceptional collection fails. (A) Seven copies of each variety, with monochromatic corners marked. (B) Targets with a trichromatic corner need both orientations; the example's bottom face is hidden. (C) Targets without trichromatic corners require eight cubes with the same repeated colour, but only seven are available; a, b, c are distinct. Reflection gives the other obstruction.

Corollary 1 (Restricted targets)**.** *Every collection of 29 four-colour cubes, and every solvable collection of 28, builds a target of type Q, R or D. These families contain just 28 of the 120 target varieties.*[1]

Proof. By Theorems 1 and 3, a 28-cube collection avoiding these families is unsolvable. The swap in the proof of Theorem 2 rules out a 29-cube collection avoiding them. There are $4\binom{3}{2} = 12$ varieties each of Q and R: choose a and the unordered pair $\{b, c\}$. There are $\binom{4}{3} = 4$ varieties of D, one for each three-colour set. □

[1]For any fixed three colours, a rotation with k face cycles fixes $3^k - 3 \cdot 2^k + 3$ colourings using all three. The 24 rotations have $6, 4, 3, 2$ face cycles with multiplicities $1, 3, 12, 8$, so Burnside's lemma gives $(540+3\cdot36+12\cdot6)/24 = 30$ varieties. Four choices of omitted colour give 120.

Remark. For the palette $\{0,1,2,3\}$, write $\delta(xyz) = n(xyz) - n(xzy)$. Applying (3) to the different colour pairs gives

$$\delta(012) = \delta(031) = \delta(023) = \delta(132).$$

This common integer is the cube's *degree*, relative to Ω. Reflection negates the degree, as does each transposition of colour labels; thus even colour permutations preserve it and odd ones negate it. Each ordinary representative in Table 2 has eight distinct corner types, so each occurs once; the two ordinary classes therefore have degrees $+1$ and -1. The checkerboard representative has neither 023 nor 032, so its degree, and that of every checkerboard, is zero.

5 Frames of arbitrary side length

An n-frame consists of the eight corner cubes and the $n-2$ further cubes on each of the twelve edges of an $n \times n \times n$ cube. Thus it contains $L = 12n - 16$ cubes. Its six exterior faces are monochromatic and use exactly three colours; face interiors are not part of the frame. This is the frame model of BES [1, Section 1], with three target colours instead of four.

Theorem 4 (Frame threshold and classification)**.** *For every $n \geq 2$, every collection of $16n-3$ four-colour cubes contains an n-frame. For a fixed palette, exactly two collections of $16n-4$ cubes fail: $4n-1$ copies of each of the four T varieties in one orientation class, or their reflections. Consequently the sharp frame threshold is $F(n) = 16n-3$.*

The corner-containment reduction also preserves edge availability: an edge occurs in a corner, so containment of corner types implies containment of edge types [1, Section 2, after Definition 2]. An edge type is an unordered pair of colours, allowing repetition. Every ordinary variety has all six bichromatic edge types. A checkerboard has five, missing precisely the pair of its singleton colours; different checkerboards miss different pairs. These facts follow from the representatives in Table 2, and also appear in BES's frame arguments [1, proofs of Lemmas 7–8 and Table 5]. A cube in bucket a has an aa edge.

Lemma 4. *Let $n \geq 2$. If a reduced collection of at least $16n-4$ cubes has a D corner solution, it has an n-frame.*

Proof. If a checkerboard variety occurs at least L times, those copies build a D-frame on either supported triple. Otherwise every checkerboard variety occurs at most $L-1$ times. Remove the eight cubes in the given $D(a,b,c)$ corner solution. The remaining positions demand $4(n-2)$ copies of each of ab, ac, bc. Each type is available on at least

$$(16n-4) - 8 - (L-1) = 4n+5$$

remaining cubes. The union of the available source sets for any two types is the entire remaining collection, since no reduced cube misses two bichromatic types. There are at least $12(n-2)$ remaining cubes. These are Hall's conditions for the three groups of edge positions, so all positions can be filled with distinct copies. □

One more target supplies the growing repeated-colour demands. For distinct a, b, c, let $U(a,b;c)$ have face tuple (a,b,a,b,a,c). Its corner and edge multisets are respectively

$$aaa + 2aab + aac + abb + bbc + abc + acb, \qquad 3aa + bb + 4ab + 2ac + 2bc.$$

Thus an ordinary class supplies a U-frame using $3n-2$ of its T_a cubes, n of its T_b cubes, and $1 + 8(n-2)$ other cubes, together with one minority cube supporting $\{a,b,c\}$. The T_a cubes fill four corners and the $3(n-2)$ monochromatic aa edge positions; the T_b cubes fill two corners and the $n-2$ monochromatic bb positions. The minority cube and one further ordinary cube fill the two trichromatic corners. Every remaining edge is bichromatic and can use any ordinary cube.

Proof of Theorem 4. We first classify a failing collection of $N = 16n - 4$ cubes. The case $n = 2$ is Theorem 3, so assume $n \geq 3$. Apply the same reduction as in Theorem 1, replacing every non-retained variety by a checkerboard. Keep the notation $A, B, C, M, t_a, p_a, n_T, n_P$ for the majority class, minority class, checkerboards and majority bucket counts.

By Lemma 4, no D corner solution exists. The first step of Theorem 1 uses only these targets, so it still applies, since $N \geq 28$ preserves its preliminary bound $A \geq 7$. It gives

$$4B + 2C \leq 12, \qquad M = B + C \leq 6, \qquad A = N - M \geq 16n - 10.$$

A majority bucket cannot contain $4n$ cubes: eight build a Q corner solution by Lemma 2, another $4(n-2)$ fill its aa edge positions, and $8(n-2)$ further majority cubes fill its bichromatic edges. The supply is sufficient since $A - 4n \geq 12n - 10 \geq 8(n-2)$. Consequently

$$t_a + p_a \leq 4n - 1, \qquad n_T + 2n_P \leq 16n - 4.$$

Together with $n_T + n_P = N - M$, this implies

$$n_P \leq M, \qquad n_T \geq 16n - 16.$$

Suppose $M > 0$. Some colour a has $t_a \geq 4n - 4 \geq 3n - 2$. At most one colour has fewer than n majority T cubes: two such colours would give

$$n_T \leq 2(n-1) + 2(4n-1) = 10n - 4 < 16n - 16.$$

Choose a minority cube. It supports a triple containing a: smaller-class ordinary cubes support every triple, and a checkerboard's two supported triples together contain every colour. At least one of the other two colours in that triple has at least n majority T cubes; call it b, and call the third c. The $U(a, b; c)$ construction now applies. After selecting its $4n - 1$ majority corner and monochromatic-edge cubes, at least $A - (4n-1) \geq 12n - 9 \geq 8(n-2)$ majority cubes remain. This contradicts failure.

Hence $M = 0$, then $n_P = 0$, and the four bucket capacities force $t_a = 4n - 1$ for every a. No checkerboards remain, so no replacement occurred. This proves the classification of possible failures.

Both stated collections do fail. A target with a trichromatic corner needs both orientations by Lemma 3. Otherwise it is $Q(a; b, c)$. Its eight corners and its $4(n-2)$ monochromatic aa edge positions all require T_a cubes, giving demand $4n$ and supply $4n - 1$.

Finally, a failing collection of $16n-3$ cubes would have every $(16n-4)$-cube subcollection of the classified form. The swap in Theorem 2 changes one variety's multiplicity to 1 or $4n$, whereas the classification allows only 0 and $4n - 1$. This contradiction proves the upper bound. □

Corollary 2 (Complete cubes)**.** *For every integer $n \geq 2$, the sharp threshold for a complete $n \times n \times n$ cube with exactly three exterior colours is*

$$G(n) = \max\{n^3, 16n - 3\}.$$

In particular, $G(2) = 29$, $G(3) = 45$, and $G(n) = n^3$ for $n \geq 4$.

Proof. Both terms are necessary: an assembly uses n^3 pieces, and contains an n-frame. Conversely, construct a frame using Theorem 4. Each face-interior position prescribes just one colour, which every unused source cube has; interior positions prescribe none. Thus arbitrary unused cubes complete the assembly whenever at least n^3 pieces are available. □

6 Formal verification and computation

The frame threshold, extremal classification, complete-cube threshold and Corollary 1 are formalised in Lean 4.35.0 [3] with mathlib v4.35.0 [4] for arbitrary indexed families of four-colour cubes. Solutions select distinct copies and proper rotations matching the exposed faces at every required position. The frame statements cover every side length $n \geq 2$; completing a frame adds only the requirement that at least n^3 pieces are available. The threshold, sharpness and restricted-target statements use no catalogues; the classification counts cubes in each rotation class. Kernel checks verify the catalogues and finite geometry, and a lemma for arbitrary palettes equates cyclic corner matching with physical placement.

Thirty-five results are restated in a standalone challenge importing only mathlib. A comparator checks their statements against the proof library and replays the proofs through Lean, NanoDa and con-ron.

The audit reports three standard axioms: `propext`, `Classical.choice` and `Quot.sound`. No proof uses holes or `native_decide`. The repository[2] provides sources, reproduction commands and details of the statement checks. The physical model and palette-independent geometry come from the shared `colour-cubes-core` library [5]. The challenge restates its definitions locally.

The Lean proof uses counting and finite Hall certificates. Its corner-case proof treats checkerboards and the smaller ordinary class separately; the frame proof combines them. Finite checks establish each target's corner and edge requirements, while the allocation and counting proofs are symbolic in the side length. Neither proof enumerates all collections.

A computational search found the corner obstruction and agreed with the corner threshold; separate scripts check the geometric tables, incidences, edge constructions and sample frame obstructions. Neither proof depends on these computations.

Literature and priority

A literature search on November 1, 2026 found no earlier answer to the corner question; targeted searches for the frame extension on November 2 also found none. These searches do not establish priority. Rosenfeld's October 2026 article [2] *MacMahon Cubes and Moshe's Insanity, Puzzle 1* remains unread, and its relevance is undetermined.

Declaration of AI-generated research and review

GPT-6-Astra (OpenAI), used through Codex, developed the original research argument, computational search, Lean formalisation and manuscript under the author's direction. Claude Opus 5.5 (Anthropic), used through Claude Code, contributed the minority-cube argument, balance proof, further mathematical simplifications, illustrations and review. Subsequent Codex revisions incorporated and extended these contributions. Detailed attribution and review scope are recorded in the repository's `formalization.yaml`. The human author initiated and directed the research and accepts responsibility for the paper's content and the accuracy of this disclosure.

References

[1] E. Berkove, A. Eickemeyer and M. Schulman, *The 4-Color Cubes Puzzle*, Integers **18** (2018), A68. Journal PDF.

[2] M. Rosenfeld, *MacMahon Cubes and Moshe's Insanity, Puzzle 1*, Geombinatorics Quarterly **36**(2) (October 2026), starts at p. 78. Journal listing; full text not examined.

[2] `https://github.com/jasondavies/four-to-three-colour-cubes`, release `v1.0.0`.

[3] L. de Moura and S. Ullrich, *The Lean 4 Theorem Prover and Programming Language*, in Automated Deduction—CADE 28, LNCS **12699**, Springer (2021), 625–635. doi:10.1007/978-3-030-79876-5_37.

[4] The mathlib Community, *The Lean Mathematical Library*, in Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), ACM (2020), 367–381. doi:10.1145/3372885.3373824. arXiv:1910.09336.

[5] J. Davies, `colour-cubes-core`: palette-independent Lean definitions and proofs for face-coloured cubes, Lean library (2026). Release `v1.0.0`, `https://github.com/jasondavies/colour-cubes-core`.