Hankel determinants and the length of finite Pierce expansions

Jason Davies*

October 23, 2026

Abstract

For integers $a > b > 0$, let $P(a, b)$ be the number of steps the iteration $x \mapsto a \bmod x$ takes to reach 0 from b; it is the length of the Pierce expansion of b/a. We prove

$$P(a, b) \ll_\varepsilon a^{\beta_* + \varepsilon}, \qquad \beta_* = \frac{41301 - 3\sqrt{82849}}{137480} = 0.2941336553\ldots,$$

improving the exponent $1/3 - 2/177$ of Chase and Pandey. The tail of an orbit carries a sequence of positive integers, the Hankel determinants of a natural moment sequence, and these are small along short orbit segments. Grouping segments by such integers places the quotients of each group on curves of bounded degree, whose integer points we count with the uniform bounds of Castryck, Cluckers, Dittmann and Nguyen. The exponent improves as the degree of these curves grows, and we raise it by adjoining further integer invariants, by an effective Hilbert irreducibility sieve, and by an exact identity for the direction of the curvature vector. We also give an elementary proof of the exponent $3/10 + \varepsilon$ and a bound for orbits whose quotient increments rarely rise. All results are formalized in Lean 4 with Mathlib. Shallit's conjectured $O((\log a)^2)$ bound remains open.

1 Introduction

Let $a > b > 0$ be integers. Put $b_0 = b$ and $b_{i+1} = a \bmod b_i$ while $b_i > 0$. The sequence decreases until it reaches 0; if $b_n = 0$ we write $P(a, b) = n$. For example, $P(35, 22) = 7$, from $22, 13, 9, 8, 3, 2, 1, 0$. This is the length of the Pierce expansion of b/a [13, 14], since

$$\frac{b}{a} = \frac{1}{q_0} - \frac{1}{q_0 q_1} + \frac{1}{q_0 q_1 q_2} - \cdots, \qquad q_i = \lfloor a/b_i \rfloor.$$

Shallit [12] asked whether

$$P(a, b) = O((\log a)^2). \tag{1}$$

Shallit [14] proved $P(a, b) \ll a^{1/2}$ by "Archimedean" arguments. Erdős and Shallit [6] combined these with a divisor argument to obtain $P(a, b) \ll_\varepsilon a^{1/3+\varepsilon}$, and showed that $\max_b P(a, b)/\log a$ does not tend to 0. Chase and Pandey [5] improved the upper exponent to $1/3 - 2/177$, using exponential sum estimates, and proved $\max_b P(a, b) \gg \log a/\log\log a$ for all large a. Baraskar and Vukusic [1] study the related sets of iterated remainders at fixed iteration depth. Shallit's talk pages [15] report a bound $n^{3/10}$, described as "LLM, unpublished so far". We have not seen that argument; our elementary bound below has the same exponent up to ε, and we make no claim about how the arguments are related.

Our main result is the following uniform bound.

*jason@jasondavies.com

Theorem 1.1. *For every $\varepsilon > 0$ there is $C_\varepsilon > 0$, independent of a, b, such that*

$$P(a,b) \le C_\varepsilon a^{\beta_*+\varepsilon} \qquad (0 < b < a), \qquad \beta_* = \frac{41301 - 3\sqrt{82849}}{137480}.$$

Here $\beta_* = 0.2941336553\ldots$. The proof depends on two fixed positive integers m, n, and β_* is the limit of the resulting exponents as n/m tends to an irrational ratio. For instance $(m,n) = (3,2)$ gives the rational exponent $4317/14677 = 0.2941336785\ldots$ (Theorem 10.8).

The curve-counting bounds of this paper (Theorems 1.1, 1.2 and 1.3) all lie just above $5/17 = 0.2941176\ldots$, and Theorem 1.1 is within $1.6 \cdot 10^{-5}$ of it. We first introduce the integer invariants of an orbit, and then explain where $5/17$ comes from, since it organizes the whole argument.

Integer invariants of an orbit

Write $k_i = q_{i+1} - q_i$ for the quotient increments and $d_i = b_i - b_{i+1}$ for the drops. The *curvature* $D_i = d_i k_{i+1} - d_{i+1} k_i$ of four consecutive terms satisfies

$$D_i b_{i+2} = d_i d_{i+2} + k_i d_i d_{i+1} + (k_i - 1) d_{i+1}^2. \tag{2}$$

Thus D_i is a positive integer, although its defining terms largely cancel, and it is small when the drops are small. A six-term identity supplies a second positive integer L, and with it the bound $q_i \le 12K^3 \max(D_{i+1}, D_{i+2})$ when the next four increments are at most K (Section 5).

These integers belong to one hierarchy. Each tail carries a finite positive measure ν_i of mass b_i/a, and

$$I_r(i) = a \det\left(\int x^{j+l}\, d\nu_i\right)_{j,l<r}$$

is a positive integer when the tail has at least r positive terms. The first four invariants are b_i, d_i, D_i, L_i, in the nonterminal ranges of Theorem 6.2.

Where the exponent $5/17$ comes from

We count the T orbit terms in a remainder window $[A, 2A)$, then sum over dyadic A. The total drop across the window is less than A, and Erdős and Shallit's divisor argument shows that a drop value can repeat only $a^{o(1)}$ times; hence $A \ge T^{2-o(1)}$. Similarly the quotients increase by at most a/A across the window, so a fixed proportion of starts have their next few increments bounded by $H \asymp a/(AT) \le a^{1+o(1)}/T^3$. For such a start the ten maximal minors of a 3×5 matrix whose rows are drops, increments and a quadratic expression in the increments, among them $I_4(i), I_4(i+1), I_4(i+2)$, are integers of absolute value at most

$$X \asymp \frac{A^4 H^6}{a^3} \asymp \frac{a^3}{A^2 T^6} \le \frac{a^{3+o(1)}}{T^{10}}.$$

Their vector is the *pattern* of the start. It is an integral point of the affine cone over the Grassmannian $\mathrm{Gr}(3,5)$, and there are $X^{5+o(1)}$ such points (Section 7); the exponent 5 is the Fano index of $\mathrm{Gr}(3,5)$, so this count cannot be improved. If every pattern were shared by $O(1)$ starts, then $T \ll X^5$ would give $T^{51} \le a^{15+o(1)}$, that is, $T \le a^{5/17+o(1)}$.

The rest of the argument bounds the number of starts sharing a pattern. For a fixed pattern, three nodes $s_j = q_{i+j} - q_i$, $j = 1,2,3$, determine the increments. The nodes satisfy a quadratic equation and, on retaining the fixed dividend a, a second equation of degree at most twelve. They therefore lie on a union of curves of total degree at most 24. The coefficient-uniform bound of

Castryck, Cluckers, Dittmann and Nguyen [3] says that an integral curve over $\mathbb{Q}$ of degree δ has at most $B^{1/\delta+o(1)}$ integer points in a box of size B, whatever its coefficients. So a pattern carries $B^{1/\delta+o(1)}$ starts. Since $B \le a^{O(1)}$, the factor $B^{1/\delta}$ becomes $a^{o(1)}$ as $\delta \to \infty$, so the exponent tends to $5/17$ as the degree grows. Every refinement below is a way of raising δ, or of shrinking B, for almost all patterns.

Raising the degree

We adjoin three integer curvature minors $W = (W_0, W_1, W_2)$ and the fifth Hankel invariant $J = I_5(i)$, whose sizes satisfy

$$|W_j| \ll S := A^3H^3/a^2, \qquad |J| \ll Y := A^5H^{10}/a^4.$$

On a component of degree d, J has a pole of total order $5d/2$ at infinity, while the W_j vanish there to total order $3d/2$; since some combination of them has a further affine zero, they have at least $3d/2 + 1$ affine poles. For fixed positive integers m, n, the graph obtained by adjoining W_0^m, W_1^m, W_2^m, J^n to the nodes therefore has degree at least $\frac{5n}{2}d + m(\frac{3d}{2} + 1)$, and the box grows only to $\max(H, S^m, Y^n)$. Conics and components on which the reconstruction fails are counted separately. This gives Theorem 1.2 below.

The pattern also determines a rank-three lattice. If λ is the smallest sup norm of a nonzero lattice vector, a reduced basis gives integer gap and curvature coordinates of sizes $O(H/\lambda)$ and $O(S/\lambda^2)$. Averaged bounds for short lattice vectors shrink the box for most patterns without changing the graph degree (Section 9, Theorem 1.3).

The polar improvement

Two further facts give Theorem 1.1. First, an effective Hilbert irreducibility sieve shows that, outside a sparse set of patterns, every relevant curve component has the maximal degree 24. Second, an exact identity, $JW = N_0^2(G\mathbf{k} + \ell)$, expresses the curvature vector through three affine-linear forms in the gaps $\mathbf{k} = (k_0, k_1, k_2)$. Here N_0 is a pattern coordinate, and $G\mathbf{k}+\ell$ arises by polarizing the pattern quadratic $f(\mathbf{k}) = \mathbf{k} \cdot (G\mathbf{k} + \ell)$: the division equations say exactly that the drop vector $\mathbf{d}$ is also orthogonal to $G\mathbf{k} + \ell$, so that $W = \mathbf{d} \times \mathbf{k}$ is parallel to it.

On a component of degree 24, J has 60 zeros, all affine. Since $W = N_0^2(G\mathbf{k}+\ell)/J$, each zero of J is a pole of W unless the three linear forms also vanish there, and outside a fixed proper algebraic exceptional set of patterns this happens at most four times, counted with multiplicity. So W has at least $60 - 4 = 56$ affine poles, and the powered graph has degree

$$\deg \Gamma \ge \frac{5n}{2} \cdot 24 + m(60 - 4) = 60n + 56m,$$

compared with $60n + 37m$ from the general bound above.

The sieve and the exceptional set together discard $O_\eta(a^\eta X^{14/3})$ patterns, which are handled by the earlier estimate. Applying the curve bound in the reduced lattice coordinates and optimizing the ratio n/m gives β_* (Section 10).

Intermediate bounds

The two curve-counting stages before the polar improvement give useful bounds without the specialization sieve. Put

$$\alpha_* = \frac{1310 - \sqrt{655}}{4365} = 0.2942513\ldots, \qquad \gamma_* = \frac{471}{1559} - \frac{6\sqrt{105}}{7795} = 0.2942294\ldots.$$

Theorem 1.2 (Joint-invariant bound)**.** *For every $\varepsilon > 0$, $P(a,b) \ll_\varepsilon a^{\alpha_*+\varepsilon}$ uniformly for $0 < b < a$.*

Theorem 1.3 (Exterior lattice bound)**.** *For every $\varepsilon > 0$, $P(a,b) \ll_\varepsilon a^{\gamma_*+\varepsilon}$ uniformly for $0 < b < a$.*

Fixed powers give the rational instances 128/435 and 2201/7480 of the joint-invariant bound, and 153/520 of the exterior lattice bound; see Corollaries 8.14 and 9.12. The table records how the exponent approached 5/17 during this work. The first four rows are superseded stages, recorded for orientation only; their proofs are formalized in the repository but omitted here.

Exponent	Exponent $-$ 5/17	Main input
23/77	$4.6 \cdot 10^{-3}$	I_4 as an integer label
11/37	$3.2 \cdot 10^{-3}$	a ternary quadratic and norm equations on slices
8/27	$2.2 \cdot 10^{-3}$	the $X^{5+o(1)}$ count of minor vectors
45/152	$1.9 \cdot 10^{-3}$	lattices of pivot gaps and their short vectors
α_*	$1.3 \cdot 10^{-4}$	adjoining curvature and rank-five invariants
γ_*	$1.1 \cdot 10^{-4}$	reducing the joint coordinates in a lattice basis
β_*	$1.6 \cdot 10^{-5}$	the Hilbert sieve and the polar identity

Elementary and structured bounds

The initial identities already yield an elementary improvement over the exponent 1/3.

Theorem 1.4. *For every $\varepsilon > 0$, $P(a,b) \ll_\varepsilon a^{3/10+\varepsilon}$ uniformly for $0 < b < a$.*

The proof uses only division equations and the divisor estimate $\tau(n) \ll_\eta n^\eta$. For a window of T remainders in $[A, 2A)$,

$$T(T-1) \le 4A\Delta, \tag{3}$$

$$T^6 A^2 \ll a^3 \Delta, \tag{4}$$

where $\Delta \ll_\eta a^\eta$ is a maximum of the divisor function (Section 4). The first is the Erdős–Shallit drop argument. For the second, there are few small triples (k_i, k_{i+1}, D_i), and a divisor argument bounds how often a triple can occur. Multiplying the second bound by the square of the first eliminates A and gives $T^{10} \ll a^3\Delta^3$. The six-step identity further gives

$$T^6 A^2 \ll a^3, \tag{5}$$

without a divisor loss. This sharper estimate is used in the later arguments.

There is also a bound sensitive to the order of the increments. Let

$$R = 1 + \#\{i : 0 \le i \le P(a,b) - 3,\ k_i < k_{i+1}\}.$$

Theorem 1.5. *$P(a,b) \ll a^{1/4}\sqrt{R}$ uniformly for $0 < b < a$. In particular, nonincreasing quotient increments give $P(a,b) \ll a^{1/4}$.*

Curvature positivity makes the drops decrease between rises. The six-step identity limits repeated equal increments, and summing the resulting window bounds proves the theorem. This argument is independent of the curve-counting refinements.

Reading guide and formal verification

Sections 3–5 establish the elementary identities and bound. Sections 6–7 construct the integer invariants and their patterns. Section 8 builds the enlarged curves, Section 9 reduces their lattice coordinates, and Section 10 completes the main theorem. The independent rise bound is proved in Section 11. The conic count, graph-degree upper bound, and three-dimensional lattice reduction are proved in Appendices A, B and C.

All the unconditional results above, including their curve-counting and specialization inputs, are formalized in Lean 4 [8] with Mathlib [9]. Section 12 identifies the proof artifact and its verification scope. Section 13 explains the present 5/17 barrier; Shallit's logarithmic conjecture remains open.

2 Preliminaries

2.1 Notation

Throughout, $a > b > 0$ are integers, $b_0 = b$, $b_{i+1} = a \bmod b_i$, and $P = P(a,b)$, so that $a > b_0 > b_1 > \cdots > b_{P-1} > 0 = b_P$. Write $q_i = \lfloor a/b_i \rfloor$, so

$$a = q_i b_i + b_{i+1}, \qquad 0 \le b_{i+1} < b_i \qquad (0 \le i < P).$$

An index i of the orbit is also called a *state*; quantities "at state i" are computed from $b_i, b_{i+1}, \ldots$ and $q_i, q_{i+1}, \ldots$, and are defined whenever the orbit terms they use exist. Let $\tau(n)$ be the number of positive divisors of n. We write $F \ll_\eta G$ when $F \le C_\eta G$, with the constant uniform in the dividend and orbit. A factor denoted $a^{o(1)}$ is bounded by $C_\eta a^\eta$ for every fixed $\eta > 0$; such factors include the divisor losses below. We write $F \asymp G$ when both $F \ll G$ and $G \ll F$ hold. Absolute constants play no role in the exponents, and we do not track them: implied constants are absolute unless their dependence is indicated, and a letter c or D with a descriptive subscript denotes a suitable absolute constant introduced where it is needed. Explicit values of all constants are recorded in the Lean development.

Lemma 2.1 (Quotients increase)**.** *If $b_{i+1} > 0$ then $q_{i+1} \ge q_i + 1$.*

Proof. Since $b_{i+1} < b_i$ we have $(q_i+1)b_{i+1} \le q_i b_i + b_{i+1} = a$, so $q_{i+1} = \lfloor a/b_{i+1} \rfloor \ge q_i + 1$. □

Lemma 2.2 (Divisor bound)**.** *For every integer $k \ge 1$ there is C_k with $\tau(n)^k \le C_k\, n$ for all $n \ge 1$.*

Proof. This is standard: $\tau(n)^k/n = \prod_{p^\alpha \| n} (\alpha+1)^k/p^\alpha$. Each factor with $p \ge 2^k$ is at most $(\alpha+1)^k/2^{k\alpha} \le 1$. Each of the fewer than 2^k factors with $p < 2^k$ is at most $\sup_\alpha (\alpha+1)^k/2^\alpha < \infty$. □

2.2 Window runs

Definition 2.3. Let $A \ge 1$ be an integer. A *window run* of length T is a sequence $x_0, x_1, \ldots, x_{T-1}$ of consecutive terms of the sequence (b_i), so $x_{j+1} = a \bmod x_j$, all lying in $[A, 2A)$.

In a window run, $x_0 > x_1 > \cdots > x_{T-1}$ and $x_j \le a$.

3 The curvature of three consecutive steps

Fix four consecutive terms

$$x = b_i, \quad y = b_{i+1} = a \bmod x, \quad z = b_{i+2} = a \bmod y, \quad w = b_{i+3} = a \bmod z,$$

with $x, y, z > 0$ (and $w \geq 0$). Write

$$q = \left\lfloor \frac{a}{x} \right\rfloor, \quad u = \left\lfloor \frac{a}{y} \right\rfloor - q, \quad v = \left\lfloor \frac{a}{z} \right\rfloor - \left\lfloor \frac{a}{y} \right\rfloor, \qquad d = x - y, \quad e = y - z, \quad f = z - w.$$

In the notation of the introduction, $q = q_i$, $u = k_i$, $v = k_{i+1}$ and $d, e, f = d_i, d_{i+1}, d_{i+2}$; the short letters keep the identities below readable. By Lemma 2.1, $u, v \geq 1$, and clearly $d, e, f \geq 1$. The division equations read

$$xq + y = a, \qquad y(q+u) + z = a, \qquad z(q+u+v) + w = a. \tag{6}$$

Definition 3.1. The *curvature* of the block is $D = dv - eu$.

Lemma 3.2 (Curvature identity)**.**

$$D\,z = df + ude + (u-1)e^2.$$

In particular $D \geq 1$.

Proof. Subtracting the first two equations of (6) gives $uy = qd+e$, hence $uz = uy-ue = qd+(1-u)e$. Subtracting the last two gives $vz = (q+u)e + f$. Therefore

$$(dv - eu)z = d\big((q+u)e + f\big) - e\big(qd + (1-u)e\big) = df + ude + (u-1)e^2.$$

The right-hand side is at least $df \geq 1$, and $z > 0$, so $D > 0$. □

Lemma 3.3 (The fiber equation)**.** *Put* $M(q) = uq^2 + (u^2+u+v)q + v$ *and* $N(q) = uq + u^2 + v$. *Then*

$$x\,M(q) = a\,N(q) + D.$$

Proof. From (6), $d = (q+1)x - a$ and $e = y - z = (q+u+1)y - a = a(q+u) - q(q+u+1)x$. Substituting these into $D = dv - eu$ gives $D = x\big(uq(q+u+1) + v(q+1)\big) - a\big(u(q+u) + v\big)$, which is the claim. □

Lemma 3.4 (Size of the parameters)**.** *If* $x \leq a$ *then*

$$u\,yx \leq a(d+e), \qquad v\,zy \leq a(e+f), \qquad D\,z \leq (u+1)(d+e+f)^2.$$

Proof. From $uy = qd + e$ we get $uyx = (qx)d + ex \leq ad + ae$, since $qx \leq a$ and $x \leq a$. Similarly $vz = (q+u)e+f$ with $(q+u)y \leq a$ and $y \leq a$ gives the second bound. For the third, by Lemma 3.2, $Dz = df + ude + (u-1)e^2 \leq df + u\,e(d+e) \leq (1+u)(d+e+f)^2$. □

4 Proof of Theorem 1.4

We bound the number T of orbit terms in a window $[A, 2A)$ in two ways. Write $B = a/A$, so that the quotients in the window are about B. The drop argument (Proposition 4.2) gives $T \ll a^{1/2+o(1)}B^{-1/2}$, which is good when the quotients are large, and the curvature argument (Proposition 4.3) gives $T \ll a^{1/6+o(1)}B^{1/3}$, which is good when they are small. The two agree at $B = a^{2/5}$, where both equal $a^{3/10+o(1)}$. Multiplying suitable powers of the two bounds eliminates A (Corollary 4.4), so no case split is needed. Throughout,

$$\Delta = \Delta(a) = \max_{1 \leq n \leq 32a^7} \tau(n) \ll_\eta a^\eta$$

by Lemma 2.2; every integer to which we apply the divisor bound is at most $32a^7$.

4.1 Few blocks share a triple (u, v, D)

Lemma 4.1 (Fiber count). *Let $1 \le u, v \le a$ and $1 \le D \le 2a$. Let $M(q) = uq^2 + pq + v$ with $p = u^2 + u + v$, and let $\Lambda(q) = cq + \ell$ with $c = au$ and $\ell = a(u^2 + v) + D$. Then the set*

$$Q = \{q \in \mathbb{Z}_{\ge 0} : \ xM(q) = \Lambda(q) \text{ for some integer } x \ge 1\}$$

has at most Δ elements.

Proof. Put $U(q) = ucq + cp - u\ell$ and $R = u\ell^2 - cp\ell + c^2 v$. Expanding shows

$$c^2 M(q) - U(q)\Lambda(q) = R \tag{7}$$

identically in q. If $q \in Q$ with witness x, substituting $\Lambda(q) = xM(q)$ into (7) gives

$$R = M(q)(c^2 - U(q)x). \tag{8}$$

We bound the sizes first. From $u, v \le a$ and $D \le 2a$ we get $\ell \le 4a^3$, $p \le 3a^2$ and $c \le a^2$, so

$$|R| \le u\ell^2 + cp\ell + c^2 v \le 16a^7 + 12a^7 + a^5 \le 32a^7, \qquad c^2 \le a^4 \le 32a^7.$$

Case $R \neq 0$. By (8), $M(q)$ is a positive divisor of $|R|$. Since $u, p \ge 1$, the map $q \mapsto M(q)$ is strictly increasing on $q \ge 0$, hence injective. Therefore $|Q| \le \tau(|R|) \le \Delta$.

Case $R = 0$. Since $M(q) > 0$, (8) gives $U(q)x = c^2$. Here $c^2 > 0$ and $x \ge 1$, so $U(q)$ is a positive divisor of c^2. Since U is linear with positive leading coefficient uc, it is injective. Therefore $|Q| \le \tau(c^2) \le \Delta$. □

The case $R = 0$ cannot be dropped: it is the case in which M and Λ might share a factor.

4.2 Counting visits to a window

In this subsection $x_0, \dots, x_{T-1}$ is a window run in $[A, 2A)$, and $q_j = \lfloor a/x_j \rfloor$ are its quotients.

Proposition 4.2 (Drop bound; after Erdős–Shallit). *If $T \ge 2$ then $T(T-1) \le 4A\Delta$.*

Proof. Let $h_j = x_j - x_{j+1} \ge 1$ for $0 \le j \le T-2$. The total drop is $\sum_j h_j = x_0 - x_{T-1} \le A - 1$. Thus fewer than $(T-1)/2$ drops exceed $2A/(T-1)$. The remaining set $\mathcal{S}$ has $2|\mathcal{S}| \ge T$.

For $j \in \mathcal{S}$ we have $1 \le h_j \le \bar{h} := \lfloor 2A/(T-1) \rfloor$. Also $a + h_j = q_j x_j + x_{j+1} + x_j - x_{j+1} = (q_j + 1)x_j$, so $x_j \mid a + h_j$. Moreover $a + h_j \le 2a \le 32a^7$. Since $j \mapsto x_j$ is injective, at most $\tau(a + h) \le \Delta$ indices in $\mathcal{S}$ have $h_j = h$. Hence $|\mathcal{S}| \le \bar{h}\Delta$, and

$$T(T-1) \le 2|\mathcal{S}|(T-1) \le 2\Delta\,\bar{h}(T-1) \le 4A\Delta. \qquad \square$$

Proposition 4.3 (Curvature bound). *If $T \ge 12$ then $T^6 A^2 \ll a^3 \Delta$.*

Proof. For $0 \le j \le T-4$ let $s_j = x_j - x_{j+3}$ be the total drop of the block $x_j, x_{j+1}, x_{j+2}, x_{j+3}$. By telescoping,

$$\sum_{j=0}^{T-4} s_j = (x_0 + x_1 + x_2) - (x_{T-3} + x_{T-2} + x_{T-1}) \le 3(2A - 1) - 3A = 3A - 3.$$

Call a block small if $s_j \le 12A/T$, and let $\mathcal{S}$ be the set of small blocks. Fewer than $T/4$ of the $T-3$ blocks fail this condition, so $|\mathcal{S}| \ge T/2$ when $T \ge 12$.

Fix $j \in \mathcal{S}$. Apply Section 3 to $x = x_j$, so that $y, z, w = x_{j+1}, x_{j+2}, x_{j+3}$, all in $[A, 2A)$. Write $s = s_j = d + e + f$. Lemma 3.4 gives $uA^2 \le uyx \le a(d+e) \le as$. Multiplying by T and using $sT \le 12A$ gives $uA^2T \le 12aA$, that is,

$$u\,AT \le 12a, \qquad \text{and likewise} \qquad v\,AT \le 12a.$$

Also $DA \le Dz \le (u+1)s^2 \le 2us^2$. Multiplying by T^2 gives $DAT^2 \le 2u(sT)^2 \le 288uA^2$, so $DT^2 \le 288uA$, and then

$$D\,T^3 \le 288\,u\,AT \le 3456\,a.$$

Thus the triple (u, v, D) of a small block lies in $[1, U^*]^2 \times [1, D^*]$, where

$$U^* = \lfloor 12a/(AT) \rfloor \le a, \qquad D^* = \lfloor 3456a/T^3 \rfloor \le 2a.$$

The inequalities $U^* \le a$ and $D^* \le 2a$ hold because $AT \ge 12$ and $T^3 \ge 1728$.

Now fix such a triple and consider the small blocks that have it. By Lemma 3.3, each one gives a point $q = q_j = \lfloor a/x_j \rfloor$ of the set Q of Lemma 4.1. The map $j \mapsto q_j$ is injective on these blocks: q_j determines $x_j = \Lambda(q_j)/M(q_j)$, and x_j determines j. By Lemma 4.1, at most Δ small blocks share a triple. Consequently

$$\frac{T}{2} \le |\mathcal{S}| \le (U^*)^2 D^* \Delta \le \left(\frac{12a}{AT}\right)^2 \frac{3456a}{T^3}\Delta,$$

which is the claimed bound. □

Corollary 4.4 (Window bound)**.** *If $T \ge 12$ then $T^{10} \ll a^3\Delta^3$.*

Proof. The drop bound and $T \le 2(T-1)$ give $T^4 \le 64A^2\Delta^2$. Multiply by the curvature bound and cancel A^2. □

4.3 Conclusion of the proof

Let $0 < b < a$. For $0 \le k \le \log_2 a$ let T_k be the number of indices $i < P$ with $2^k \le b_i < 2^{k+1}$. Every b_i with $i < P$ lies in $[1, a)$, so $P = \sum_{0 \le k \le \log_2 a} T_k$. Since (b_i) is decreasing, the indices counted by T_k are consecutive. So when $T_k > 0$ they form a window run in $[A, 2A)$ with $A = 2^k$.

Inserting $\Delta \ll_\eta a^\eta$ with $\eta = 5\varepsilon/3$ into Corollary 4.4 yields $T_k \ll_\varepsilon a^{3/10+\varepsilon/2}$; the windows with $T_k < 12$ satisfy the same estimate. There are $O(\log a) \ll_\varepsilon a^{\varepsilon/2}$ windows, so summing proves Theorem 1.4. □

5 Six consecutive steps

Fix six consecutive terms $x_0 > x_1 > \cdots > x_5 \ge 0$ of the sequence (b_i) with $x_4 > 0$. Write $q = \lfloor a/x_0 \rfloor$ and let $k_0, \dots, k_3$ be the next four quotient increments, so that

$$\lfloor a/x_j \rfloor = q + k_0 + \cdots + k_{j-1} \qquad (0 \le j \le 4).$$

By Lemma 2.1, $k_0, \dots, k_3 \ge 1$. Let $f_j = x_j - x_{j+1} \ge 1$ be the drops, and let

$$D = f_0k_1 - f_1k_0, \qquad E = f_1k_2 - f_2k_1, \qquad F = f_2k_3 - f_3k_2$$

be the curvatures of the blocks starting at x_0, x_1 and x_2. Write $\mathcal{E}_j$ for the division equation $x_jq_j + x_{j+1} - a = 0$, where $q_j = \lfloor a/x_j \rfloor$. Define polynomials

$$\begin{aligned}\sigma(y_1,y_2,y_3) &= y_1^2y_2 + (y_1-1)y_2^2 + y_1y_3,\\ \chi(y_1,y_2,y_3,y_4) &= (y_1^2 + (y_1-1)y_2)y_4 + y_3(y_1^3 + y_1y_2(2y_1-1) + (y_1-1)y_2^2)\\ &\quad + y_3^2((y_1-1)(y_1+y_2)+1),\end{aligned}$$

and put

$$L = (k_1k_2 + k_2^2 + k_3)\,D - (k_0^2 + (k_0-1)k_1)E - k_2\,(k_2f_0 - k_0f_2).$$

Lemma 5.1 (The six-step identity). *1.* $\sigma(k_1,k_2,k_3)\,D - \sigma(k_0,k_1,k_2)\,E = k_1L$.

2. $q\,k_2L = \chi(k_0,k_1,k_2,k_3)\,E + \sigma(k_0,k_1,k_2)\,F$.

Proof. Part (1) is a polynomial identity in the drops and increments. Modulo k_1 we have $\sigma(k_1,k_2,k_3) \equiv -k_2^2$ and $\sigma(k_0,k_1,k_2) \equiv k_0k_2$, and $k_2D + k_0E = k_1(k_2f_0 - k_0f_2)$; so the left side is divisible by k_1, and expanding the quotient gives L. Part (2) holds modulo the division equations: $qk_2L - \chi E - \sigma(k_0,k_1,k_2)F = k_2(c_0\mathcal{E}_0 - c_1\mathcal{E}_1 + c_2\mathcal{E}_2 - c_3\mathcal{E}_3)$, where

$$\begin{aligned}c_0 &= k_1^2k_2 + k_1k_2^2 + k_1k_3 - k_2^2,\\ c_1 &= k_0^2k_2 + 2k_0k_1k_2 + k_0k_2^2 + k_0k_3 + k_1^2k_2 + k_1k_2^2 - k_1k_2 + k_1k_3 - k_2^2,\\ c_2 &= k_0^2k_1 + k_0^2k_2 + k_0k_1^2 + 2k_0k_1k_2 + k_0k_2^2 + k_0k_2 + k_0k_3 - k_1^2 - k_1k_2,\\ c_3 &= k_0^2k_1 + k_0k_1^2 + k_0k_2 - k_1^2.\end{aligned}$$

This is an identity of polynomials in a, q, the x_j and the increments. □

Corollary 5.2. *Suppose $q \geq 1$ and $k_0,\dots,k_3 \leq K$. Then $E,F \geq 1$, $L \geq 1$, and*

$$q \leq 9K^3E + 3K^3F.$$

Proof. Lemma 3.2, applied to the blocks starting at x_1 and x_2, gives $E,F \geq 1$. Clearly $\chi > 0$ and $\sigma \geq 0$, so by Lemma 5.1(2) $qk_2L > 0$. Since L is an integer, $L \geq 1$ and $qk_2 \leq \chi E + \sigma F$. Bounding each monomial gives $\chi \leq 9K^3k_2$ and $\sigma(k_0,k_1,k_2) \leq 3K^3k_2$. Dividing by k_2 gives the claim. □

So the quotient q, which is about a/A, is controlled by the small parameters of a single block of six terms. No counting over many blocks is needed.

Proposition 5.3 (Six-step window bound). *For a window run of length $T \geq 10$,*

$$T^6A^2 \ll a^3.$$

Proof. For $0 \leq j \leq T-6$ let $s_j = x_j - x_{j+5}$. By telescoping, $\sum_{j\leq T-6} s_j \leq 5(2A-1) - 5A = 5A-5$. So some j has $s_j(T-5) \leq 5A-5$, and since $T \leq 2(T-5)$, we get $s := s_j$ with $sT \leq 10A$. Apply Corollary 5.2 to $x_j,\dots,x_{j+5}$, all of which lie in $[A,2A)$, with $K = \max(k_0,\dots,k_3)$.

By Lemma 3.4 at x_j, x_{j+1} and x_{j+2}, each increment k satisfies $kA^2 \leq as$. Multiplying by T gives $KAT \leq 10a$. By Lemma 3.4 again, $EA \leq Ex_{j+3} \leq (k_1+1)s^2 \leq 2Ks^2$, and likewise $FA \leq 2Ks^2$. Hence $qA \leq 24K^4s^2$. Also $a = qx_j + x_{j+1} < (q+1)x_j < 2A(q+1) \leq 4Aq$. Therefore

$$qA \cdot A^4T^6 \leq 24\,(KAT)^4(sT)^2 \ll a^4A^2,$$

so $qA^3T^6 \ll a^4$, and $aA^2T^6 \leq 4qA^3T^6 \ll a^4$. □

Remark 5.4. In terms of $B = a/A$, Proposition 5.3 reads $T \ll a^{1/6}B^{1/3}$. This is Proposition 4.3 without the factor $\Delta^{1/6}$. Combined with Proposition 4.2 as in Corollary 4.4, it gives $T^{10} \ll a^3\Delta^2$ and another proof of Theorem 1.4.

6 The Hankel hierarchy

From now on we write $N = P(a,b)$, freeing the letter P for polynomials, so that $b_0 > \cdots > b_{N-1} > 0 = b_N$, and $k_i = q_{i+1} - q_i$. The division equation gives $b_i/a = (1 - b_{i+1}/a)/q_i$, and iterating it expresses b_i/a as the alternating Pierce series in the tail quotients $q_i, q_{i+1}, \ldots$. The construction below rewrites this series as a Stieltjes transform,

$$\frac{b_i}{a} = \int \frac{d\mu_i(x)}{q_i + x},$$

of a probability measure μ_i on the quotient gaps $q_{i+j} - q_i$ (Lemma 6.1(2)). The measure $\nu_i = \mu_i/(q_i + x)$ then has total mass b_i/a, and the Hankel determinants of its moments are the integer invariants we need.

Put $p_i(j) = q_{i+j} - q_i$ for $0 \le j < N - i$; the symbols p_i, w_i are local to this construction. Define the weights backwards from $w_{N-1}(0) = 1$ by

$$w_i(0) = 1 - \sum_{j=0}^{N-i-2} \frac{w_{i+1}(j)}{p_i(j+1)}, \qquad w_i(j+1) = \frac{w_{i+1}(j)}{p_i(j+1)} \quad (0 \le j < N - i - 1).$$

Let μ_i be the measure with mass $w_i(j)$ at $p_i(j)$, and let

$$d\nu_i(x) = \frac{d\mu_i(x)}{q_i + x}, \qquad h_i(s) = \int x^s \, d\nu_i = \sum_{j=0}^{N-i-1} \frac{w_i(j) p_i(j)^s}{q_{i+j}}.$$

The recursion is chosen to reproduce the division equation $q_i h_i(0) + h_{i+1}(0) = 1$ for $i + 1 < N$.

Lemma 6.1 (The moment model). *1. The weights are positive and sum to* 1.

2. $h_i(0) = b_i/a$, *that is,* $b_i/a = \mathbb{E}[1/(q_i + X_i)]$ *for* $X_i \sim \mu_i$; *and* $h_{N-1}(s) = 0$ *for* $s \ge 1$.

3. $h_i(s+1) = \sum_{t \le s} \binom{s}{t} k_i^{s-t} h_{i+1}(t)$ *for* $i + 1 < N$.

4. $h_i(s) = (-q_i)^s b_i/a + z$ *with* $z \in \mathbb{Z}$.

Proof. (1) Since $p_i(j+1) \ge 1$, the sum defining $w_i(0)$ is at most 1. It is strictly less: if $N - i - 1 \ge 2$, the term $j = 1$ has $p_i(2) \ge 2$; if $i = N - 2$, the last gap satisfies $q_{N-1} \ge q_{N-2} + 2$, because $b_{N-1} \mid a$ gives $q_{N-1} - q_{N-2} - 1 = q_{N-2}(b_{N-2} - b_{N-1})/b_{N-1} \ge 1$. (2) Since $p_i(j+1) = q_{i+j+1} - q_i$, the definitions give $q_i h_i(0) + h_{i+1}(0) = 1$, which is the normalized division equation; and $h_{N-1}(0) = 1/q_{N-1} = b_{N-1}/a$; since $\mu_{N-1} = \delta_0$, also $h_{N-1}(s) = 0$ for $s \ge 1$. (3) Use $p_i(j+1) = k_i + p_{i+1}(j)$ and the binomial theorem. (4) For $i = N - 1$ this holds with $z = -(-q_{N-1})^s b_{N-1}/a = -(-1)^s q_{N-1}^{s-1}$ for $s \ge 1$, since $q_{N-1} b_{N-1} = a$. Then induct backwards on i using (3), $b_{i+1}/a = 1 - q_i b_i/a$ and $k_i - q_{i+1} = -q_i$. □

The weights also explain the role of the Newton polynomials below. Write $\mu_i(g) = \int g \, d\mu_i$. For $i+1 < N$ and any polynomial g, the zero atom contributes nothing to $\mu_i(xg)$, and $p_i(j+1) w_i(j+1) = w_{i+1}(j)$ gives

$$\mu_i(xg(x)) = \mu_{i+1}(g(k_i + x)).$$

Put $s_j = q_{i+j} - q_i$ and $F_j(x) = \prod_{0 \le l < j} (x - s_l)$, with $F_0 = 1$. Iterating this identity gives the *shift rule*

$$\mu_i(F_j(x) g(x)) = \mu_{i+j}(g(x + s_j)) \qquad (i + j < N). \tag{9}$$

Indeed, each step removes the first factor of F_j and translates the remaining nodes by the next quotient gap. Thus multiplication by a Newton polynomial advances along the orbit; this is why this basis is useful for bounding its Hankel determinants.

Theorem 6.2 (Hankel determinants)**.** *For $1 \le r \le N-i$, the number $I_r(i) = a\det(h_i(j+l))_{0\le j,l<r}$ is a positive integer. Moreover $I_1(i) = b_i$, $I_2(i) = d_i$ (if $i+1<N$), $I_3(i) = D_i$ (if $i+3<N$), and $I_4(i) = L_i$ (if $i+5<N$), where D_i and L_i are the curvature and the six-step integer of the blocks starting at b_i.*

Proof. Integrality. By Lemma 6.1(4), $h_i(s+1) + q_i h_i(s) \in \mathbb{Z}$. Multiply the Hankel matrix on the left by the unit lower triangular matrix that adds q_i times row $j-1$ to row j. This makes rows $1,\dots,r-1$ integral and leaves the determinant unchanged. Expand along row 0, whose entries are $(-q_i)^l b_i/a + (\text{integer})$; then $a\det$ is an integer.

Positivity. The Hankel matrix is the Gram matrix of $1, x, \dots, x^{r-1}$ in $L^2(\nu_i)$. Its quadratic form is $\sum_j w_i(j) q_{i+j}^{-1} \big(\sum_l c_l p_i(j)^l\big)^2$. If this vanishes, the polynomial $\sum_l c_l x^l$ vanishes at the $N-i \ge r$ distinct points $p_i(j)$, so it is zero, by the invertibility of the Vandermonde matrix. So the matrix is positive definite.

The first four. First, $I_1(i) = a h_i(0) = b_i$. For $i+1<N$, write $y_j = b_{i+j}/a$. Then $h_i(0) = y_0$, $h_i(1) = y_1$ and $h_i(2) = k_i y_1 + y_2$. Hence

$$I_2(i) = a(y_0(k_i y_1 + y_2) - y_1^2) = a(y_0 - y_1) = d_i,$$

using $q_i y_0 + y_1 = q_{i+1} y_1 + y_2 = 1$. For the larger determinants we use the Newton basis. By Lemma 6.1(3), $\nu_i(xg(x)) = \nu_{i+1}(g(x+k_i))$, so ν_i obeys the same shift rule (9) as μ_i, and for $j \le l$

$$\nu_i(F_j F_l) = \nu_{i+l}(F_j(x+s_l)) = \nu_{i+l}\Big(\prod_{m<j}(x+s_l-s_m)\Big).$$

Since $F_0,\dots,F_{r-1}$ are monic of degrees $0,\dots,r-1$, the Gram matrix $(\nu_i(F_jF_l))_{j,l<r}$ has the same determinant as the Hankel matrix. Its entries are explicit: $\nu_{i+l}(1) = y_l$, and higher moments of ν_{i+l} follow from the same rule. For $r=3$ the matrix is

$$\begin{pmatrix} y_0 & y_1 & y_2 \\ y_1 & y_2 + k_i y_1 & y_3 + s_2 y_2 \\ y_2 & y_3 + s_2 y_2 & y_4 + (s_2 + k_{i+1} + k_{i+2}) y_3 + s_2 k_{i+1} y_2 \end{pmatrix}.$$

Using $y_{j+1} = 1 - q_{i+j} y_j$ for $0 \le j \le 3$ to express every entry in terms of y_0, q_i and the gaps, and expanding, gives a times its determinant as $(b_i - b_{i+1})k_{i+1} - (b_{i+1} - b_{i+2})k_i = D_i$. The case $r=4$ is the same computation with $F_0,\dots,F_3$; its entries involve $\beta_0,\dots,\beta_6$, and the same reduction to β_0, q_i and the gaps gives L_i. At the end of the orbit these computations use $h_{N-1}(s) = 0$ for $s \ge 1$, that is, $b_N = 0$. □

The ranges $i+3<N$ and $i+5<N$ cannot be relaxed to all i at which D_i or L_i is defined. For $a = 926$ and $b = 229$ the orbit is $229, 10, 6, 2$, so $N = 4$, and $I_3(1) = 984$ while $D_1 = 988$.

Remark 6.3. In these index ranges, the positivity of D and L, proved earlier by explicit identities, is an instance of the positivity of Gram determinants. The integrality of L (Lemma 5.1(1)) is an instance of the integrality of I_4. A priori $I_r(i)$ involves the moments $h_i(s)$ for $s \le 2r-2$, and hence $b_i,\dots,b_{i+2r-2}$. In fact $I_3(i)$ depends only on b_i, b_{i+1}, b_{i+2} and q_i, q_{i+1}, q_{i+2}. Similarly $I_4(i)$ depends only on $b_i,\dots,b_{i+3}$ and $q_i,\dots,q_{i+4}$: the later remainders cancel.

7 Patterns

In this section we develop the patterns used in Section 8, together with the bounds on their size and number. We keep the notation of Section 6: $b_0 > b_1 > \cdots > b_{N-1} > 0$ is the positive part of the orbit, $N = P(a,b)$, and $q_i = \lfloor a/b_i \rfloor$.

7.1 Heuristics

The exponent $3/10$ of Theorem 1.4 comes from a single balanced configuration: $T \asymp a^{3/10}$ remainders in a window $[A, 2A)$ with $A \asymp a^{3/5}$, quotients $q \asymp a^{2/5}$ and typical increments $k \asymp a^{1/10}$. At this scale the four-term Hankel invariant $I_4(i) = L(i)$ has size $\asymp aH^6/q^4 \asymp 1$, where H bounds the increments. The idea is that a positive integer of bounded size carries information. The labels $I_4(i), I_4(i+1), I_4(i+2)$, together with four further integers of the same size, form a *pattern* (made precise in Section 7.6). States with a given pattern satisfy two linear equations and one quadratic equation in five consecutive increments (Lemma 7.1), and the quotient q_i is a root of a quartic fixed by the pattern and the increments (Lemma 7.3). So the problem becomes one of counting patterns, and the states belonging to each pattern. Counting the latter with these equations alone already improves on $3/10$ (see the intermediate bounds in Section 1). Section 8 instead keeps the fixed dividend a in the equations, and this is essential: the linear and quadratic pattern relations alone admit free families. For example, the Newton coefficients $(\alpha, -3\alpha - 1, 3\alpha + 3, -\alpha - 3, 1)$ at every state, with coefficient polynomial $(z-1)^3(z-\alpha)$, and constant increments $k_j = t$ satisfy all the linear and quadratic relations for every t. The remainder equations, which involve a, exclude such a family.

7.2 The pattern matrix

Fix a state i with $i + 6 < N$ and write, for $m \geq 0$,

$$k_m = q_{i+m+1} - q_{i+m}, \qquad d_m = b_{i+m} - b_{i+m+1},$$
$$s_m = k_0 + \cdots + k_{m-1}, \qquad r_m = k_m(s_m + s_{m+1}) + k_{m+1}.$$

All indices here are local to the chosen state, and the definitions apply whenever the indicated orbit entries exist; the empty sum gives $s_0 = 0$. So $s_m = q_{i+m} - q_i$, $r_0 = k_0^2 + k_1$, $r_1 = k_1(2k_0 + k_1) + k_2$, and so on. Consider the 3×4 integer matrix

$$\mathcal{M}_i = \begin{pmatrix} d_0 & d_1 & d_2 & d_3 \\ k_0 & k_1 & k_2 & k_3 \\ r_0 & r_1 & r_2 & r_3 \end{pmatrix},$$

and let $U_m(i) = (-1)^{m+1} \det(\mathcal{M}_i \text{ with column } m \text{ deleted})$ for $0 \leq m \leq 3$.

Lemma 7.1 (Cofactor relations). *1. $U_3(i) = L(i) = I_4(i)$ and $U_0(i) = -L(i+1) = -I_4(i+1)$, where L is the six-step integer of Lemma 5.1.*

2. $\sum_m U_m(i)\, d_m = \sum_m U_m(i)\, k_m = \sum_m U_m(i)\, r_m = 0$.

Proof. (2) is the Laplace expansion of a 4×4 determinant with a repeated row. For (1), the identity $\det(\text{columns } 0, 1, 2) = L(i)$ holds as a polynomial identity in $d_0, d_1, d_2, k_0, \ldots, k_3$, by direct expansion; and $L(i) = I_4(i)$ by Theorem 6.2. Passing from i to $i+1$ shifts the d- and k-rows by one column, and, since $s_m(i+1) = s_{m+1}(i) - k_0$, it replaces r_m by $r_{m+1} - 2k_0k_{m+1}$. This row operation does not change the determinant, so $\det(\text{columns } 1, 2, 3) = L(i+1)$. □

For example, the positive part of the orbit with $(a, b) = (95, 61)$ is $61, 34, 27, 14, 11, 7, 4, 3, 2, 1$. Its first pattern matrix and cofactor vector are

$$\mathcal{M}_0 = \begin{pmatrix} 27 & 7 & 13 & 3 \\ 1 & 1 & 3 & 2 \\ 2 & 6 & 23 & 29 \end{pmatrix}, \qquad U(0) = (-81, 833, -296, 68).$$

Thus $L(0) = 68$, $L(1) = 81$, and each row of $\mathcal{M}_0$ is orthogonal to $U(0)$.

The cofactor vector has a meaning, which is the source of everything that follows. Recall the Newton polynomials $F_m(x) = \prod_{l<m}(x - s_l)$ and the measures μ_i, ν_i of Section 6.

Lemma 7.2 (The orthogonal quartic)**.** *Let $i + 6 < N$ and $C_m = U_m(i)/U_3(i)$. Then*

$$\pi_i = \sum_{m=0}^{3} C_m \, (F_{m+1} - F_m)$$

is the monic orthogonal polynomial of degree four for ν_i. Writing $\pi_i = \sum_{l\le 4} \gamma_l F_l$, one has $U_m(i)/I_4(i) = \gamma_{m+1} + \cdots + \gamma_4$ for $0 \le m \le 3$, and $\gamma_0 = I_4(i+1)/I_4(i)$.

Proof. Note that $F_{m+1} - F_m = F_m(x)(x - s_m - 1)$. The shift rule (9) holds for ν_i as well as for μ_i (proof of Theorem 6.2). Together with $\nu_{i+m}(1) = b_{i+m}/a$, $\nu_{i+m}(x) = b_{i+m+1}/a$ and, for $m \le 3$, the moments

$$\mu_{i+m}(1) = \mu_{i+m}(x) = 1, \quad \mu_{i+m}(x^2) = k_m + 1, \quad \mu_{i+m}(x^3) = k_m^2 + 2k_m + k_{m+1} + 1$$

(the first by Lemma 6.1(1), the others from $\mu_j(xg(x)) = \mu_{j+1}(g(x + k))$ with k the gap at state j), it gives, for $m \le 3$,

$$\begin{aligned}
\nu_i(F_m(x)(x - s_m - 1)) &= \nu_{i+m}(x - 1) = -d_m/a, \\
\mu_i(F_m(x)(x - s_m - 1)) &= \mu_{i+m}(x - 1) = 0, \\
\mu_i(xF_m(x)(x - s_m - 1)) &= \mu_{i+m}((x + s_m)(x - 1)) = k_m, \\
\mu_i(x^2F_m(x)(x - s_m - 1)) &= \mu_{i+m}((x + s_m)^2(x - 1)) = r_m + k_m.
\end{aligned}$$

By Lemma 7.1(2), the three rows of $\mathcal{M}_i$ are orthogonal to (C_m). Hence $\nu_i(\pi_i) = \mu_i(\pi_i) = \mu_i(x\pi_i) = \mu_i(x^2\pi_i) = 0$. Since $\mu_i = (q_i + x)\nu_i$, we have $\nu_i(xg) = \mu_i(g) - q_i\nu_i(g)$, and therefore $\nu_i(x^j\pi_i) = 0$ for $j \le 3$, inductively in j. The coefficient of F_4 in π_i is $C_3 = 1$, so π_i is monic of degree four. Finally $\gamma_l = C_{l-1} - C_l$ with $C_{-1} = C_4 = 0$, which gives the tail sums, and $\gamma_0 = -C_0 = I_4(i+1)/I_4(i)$ by Lemma 7.1(1). □

The *pattern integers* of i are the seven integers

$$\theta(i) = \big(L(i),\, L(i+1),\, L(i+2),\, U_1(i),\, U_2(i),\, U_1(i+1),\, U_2(i+1)\big),$$

and we write $\theta(i) = (N_0, N_1, N_2, u_1, u_2, v_1, v_2)$. By Lemma 7.1, $\theta(i)$ determines $U(i) = (-N_1, u_1, u_2, N_0)$ and $U(i+1) = (-N_2, v_1, v_2, N_1)$ completely, and the k-relations at i and $i + 1$ read

$$N_1k_0 = u_1k_1 + u_2k_2 + N_0k_3, \qquad N_1k_4 = N_2k_1 - v_1k_2 - v_2k_3.$$

7.3 The normalization identity and the label bound

Lemma 7.3 (The normalization identity)**.** *Let $i + 6 < N$, let π_i be as in Lemma 7.2, and let $H_4 = \det(\mu_i(x^{j+l}))_{j,l<4}$, a polynomial in $k_0, \ldots, k_4$ independent of i. Then*

$$I_4(i)\, \pi_i(-q_i) = a\, H_4(k_0, \ldots, k_4).$$

As a polynomial in q, $I_4(i)\pi_i(-q)$ has degree four and leading coefficient $I_4(i) = L(i) \ge 1$.

This is Christoffel's formula for the modification $\mu_i = (q_i + x)\nu_i$ of the measure; we include the short proof.

Proof. Write $\nu_s = \nu_i(x^s)$ and $\mu_s = \mu_i(x^s)$, so that $\nu_{s+1} + q_i\nu_s = \mu_s$. Expanding along the last row shows that

$$\det(\nu_{j+l})_{j,l<4}\,\pi_i(x) = \det\begin{pmatrix} \nu_0 & \nu_1 & \cdots & \nu_4 \\ \vdots & & & \vdots \\ \nu_3 & \nu_4 & \cdots & \nu_7 \\ 1 & x & \cdots & x^4 \end{pmatrix},$$

since the right side is a polynomial of degree four with leading coefficient $\det(\nu_{j+l})_{j,l<4}$ that is orthogonal to $1, x, x^2, x^3$ (replacing the last row by a moment row gives a repeated row). Put $x = -q_i$ and, for $l = 4, 3, 2, 1$ in turn, add q_i times column $l-1$ to column l. The last row becomes $(1, 0, 0, 0, 0)$, and the entry in row $j < 4$ and column $l \geq 1$ becomes $\nu_{j+l} + q_i\nu_{j+l-1} = \mu_{j+l-1}$. Expanding along the last row gives $\det(\nu_{j+l})\,\pi_i(-q_i) = \det(\mu_{j+l})_{j,l<4} = H_4$. Multiply by a and use $a\det(\nu_{j+l}) = I_4(i)$. The leading coefficient is that of the monic π_i. □

Lemma 7.4 (Labels). *Let $i + 6 < N$ and put $\mathcal{G} = \prod_{j<4}\mu_i(F_j^2)$. Explicitly,*

$$\mathcal{G}(k_0, \ldots, k_4) = (k_0+1)\big(k_1s_2 + s_2 + k_1 + k_2 + 1\big)\big(k_2(k_1+k_2)s_3 + g_3\big),$$

where $g_3 = s_3(k_1+k_2) + s_3k_2 + (k_1+k_2)k_2 + (k_0+2k_1+3k_2)(k_3+1) + k_3^2 + 2k_3 + k_4 + 1$. Then $L(i)\,q_i^4 \leq a\,\mathcal{G}(k_0, \ldots, k_4)$. If moreover $b_i < 2A$, then $L(i)\,a^3 \leq 256A^4\,\mathcal{G}(k_0, \ldots, k_4)$.

Proof. The Newton polynomials $F_0, \ldots, F_3$ are monic of degrees $0, \ldots, 3$. So the matrix $(\nu_i(F_jF_l))$ is $\mathcal{U}\mathcal{H}\mathcal{U}^{\mathsf{T}}$, where $\mathcal{H}$ is the Hankel matrix of ν_i and $\mathcal{U}$ is unit lower triangular. Hence it has determinant $I_4(i)/a$, and it is positive definite, as in the proof of Theorem 6.2. By Hadamard's inequality,

$$\frac{I_4(i)}{a} \leq \prod_{j<4}\nu_i(F_j^2) \leq \prod_{j<4}\frac{\mu_i(F_j^2)}{q_i},$$

since $\nu_i = \mu_i/(q_i + x) \leq \mu_i/q_i$ on the nonnegative support. By the shift rule (9) and the moments $\mu_{i+m}(1) = \mu_{i+m}(x) = 1$, $\mu_{i+m}(x^2) = k_m + 1$, $\mu_{i+m}(x^3) = k_m^2 + 2k_m + k_{m+1} + 1$, the product $\prod_{j<4}\mu_i(F_j^2)$ expands to the displayed formula. For the second claim, $a = q_ib_i + b_{i+1} < 2A(q_i+1) \leq 4Aq_i$, so $L(i)a^4 \leq 256A^4L(i)q_i^4 \leq 256A^4a\mathcal{G}$. □

7.4 Signs

Throughout this subsection the pattern and gaps arise from the same state, so the cofactor relations of Lemma 7.1 remain in force. Besides the polynomial σ of Section 5, put $\tilde{\sigma}(y_1, y_2, y_3, y_4) = y_1^2y_3 + (2y_1 - 1)y_2y_3 + y_1y_3^2 + y_1y_4$. Both σ and $\tilde{\sigma}$ are positive at positive integers.

Lemma 7.5 (Signs). *If $N_0, N_1 \geq 1$ and $k_0, \ldots, k_4 \geq 1$, then $u_1 > 0 > u_2$.*

Proof. Solving the k- and r-relations of Lemma 7.1 for (u_1, u_2) by Cramer's rule gives the exact identities

$$\begin{aligned} u_1\sigma(k_1, k_2, k_3) &= N_1\tilde{\sigma}(k_0, k_1, k_2, k_3) + N_0\sigma(k_2, k_3, k_4), \\ -u_2\sigma(k_1, k_2, k_3) &= N_1\sigma(k_0, k_1, k_2) + N_0\tilde{\sigma}(k_1, k_2, k_3, k_4), \end{aligned}$$

since the Cramer determinant is $k_1r_2 - k_2r_1 = \sigma(k_1, k_2, k_3)$. □

7.5 Good starts

Lemma 7.6 (Good starts). *Let $b_s, \dots, b_{s+T-1}$ be a window run in $[A, 2A)$ with $T \geq 36$, and put $H = \lfloor 28a/(AT) \rfloor + 1$. Call a start i* good *if $s \leq i$, $i+9 \leq s+T$ and $q_{j+1} - q_j \leq H$ for $i \leq j \leq i+6$, that is, $k_0, \dots, k_6 \leq H$ at state i. Then at least $T/2$ starts are good, and every good start satisfies $i + 8 < N$ and $b_i, b_{i+1}, b_{i+2} < 2A$.*

Proof. Let $\mathcal{B}$ be the set of $j \in [s, s+T-1)$ with $q_{j+1} - q_j > H$. Then $|\mathcal{B}|(H+1) \leq q_{s+T-1} - q_s \leq a/A$. Since $H \cdot AT > 28a$, this gives $28|\mathcal{B}| \leq T$. Each $j \in \mathcal{B}$ spoils at most seven starts, so at least $T - 9 - 7|\mathcal{B}| \geq T/2$ starts are good, as $T \geq 36$. The last assertion is immediate, since the states $i, \dots, i+8$ lie in the window. □

7.6 The full minor vector

The seven pattern integers are not independent: they are coordinates of a single vector of 3×3 minors. If all ten minors are at most X in absolute value, there are only $X^{5+o(1)}$ such vectors (Proposition 7.9), rather than the X^7 choices of seven independent integers. We therefore define the *pattern* of a state i to be the vector $p(i)$ of the ten 3×3 minors of the matrix $\widehat{\mathcal{M}}_i$ below; it determines the pattern integers $\theta(i)$.

Let $\widehat{\mathcal{M}}_i$ be the 3×5 matrix with rows $(d_0, \dots, d_4)$, $(k_0, \dots, k_4)$ and $(r_0, \dots, r_4)$, and let $p_{c_1 c_2 c_3}$ be its minor on columns $c_1 < c_2 < c_3$. Expanding the cofactors above gives

$$\theta(i) = (p_{012},\, p_{123},\, p_{234},\, p_{023},\, -p_{013},\, p_{134},\, -p_{124}).$$

Lemma 7.7 (Expansion). *Let $q = q_i$ and $s_j = q_{i+j} - q$. If $i + j + 3 < N$ then*

$$\frac{b_{i+j}}{a} = \frac{1}{q} - \frac{1+s_j}{q^2} + \frac{s_j^2 + 2s_j + k_j + 1}{q^3} - e_j, \qquad 0 \leq e_j \leq \frac{\mu_{i+j}((x+s_j)^3)}{q^4}.$$

Proof. Apply μ_{i+j} to $\frac{1}{q+y} = \frac{1}{q} - \frac{y}{q^2} + \frac{y^2}{q^3} - \frac{y^3}{q^3(q+y)}$ with $y = s_j + x \geq 0$, using $b_{i+j}/a = \nu_{i+j}(1) = \mu_{i+j}(1/(q + s_j + x))$, $\mu(1) = \mu(x) = 1$ and $\mu(x^2) = k + 1$. □

Proposition 7.8 (All minors). *Let $i + 8 < N$, $b_i < 2A$ and $k_0, \dots, k_6 \leq H$ at state i. Then every minor of $\widehat{\mathcal{M}}_i$ satisfies $|p| \leq c_X A^4 H^6 / a^3$, for an absolute constant c_X.*

Proof. Subtracting the expansions of Lemma 7.7 for j and $j+1$ gives $d_j/a = k_j/q^2 - (r_j + k_j)/q^3 + e_{j+1} - e_j$. So replacing the first row by $d'_j = d_j - a(q^{-2} - q^{-3})k_j + aq^{-3}r_j = a(e_{j+1} - e_j)$ changes no minor. Now $|d'_j| \ll aH^3/q^4$ by the error bound of Lemma 7.7, while $|k_j| \leq H$ and $|r_j| \leq 10H^2$, so $|p| \ll aH^6/q^4$. Finally $a \leq 4Aq$. □

7.7 Counting minor vectors

Proposition 7.9 (Minor vectors). *The number of vectors of minors of integer 3×5 matrices with all coordinates in $[-X, X]$ is $\ll 1 + \sum_{\kappa \leq X} \kappa^4 \tau(\kappa) \ll X^{5+o(1)}$.*

The minor vectors are the integral points of the affine cone over the Grassmannian $\mathrm{Gr}(3,5)$ in its Plücker embedding, which we call the *Plücker cone*. This cone has dimension seven, but its integral points of height at most X number about X^5, as for rational points of bounded height on $\mathrm{Gr}(3,5)$, whose anticanonical class is five times the hyperplane class [7]. So the exponent 5 is optimal. The proof uses the standard affine chart of the Grassmannian.

Proof. For a nonzero vector p choose a pivot π with $\kappa = |p_\pi|$ maximal, and let c_4, c_5 be the two columns outside π. Let $\mathbf{u}_r$ ($\mathbf{v}_r$) be the minor with the r-th pivot column replaced by c_4 (c_5), with the sign of the column permutation. Write $\mathcal{M} = \mathcal{M}_\pi(I \mid B)$ up to a permutation of columns, where $\mathcal{M}_\pi$ is the pivot block; by Cramer's rule the two columns of B are $\mathbf{u}/p_\pi$ and $\mathbf{v}/p_\pi$. The minors of $(I \mid B)$ are ± 1, the entries of B, and the 2×2 minors of B; multiplying by $\det \mathcal{M}_\pi = p_\pi$ shows that the three minors p_{w_r} using both c_4 and c_5, where w_r consists of c_4, c_5 and the rth pivot column, satisfy $p_\pi\, p_{w_r} = \pm(\mathbf{u} \times \mathbf{v})_r$. So p is determined by $(\pi, \operatorname{sgn} p_\pi, \kappa, \mathbf{u}, \mathbf{v})$, where $|\mathbf{u}_r|, |\mathbf{v}_r| \le \kappa$ and κ divides every coordinate of $\mathbf{u} \times \mathbf{v}$.

Fix $\kappa \ge 1$ and $\mathbf{u}$, and let $g = \gcd(\kappa, \mathbf{u}_1, \mathbf{u}_2, \mathbf{u}_3)$, $\ell = \kappa/g$, $\mathbf{u} = g\mathbf{u}_0$. By Bézout there are $z \in \mathbb{Z}^3$ and $t \in \mathbb{Z}$ with $z \cdot \mathbf{u}_0 + t\ell = 1$. If $\kappa \mid \mathbf{u} \times \mathbf{v}$, then $\ell \mid \mathbf{u}_0 \times \mathbf{v}$, and the identity $\mathbf{v} - (z \cdot \mathbf{v})\mathbf{u}_0 = t\ell\, \mathbf{v} + z \times (\mathbf{v} \times \mathbf{u}_0)$ shows $\mathbf{v} = c\mathbf{u}_0 + \ell y$ with $c = z \cdot \mathbf{v}$ and $y \in \mathbb{Z}^3$. Hence $\mathbf{v}$ lies in one of ℓ classes modulo ℓ, each meeting $[-\kappa, \kappa]^3$ in at most $(3g)^3$ points: at most $27\kappa g^2$ choices. Summing over $\mathbf{u}$, grouped by $g = e \mid \kappa$, gives $\sum_{\mathbf{u}} g^2 \le \sum_{e \mid \kappa} e^2 (3\kappa/e)^3 \le 27\kappa^3 \tau(\kappa)$, so at most $729\kappa^4\tau(\kappa)$ pairs $(\mathbf{u}, \mathbf{v})$. There are ten pivots and two signs. □

8 Joint integer invariants and the optimized bound

We now prove Theorem 1.2. Fix a window run in $[A, 2A)$ with $T \ge 36$ terms. Throughout this section constants may depend on fixed positive integers m, n (the powers of Section 8.9) and on a loss $\eta > 0$, but never on the dividend, the orbit or the pattern. The argument has six steps.

1. For a fixed pattern, the nodes of every good start lie on at most 24 curves (Section 8.1).
2. On these curves the quotient q_i and the curvature $I_3(i)$ are rational functions of the nodes. Components on which the reconstruction identities fail contribute $O(1)$ starts (Section 8.2).
3. The directions at infinity of the curves are controlled by the Newton coefficients of the pattern. This makes nonexceptional components have even degree (Section 8.3), shows that exceptional patterns are few, and excludes lines (Section 8.4).
4. The reconstructed functions, and hence two families of new integer coordinates W and J, have prescribed orders at infinity (Sections 8.5 and 8.6).
5. Each nonexceptional conic contributes $a^{o(1)}$ starts (Section 8.7).
6. On the remaining components, adjoining W_j^m and J^n raises the degree. The curve-counting theorem and a balancing of heights then give the exponent (Sections 8.8 and 8.9).

Notation. In this section

- $\theta = (N_0, N_1, N_2, u_1, u_2, v_1, v_2)$ are the pattern integers of Section 7, with $N_0 = I_4(i)$, and bars denote their normalized versions (10);
- s_j are the nodes, k_j the gaps, P_4, R_4 the two orthogonal quartics, Ψ the quartic norm and f the pattern quadratic;
- $D_{jl} = d_j k_l - d_l k_j$ are the 2×2 minors of drops and gaps, so that $D_{01} = I_3(i)$ is the curvature of Section 3;
- Q and E_0 are the reconstructed quotient and curvature, $W = (W_0, W_1, W_2)$ and J the joint coordinates, and S, Y their height bounds;
- e is a multiplicity at infinity, d the degree of a component, δ the degree of a graph and $\delta_{\min} = 10n + 7m$ its lower bound (Lemma 8.12).

8.1 Good starts and the pattern curve

Put

$$H = \left\lfloor \frac{28a}{AT} \right\rfloor + 1, \qquad X = \left\lfloor \frac{c_X A^4 H^6}{a^3} \right\rfloor,$$

with c_X the constant of Proposition 7.8. Let the good starts be as in Lemma 7.6; at least $T/2$ starts are good. By Proposition 7.8, the minor vector of a good start has all coordinates at most X, and $X \geq 1$ because its coordinate $p_{012} = L(i)$ is a positive integer. The pattern of a good start (Section 7.6) therefore has height at most X, and it determines $\theta(i)$. By Proposition 7.9, there are $O_\eta(a^\eta X^5)$ patterns.

The *nodes* of state i are $s_j = q_{i+j} - q_i$. They are the support points of the measure μ_i of Section 6, and the interpolation nodes of the Newton polynomials of Section 7. The three *free nodes* s_1, s_2, s_3 of a good start are integers in $[0, 3H]$. Distinct good starts with the same pattern have distinct free-node triples (Lemma 8.2 below).

For the pattern integers θ, put

$$(\bar N_1, \bar u_1, \bar u_2) = \frac{(N_1, u_1, u_2)}{N_0}, \qquad (\bar N_2, \bar v_1, \bar v_2) = \frac{(N_2, v_1, v_2)}{N_1}. \tag{10}$$

By Lemma 7.5, $\bar N_1 > 0$, $\bar u_1 > 0$ and $\bar u_2 < 0$. Treat the free nodes s_1, s_2, s_3 as coordinates and define

$$s_0 = 0, \quad s_4 = (\bar N_1 + \bar u_1)s_1 - (\bar u_1 - \bar u_2)s_2 - (\bar u_2 - 1)s_3, \quad k_j = s_{j+1} - s_j \quad (0 \leq j \leq 3),$$

$$k_4 = \bar N_2 k_1 - \bar v_1 k_2 - \bar v_2 k_3, \qquad s_5 = s_4 + k_4.$$

At a good start these are the actual nodes and gaps, by the two k-relations of Lemma 7.1; the first is equivalent to $\bar N_1 k_0 = \bar u_1 k_1 + \bar u_2 k_2 + k_3$. Let

$$\gamma = (\bar N_1,\ -\bar N_1 - \bar u_1,\ \bar u_1 - \bar u_2,\ \bar u_2 - 1,\ 1),$$

and let γ' be the same expression in $\bar N_2, \bar v_1, \bar v_2$. Put

$$P_4(x) = \sum_{j=0}^{4} \gamma_j \prod_{\ell=0}^{j-1} (x - s_\ell), \qquad R_4(x) = \sum_{j=0}^{4} \gamma'_j \prod_{\ell=1}^{j} (x - s_\ell).$$

By Lemma 7.2, at a good start P_4 is the monic orthogonal polynomial π_i of degree four for ν_i, and $R_4(x) = \pi_{i+1}(x - k_0)$ is the corresponding polynomial of the next state, written in the variable of the current state. Put $Z = xR_4$ and $f = \sum_{j=0}^{4} \gamma_j (s_j^2 + k_j)$, the *pattern quadratic*, and let

$$\begin{aligned} \Psi = {} & \sum_{j=0}^{4} \gamma_j (s_j^3 + 3s_j^2 + 3s_j(1 + k_j) + (1 + k_j)^2) + \sum_{j=0}^{3} \gamma_j k_{j+1} \\ & + \bar N_2 r_1 - \bar v_1 r_2 - \bar v_2 r_3 - k_4(2s_4 + k_4), \end{aligned}$$

where $r_j = k_j(s_j + s_{j+1}) + k_{j+1}$ for $j = 1, 2, 3$. This is a cubic polynomial in the free nodes, the *quartic norm*: at a good start it is the squared norm $\nu_i(P_4^2)$ of the orthogonal quartic (Lemma 8.1).

The degree of an affine curve is that of its projective closure, equivalently the leading coefficient of its affine Hilbert function. "Integral" means that the defining ideal is prime over $\mathbb{Q}$; it does not assert geometric irreducibility or a rational parametrization.

Lemma 8.1 (The pattern curve). *1. At a good start i, $\Psi = \nu_i(P_4^2) > 0$, and the free nodes satisfy*

$$f = 0, \qquad N_0 \operatorname{Res}(P_4, Z) = a\Psi^4. \tag{11}$$

2. For every pattern with $\bar{N}_1, \bar{u}_1 > 0 > \bar{u}_2$, the two equations (11) have degrees two and at most twelve and no common factor. Their common zero set in $\mathbb{A}^3$ is the union of at most twenty-four integral affine curves over $\mathbb{Q}$, of total degree at most twenty-four.

Proof. (1) Since $r_m = s_{m+1}^2 - s_m^2 + k_{m+1}$, summation by parts and the k-relation at state i show that $N_0 f$ is the r-relation $\sum_m U_m(i) r_m = 0$ of Lemma 7.1. Next, the shift rule $\mu_i(g \prod_{\ell<j}(x - s_\ell)) = \mu_{i+j}(g(x + s_j))$ and the moments listed in the proof of Lemma 7.4 give $\mu_i(x^3 P_4) = \sum_j \gamma_j (s_j^3 + 3s_j^2 + 3s_j(1 + k_j) + (1 + k_j)^2 + k_{j+1})$. Eliminating k_5 with the r-relation at state $i + 1$ turns this into Ψ. Since $\nu_i(xg) = \mu_i(g) - q_i \nu_i(g)$ and P_4 is orthogonal to x^3, $\nu_i(P_4^2) = \nu_i(x^4 P_4) = \mu_i(x^3 P_4) = \Psi$, and this is positive as in the proof of Theorem 6.2.

For the resultant, Lemma 6.1(3) gives $\nu_i(xg(x)) = \nu_{i+1}(g(x + k_0))$. Hence, for $j < 4$, $\nu_i(x^j Z) = \nu_{i+1}((x + k_0)^j \pi_{i+1}(x)) = 0$. So the monic quintic Z differs from the monic orthogonal polynomial $\pi^{(5)}$ of degree five for ν_i by a multiple of P_4, and $\operatorname{Res}(P_4, Z) = \operatorname{Res}(P_4, \pi^{(5)})$. Let n_j be the squared ν_i-norm of the monic orthogonal polynomial of degree j. The three-term recurrence, whose coefficients are the ratios n_j / n_{j-1}, gives

$$\operatorname{Res}(P_4, \pi^{(5)}) = \prod_{j=1}^{4} \Big(\frac{n_j}{n_{j-1}}\Big)^j = \frac{n_4^4}{n_0 n_1 n_2 n_3} = \frac{\Psi^4}{\det(\nu_i(x^{r+s}))_{r,s<4}} = \frac{a\Psi^4}{I_4(i)}.$$

Since $N_0 = I_4(i)$, this is the second equation.

(2) The degree bounds are immediate, since the resultant has total degree at most ten in the nodes. Note that $\gamma_1 < 0 < \gamma_2$, $\gamma_3 = \bar{u}_2 - 1 < -1$ and $\gamma_0, \gamma_4 > 0$. Since $s_0 = 0$ and $\sum_j \gamma_j s_j = 0$, the quadratic part of f is

$$f_2 = \gamma_1 s_1^2 + \gamma_2 s_2^2 + \gamma_3 s_3^2 + (\gamma_1 s_1 + \gamma_2 s_2 + \gamma_3 s_3)^2,$$

whose Hessian determinant is $8\gamma_1\gamma_2\gamma_3(1 + \gamma_1 + \gamma_2 + \gamma_3) = -8\gamma_0\gamma_1\gamma_2\gamma_3 \neq 0$. A ternary quadratic form of rank three is not a product of linear forms, so f_2, and hence f, is irreducible. The degree-twelve part of the second equation is $-ac^4$ with $c = \sum_j \gamma_j s_j^3$. If f divided the second equation, the irreducible form f_2 would divide c^4, hence c. Divide c by f_2 as polynomials in s_3; the s_3^2 coefficient of f_2 is $\gamma_3(1 + \gamma_3) \neq 0$. The coefficients of the remainder vanish only if $\gamma_1\gamma_2\gamma_3^2(\gamma_3^2 - 1) = 0$, which the signs above exclude. So the two equations have no common factor, their common zero set is a curve, and by Bézout's theorem its components over $\mathbb{Q}$ have total degree at most $2 \cdot 12$. □

8.2 Reconstruction and retained components

Two further identities single out the components on which the later rational functions behave as required. The idea is that the two orthogonal polynomials P_4 and Z determine the moment vector of ν_i up to scale, and the two normalized division equations fix the scale; this recovers q_i and $I_3(i)$ as rational functions of the free nodes. Concretely, form the 8×9 matrix of coefficients of $P_4, xP_4, x^2P_4, x^3P_4, Z, xZ, x^2Z, x^3Z$ in the basis $1, x, \dots, x^8$. Let c_j be $(-1)^j$ times the minor obtained by deleting column j, and define

$$Q = \frac{c_1 - c_2}{c_1 - c_0}, \qquad E_0 = a\,\frac{\det(c_{r+s})_{0 \le r,s<3}(c_1 - c_0)^3}{(c_1^2 - c_0 c_2)^3}.$$

At a good start the matrix annihilates the moment vector $(h_i(0),\dots,h_i(8))$ and has rank eight. Indeed, the resultant in (11) is positive, so P_4 and Z are coprime. The identities $q_ih_i(0)+h_i(1)=q_ih_i(1)+h_i(2)=1$, together with $h_i(0)h_i(2)>h_i(1)^2$, then give

$$h_i(j)=\frac{c_j(c_1-c_0)}{c_1^2-c_0c_2}\quad(0\le j\le 8),\qquad Q=q_i,\qquad E_0=I_3(i).$$

In particular, all displayed denominators are nonzero at good starts.

To state the identities as equations on the curve, use the raw-moment polynomials

$$M_0=1,\qquad M_{r+1}(k_0,k_1,\dots)=\sum_{t=0}^{r}\binom{r}{t}k_0^{r-t}M_t(k_1,k_2,\dots).$$

Only $M_0,\dots,M_6$ are needed; they involve $k_0,\dots,k_4$. At good starts, $M_r=\mu_i(x^r)$ for $0\le r\le 6$, by the weight recursion in Section 6. Set

$$H_3=\det(M_{r+s})_{0\le r,s<3},\qquad H_4=\det(M_{r+s})_{0\le r,s<4},\qquad H_{013}=\det(M_{r+s})_{\substack{r=0,1,2\\ s=0,1,3}}.$$

This H_4 is the polynomial of Lemma 7.3. With $p_3=\bar u_2-1-s_1-s_2-s_3$, the coefficient of x^3 in P_4, consider the *normalization* and *resolvent identities*

$$N_0P_4(-Q)=aH_4,\qquad E_0P_4(-Q)=a((Q-p_3)H_3-H_{013}).\tag{12}$$

Both hold at good starts, where $Q=q_i$, $E_0=I_3(i)$ and $P_4=\pi_i$. The first is Lemma 7.3. The second follows from the same bordering argument. Write $v_j=h_i(j)$, $M_j=\mu_i(x^j)=q_iv_j+v_{j+1}$ and $P_4(x)=x^4+p_3x^3+p_2x^2+p_1x+p_0$. Since P_4 is orthogonal to $1,x,x^2$, expanding along the last row gives

$$\det(v_{r+s})_{r,s<3}\,P_4(x)=\det\begin{pmatrix}v_j & v_{j+1} & v_{j+2} & v_{j+4}+p_3v_{j+3}\\ 1 & x & x^2 & x^4+p_3x^3\end{pmatrix}_{j=0,1,2}.$$

At $x=-q_i$, add q_i times column 1 to column 2, then q_i times column 0 to column 1; the first three columns of row j become v_j,M_j,M_{j+1}, and the last row becomes $(1,0,0,q_i^4-p_3q_i^3)$. Using $v_{s+1}=M_s-q_iv_s$ repeatedly, the last column equals $M_{j+3}+(p_3-q_i)M_{j+2}$ plus a combination of the columns M_j, M_{j+1} and $q_i^3(q_i-p_3)$ times the column $(v_j;1)$; removing these leaves $(M_{j+3}+(p_3-q_i)M_{j+2};0)$. Expanding along the last row gives

$$\det(v_{r+s})_{r,s<3}\,P_4(-q_i)=(q_i-p_3)H_3-H_{013},$$

and $a\det(v_{r+s})_{r,s<3}=I_3(i)$. Among components containing a good start, we call a component *retained* if both equations, after clearing denominators, hold identically on it.

Lemma 8.2 (Removing components by polynomial equations)**.** *After clearing denominators, the two identities in (12) are polynomial equations in the free nodes of degree at most an absolute constant D_{cut}. On a degree-d component, each identity either holds everywhere or holds at no more than $D_{\mathrm{cut}}d$ good starts. Consequently each component that is not retained contributes $O(1)$ good starts.*

Proof. Each entry of the 8×9 coefficient matrix has degree at most four in the nodes, so the cofactors c_j have bounded degree. Write $Q=Q_{\mathrm{n}}/Q_{\mathrm{d}}$ and $E_0=E_{\mathrm{n}}/E_{\mathrm{d}}$ using the formulas above, and clear denominators:

$$\Phi_N=N_0\widehat P-aH_4Q_{\mathrm{d}}^4,$$
$$\Phi_R=E_{\mathrm{n}}\widehat P-a((Q_{\mathrm{n}}-p_3Q_{\mathrm{d}})H_3-H_{013}Q_{\mathrm{d}})Q_{\mathrm{d}}^3E_{\mathrm{d}},$$

where $\widehat{P} = Q_{\mathrm{d}}^4 P_4(-Q_{\mathrm{n}}/Q_{\mathrm{d}})$. Every factor here is a fixed polynomial expression in the nodes and the pattern coordinates, so both have bounded degree in the nodes.

If a polynomial of degree r does not vanish identically on an integral curve of degree d, its zero set on that curve has at most rd distinct rational points, by Bézout's theorem. Every good start satisfies both cleared equations. Moreover, the free nodes and pattern recover $q_i = Q$, and q_i strictly increases along the fixed orbit, so distinct good starts of this pattern give distinct free-node triples. Applying the point bound to whichever equation is not an identity proves the claim; the bound $d \le 24$ makes it uniform. □

8.3 Directions at infinity

Call a pattern *exceptional* if two of the five Newton coefficients γ_j sum to zero. Since $\bar{N}_1 > 0$, $\bar{u}_1 > 0$ and $\bar{u}_2 < 0$, every γ_j is nonzero, and a pair sum can vanish only in one of the four cases

$$\bar{N}_1 + \bar{u}_2 = 1, \qquad \bar{N}_1 + \bar{u}_2 = 0, \qquad \bar{u}_1 = 1, \qquad \bar{N}_1 + \bar{u}_1 = 1. \tag{13}$$

Equivalently, the integer pattern coordinates satisfy $N_1 + u_2 = N_0$, $N_1 + u_2 = 0$, $u_1 = N_0$ or $N_1 + u_1 = N_0$.

Lemma 8.3 (Leading nodes and determinant orders)**.** *Let C be an integral component of positive degree of (11). Let P be a closed point above infinity on the normalization of the projective closure of C, write $\mathrm{ord} = \mathrm{ord}_P$, and let $e > 0$ be the multiplicity of the hyperplane at infinity at P. For a local parameter ϖ at P, let $z_j \in \mathbb{Q}(P)$ be the value at P of the regular function $\varpi^e s_j$, $0 \le j \le 4$; we call z_j the* leading node*. Then the z_j are not all equal and satisfy*

$$\sum_{j=0}^{4} \gamma_j z_j^r = 0 \qquad (0 \le r \le 3).$$

If two of these five leading nodes coincide, the pattern is exceptional. For a nonexceptional pattern, therefore,

$$\begin{aligned} \mathrm{ord}(s_l - s_j) &= -e \quad (0 \le j < l \le 4), \\ \mathrm{ord}\,\mathrm{Res}(P_4, Z) &= -10e, \\ \mathrm{ord}\, H_3 = -3e, &\qquad \mathrm{ord}\, H_4 = -6e. \end{aligned}$$

Positive order denotes a zero and negative order a pole.

Proof. All nonzero rational constants have order zero. The multiplicity e can equivalently be written as

$$e = \max_{1 \le j \le 3} \{-\mathrm{ord}\, s_j\} > 0.$$

Thus a polynomial of total node degree at most r has order at least $-re$. The values z_j lie in the residue field $\mathbb{Q}(P)$; we do not require P to be rational. We have $z_0 = 0$, and at least one of z_1, z_2, z_3 is nonzero.

The leading nodes are distinct. The definitions of the nodes give the moment equations of orders zero and one. The highest-degree part of $f = 0$ gives the equation of order two. Finally, the highest-degree part of $N_0\,\mathrm{Res}(P_4, Z) = a\Psi^4$ gives the equation of order three: the resultant has total degree at most ten, whereas the cubic part of Ψ is $\sum_j \gamma_j s_j^3$. Consequently

$$\sum_{j=0}^{4} \gamma_j z_j^r = 0 \qquad (0 \le r \le 3). \tag{14}$$

For the last of these, multiply the fixed-dividend equation by ϖ^{12e} and evaluate at P; its left side vanishes and its right side is $a(\sum_j \gamma_j z_j^3)^4$.

If two nodes coincided, there would be at most four distinct values among the z_j. Grouping the sums in (14) by those values and using the corresponding Vandermonde matrix shows that the sum of the γ_j in each group is zero. Since there are five indices and at least two groups, some group has at most two elements. It cannot have one, because every γ_j is nonzero; it must therefore give a vanishing pair sum. This proves the collision assertion. For a nonexceptional pattern it is impossible, so all five nodes are distinct, and

$$\operatorname{ord}(s_l - s_j) = -e \qquad (0 \le j < l \le 4). \tag{15}$$

The determinant orders. Write $\mathcal{R} = \operatorname{Res}(P_4, Z)$. The leading homogeneous parts needed below are

$$[\mathcal{R}]_{10} = \gamma_0\gamma_1\gamma_2\gamma_3 \prod_{0\le j<l\le 4} (s_l - s_j), \tag{16}$$

$$[H_3]_3 = k_0k_1(k_0 + k_1),$$

$$[H_4]_6 = k_0k_1k_2(k_0 + k_1)(k_1 + k_2)(k_0 + k_1 + k_2). \tag{17}$$

Here $[F]_r$ denotes the homogeneous part of total node degree r. For completeness, the degree-ten formula can be checked without expanding the resultant. In the Newton basis beginning at s_1, s_2, s_3, its 4×4 Bezout matrix for P_4, R_4 has entry degrees at most $4-\max(i,j)$, for $0 \le i, j < 4$. Only the diagonal term of its determinant can have degree ten. Its four leading diagonal entries are

$$\begin{aligned} &-\gamma_1 s_1(s_1 - s_2)(s_1 - s_3)(s_1 - s_4),\\ &-\gamma_2 s_2(s_2 - s_3)(s_2 - s_4),\\ &-\gamma_3 s_3(s_3 - s_4), \qquad -s_4. \end{aligned}$$

Their product gives the stated Vandermonde; the additional factor x in the second polynomial contributes $P_4(0) = \gamma_0$. The raw-moment recurrence gives

$$H_3 = k_0k_1(k_0 + k_1) + k_0k_2 - k_1^2,$$

and gives the displayed degree-six part of H_4, with remainder of degree at most five. These are finite polynomial identities.

Every factor in these leading products has order $-e$ by (15). The lower-degree terms have strictly larger order, so they cannot cancel the leading terms. Thus

$$\operatorname{ord}\mathcal{R} = -10e, \qquad \operatorname{ord} H_3 = -3e, \qquad \operatorname{ord} H_4 = -6e. \tag{18}$$

□

Corollary 8.4 (Even component degrees)**.** *Every positive-degree nonexceptional component of* (11) *has even degree. At each point above infinity its hyperplane multiplicity e is even, and* $\operatorname{ord}(N_0\Psi) = -5e/2$.

Proof. Taking orders in the fixed-dividend equation and using Lemma 8.3 gives $4\operatorname{ord}\Psi = -10e$. Since orders are integers and $N_0 \neq 0$, this proves both local assertions. The degree of the curve is the sum of the multiplicities e, weighted by the residue-field degrees, so it is even. The retained identities are not needed here. □

8.4 Exceptional patterns and line components

Exceptional patterns may have components of odd degree, including lines, but there are fewer of them.

Lemma 8.5 (Exceptional patterns and lines)**.** *There are $O(X^4 \log(2X)) = O_\eta(a^\eta X^4)$ exceptional patterns. In the range*

$$c_{\mathrm{L}} A^4 H^5 X < a^3, \tag{19}$$

where c_{L} is a suitable absolute constant, no line component contains a good start.

Proof. Each relation in (13) becomes a vanishing maximal minor after a fixed invertible integer column change. The transformed minor vector still has height $O(X)$. For example, if the original columns are $C_0, \dots, C_4$, then

$$\det(C_1, C_2 + C_0, C_3 - C_0) = p_{123} - p_{013} - p_{012} = N_1 + u_2 - N_0.$$

The other three relations follow by the same column additions and permutations.

A fixed zero minor reduces the Plücker relations to a simple shape by Grassmann duality. Up to sign, the ten minors p_{abc} are the Plücker coordinates q_{de}, $\{a, b, c, d, e\} = \{0, \dots, 4\}$, of the orthogonal complement of the row space, a 2-plane in $\mathbb{Q}^5$. After permuting columns we may take the vanishing minor to be $q_{34} = 0$. The Plücker relations $q_{ij}q_{kl} - q_{ik}q_{jl} + q_{il}q_{jk} = 0$ then say exactly that the integer vectors

$$\mathbf{x} = (q_{03}, q_{13}, q_{23}), \qquad \mathbf{y} = (q_{04}, q_{14}, q_{24}), \qquad \mathbf{z} = (q_{12}, -q_{02}, q_{01})$$

satisfy $\mathbf{x} \times \mathbf{y} = 0$ and $\mathbf{x} \cdot \mathbf{z} = \mathbf{y} \cdot \mathbf{z} = 0$. They have height $O(X)$ and are signed rearrangements of the nine remaining minors, so they determine the minor vector. The case $\mathbf{x} = \mathbf{y} = 0$ has $O(X^3)$ choices. Otherwise write $\mathbf{x} = r\mathbf{v}$, $\mathbf{y} = s\mathbf{v}$ with $\mathbf{v}$ primitive. In a dyadic range $R \le |\mathbf{v}|_\infty < 2R$, there are $O(R^3)$ choices for $\mathbf{v}$ and $O((X/R)^2)$ for (r, s). The equation $\mathbf{v} \cdot \mathbf{z} = 0$ has $O(X^2/R)$ solutions in the box. Indeed, take a coordinate of absolute value $n = |\mathbf{v}|_\infty$. Primitivity gives n admissible pairs of the other two coordinates modulo n, and the chosen coordinate is then determined. Since $R \le n \ll X$, this gives $O(X^2/n)$ choices. Each range therefore contributes $O(X^4)$, and summing over $O(\log(2X))$ ranges proves the pattern count. Since $X \le a^{O(1)}$, the logarithm is absorbed into a^η.

Now suppose that a line contained in the zero set of (11) passes through the free-node triple s of a good start, with direction w. Extend w to the five nodes linearly, as in Section 8.1, and write $s(t) = s + tw$. Both equations of (11) hold identically in t.

1. The resultant has degree at most ten in t, so the t^{12} coefficient of the second equation, $-a\big(\sum_j \gamma_j w_j^3\big)^4$, vanishes. Then $\Psi(s(t))$ has degree at most two, so Ψ^4 has degree at most eight, and the t^{10} coefficient $[\mathcal{R}]_{10}(w)$ of the resultant, given by (16), vanishes too: two coordinates of w coincide.

2. The direction satisfies the four moment equations of Lemma 8.3: those of orders zero and one by the definition of the nodes, order two from the t^2 coefficient of f, and order three from step (1). The grouping argument in the proof of Lemma 8.3 then shows that the pattern is exceptional, with exactly two groups of equal coordinates, of sizes two and three. After scaling, and in the order of (13), w is one of

$$(0, 1, 1, 0, 1), \quad (0, 1, 1, 0, 0), \quad (0, 0, 1, 1, 0), \quad (0, 1, 0, 0, 1).$$

3. For each of these four directions the resultant has degree at most six in t. This follows from a row and column degree count in the Newton-basis Bezout matrix used in the proof of Lemma 8.3, in which coinciding leading nodes lower the degrees of the corresponding entries. Hence Ψ^4 has degree at most six, $\Psi(s(t))$ is affine in t, and its second difference $\Psi(s+w)+\Psi(s-w)-2\Psi(s)$ vanishes.

4. Expanding this second difference, and using $f=0$ at $s\pm w$, gives respectively

$$-2\bar{N}_1 s_3, \qquad 2(\bar{N}_1+\bar{u}_1)(k_1-1), \qquad 2(\bar{u}_2-1)(k_2-1), \qquad 2((\bar{u}_1-\bar{u}_2)-(s_4-s_1)).$$

The factors $\bar{N}_1$, $\bar{N}_1+\bar{u}_1$ and $\bar{u}_2-1$ are nonzero, so respectively

$$s_3=0, \qquad k_1=1, \qquad k_2=1, \qquad s_4-s_1=\bar{u}_1-\bar{u}_2.$$

The first case is impossible at a good start, where $s_3\geq 3$. The polynomial $\mathcal{G}$ in Lemma 7.4 satisfies, by expansion,

$$\mathcal{G}\leq H^5(12k_1+464), \qquad \mathcal{G}\leq H^5(12k_2+464).$$

In the last obstruction, $k_1+k_2+k_3=\bar{u}_1-\bar{u}_2=(u_1-u_2)/N_0\leq 2X$. Thus each surviving case gives $k_1\leq 2X$ or $k_2\leq 2X$. The label bound $N_0 q_i^4\leq a\mathcal{G}$, with $N_0\geq 1$, now gives

$$q_i^4\leq aH^5(24X+464).$$

But $q_i\geq a/(4A)$ and $24X+464\leq 488X$ for $X\geq 1$, so $a^3\leq 4^4\cdot 488\,A^4H^5X$. Condition (19) with $c_{\mathrm{L}}=4^4\cdot 488$ gives the strict reverse inequality. □

8.5 Orders of the reconstructed functions

Lemma 8.6 (Orders of the reconstructed functions)**.** *Let C be a retained nonexceptional component of positive degree containing a good start. At every point above infinity, with hyperplane multiplicity e, one has*

$$\operatorname{ord} Q=-\frac{3e}{2}, \qquad \operatorname{ord} E_0=\frac{3e}{2}.$$

In particular, E_0 is a nonzero rational function vanishing at every point above infinity.

Proof. The good start guarantees that the reconstruction denominators are not identically zero on C. Thus Q and E_0 are rational functions on C, and retention makes both identities (12) identities in its function field.

The quotient order. Write $P_4(x)=x^4+p_3x^3+p_2x^2+p_1x+p_0$. Its Newton expression gives $\deg p_j\leq 4-j$, and hence $\operatorname{ord} p_j\geq -(4-j)e$. The first retained identity gives

$$\operatorname{ord} P_4(-Q)=\operatorname{ord}(aH_4/N_0)=-6e.$$

If $\operatorname{ord} Q\geq -e$, every term of $P_4(-Q)$ would have order at least $-4e$, a contradiction. Put $r=\operatorname{ord} Q<-e$. For $0\leq j<4$,

$$\operatorname{ord}(p_j(-Q)^j)-4r \;\geq\; (4-j)(-e-r)>0.$$

Thus Q^4 is the unique term of lowest order. It follows that $4r=-6e$, or $\operatorname{ord} Q=-3e/2$.

The reconstructed curvature order. The coefficient $p_3=\bar{u}_2-1-s_1-s_2-s_3$ has order at least $-e$. The raw-moment recurrence also shows that H_{013} has degree at most four: its degree-four part is $(2k_0+k_1)k_0k_1(k_0+k_1)$. Consequently

$$\operatorname{ord}(Q-p_3)=-3e/2, \qquad \operatorname{ord}((Q-p_3)H_3)=-9e/2<-4e\leq \operatorname{ord} H_{013}.$$

There is again a unique term of lowest order. Dividing the second retained identity by $P_4(-Q)$ therefore gives

$$\operatorname{ord} E_0 = -9e/2 - (-6e) = 3e/2.$$

The argument uses the discrete valuation on the normalization, so it also covers branches above singular points of the original curve. □

8.6 The joint coordinates

Put $d_j = b_{i+j} - b_{i+j+1}$ and

$$D_{01} = d_0k_1 - d_1k_0, \quad D_{02} = d_0k_2 - d_2k_0, \quad D_{12} = d_1k_2 - d_2k_1.$$

The first of these, D_{01}, is the curvature $I_3(i)$. With $s_j = q_{i+j} - q_i$ and $r_j = k_j(s_j + s_{j+1}) + k_{j+1}$, the Hankel polynomial H_3 of Section 8.2 and its shift are

$$H_3 = k_0r_1 - k_1r_0 = \sigma(k_0, k_1, k_2),$$
$$H_3' = k_1r_2 - k_2r_1 = \sigma(k_1, k_2, k_3),$$

with σ as in Section 5. They are positive at every state at which they are defined. On a retained component, use the reconstructed function E_0 to define

$$E_{12} = \frac{H_3'E_0 - N_0k_1}{H_3}, \qquad E_{02} = \frac{k_0E_{12} + k_2E_0}{k_1}, \qquad W = (E_{12}, -E_{02}, E_0), \qquad J = N_0\Psi.$$

The determinant and cross-product identities give, identically,

$$W \cdot (k_0, k_1, k_2) = 0, \qquad W \cdot (r_0, r_1, r_2) = N_0. \tag{20}$$

At a good start these rational functions take the integer values $W = (D_{12}, -D_{02}, D_{01})$. Changing the rank-five Gram basis to $1, x, x^2, x^3, P_4$ gives

$$\det\big(\nu_i(x^{r+s})\big)_{r,s<5} = \det\big(\nu_i(x^{r+s})\big)_{r,s<4}\,\nu_i(P_4^2).$$

Consequently $J = I_5(i)$ is a positive integer at good starts. The heights of these integers are controlled by

$$S = \left\lfloor \frac{c_S A^3 H^3}{a^2} \right\rfloor, \qquad Y = \left\lfloor \frac{c_Y A^5 H^{10}}{a^4} \right\rfloor,$$

for suitable absolute constants c_S, c_Y.

Lemma 8.7 (Heights of the joint coordinates)**.** *At good starts, $|W_j| \le S$ and $1 \le J \le Y$.*

Proof. Write $q = q_i$, and recall $q \ge a/(4A)$ (proof of Lemma 7.4). For $j \le 3$, apply μ_{i+j} to the identity $(q+y)^{-1} = q^{-1} - yq^{-2} + y^2/(q^2(q+y))$ with $y = s_j + x$, as in Lemma 7.7. Since $\mu_{i+j}((s_j+x)^2) = s_j^2 + 2s_j + k_j + 1 \ll H^2$, this gives

$$\frac{b_{i+j}}{a} = \frac{1}{q} - \frac{s_j + 1}{q^2} + \epsilon_j, \qquad 0 \le \epsilon_j \ll H^2/q^3.$$

Subtracting consecutive expansions, $d_j/a = k_j/q^2 + \delta_j$ with $|\delta_j| \ll H^2/q^3$. In $D_{jl} = d_jk_l - d_lk_j$ the leading terms cancel, so $|D_{jl}|q^3 \ll aH^3$, and $q \ge a/(4A)$ gives $|D_{jl}| \le S$ for suitable c_S, including the floor because the minors are integers.

For $J = I_5(i)$, argue as in the proof of Lemma 7.4 with five Newton polynomials instead of four: Hadamard's inequality and $\nu_i \le \mu_i/q$ give $Jq^5 \le a\prod_{j<5}\mu_i(F_j^2)$. By the shift rule, $\mu_i(F_j^2) = \mu_{i+j}(F_j(x+s_j))$ is a polynomial of degree j in the gaps, and the gap bound gives $\mu_i(F_j^2) \ll H^j$. Hence $Jq^5 \ll aH^{10}$, and $q \ge a/(4A)$ gives $J \le Y$ for suitable c_Y. □

Corollary 8.8 (Orders of the joint coordinates)**.** *In the setting of Lemma 8.6, the joint coordinates satisfy*

$$\operatorname{ord} W_0 = \operatorname{ord} W_2 = 3e/2, \qquad \operatorname{ord} W_1 \geq 3e/2, \qquad \operatorname{ord} J = -5e/2. \tag{21}$$

Proof. The cubic leading part of H_3' is $k_1k_2(k_1+k_2)$. By (15), it has order $-3e$, whereas the quadratic remainder $k_1k_3 - k_2^2$ has order at least $-2e$. Thus $\operatorname{ord} H_3 = \operatorname{ord} H_3' = -3e$ and $\operatorname{ord} k_1 = -e$. In the numerator defining E_{12} the two terms have different orders:

$$\operatorname{ord}(H_3' E_0) = -3e/2 < -e = \operatorname{ord}(N_0 k_1).$$

Therefore $\operatorname{ord} E_{12} = 3e/2$. Each term in $k_0E_{12} + k_2E_0$ has order $e/2$, so their sum has order at least $e/2$. Dividing by k_1 gives $\operatorname{ord} E_{02} \geq 3e/2$. Cancellation here is harmless: equality for the first and third coordinates already gives $\min_j \operatorname{ord} W_j = 3e/2$. The formula for $J = N_0\Psi$ is Corollary 8.4. □

8.7 Counting the nonexceptional conics

Proposition 8.9 (Conics)**.** *A nonexceptional component of degree two contains the free-node triples of $O_\eta(a^\eta)$ good starts, with an implied constant independent of the pattern and the component. Consequently all nonexceptional conics together contribute $O_\eta(a^\eta X^5)$ good starts.*

The idea is short. A component that is not retained contributes $O(1)$ good starts (Lemma 8.2). A retained conic through three free-node triples of good starts lies in a plane, and Corollary 8.4 leaves it a single rational branch at infinity, of multiplicity two. So it is a parabola, the image of a quadratic polynomial map $t \mapsto (\xi_0(t), \xi_1(t), \xi_2(t))/\Delta_c$ whose parameter t is an integer at every integer point. By Lemma 8.6, the reconstructed curvature E_0 is a rational function of t with a zero of order three at infinity, and it takes the integer value $I_3(i)$ at each good start. Its nonconstant denominator must then divide a fixed nonzero resultant at each such t, which leaves $a^{o(1)}$ possibilities. Appendix A gives the details, including the uniform bounds on the coefficients.

8.8 The extra degree and the powered graph

The conic count leaves retained nonexceptional components of degree at least four. Fix such a component C of degree d. The additional integer coordinates will increase its degree without increasing their heights too much. The key is one extra zero over the affine part of C.

All divisors in this argument are on the normalization $\widetilde{C}$ of the projective closure of C, over $\mathbb{Q}$. For a closed point P of $\widetilde{C}$, write $\deg P = [\mathbb{Q}(P) : \mathbb{Q}]$, where $\mathbb{Q}(P)$ is the residue field; thus $\deg(\sum_P a_P[P]) = \sum_P a_P \deg P$. Let $E_\infty = \sum_{P|\infty} e_P[P]$ be the pullback of the original hyperplane at infinity. Then $\sum_{P|\infty} e_P \deg P = d$. We do not require the closed points, or the branches over a rational point of C, to be rational.

Let $\mathcal{F}$ be the pole envelope of the curvature coordinates:

$$\mathcal{F} = \sum_P \max\{0, -\operatorname{ord}_P(W_0), -\operatorname{ord}_P(W_1), -\operatorname{ord}_P(W_2)\}[P],$$

with $\operatorname{ord}_P(0) = +\infty$. The orders in (21) show that $\mathcal{F}$ is supported over the affine part of C, and hence is disjoint from E_∞.

Lemma 8.10 (An additional affine zero)**.** *There is a rational linear combination ϕ of W_0, W_1, W_2 which is not the zero function and has a positive-order zero at a closed point P_0 of $\widetilde{C}$ outside E_∞.*

Proof. First, the W_j cannot all be scalar multiples, in $\mathbb{Q}(C)$, of a fixed nonzero vector $v \in \mathbb{Q}^3$. Such a relation, together with $W \cdot (k_0, k_1, k_2) = 0$, would put C in the plane $v_0k_0 + v_1k_1 + v_2k_2 = 0$. This is a genuine plane: the first three gaps are an invertible linear change of the three node coordinates. The pattern quadratic is irreducible, so its restriction to this plane is nonzero and has degree at most two. Every curve contained in this plane section therefore has degree at most two, contrary to $d \geq 4$.

Choose the free-node triple $\mathbf{s}_* \in C(\mathbb{Q})$ of a good start. All the W_j are regular there, because their common reconstruction denominator does not vanish at a good start. Write $w_j = W_j(\mathbf{s}_*)$. The identity $W \cdot (r_0, r_1, r_2) = N_0 \neq 0$ gives $w \neq 0$. Choose l with $w_l \neq 0$. At least one function $w_lW_j - w_jW_l$ is nonzero: otherwise $W = (W_l/w_l)w$, contradicting the preceding paragraph. Take such a function for ϕ. It has rational coefficients and satisfies $\phi(\mathbf{s}_*) = 0$.

This zero survives normalization even if $\mathbf{s}_*$ is singular. Indeed, write $\phi = g/g_0$ near $\mathbf{s}_*$, with g not identically zero on C, $g(\mathbf{s}_*) = 0$ and $g_0(\mathbf{s}_*) \neq 0$. The finite normalization has a closed point P_0 above $\mathbf{s}_*$. In its discrete valuation ring the numerator is in the maximal ideal and the denominator is a unit, so $\operatorname{ord}_{P_0}(\phi) > 0$. Since $\mathbf{s}_*$ is affine, P_0 does not lie in E_∞. □

Lemma 8.11 (The curvature pole degree)**.** *The pole envelope satisfies*

$$\deg \mathcal{F} \geq \frac{3d}{2} + 1. \tag{22}$$

Proof. Use the function ϕ from Lemma 8.10. At every point over infinity, each W_j has order at least $3e_P/2$. The same is true of a rational linear combination; cancellation can only increase its zero order. Consequently

$$(\phi)_0 \;\geq\; \frac{3}{2}E_\infty + [P_0], \qquad (\phi)_\infty \;\leq\; \mathcal{F}.$$

The first divisor is integral because every e_P is even. The function ϕ is nonzero and has a zero, so its principal divisor has equal zero and pole degrees. Hence

$$\deg \mathcal{F} \;\geq\; \deg(\phi)_\infty = \deg(\phi)_0 \;\geq\; \frac{3}{2}\sum_{P|\infty} e_P \deg P + \deg P_0 \;\geq\; \frac{3d}{2} + 1.$$

The last inequality uses only $\deg P_0 \geq 1$, not that P_0 is rational. □

For positive integers m, n, let $\Gamma_{m,n} \subseteq \mathbb{A}^7$ be the closure of the graph obtained by adjoining W_0^m, W_1^m, W_2^m, J^n to the three original node coordinates. The graph is taken on the common domain of these rational functions. It is an integral curve, and projection to the first three coordinates is birational: on that domain the four remaining coordinates are uniquely determined. Thus $\Gamma_{m,n}$ and C have the same function field. Taking powers does not lose this property, because the original coordinates remain in the graph.

Lemma 8.12 (Degree of the powered graph)**.** *If $\delta = \deg \Gamma_{m,n}$, then*

$$2\delta \geq (5n + 3m)d + 2m, \qquad \delta \geq 10n + 7m =: \delta_{\min}. \tag{23}$$

Moreover $\delta \leq D_{\mathrm{gr}}(m + n)d$ for an absolute constant D_{gr}. For $m = n = 1$ the lower bound reads $\delta \geq 4d + 1 \geq 17$.

Proof. Lower bound. At a point P over the old infinity hyperplane, the original node coordinates have pole orders at most e_P, while J^n has pole order $5ne_P/2$ and all W_j^m vanish. Since $n \geq 1$, the pole envelope of the seven graph coordinates at P is exactly $5ne_P/2$. Over the affine part, the node

coordinates and J are regular: $J = N_0\Psi$ is polynomial in the nodes. There the envelope is exactly m times the curvature envelope. The complete envelope of the graph coordinates is therefore

$$\mathcal{F}_{m,n} = \frac{5n}{2}E_\infty + m\mathcal{F}.$$

There is a single rational linear combination λ of the seven coordinates whose pole divisor is $\mathcal{F}_{m,n}$. To see this, fix a point P in the finite support of $\mathcal{F}_{m,n}$ and multiply all seven coordinates by a local uniformizer to the power $\mathcal{F}_{m,n}(P)$. Their values in $\mathbb{Q}(P)$ are not all zero. Cancellation of the leading pole is therefore a proper $\mathbb{Q}$-linear condition on the seven coefficients. The finitely many such proper subspaces do not cover $\mathbb{Q}^7$. Choosing coefficients outside their union gives the asserted simultaneous absence of cancellation, also when $\mathbb{Q}(P) \neq \mathbb{Q}$.

As a linear function on $\Gamma_{m,n}$, λ has degree at most δ. For example, homogenize it on the projective closure of the graph: its pencil with the homogenizing coordinate consists of hyperplane sections of degree δ. Removing common zeros can only decrease the degree of the resulting map to $\mathbb{P}^1$. Birational projection identifies its function field with $\mathbb{Q}(C)$, so this degree is $\deg(\lambda)_\infty = \deg \mathcal{F}_{m,n}$. It follows that

$$\delta \;\geq\; \frac{5n}{2}d + m\deg\mathcal{F} \;\geq\; \frac{5n+3m}{2}d + m.$$

This is the first inequality in (23); $d \geq 4$ gives the second. The auxiliary linear combination is used only to measure degree: no bound on its coefficients enters the integer-point box.

Upper bound. This is a routine count of polynomial degrees, given in Appendix B. □

Every good start whose free-node triple lies on C gives an integer point on this graph. By Lemma 8.7, its coordinates have absolute values at most

$$\max(3H, S^m, Y^n).$$

The original node coordinates also show that distinct such node triples remain distinct after adjoining the powers.

8.9 The window bound and the final exponents

We use the coefficient-uniform affine-curve theorem of Castryck–Cluckers–Dittmann–Nguyen [3, Theorem 3], also stated as [4, Corollary 2.2]: an integral affine curve C over $\mathbb{Q}$ of degree δ in $\mathbb{A}^r$, with $r \geq 2$ fixed, has

$$N_{\mathrm{aff}}(C, B) \ll_r \delta^3 B^{1/\delta}(\log B + \delta), \qquad B \geq 1, \tag{24}$$

integer points in the box $[-B, B]^r$, with an implied constant independent of the coefficients of C. Its formal proof is discussed in Section 12.

Proposition 8.13 (Window bound)**.** *Fix positive integers m, n and $\eta > 0$, and put $\mathcal{B} = \max(H, S^m, Y^n)$. For a window run in $[A, 2A)$ with $T \geq 36$ satisfying (19),*

$$T \ll_{m,n,\eta} a^\eta \left(X^5 + X^4 H^{1/2} + X^5 \mathcal{B}^{1/\delta_{\min}}\right). \tag{25}$$

Proof. At least $T/2$ starts are good, and for each pattern the free-node triples of its good starts are distinct (Lemma 8.2). By Lemma 8.1, these triples lie on at most twenty-four integral curves of degree at most twenty-four, and we bound the good starts on each curve, allowing overlaps. All the heights H, S, Y and X are at most a fixed power of a, so the logarithm in (24) is $O_\eta(a^\eta)$.

- *Exceptional patterns.* There are $O_\eta(a^\eta X^4)$ of them (Lemma 8.5). Under (19) no line contains a good start, so every relevant curve has degree $2 \le \delta \le 24$, and (24) in $\mathbb{A}^3$ with $B = 3H$ gives $O_\eta(a^\eta H^{1/2})$ good starts per pattern. This gives the term $X^4 H^{1/2}$.

- *Nonexceptional patterns: components that are not retained, and conics.* By Lemma 8.2 and Section 8.7, these contribute $O_\eta(a^\eta)$ per pattern, hence $O_\eta(a^\eta X^5)$ in all.

- *Remaining nonexceptional components.* Lines do not occur, by Lemma 8.5, and odd degrees do not occur, by Corollary 8.4, so these components have degree $d \ge 4$. Their good starts give distinct integer points of $\Gamma_{m,n} \subseteq \mathbb{A}^7$ in a box of side $\max(3H, S^m, Y^n) \le 3\mathcal{B}$. By Lemma 8.12, the degree δ of $\Gamma_{m,n}$ satisfies $\delta_{\min} \le \delta \le 24 D_{\mathrm{gr}}(m+n)$, so (24) gives $O_{m,n,\eta}(a^\eta \mathcal{B}^{1/\delta_{\min}})$ good starts per component. This gives the term $X^5 \mathcal{B}^{1/\delta_{\min}}$.

The added coordinates W_j and J are functions of the free nodes on each component. They therefore introduce no further freely counted coordinates, and the pattern count is not enlarged. □

Corollary 8.14. *For fixed positive integers m, n and every $\varepsilon > 0$,*

$$P(a,b) \ll_{m,n,\varepsilon} a^{E(m,n)+\varepsilon}, \qquad E(m,n) = \max\left(\frac{15\delta_{\min} + m}{51\delta_{\min} + 3m}, \frac{15\delta_{\min} + 6n}{51\delta_{\min} + 20n} \right),$$

where $\delta_{\min} = 10n + 7m$. In particular $E(1,1) = 128/435$ and $E(50,51) = 2201/7480$.

Proof. We bound the number T of terms in each dyadic window, eliminating the window scale A. If $T < 36$ there is nothing to prove. If $AT > 28a$, then Proposition 5.3 gives $T^4 \ll a$. Otherwise $HAT \le 56a$, and Proposition 4.2 gives $T^2 \le 8\Delta A$. Substituting $H \le 56a/(AT)$ and then $A \ge T^2/(8\Delta)$ into the definitions of X, S and Y gives

$$H \ll \frac{\Delta a}{T^3}, \qquad X \ll \frac{\Delta^2 a^3}{T^{10}}, \qquad S \ll \frac{a}{T^3}, \qquad Y \ll \frac{\Delta^5 a^6}{T^{20}}. \tag{26}$$

Every power of Δ is $O(a^\eta)$, and we absorb such factors into $a^{O(\eta)}$.

- If (19) fails, then $c_{\mathrm{L}} A^4 H^5 X \ge a^3$. With $Xa^3 \le c_X A^4 H^6$ this gives $A^8 H^{11} \gg a^6$, and $H \le 56a/(AT)$ turns it into $A^3 T^{11} \ll a^5$, so $T^{17} \ll \Delta^3 a^5$.

- Otherwise Proposition 8.13 applies. By (26), its first term gives $T \ll a^{O(\eta)} a^{15}/T^{50}$, so $T^{51} \ll a^{15+O(\eta)}$. Its second term gives $T \ll a^{O(\eta)} a^{12+1/2}/T^{40+3/2}$, so $T^{85} \ll a^{25+O(\eta)}$.

- For the graph term, $T \ll a^{O(\eta)} X^5 \mathcal{B}^{1/\delta_{\min}}$ gives $T^{51\delta_{\min}} \ll a^{O(\eta)} a^{15\delta_{\min}} \mathcal{B}$. Since $H \ge 1$, $\max(H, S^m) \le \max(H,S)^m \ll a^{O(\eta)} (a/T^3)^m$, so if $\mathcal{B} = \max(H, S^m)$ then $T^{51\delta_{\min}+3m} \ll a^{15\delta_{\min}+m+O(\eta)}$. If $\mathcal{B} = Y^n$ then $T^{51\delta_{\min}+20n} \ll a^{15\delta_{\min}+6n+O(\eta)}$.

The first three exponents all equal $5/17$, both fractions in $E(m,n)$ exceed $5/17$, and $T^4 \ll a$ is stronger still. Choosing η small in terms of ε, m and n, every window has $T \ll a^{E(m,n)+\varepsilon/2}$, and summing over the $O(\log a)$ windows proves the bound. For $m = n = 1$ the two fractions are $256/870 = 128/435$ and $261/887$, and the first is larger. For $(m,n) = (50,51)$ the larger fraction is $2201/7480$. □

Proof of Theorem 1.2. Set $r = n/m$. The two fractions in $E(m,n)$ become

$$E_S(r) = \frac{150r + 106}{510r + 360}, \qquad E_Y(r) = \frac{156r + 105}{530r + 357}.$$

The first decreases and the second increases on $r > 0$. Their crossing satisfies $30r^2 - 10r - 21 = 0$, so $r_* = (5 + \sqrt{655})/30$ and their common value is $\alpha_* = (1310 - \sqrt{655})/4365$. Positive rational ratios approximate r_*; choose the fixed positive integers m, n to make $E(m, n) < \alpha_* + \varepsilon/2$, and apply Corollary 8.14 with loss $\varepsilon/2$. The powers and all constants depend only on ε, not on a, b. □

Remark 8.15. In terms of the scales, at the critical point $T \approx a^{\alpha_*}$ one has $A \approx a^{2\alpha_*}$, $H \approx S \approx a^{1-3\alpha_*} \approx a^{0.117}$, $X \approx a^{3-10\alpha_*} \approx a^{0.057}$ and $Y \approx a^{6-20\alpha_*} \approx a^{0.115}$. The pattern count $X^5 \approx a^{0.287}$ alone would give the exponent $5/17$. The graph term multiplies it by $\mathcal{B}^{1/\delta_{\min}}$, and at $n/m \approx r_*$ the two box radii S^m and Y^n balance, at $\mathcal{B} \approx a^{0.117m}$. This gives the extra factor $a^{0.117/(10r_*+7)} \approx a^{0.0068}$.

9 Exterior lattice coordinates

We now prove Theorem 1.3. We keep the notation and the conventions on constants of Section 8, and write $\delta_{\min} = 10n + 7m$ as there. Only the count of good starts on retained nonexceptional components of degree at least four changes. Proposition 9.11 replaces the term $X^5\mathcal{B}^{1/\delta_{\min}}$ of Proposition 8.13 by

$$X^{5-1/(3\delta_{\min})}H^{1/\delta_{\min}} + X^{5-2m/(3\delta_{\min})}S'^{\,m/\delta_{\min}} + X^5Y^{n/\delta_{\min}},$$

where S' differs from S only in its constant (Lemma 9.8). In effect H is replaced by $H/X^{1/3}$ and S by $S/X^{2/3}$, while Y is unchanged. The node term stays below the critical exponent, so of the two fractions balanced in the proof of Theorem 1.2 only the curvature fraction improves, and the crossing moves from $\alpha_* = 0.2942513\ldots$ to $\gamma_* = 0.2942294\ldots$. Every pattern has $\lambda(p) \ll X^{1/3}$ (Lemma 9.2), and the weighted count of Lemma 9.6 shows that this saving is available on average: the patterns with an unusually short vector are few enough to pay for their larger boxes.

Section 9.1 attaches to each pattern p a rank-three lattice $\Lambda_p \subseteq \mathbb{Z}^3$ that contains the restrictions of the rows of $\widehat{\mathcal{M}}_i$ to three pivot columns, and gives it a reduced basis. Section 9.2 shows that few patterns have a short vector in Λ_p. Section 9.3 writes the nodes and the curvature coordinates in the reduced basis. Section 9.4 checks that this linear change of coordinates keeps the degree bounds of Lemma 8.12, and Section 9.5 sums over patterns and optimizes the exponent.

9.1 The lattice of a pattern

Let $p \neq 0$ be the minor vector of a 3×5 integer matrix $\mathcal{M}$, with pivot π as in the proof of Proposition 7.9; so $|p_\pi|$ is maximal among the coordinates of p. Put $\kappa = |p_\pi| \geq 1$, let $j_1 < j_2 < j_3$ be the pivot columns and $j_4 < j_5$ the other two, and call $x_\pi = (x_{j_1}, x_{j_2}, x_{j_3})$ the *pivot part* of a row x of $\mathcal{M}$. Let $\mathbf{u}, \mathbf{v} \in \mathbb{Z}^3$ be as in that proof: $\mathbf{u}_r$ ($\mathbf{v}_r$) is the minor with the r-th pivot column replaced by column j_4 (j_5), with the sign of the column permutation. So the coordinates of $\mathbf{u}$ and $\mathbf{v}$ are, up to sign, coordinates of p, and $p_\pi x_{j_4} = \mathbf{u} \cdot x_\pi$ and $p_\pi x_{j_5} = \mathbf{v} \cdot x_\pi$ for every row x (Lemma 9.1). Let $\Lambda_p \subseteq \mathbb{Z}^3$ be the subgroup of x with $\kappa \mid \mathbf{u} \cdot x$ and $\kappa \mid \mathbf{v} \cdot x$. It contains $\kappa\mathbb{Z}^3$. Write $|x|$ for the Euclidean norm and $|x|_\infty$ for the sup norm, and let the *minimum* $\lambda(p)$ be the least sup norm of a nonzero vector of Λ_p. Then $|x| \geq \lambda(p)$ for every nonzero $x \in \Lambda_p$.

Lemma 9.1 (Lifting)**.** *For a column l, let $L_l \in \mathbb{Z}^3$ be $p_\pi \mathbf{e}_r$ if $l = j_r$ is a pivot column, $\mathbf{u}$ if $l = j_4$ and $\mathbf{v}$ if $l = j_5$, where $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ is the standard basis. For every row x of $\mathcal{M}$ and every column l, $p_\pi x_l = L_l \cdot x_\pi$. Moreover $\det(L_0, L_1, L_2) = p_\pi^2 p_{012}$, and $|L_l|_\infty \leq \kappa$.*

Proof. The vectors L_l are the rows of $\mathcal{M}^{\mathsf{T}} \operatorname{adj}(\mathcal{M}_\pi^{\mathsf{T}})$, where $\mathcal{M}_\pi$ is the pivot 3×3 block, and the first identity is Cramer's rule for the system $\mathcal{M}_\pi c = (\text{column } l)$, whose solution is $c = L_l/p_\pi$. Taking

the rows $l = 0, 1, 2$ gives $\det(L_0, L_1, L_2) = p_{012} \det \operatorname{adj}(\mathcal{M}_\pi^{\mathsf{T}}) = p_{012} p_\pi^2$. The bound holds because the coordinates of $\mathbf{u}$ and $\mathbf{v}$ are, up to sign, coordinates of p, and $|p_\pi|$ is maximal. □

So the pivot parts x_π of the rows of $\mathcal{M}$ lie in Λ_p. They are the rows of $\mathcal{M}_\pi$, which span a sublattice of index $|\det \mathcal{M}_\pi| = \kappa$, so $[\mathbb{Z}^3 : \Lambda_p] \le \kappa$.

Lemma 9.2 (Index). *If K is a nonnegative integer and $(K+1)^3 > \kappa$, then $\lambda(p) \le K$.*

Proof. The box $[0, K]^3$ has more than $\kappa \ge [\mathbb{Z}^3 : \Lambda_p]$ points, so two of them lie in the same coset of Λ_p, and their difference is a nonzero vector of Λ_p with sup norm at most K. □

The next lemma is the classical reduction of a three-dimensional lattice: any basis with $|\mathbf{f}_1||\mathbf{f}_2||\mathbf{f}_3| \ll V$ would do, and the constant does not affect the exponents. For completeness we give the elementary proof in Appendix C.

Lemma 9.3 (Reduction). *Let $\Lambda \subseteq \mathbb{Z}^3$ be a subgroup containing $\kappa\mathbb{Z}^3$, $\kappa \ge 1$. There is a basis $\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3$ of Λ with $V = \det(\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3)$, $0 < V \le \kappa^3$ and*

$$27\,|\mathbf{f}_1|^2 |\mathbf{f}_2|^2 |\mathbf{f}_3|^2 \le 64\, V^2.$$

Hence every $x \in \Lambda$ is $\sum \xi_i \mathbf{f}_i$ with $\xi \in \mathbb{Z}^3$ and $27|\mathbf{f}_i|^2 \xi_i^2 \le 64|x|^2$.

9.2 Few patterns have a short vector

Lemma 9.4 (Divisibility count). *For $\ell \ge 1$ and $x \in \mathbb{Z}^3$, at most $27\ell^2 \gcd(\ell, x)$ vectors $y \in [-\ell, \ell]^3$ have $\ell \mid y \cdot x$. Here $\gcd(\ell, x) = \gcd(\ell, x_1, x_2, x_3)$.*

Proof. Each residue class modulo ℓ meets $[-\ell, \ell]^3$ in at most 3^3 points, so it suffices to show that the set $\mathcal{K}$ of solutions in $[0, \ell)^3$ has at most $\ell^2 e$ elements, where $e = \gcd(\ell, x)$. By Bézout choose $z \in \mathbb{Z}^3$ with $z \cdot x \equiv e \pmod{\ell}$. The map $(y, j) \mapsto (y + jz) \bmod \ell$ is injective on $\mathcal{K} \times [0, \ell/e)$: equal images give $\ell \mid (j - j')e$, so $j = j'$, and then $y = y'$. Hence $|\mathcal{K}|\, \ell/e \le \ell^3$. □

Proposition 9.5 (Short vectors). *Let $\mathcal{S}$ be a set of nonzero minor vectors with coordinates in $[-X, X]$, and let $\tau(j) \le \tau_0$ for $1 \le j \le X$. Then for every $R \ge 1$,*

$$\#\{p \in \mathcal{S} : \lambda(p) \le R\} \ll R^3 X^4 \tau_0^2.$$

Proof. If $\lambda(p) \le R$, some nonzero $x \in [-R, R]^3$ lies in Λ_p, so $\kappa \mid \mathbf{u} \cdot x$, where $\kappa = |p_\pi| \le X$; we drop the condition $\kappa \mid \mathbf{v} \cdot x$. For fixed x, the encoding $p \mapsto (\pi, \operatorname{sgn} p_\pi, \kappa, \mathbf{u}, \mathbf{v})$ of the proof of Proposition 7.9 gives at most $20 \sum_{\kappa \le X} N(\kappa, x)$ such p, where $N(\kappa, x)$ counts pairs $(\mathbf{u}, \mathbf{v})$ in $[-\kappa, \kappa]^3$ with $\kappa \mid \mathbf{u} \times \mathbf{v}$ and $\kappa \mid \mathbf{u} \cdot x$. By the proof of Proposition 7.9, for each $\mathbf{u}$ there are at most $27\kappa \gcd(\kappa, \mathbf{u})^2$ vectors $\mathbf{v}$. Group the $\mathbf{u}$ by $g = \gcd(\kappa, \mathbf{u})$ and write $\mathbf{u} = g\mathbf{u}_0$ with $(\kappa/g) \mid \mathbf{u}_0 \cdot x$. Lemma 9.4 with $\ell = \kappa/g$, and $\gcd(\kappa/g, x) \le \gcd(\kappa, x)$, give

$$\sum_{\mathbf{u}} g^2 \le \sum_{g \mid \kappa} g^2 \cdot 27 \Big(\frac{\kappa}{g}\Big)^2 \gcd(\kappa, x) = 27\kappa^2 \tau(\kappa) \gcd(\kappa, x),$$

so $N(\kappa, x) \le 729 \kappa^3 \tau(\kappa) \gcd(\kappa, x)$. Next, $\gcd(\kappa, x) = \sum_{e \mid \gcd(\kappa, x)} \varphi(e) \le \sum_{e \mid \kappa,\, e \mid x} e$, and swapping the sums,

$$\sum_{0 < |x|_\infty \le R} \gcd(\kappa, x) \le \sum_{e \mid \kappa} e\, \#\{x \ne 0 : |x|_\infty \le R,\ e \mid x\} \le 27 R^3 \tau(\kappa).$$

Indeed, a divisor $e > R$ of κ divides no nonzero x in the box, and for $e \le R$ the multiples of e number at most $(2R/e + 1)^3$, with $e(2R/e + 1)^3 \le 27R^3$. Sum over x and then over $\kappa \le X$, using $\sum_{\kappa \le X} \kappa^3 \le X^4$. □

Lemma 9.6 (Fractional weights). *Let $\mathcal{S}$ be a set of nonzero minor vectors with coordinates in $[-X, X]$, $X \geq 1$, and let $\tau_0 \geq 1$ satisfy $\tau(j) \leq \tau_0$ for $1 \leq j \leq X$. Then for $0 \leq t \leq 1$,*

$$\sum_{p \in \mathcal{S}} \lambda(p)^{-t} \ll \tau_0^2 X^{5-t/3}.$$

Proof. Choose an integer K with $X \leq K^3 \leq 8X$. By Lemma 9.2, $1 \leq \lambda(p) \leq K$ for $p \in \mathcal{S}$. Writing $1/\lambda = 1/K + \sum_{\lambda \leq j < K} 1/(j(j+1))$ and using Proposition 9.5 with $R = j$,

$$K \sum_{p \in \mathcal{S}} \frac{1}{\lambda(p)} \leq \#\mathcal{S} + O\Big(K X^4 \tau_0^2 \sum_{1 \leq j < K} \frac{j^2}{j+1}\Big) \leq \#\mathcal{S} + O(X^4 \tau_0^2 K^3).$$

The proof of Proposition 7.9 gives $\#\mathcal{S} \ll \tau_0 X^5$. Since $K^3 \leq 8X$ and $\tau_0 \geq 1$, the right side is $O(\tau_0^2 X^5)$. Dividing by $K \geq X^{1/3}$ proves the case $t = 1$. For $0 < t < 1$, Hölder's inequality with exponents $1/(1-t)$ and $1/t$ gives $\sum \lambda^{-t} \leq (\#\mathcal{S})^{1-t} (\sum \lambda^{-1})^t$, which is the claim; the cases $t = 0$ and $t = 1$ are the two bounds above. □

9.3 Exterior coordinates

Lemma 9.7 (Exterior coordinates). *Let $\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3$ be a basis of $\Lambda \subseteq \mathbb{Z}^3$ as in Lemma 9.3, with $|\mathbf{f}_i| \geq \lambda > 0$. Let $x, x' \in \Lambda$ have coordinates $\xi, \xi' \in \mathbb{Z}^3$, and put $w = x \times x'$ and $c = \xi \times \xi' \in \mathbb{Z}^3$. Then*

$$w = c_1\, \mathbf{f}_2 \times \mathbf{f}_3 + c_2\, \mathbf{f}_3 \times \mathbf{f}_1 + c_3\, \mathbf{f}_1 \times \mathbf{f}_2, \qquad \mathbf{f}_i \cdot w = V c_i, \qquad 27 \lambda^4 c_i^2 \leq 64 |w|^2.$$

Proof. The first identity follows by expanding $(\sum \xi_i \mathbf{f}_i) \times (\sum \xi_i' \mathbf{f}_i)$ bilinearly, and taking the dot product with $\mathbf{f}_i$ gives the second. By Cauchy–Schwarz and Lemma 9.3,

$$27 |\mathbf{f}_1|^2 |\mathbf{f}_2|^2 |\mathbf{f}_3|^2 c_i^2 \leq 64 V^2 c_i^2 = 64 (\mathbf{f}_i \cdot w)^2 \leq 64 |\mathbf{f}_i|^2 |w|^2.$$

Cancelling $|\mathbf{f}_i|^2$ and using $|\mathbf{f}_j| \geq \lambda$ for the other two vectors gives the claim. □

The lemma replaces the bounds $|\xi| \ll |x|/\lambda$ and $|\xi'| \ll |x'|/\lambda$, which would give $|c| \ll |x||x'|/\lambda^2$, by $|c| \ll |x \times x'|/\lambda^2$. This matters because the drop row is large while the curvature minors are small.

The pivot columns of a pattern may include columns 3 and 4, so the cross product of the pivot parts involves minors D_{jl} with $l \geq 3$. We therefore bound all ten minors, not only the three of Lemma 8.7.

Lemma 9.8 (All curvature minors). *At a good start, all ten minors $D_{jl} = d_j k_l - d_l k_j$ with $0 \leq j < l \leq 4$ satisfy*

$$|D_{jl}| \leq S' := \left\lfloor \frac{c_{S'} A^3 H^3}{a^2} \right\rfloor,$$

for a suitable absolute constant $c_{S'}$.

Proof. Argue as in Lemma 8.7, using the expansions for $j \leq 5$, hence d_j for $j \leq 4$. The gaps $k_0, \dots, k_5$ are at most H, so $s_j \leq 5H$ and $\mu_{i+j}((s_j + x)^2) = s_j^2 + 2s_j + k_j + 1 \ll H^2$. This gives $d_j/a = k_j/q^2 + \delta_j$ with $|\delta_j| \ll H^2/q^3$ for $j \leq 4$, hence $|D_{jl}| q^3 \ll a H^3$, and $q \geq a/(4A)$ gives $|D_{jl}| \ll A^3 H^3 / a^2$. □

Now fix a pattern p that occurs at a good start, and let $\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3$ be a basis of Λ_p as in Lemma 9.3. Write $\lambda = \lambda(p)$. At a good start i with pattern p, the pivot parts d_π and k_π of the drop and gap rows of $\widehat{\mathcal{M}}_i$ lie in Λ_p, with coordinates ξ and ζ. Put

$$W' = \xi \times \zeta, \qquad w = d_\pi \times k_\pi.$$

The entries of w are, up to sign, the minors D_{jl} with j, l pivot columns. The next lemma shows that $W' = G_p W$ for a constant matrix G_p. We use W' rather than $G_p W$ because W' is visibly integral, while G_p has rational entries.

Lemma 9.9 (Coordinates in the reduced basis)**.** *There are matrices $B_p, G_p \in \mathrm{GL}_3(\mathbb{Q})$, depending only on p, such that at every good start with pattern p*

$$(s_1, s_2, s_3) = B_p \zeta, \qquad W' = G_p W, \qquad |\zeta|_\infty \le \frac{3H}{\lambda}, \qquad |W'|_\infty \le \frac{8S'}{3\lambda^2}.$$

Proof. By Lemma 9.1, $p_\pi(k_0, k_1, k_2) = \mathcal{L} k_\pi$, where $\mathcal{L}$ has rows L_0, L_1, L_2 and $\det \mathcal{L} = p_\pi^2 p_{012} \neq 0$, since $p_{012} = N_0 = I_4(i) \ge 1$ at a good start. With $k_\pi = (\mathbf{f}_1\, \mathbf{f}_2\, \mathbf{f}_3)\zeta$ and the cumulative sums $s_j = k_0 + \cdots + k_{j-1}$, this gives B_p. Since $|k_\pi|^2 \le 3H^2$ and $|\mathbf{f}_i| \ge \lambda$, Lemma 9.3 gives $|\zeta_i| \le (8/3)H/\lambda \le 3H/\lambda$.

For the curvature, $W = (D_{12}, -D_{02}, D_{01}) = (d_0, d_1, d_2) \times (k_0, k_1, k_2)$, and $p_\pi(d_0, d_1, d_2) = \mathcal{L} d_\pi$. Let $\mathrm{Cof}(\mathcal{L})$ have rows $L_1 \times L_2$, $L_2 \times L_0$, $L_0 \times L_1$; its determinant is $(\det \mathcal{L})^2 \neq 0$. The Binet–Cauchy identity $(L \cdot x)(L' \cdot y) - (L' \cdot x)(L \cdot y) = (L \times L') \cdot (x \times y)$, applied to pairs of rows of $\mathcal{L}$, gives $(\mathcal{L}x) \times (\mathcal{L}y) = \mathrm{Cof}(\mathcal{L})(x \times y)$, so $p_\pi^2 W = \mathrm{Cof}(\mathcal{L})\, w$. Lemma 9.7 gives $VW' = (\mathbf{f}_1\, \mathbf{f}_2\, \mathbf{f}_3)^{\mathsf{T}} w$. Together, $W' = p_\pi^2 V^{-1} (\mathbf{f}_1\, \mathbf{f}_2\, \mathbf{f}_3)^{\mathsf{T}} \mathrm{Cof}(\mathcal{L})^{-1} W$, which defines G_p. Finally $|w|^2 \le 3S'^2$ by Lemma 9.8, and Lemma 9.7 gives $3\lambda^2 |W_i'| \le 8S'$. □

9.4 The lattice graph

On a retained nonexceptional component C of degree $d \ge 4$ of a pattern p, regard $W' = G_p W$ as a vector of rational functions on C, and let $\Gamma'_{m,n}$ be the image of the graph of

$$(s_1, s_2, s_3, (W_1')^m, (W_2')^m, (W_3')^m, J^n)$$

under the linear map that replaces the nodes s by $\zeta = B_p^{-1} s$.

Lemma 9.10 (Degree of the lattice graph)**.** *The degree δ' of $\Gamma'_{m,n}$ satisfies $\delta_{\min} \le \delta' \le D_{\mathrm{gr}}(m+n)d$.*

Proof. An invertible linear change of the node coordinates is a linear automorphism of $\mathbb{A}^7$, so it preserves the affine Hilbert function, and we may keep the nodes s. The proof of Lemma 8.12, with Appendix B, uses the curvature coordinates only through the following facts.

1. At every point P over infinity, $\mathrm{ord}_P W_j \ge 3e_P/2 > 0$ for each j (Corollary 8.8).
2. The pole envelope $\mathcal{F}$ is supported over the affine part of C, and $\deg \mathcal{F} \ge 3d/2 + 1$ (Lemma 8.11).
3. Some rational linear combination of the seven graph coordinates has pole divisor $\frac{5n}{2} E_\infty + m\mathcal{F}$; its construction uses only this envelope.
4. The W_j have a common denominator Ω and numerators of degree at most D_W (Appendix B).

Each of these is unchanged when W is replaced by $W' = G_p W$ for a constant invertible matrix G_p. Each W'_j is a constant combination of the W_l, and each W_l is a constant combination of the W'_j, so for every point P

$$\min_j \operatorname{ord}_P W'_j = \min_l \operatorname{ord}_P W_l.$$

This gives (1) for W', and it shows that the pole envelope of W' is $\mathcal{F}$, which gives (2) and (3). The numerators of W' over Ω are constant combinations of those of W, which gives (4). The proof of Lemma 8.12 therefore applies unchanged. □

9.5 The window bound and the optimized exponent

Proposition 9.11 (Lattice window bound)**.** *Fix positive integers m, n and $\eta > 0$. For a window run in $[A, 2A)$ with $T \geq 36$ satisfying (19),*

$$T \ll_{m,n,\eta} a^{\eta} \left(X^5 + X^4 H^{1/2} + X^{5-1/(3\delta_{\min})} H^{1/\delta_{\min}} + X^{5-2m/(3\delta_{\min})} {S'}^{m/\delta_{\min}} + X^5 Y^{n/\delta_{\min}} \right).$$

Proof. The proof of Proposition 8.13 applies, except for the remaining components of the nonexceptional patterns. For such a component of a pattern p, the good starts give integer points $(\zeta, (W'_1)^m, (W'_2)^m, (W'_3)^m, J^n)$ of $\Gamma'_{m,n}$, and distinct node triples give distinct points because ζ determines s. By Lemma 9.9 they lie in the box of side

$$\mathcal{B}_p = \max\Big(1, \frac{3H}{\lambda(p)}, \Big(\frac{8S'}{3\lambda(p)^2}\Big)^m, Y^n\Big).$$

By Lemma 9.10 and (24), there are $O_{m,n,\eta}(a^{\eta} \mathcal{B}_p^{1/\delta_{\min}})$ of them, and

$$\mathcal{B}_p^{1/\delta_{\min}} \leq 1 + 3H^{1/\delta_{\min}} \lambda(p)^{-1/\delta_{\min}} + 8{S'}^{m/\delta_{\min}} \lambda(p)^{-2m/\delta_{\min}} + Y^{n/\delta_{\min}}.$$

Sum over the $O_{\eta}(a^{\eta} X^5)$ patterns. Since $\delta_{\min} \geq 2m$, both $1/\delta_{\min}$ and $2m/\delta_{\min}$ lie in $[0,1]$, and Lemma 9.6, with $\tau_0 = a^{o(1)}$, gives $\sum_p \lambda(p)^{-1/\delta_{\min}} \ll a^{\eta} X^{5-1/(3\delta_{\min})}$ and $\sum_p \lambda(p)^{-2m/\delta_{\min}} \ll a^{\eta} X^{5-2m/(3\delta_{\min})}$. □

Corollary 9.12. *For fixed positive integers m, n and every $\varepsilon > 0$,*

$$P(a,b) \ll_{m,n,\varepsilon} a^{E'(m,n)+\varepsilon},$$
$$E'(m,n) = \max\left(\frac{5}{17}, \frac{45\delta_{\min}}{153\delta_{\min} - 1}, \frac{45\delta_{\min} - 3m}{153\delta_{\min} - 11m}, \frac{15\delta_{\min} + 6n}{51\delta_{\min} + 20n} \right),$$

where $\delta_{\min} = 10n + 7m$. In particular $E'(3,2) = 153/520$.

Proof. Argue as in the proof of Corollary 8.14, with $S' \ll a/T^3$ in place of S in (26). The first two terms and the windows in which (19) fails give the exponent $5/17$. For the third term, $T^{3\delta_{\min}} \ll a^{O(\eta)} X^{15\delta_{\min}-1} H^3$, and (26) gives $T^{153\delta_{\min}-1} \ll a^{45\delta_{\min}+O(\eta)}$. For the fourth, $T^{3\delta_{\min}} \ll a^{O(\eta)} X^{15\delta_{\min}-2m} {S'}^{3m}$ gives $T^{153\delta_{\min}-11m} \ll a^{45\delta_{\min}-3m+O(\eta)}$. For the fifth, $T^{\delta_{\min}} \ll a^{O(\eta)} X^{5\delta_{\min}} Y^n$ gives $T^{51\delta_{\min}+20n} \ll a^{15\delta_{\min}+6n+O(\eta)}$. For $(m,n) = (3,2)$, $\delta_{\min} = 41$ and the last three fractions are $1845/6272$, $153/520$ and $627/2131$. □

Proof of Theorem 1.3. Set $r = n/m$. The last two fractions in $E'(m,n)$ become

$$E'_S(r) = \frac{450r + 312}{1530r + 1060}, \qquad E_Y(r) = \frac{156r + 105}{530r + 357},$$

where E_Y is the same function as in the proof of Theorem 1.2. The first decreases and the second increases on $r > 0$. Their crossing satisfies $15r^2 = 7$, so $r_* = \sqrt{105}/15$, and their common value is $\gamma_* = 471/1559 - 6\sqrt{105}/7795$. The fraction $45\delta_{\min}/(153\delta_{\min} - 1)$ decreases in $\delta_{\min}$, and it is below γ_* once $\delta_{\min} > \gamma_*/(153\gamma_* - 45) \approx 17.2$; also $5/17 < \gamma_*$. Choose a positive rational r close to r_* with $\max(E'_S(r), E_Y(r)) < \gamma_* + \varepsilon/2$, and write $r = n/m$ with $\delta_{\min} = 10n + 7m \geq 18$, scaling m and n by a common factor if necessary. Then $45\delta_{\min}/(153\delta_{\min} - 1) < \gamma_*$, so $E'(m, n) < \gamma_* + \varepsilon/2$, and Corollary 9.12 with loss $\varepsilon/2$ gives the theorem. □

10 Exact polar projection and the branch sieve

The curvature vector W is a multiple of an affine-linear vector $G\mathbf{k}+\ell$ in the gaps (Proposition 10.1). Away from the single common zero of $G\mathbf{k} + \ell$, every zero of J is therefore a pole of W, and this raises the graph degree provided that (i) little cancellation occurs at that base point and (ii) the component has the full degree 24. A local intersection calculation gives (i), and a sieve for branches at infinity gives (ii). We then count the additional exceptional patterns and obtain Theorem 1.1.

10.1 An exact identity at orbit states

For this section abbreviate the normalized parameters of (10) by

$$(\rho, \varphi, \chi) = (\bar{N}_1, \bar{u}_1, \bar{u}_2), \qquad (\rho', \varphi', \chi') = (\bar{N}_2, \bar{v}_1, \bar{v}_2).$$

Thus $\rho, \varphi > 0 > \chi$ and $\gamma = (\rho, -\rho - \varphi, \varphi - \chi, \chi - 1, 1)$; primes indicate the next state. We retain the notation N_0, Ψ, P_4, R_4 and D_{jl} from Section 8. Write $\mathbf{k} = (k_0, k_1, k_2)^{\mathsf{T}}$ and $\mathbf{d} = (d_0, d_1, d_2)^{\mathsf{T}}$, so that $W = (D_{12}, -D_{02}, D_{01})^{\mathsf{T}} = \mathbf{d} \times \mathbf{k}$.

The gap and drop rows of the pattern matrix satisfy the same linear relations (Lemma 7.1): the cofactor relations at i and $i + 1$ give

$$k_3 = \rho k_0 - \varphi k_1 - \chi k_2, \qquad k_4 = \rho' k_1 - \varphi' k_2 - \chi' k_3,$$

and the same formulas express d_3, d_4 in terms of $\mathbf{d}$. For $\mathbf{y} \in \mathbb{Q}^3$ extend $\mathbf{y}$ to $(y_0, \dots, y_4)$ by these formulas, and let $S_j(\mathbf{y}) = y_0 + \cdots + y_{j-1}$, so that $S_j(\mathbf{k}) = s_j$ and $S_j(\mathbf{d}) = b_0 - b_j$, with local indices. Define G and ℓ by

$$\sum_{j=0}^{4} \gamma_j S_j(\mathbf{y})^2 = \mathbf{y}^{\mathsf{T}} G \mathbf{y}, \qquad \sum_{j=0}^{4} \gamma_j y_j = \ell^{\mathsf{T}} \mathbf{y}.$$

Explicitly,

$$G = \begin{pmatrix} \rho(\rho+1) & -\rho(\varphi-1) & -\rho(\chi-1) \\ -\rho(\varphi-1) & \varphi(\varphi-1) & \varphi(\chi-1) \\ -\rho(\chi-1) & \varphi(\chi-1) & \chi(\chi-1) \end{pmatrix}, \qquad \ell = \begin{pmatrix} \rho(\chi - \chi') \\ \rho' - \rho - \varphi(\chi - \chi') \\ \varphi - \varphi' - \chi(\chi - \chi') \end{pmatrix},$$

and $\det G = \rho(\rho+\varphi)(\varphi-\chi)(\chi-1) \neq 0$. The pattern quadratic is $f = \sum_j \gamma_j(s_j^2 + k_j) = \mathbf{k}^{\mathsf{T}} G \mathbf{k} + \ell^{\mathsf{T}} \mathbf{k}$. Put

$$V = G\mathbf{k} + \ell, \qquad \text{so that} \qquad f = \mathbf{k} \cdot V.$$

The linear form $\mathbf{y} \mapsto \mathbf{y} \cdot V$ is the polarization of f at $\mathbf{k}$: it is $\mathbf{y}^{\mathsf{T}} G \mathbf{k} + \ell^{\mathsf{T}} \mathbf{y}$, and it recovers f at $\mathbf{y} = \mathbf{k}$. We call V the *polar vector*.

Proposition 10.1 (Polar identity)**.** *At every orbit state with initial index i satisfying $i+7<N$,*

$$\Psi W = N_0 V, \qquad JW = N_0^2 V.$$

In particular, $[W] = [G\mathbf{k} + \ell]$.

Proof. The vector V is orthogonal to $\mathbf{k}$, since $\mathbf{k}\cdot V = f = 0$. It is also orthogonal to $\mathbf{d}$; this is where the division equations enter. By polarization,

$$\mathbf{d}\cdot V = \sum_{j=0}^{4} \gamma_j s_j t_j + \sum_{j=0}^{4} \gamma_j d_j, \qquad t_j = b_0 - b_j.$$

The relations of Lemma 7.1 give $\sum_j \gamma_j = 0$ and $\sum_j \gamma_j S_j(\mathbf{y}) = \sum_l y_l(\gamma_{l+1}+\cdots+\gamma_4) = 0$ for $\mathbf{y} = \mathbf{k}, \mathbf{d}$ (Lemma 7.2), that is, $\sum_j \gamma_j s_j = \sum_j \gamma_j b_j = 0$. Multiply the division equations $(q_i + s_j)b_j + b_{j+1} = a$ by γ_j and sum over $j \le 4$. The terms in a and in q_i cancel, and writing $b_j = b_0 - t_j$ and $b_{j+1} = b_j - d_j$ leaves $-\sum_j \gamma_j s_j t_j - \sum_j \gamma_j d_j = 0$. Hence $\mathbf{d}\cdot V = 0$.

Since $W\cdot(r_0, r_1, r_2) = N_0 > 0$ by (20), $W \neq 0$, so $\mathbf{d}$ and $\mathbf{k}$ are independent and V is a multiple of $W = \mathbf{d}\times\mathbf{k}$. To find the multiple, expand: with $r_j = k_j(s_j + s_{j+1}) + k_{j+1}$,

$$V\cdot(r_0, r_1, r_2) - \Psi = -(s_4 + 1 + \chi - \chi')f,$$

so $V\cdot r = \Psi$ while $W\cdot r = N_0$. Thus $\Psi W = N_0 V$. Now use $J = N_0\Psi$ and $\Psi > 0$. □

In projective coordinates $[\mathbf{k} : k_h]$, the three homogeneous forms $G\mathbf{k} + \ell k_h$ have a unique common zero, the finite point $\mathbf{k}_* = -G^{-1}\ell$. It lies on $f = \Psi = 0$, and also on the fixed-dividend curve for every a. Indeed, its nodes are

$$s_0^* = 0, \qquad s_j^* = \frac{\gamma'_{j-1} - \gamma_{j-1}}{\gamma_j}\ (1 \le j \le 3), \qquad s_4^* = \chi' - \chi.$$

Dividing $Z(x) = xR_4(x)$ by $P_4(x)$ gives quotient $x + c_*$, where $c_* = \chi' - \chi - s_4$, and a remainder with Newton coefficients

$$-c_*\gamma_0, \qquad \gamma'_{j-1} - \gamma_{j-1} - \gamma_j(s_j + c_*) \quad (1 \le j \le 3).$$

They all vanish at $\mathbf{k}_*$, so $Z_* = xP_{4,*}$ and $N_0 \operatorname{Res}(P_{4,*}, Z_*) - a\Psi(\mathbf{k}_*)^4 = 0$.

We add the three denominator-cleared polar identities to the retained cuts of Section 8.2. They have bounded degree, so, as in Lemma 8.2, a cut that is not an identity on a component removes all but $O(1)$ of its points. Throughout this section, a *retained* component satisfies the earlier cuts and these three new ones; on it,

$$JW = N_0^2(G\mathbf{k} + \ell)$$

holds in the function field.

10.2 The base point and the graph degree

For an integral component C of a pattern curve, the *base divisor* of the forms $G\mathbf{k} + \ell$ is

$$\sum_P \Big(\min_j \operatorname{ord}_P(G\mathbf{k} + \ell)_j\Big) P,$$

summed over the affine points P of the normalization of C; its degree weights P by $[\mathbb{Q}(P) : \mathbb{Q}]$. Since $G\mathbf{k} + \ell = G(\mathbf{k} - \mathbf{k}_*)$, it is supported over $\mathbf{k}_*$, and since G is invertible the homogenized forms $G\mathbf{k} + \ell k_h$ have no common zero at infinity.

Lemma 10.2 (The base divisor)**.** *There is a fixed rational polynomial Θ in the ten minor coordinates, independent of a and not identically zero on the Plücker cone, with the following property. For every pattern with $\Theta \neq 0$ and every integral component C of the pattern curve, the base divisor of $G\mathbf{k} + \ell$ on C has degree at most four.*

Proof. Step 1: the quartic initial form. We work on the chart where $N_0 N_1 \neq 0$, so that the normalized parameters are defined, and

$$\gamma_1\gamma_2\gamma_3 \det G \, G_{00} \, \ell_0 \neq 0.$$

These conditions make the base point and its nodes well defined. Since $\nabla f(\mathbf{k}_*) = -\ell$ and $\ell_0 \neq 0$, the quadric is smooth at $\mathbf{k}_*$. Take $s = k_1 - k_{*,1}$ and $t = k_2 - k_{*,2}$ as local coordinates and solve $f = 0$ for k_0 by formal power series. Let $R_1(x; s, t)$ be the linear part of the Newton remainder just computed, and let $\Psi_1(s, t)$ be the linear part of Ψ on this tangent plane. The curve equation has order at least four and quartic initial form

$$N_0 N_4(s,t) - a\Psi_1(s,t)^4, \qquad N_4(s,t) = \operatorname{Res}_x(P_{4,*}, R_1). \tag{27}$$

To see that only first derivatives enter, compute the resultant as a 4×4 Bézout determinant. Each row vanishes at $\mathbf{k}_*$, so its term of degree four uses exactly the linear part of every row. Higher terms in the formal solution of $f = 0$ cannot change (27).

Step 2: the polynomial Θ. Consider the rational function of the pattern parameters

$$M = N_4(1,0)\Psi_1(0,1)^4 - N_4(0,1)\Psi_1(1,0)^4.$$

If $M \neq 0$ and $N_0 \neq 0$, the two axis coefficients in (27) cannot both vanish, for any a. Now write M, the chart conditions and $N_0 N_1$ as rational functions of the ten minor coordinates, by (10), each in lowest terms, and let Θ be the product of all their numerators and denominators. Then $\Theta \neq 0$ implies that each of these functions is defined and nonzero. To see that Θ is not identically zero on the Plücker cone, take the maximal minors of

$$\begin{pmatrix} 1 & 0 & 0 & 2 & 10 \\ 0 & 1 & 0 & -5 & -22 \\ 0 & 0 & 1 & 3 & 9 \end{pmatrix},$$

which give $(\rho, \varphi, \chi, \rho', \varphi', \chi') = (2, 5, -3, 3, 6, -5)$ and $\mathbf{k}_* = (-1/7, -3/28, -1/2)$. There the chart conditions hold, and Appendix D computes the tangent frame, the quartic and the first derivatives, giving

$$M = \frac{26291112152499}{60236288} \neq 0.$$

Step 3: from the initial form to the base divisor. Let $\mathcal{O}$ be the local ring of the smooth quadric at $\mathbf{k}_*$ and let F be the curve equation. Choose $g = s$ or $g = t$ so that the complementary axis in (27) has nonzero coefficient. The completion of $\mathcal{O}/(g)$ is a power-series ring in one variable, in which F has order exactly four. Consequently $\dim_{\mathbb{Q}} \mathcal{O}/(F, g) = 4$. If C passes through $\mathbf{k}_*$, its local ring $\mathcal{O}_C = \mathcal{O}/I_C$ is a quotient with $F \in I_C$, so

$$\dim_{\mathbb{Q}} \mathcal{O}_C/g\mathcal{O}_C \leq 4.$$

In particular, g is nonzero on this one-dimensional integral local ring.

Let $\overline{\mathcal{O}}_C$ be its finite normalization. The quotient $\overline{\mathcal{O}}_C/\mathcal{O}_C$ is finite-dimensional over $\mathbb{Q}$. Apply multiplication by g to $0 \to \mathcal{O}_C \to \overline{\mathcal{O}}_C \to \overline{\mathcal{O}}_C/\mathcal{O}_C \to 0$. Multiplication is injective on the first two

rings, and its kernel and cokernel on the finite-dimensional quotient have equal dimensions. The resulting exact sequence therefore gives

$$\dim_{\mathbb{Q}} \overline{\mathcal{O}}_C/g\overline{\mathcal{O}}_C = \dim_{\mathbb{Q}} \mathcal{O}_C/g\mathcal{O}_C = \sum_{P|\mathbf{k}_*} [\mathbb{Q}(P):\mathbb{Q}] \operatorname{ord}_P g.$$

The final equality follows by factoring the ideal (g) in the semilocal Dedekind ring $\overline{\mathcal{O}}_C$. Since G is invertible, the coordinates of $G\mathbf{k}+\ell = G(\mathbf{k}-\mathbf{k}_*)$ generate the centered maximal ideal. Their minimum order at each P is at most $\operatorname{ord}_P g$. The weighted sum of these minimum orders, which is the degree of the base divisor, is therefore at most four. A component missing $\mathbf{k}_*$ has zero base divisor. □

Suppose now that a retained nonexceptional component has degree 24 and $\Theta \neq 0$. At each point at infinity, $\operatorname{ord}_P J = -5e_P/2$, by Corollary 8.4, and $\sum e_P[\mathbb{Q}(P):\mathbb{Q}] = 24$. Thus J has pole divisor, and hence zero divisor, of degree 60; all its zeros are affine. At an affine point P the forms $G\mathbf{k}+\ell$ are regular, and $W = N_0^2(G\mathbf{k}+\ell)/J$ gives

$$\max_j(-\operatorname{ord}_P W_j) = \operatorname{ord}_P J - \min_j \operatorname{ord}_P (G\mathbf{k}+\ell)_j,$$

where the subtracted term is the multiplicity of P in the base divisor. Summing over affine points, Lemma 10.2 gives an affine curvature pole envelope of degree at least $60-4=56$, in place of the bound $3\cdot 24/2+1=37$ of Lemma 8.11. These poles are disjoint from the poles of J at infinity, so the pole formula of Lemma 8.12 gives

$$\deg(\text{powered joint graph}) \geq \frac{5n}{2}\cdot 24 + 56m = \delta_{\text{pol}} := 60n+56m. \tag{28}$$

An invertible constant change of curvature coordinates preserves the minimum valuation, so this also holds for the lattice coordinates of Section 9. It remains to show that components of degree less than 24 occur for relatively few patterns.

10.3 An explicit label for a branch at infinity

The plan is as follows. At a branch at infinity of a nonexceptional component, the residue field contains the ratio t of two leading nodes, which is a root of a fixed sextic, and a square root β coming from the fixed-dividend equation. If the minimal polynomial of $t+c\beta$ has degree twelve for some slope c, every branch at infinity has residue degree at least twelve; since its multiplicity e is even, the component then has the maximal degree 24. The *label* below is a polynomial whose irreducibility certifies this, and which can be sieved over patterns.

Let P be a point above infinity on the normalization of a nonexceptional component. The five leading nodes are distinct by Lemma 8.3. Scale them to $(0,1,t,y,z)$, and put

$$z = -\gamma_1 - \gamma_2 t - \gamma_3 y.$$

The second and third moment equations become

$$Q_2(t,y) = \gamma_1 + \gamma_2 t^2 + \gamma_3 y^2 + z^2 = 0, \qquad Q_3(t,y) = \gamma_1 + \gamma_2 t^3 + \gamma_3 y^3 + z^3 = 0.$$

Define the universal sextic $E_\gamma(t)$ by the polynomial identity

$$\operatorname{Res}_y(Q_2,Q_3) = \gamma_3^2(\gamma_3+1)E_\gamma(t). \tag{29}$$

The quotient on the right is a polynomial over $\mathbb{Z}[\gamma_1,\gamma_2,\gamma_3]$, of degree six in t. At an orbit pattern $\gamma_3=\chi-1<-1$, so $\gamma_3^2(\gamma_3+1)\neq 0$. Cancelling this factor in the resultant and linear subresultant identities gives $E_\gamma(t)=0$ and $d(t)y+n(t)=0$, where

$$\begin{aligned} d(t) &= (\gamma_3-1)\big(\gamma_2(\gamma_2+\gamma_3+1)t^2+2\gamma_1\gamma_2 t+\gamma_1(\gamma_1+\gamma_3+1)\big),\\ n(t) &= \gamma_2(\gamma_2+1)(\gamma_2+\gamma_3+1)t^3+\gamma_1\gamma_2(3\gamma_2+\gamma_3+2)t^2\\ &\quad+\gamma_1\gamma_2(3\gamma_1+\gamma_3+2)t+\gamma_1(\gamma_1+1)(\gamma_1+\gamma_3+1). \end{aligned}$$

These are polynomial identities, so the cleared relation remains valid when a specialized denominator vanishes.

Write

$$V_4=ty(t-1)(y-1)(y-t),\qquad V_5=z(z-1)(z-t)(z-y).$$

By (16), the degree-ten leading resultant at these nodes is $R_0=\gamma_0\gamma_1\gamma_2\gamma_3V_4V_5$. The four moment equations give the barycentric identity

$$R_0V_4=V_5^4. \tag{30}$$

For example, the identities $\gamma_j\prod_{l\neq j}(z_j-z_l)=V_5$ for $0\le j<4$ follow by evaluating the first four moment relations; their product is (30). Let $y_\infty=1/s_1$. The value $\eta_P=(y_\infty^5\Psi^2)(P)\in\mathbb{Q}(P)$ is defined and nonzero, because $\operatorname{ord}_P\Psi=-5e_P/2$. Taking leading terms in the curve equation gives $N_0R_0=a\eta_P^2$. Thus the residue field $\mathbb{Q}(P)$ contains

$$\beta=V_5^2/\eta_P,\qquad N_0\beta^2=aV_4. \tag{31}$$

So $\mathbb{Q}(P)$ contains $\mathbb{Q}(t,\beta)$, which generically has degree twelve, not just six.

For a slope c define

$$D_\gamma(t)=-t(t-1)\,n(t)\big(n(t)+d(t)\big)\big(n(t)+td(t)\big)$$

and the label polynomial

$$L_{a,N_0,c,\gamma}(Z)=\operatorname{Res}_t\big(E_\gamma(t),N_0d(t)^3(Z-t)^2-c^2aD_\gamma(t)\big). \tag{32}$$

Here the resultant means the Sylvester determinant with degree bounds 6 and 11, padding missing coefficients with zero. This convention makes (32) a polynomial family under every specialization. The relation $d(t)y+n(t)=0$ gives $D_\gamma(t)=d(t)^3V_4$, so (31) implies

$$L_{a,N_0,c,\gamma}(t+c\beta)=0$$

in every such residue field, including when $d(t)=0$.

Lemma 10.3 (The degree-twelve label)**.** *Fix $a\in\mathbb{Q}^\times$, and regard $N_0,\gamma_1,\gamma_2,\gamma_3$ as independent parameters. For some integer c with $0\le c\le 144$, $L_{a,N_0,c,\gamma}$ is irreducible of degree twelve over $\mathbb{Q}(N_0,\gamma_1,\gamma_2,\gamma_3)$. For a specialized nonexceptional pattern, if one such label polynomial is irreducible of degree twelve over $\mathbb{Q}$, every integral component of its pattern curve has degree twenty-four.*

Proof. At $(\gamma_1,\gamma_2,\gamma_3)=(-7,8,-4)$, the sextic in (29) equals $24E_\star(t)$, where

$$E_\star(t)=2400t^6-20160t^5+69776t^4-126448t^3+125636t^2-64680t+13475.$$

Appendix D proves $E_\star$ irreducible modulo 71. This specialization preserves degree six. Gauss's lemma therefore shows that E_γ is irreducible over $K=\mathbb{Q}(\gamma_1,\gamma_2,\gamma_3)$: a factorization into two

factors of positive t-degree over the parameter ring would specialize to two factors of positive degree. Choose a root t of E_γ, so $[K(t):K]=6$. In this field $d(t)\neq 0$ since its degree is less than six, and $y=-n(t)/d(t)$. The same degree comparison shows $V_4\neq 0$.

Adjoin the independent parameter N_0. The element aV_4/N_0 has odd valuation at $N_0=0$, so it is not a square in $K(t)(N_0)$. Thus (t,β) generates an extension of degree twelve over $K(N_0)$. Among its twelve embeddings, equality of two values $t+c\beta$ excludes at most one slope per pair, so at most $12^2=144$ slopes are excluded and one of $0,\dots,144$ gives a primitive element. (Counting only unordered pairs of distinct embeddings shows that one of $0,\dots,66$ already suffices; we state the cruder range because it is the one formalized.) Its minimal polynomial divides (32), whose degree is at most twelve; hence that label polynomial is irreducible of degree twelve. This argument holds for each fixed nonzero a; the chosen slope may depend on a.

For an actual pattern, the residue field at any branch at infinity contains the root $t+c\beta$. Irreducibility of the specialized degree-twelve label forces $[\mathbb{Q}(P):\mathbb{Q}]\geq 12$. Corollary 8.4 gives $e_P\geq 2$, so

$$\deg C=\sum_{P\text{ at infinity}} e_P[\mathbb{Q}(P):\mathbb{Q}]\geq 24.$$

Lemma 8.1 gives the reverse inequality. □

For each a, fix a slope $c(a)\in\{0,\dots,144\}$ given by Lemma 10.3. Call a pattern *branch-good* if $L_{a,N_0,c(a),\gamma}$ is irreducible of degree twelve over $\mathbb{Q}$, and *branch-bad* otherwise.

10.4 The arithmetic sieve

We prove a monic variant of the effective Hilbert irreducibility theorem of [4, Theorem 1.10] along the same lines, with explicit logarithmic powers.

Theorem 10.4 (Monic specialization estimate)**.** *For fixed integers $r,d\geq 1$ there is a constant $C_{r,d}$ with the following property. Let $F\in\mathbb{Z}[T_1,\dots,T_r][Y]$ be monic in Y and irreducible over $\mathbb{Q}(T_1,\dots,T_r)$. Suppose $\deg_Y F\leq d$, each coefficient of F has total parameter degree at most d, and every integer coefficient has absolute value at most $H\geq 1$. For every integer $B\geq 1$,*

$$\#\{t\in[-B,B]^r\cap\mathbb{Z}^r : F(t,Y)\text{ is reducible over }\mathbb{Q}\}\leq C_{r,d}(1+\log(H+1))^4B^{r-1/2}(1+\log B)^{4r+1}.$$

Proof. Write $\lg(x)=1+\log(x+1)$. Implied constants depend only on r,d.

Step 1: integer roots in one parameter. Let $\mathcal{A}\in\mathbb{Z}[T][Z]$ be monic in Z, of Z-degree $e\geq 2$, irreducible over $\mathbb{Q}(T)$, of total degree at most D and height at most H_0. We claim that

$$\#\{t\in[-B,B]\cap\mathbb{Z}:\ \mathcal{A}(t,Z)\text{ has an integer root}\}\ll_D \lg(H_0)^4B^{1/e}(1+\log B).$$

An integer root z of $\mathcal{A}(t,Z)$ satisfies $|z|\leq 1+(D+1)H_0B^D$. Choose an integer $k>D$ with $k\asymp_D\lg(H_0)$, and apply the shear $(T,Z)\mapsto(T,Z+T^k)$. The polynomial $\mathcal{A}_k(T,Z)=\mathcal{A}(T,Z-T^k)$ is irreducible in $\mathbb{Q}[T,Z]$, and its total degree is exactly ke: the term $(Z-T^k)^e$ contributes $\pm T^{ke}$, while every other term has degree at most $D+k(e-1)<ke$. The points $(t,z+t^k)$ are integer points of the plane curve $\mathcal{A}_k=0$ in a box of size $B'\ll_D H_0B^k$. By (24) with $\delta=ke$ there are $\ll(ke)^3B'^{1/(ke)}(\log B'+ke)$ of them. Here $B'^{1/(ke)}\ll_D H_0^{1/(ke)}B^{1/e}\ll B^{1/e}$ by the choice of k, while $ke\ll\lg(H_0)$ and $\log B'\ll\lg(H_0)(1+\log B)$. This proves the claim.

Step 2: one parameter. Let $r=1$, and let $y_1,\dots,y_e$ be the roots of F in an algebraic closure of $\mathbb{Q}(T)$; they are integral over $\mathbb{Z}[T]$. For a nonempty proper subset $I\subset\{1,\dots,e\}$ and $1\leq j\leq|I|$, let $\sigma_{I,j}$ be the jth elementary symmetric function of $(y_i)_{i\in I}$. Since F is irreducible, $\prod_{i\in I}(Y-y_i)\notin$

$\mathbb{Q}(T)[Y]$, so some $\sigma_I := \sigma_{I,j(I)}$ lies outside $\mathbb{Q}(T)$. The resolvent $\prod_{|I'|=|I|}(Z - \sigma_{I',j(I)})$ is monic in Z, and its coefficients are fixed integer polynomials in the coefficients of F; so it lies in $\mathbb{Z}[T][Z]$ with degree $O(1)$ and height $O(H^{O(1)})$. Let $\mathcal{A}_I$ be its monic irreducible factor over $\mathbb{Q}(T)$ that vanishes at σ_I. By Gauss's lemma $\mathcal{A}_I \in \mathbb{Z}[T][Z]$; by Gelfond's factor-height inequality its height is $O(H^{O(1)})$; and its Z-degree is at least two, because $\sigma_I \notin \mathbb{Q}(T)$.

Now suppose that $F(t, Y)$ is reducible over $\mathbb{Q}$. By Gauss's lemma it has a monic factor $g \in \mathbb{Z}[Y]$ of degree e' with $1 \le e' < e$. Extend the specialization $T \mapsto t$ to a ring homomorphism ϕ from $\mathbb{Z}[T][y_1, \dots, y_e]$ to $\overline{\mathbb{Q}}$, which is possible because this ring is integral over $\mathbb{Z}[T]$. Then $F(t, Y) = \prod_i (Y - \phi(y_i))$, so the roots of g, with multiplicity, are $\phi(y_i)$ for i in some I with $|I| = e'$, and the coefficients of g are $\pm\phi(\sigma_{I,j})$. In particular $\phi(\sigma_I)$ is an integer root of $\mathcal{A}_I(t, Z)$. Step 1, applied to the $O(1)$ polynomials $\mathcal{A}_I$, bounds the number of such t by $O(\lg(H)^4 B^{1/2}(1 + \log B))$.

Step 3: one parameter, the others free. Let $r \ge 2$. We claim that for all but $O(\lg(H)^4 B^{1/2}(1 + \log B))$ integers $t_1 \in [-B, B]$, the polynomial $F(t_1, T_2, \dots, T_r, Y)$ is irreducible over $\mathbb{Q}(T_2, \dots, T_r)$. Fix an integer $b > d$ and let $\mathcal{K}$ be the Kronecker map

$$T_2^{a_2} \cdots T_r^{a_r} Y^{a_{r+1}} \longmapsto \omega^{a_2 + a_3 b + \cdots + a_{r+1} b^{r-1}}.$$

It is a linear bijection from polynomials with all partial degrees less than b onto polynomials in ω of degree less than b^r, it commutes with specializing T_1, and it is multiplicative on products whose partial degrees stay below b. Since Y occupies the highest position, $\mathcal{K}(F) \in \mathbb{Z}[T_1][\omega]$ is monic. Factor $\mathcal{K}(F) = \prod_\nu \mathcal{A}_\nu$ into monic irreducible factors over $\mathbb{Q}(T_1)$. Again $\mathcal{A}_\nu$ lies in $\mathbb{Z}[T_1][\omega]$, with bounded degree and height $O(H^{O(1)})$. By Step 2, outside an exceptional set of the stated size, every $\mathcal{A}_\nu(t_1, \omega)$ is irreducible over $\mathbb{Q}$.

Let t_1 be outside this set, and suppose $F(t_1, \cdot) = gh$ with $g, h \in \mathbb{Q}[T_2, \dots, T_r][Y]$ monic in Y of positive Y-degree. All partial degrees of g and h are at most those of F, hence less than b, so $\mathcal{K}(g)\mathcal{K}(h) = \prod_\nu \mathcal{A}_\nu(t_1, \omega)$, and by unique factorization $\mathcal{K}(g) = \prod_{\nu \in S} \mathcal{A}_\nu(t_1, \omega)$ for some set S of indices. Put $g_S = \mathcal{K}^{-1}(\prod_{\nu \in S} \mathcal{A}_\nu)$ and define g_{S^c} likewise, so that $g_S(t_1) = g$ and $g_{S^c}(t_1) = h$. If $g_S g_{S^c} = F$ identically, this is a factorization of F over $\mathbb{Q}(T_1, \dots, T_r)$ into factors of positive Y-degree, which is impossible. Otherwise some coefficient of $g_S g_{S^c} - F$ is a nonzero polynomial in T_1 of bounded degree vanishing at t_1. There are $O(1)$ sets S, so this adds $O(1)$ exceptional values.

Step 4: induction on r. The case $r = 1$ is Step 2. For $r \ge 2$, the exceptional values t_1 of Step 3 contribute at most their number times $(2B + 1)^{r-1}$. For every other $t_1 \in [-B, B]$, the family $F(t_1, T_2, \dots, T_r, Y)$ is monic, irreducible over $\mathbb{Q}(T_2, \dots, T_r)$, of the same degree bounds, and of height at most $(d+1)HB^d$. Since $\lg((d+1)HB^d) \ll \lg(H)(1 + \log B)$, the induction hypothesis bounds its reducible specializations by $O(\lg(H)^4 B^{r-3/2}(1 + \log B)^{4r+1})$. Summing over the $2B + 1$ values of t_1 gives the theorem. □

To apply the theorem with a fixed, use the four integer minors of the first four columns,

$$\mathbf{v} = (v_0, v_1, v_2, v_3) = (p_{012}, p_{013}, p_{023}, p_{123}).$$

For $v_0 = N_0 \neq 0$ their relation to the independent parameters above is

$$\gamma_1 = -\frac{v_3 + v_2}{v_0}, \qquad \gamma_2 = \frac{v_2 + v_1}{v_0}, \qquad \gamma_3 = -\frac{v_1 + v_0}{v_0}. \tag{33}$$

This change is birational. Substitute it in (32) and clear denominators by a fixed power of v_0. To make the resulting degree-twelve polynomial monic without changing its generic irreducibility, use the following polynomial operation. For $F(Y) = \sum_{i=0}^{e} b_i Y^i$ put

$$\mathrm{mon}_e(F)(Y) = Y^e + \sum_{i<e} b_i b_e^{e-1-i} Y^i.$$

When $b_e \neq 0$, this is $b_e^{e-1}F(Y/b_e)$. When $b_e = 0$ and $e \geq 2$, it is $Y^{e-1}(Y + b_{e-1})$, hence reducible. Thus irreducibility of a specialized monic normalization implies both preservation of degree and irreducibility of the original label.

For each fixed positive integer a, use the slope $c(a)$ and set $e = 12$. We obtain a generically irreducible monic integer family in $\mathbf{v}$, with uniformly bounded degree and coefficient height at most $C(a+1)^D$, for absolute constants C, D. The same is true after dilation $\mathbf{v} = g\mathbf{z}$, with height at most $C(a+1)^D g^D$. The slope may depend on a, but the bounds are uniform because there are only 145 choices. Counting vertices with $v_0 = 0$ as bad, which adds the $O(B^3)$ points of that slice, Theorem 10.4 gives

$$\#\{\mathbf{z} \in [-B, B]^4 \cap \mathbb{Z}^4 : g\mathbf{z} \text{ is branch-bad}\} \ll_\eta a^\eta B^{7/2} \tag{34}$$

whenever $1 \leq g, B \ll a^7$. This range contains all the dilations and boxes used for orbit patterns below.

Lemma 10.5 (Branch-bad full patterns)**.** *Among the selected orbit patterns of coordinate height at most X, the number that are branch-bad is $O_\eta(a^\eta X^{9/2})$.*

Proof. Partition the first-four-column vertices into dyadic bands with $\max_j |v_j| \asymp M$, and choose a maximal coordinate as signed pivot κ. The other three vertex coordinates form $\mathbf{u}$; put $g = \gcd(\kappa, \mathbf{u})$. Completing the fifth column amounts to choosing $\mathbf{w} \in [-X, X]^3$ with $\kappa \mid \mathbf{u} \times \mathbf{w}$, as in the proof of Proposition 7.9. Since κ is maximal only among the first four columns, $\mathbf{w}$ ranges over the full X-box rather than $[-\kappa, \kappa]^3$, and the congruence count is

$$O(X^3 g^2 / M^2).$$

Apply (34) after dilation by g. There are $O_\eta(a^\eta (M/g)^{7/2})$ bad vertices in this common-factor class; imposing the exact gcd only reduces this count. Their total contribution is at most

$$O_\eta\left(a^\eta X^3 M^{3/2} \sum_{g \geq 1} g^{-3/2}\right).$$

The series converges, and summing the bands $M \leq X$ proves the claim. The finitely many pivot choices only change the constant. □

10.5 Counting the geometric exceptional locus

The patterns on the hypersurface $\Theta = 0$ must be counted with a saving over the $X^{5+o(1)}$ patterns in total. The following estimate saves $X^{1/3}$.

Proposition 10.6 (Hypersurface pattern count)**.** *Let Θ be a rational polynomial in the ten minor coordinates, with $\Theta \neq 0$ at the minor vector of some rational 3×5 matrix $\mathcal{M}_0$. For every $\eta > 0$, the number of integral points of the Plücker cone of coordinate height at most $X \geq 1$ satisfying $\Theta = 0$ is*

$$O_{\Theta,\eta}(X^{14/3+\eta}).$$

The count includes nonprimitive vectors and the zero vector.

Proof. Choose a minor of maximal absolute value $\kappa > 0$ and record its sign. The six one-column replacements form $\mathbf{u}, \mathbf{w} \in [-\kappa, \kappa]^3$, and the other coordinates are signed entries of $(\mathbf{u} \times \mathbf{w})/\kappa$. There are twenty signed pivot charts. In each chart, clearing denominators in Θ gives a nonzero

polynomial $\Theta_\pi(\kappa, \mathbf{u}, \mathbf{w})$. Indeed, the product of the pivot minor and Θ at the minor vector of $\mathcal{M}_0$ is a nonzero polynomial in the matrix entries, so it is nonzero at some rational matrix; every pivot chart meets the nonvanishing locus.

Put $g = \gcd(\kappa, \mathbf{u})$, $n = \kappa/g$ and $\mathbf{u}_0 = \mathbf{u}/g$. Since $\gcd(n, \mathbf{u}_0) = 1$, the congruence $\mathbf{u}_0 \times \mathbf{w} \equiv 0 \pmod{n}$ is equivalent to

$$\mathbf{w} \in \Lambda := \mathbb{Z}\mathbf{u}_0 + n\mathbb{Z}^3.$$

This lattice contains $n\mathbb{Z}^3$, so Lemma 9.3 gives a basis $\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3$ of Λ; write $\ell_i = |\mathbf{f}_i|$. We need two bounds on these lengths.

(a) $\ell_i \ll n$. The vectors $n\mathbf{e}_1, n\mathbf{e}_2, n\mathbf{e}_3$ lie in Λ and span a sublattice of full rank, so for each i one of them has a nonzero ith coordinate in the basis, and the coordinate bound of Lemma 9.3 gives $27\ell_i^2 \le 64n^2$.

(b) $\ell_i\ell_j \ge n\lambda^\perp(\mathbf{u}_0)$ for $i \ne j$, where $\lambda^\perp(\mathbf{u}_0)$ is the least sup norm of a nonzero vector of the normal lattice $\Lambda_n^\perp = \{z \in \mathbb{Z}^3 : n \mid z \cdot \mathbf{u}_0\}$. Indeed, if $x, y \in \Lambda$ are independent, then $x \times y = nz$ for a nonzero $z \in \Lambda_n^\perp$, so $|x||y| \ge |x \times y| \ge n\lambda^\perp(\mathbf{u}_0)$.

By Lemma 9.3, the vectors $\mathbf{w} \in \Lambda$ with $|\mathbf{w}|_\infty \le \kappa$ have basis coordinates in a box with sides $N_i = O(\kappa/\ell_i)$, and by (a) and $n \le \kappa$ these sides are $\gg 1$. A nonzero polynomial of bounded degree vanishes at $\ll N_1N_2N_3/\min_i N_i$ integer points of such a box (the polynomial grid-zero bound). Applied to a nonzero fiber $\Theta_\pi(\kappa, \mathbf{u}, \cdot)$ in the basis coordinates, this and (b) give

$$\#\{\mathbf{w} \in \Lambda \cap [-\kappa, \kappa]^3 : \Theta_\pi(\kappa, \mathbf{u}, \mathbf{w}) = 0\} \ll_\Theta \sum_{i<j} \frac{\kappa^2}{\ell_i\ell_j} \ll_\Theta \frac{\kappa^2}{n\lambda^\perp(\mathbf{u}_0)}. \tag{35}$$

Here is the reciprocal estimate that produces the exponent 14/3:

$$\sum_{\mathbf{u}_0 \in [-n,n]^3 \cap \mathbb{Z}^3} \frac{1}{\lambda^\perp(\mathbf{u}_0)} \ll \tau(n)n^{8/3}. \tag{36}$$

It holds for all $\mathbf{u}_0$, without a primitivity restriction. Pigeonholing $z \mapsto \mathbf{u}_0 \cdot z \pmod{n}$ gives $\lambda^\perp(\mathbf{u}_0) \ll n^{1/3}$. For $R \ge 1$, Lemma 9.4 and a divisor sum give

$$\begin{aligned}\#\{\mathbf{u}_0 : \lambda^\perp(\mathbf{u}_0) \le R\} &\le \sum_{0<|z|_\infty \le R} \#\{\mathbf{u}_0 \in [-n, n]^3 : n \mid \mathbf{u}_0 \cdot z\} \\ &\ll n^2 \sum_{0<|z|_\infty \le R} \gcd(n, z_1, z_2, z_3) \ll \tau(n)n^2R^3.\end{aligned}$$

For the last bound, expand the gcd as a divisor sum and note that a nonzero z of size at most R has no common divisor greater than R. Partial summation up to $K \asymp n^{1/3}$ now bounds the reciprocal sum by $O(n^3/K + \tau(n)n^2K^2)$, proving (36).

For fixed κ, g, equations (35) and (36) bound the nonzero fibers by $O_\Theta(\tau(n)\kappa^2n^{5/3})$. Sum over $g \mid \kappa$ and $\kappa \le X$ to obtain

$$O_\Theta(\tau_{\max}(X)^2X^{14/3}), \qquad \tau_{\max}(X) = \max_{1\le j\le X} \tau(j).$$

Finally, an identically zero fiber forces one fixed nonzero coefficient polynomial $\Theta_0(\kappa, \mathbf{u})$ to vanish. For fixed g, the polynomial $\Theta_0(g\kappa', g\mathbf{u}')$ is still nonzero. The four-variable grid-zero bound

supplies $O_\Theta((X/g)^3)$ possible scaled vertices. Each has $O(\kappa g^2) \le O(Xg^2)$ completions in its κ-box, so these fibers total

$$O_\Theta\left(X^4 \sum_{g \le X} \frac{1}{g}\right) = O_\Theta(X^4 \log(2X)).$$

The divisor estimate absorbs $\tau_{\max}(X)^2$ into X^η. Summing the charts and adding the zero vector completes the proof. □

10.6 The window estimate and the polar exponent

Retain the window notation $[A, 2A)$, $T \ge 36$, and the heights H, X, S' and Y of Sections 8 and 9. Only windows satisfying (19) need the new estimate.

Proposition 10.7 (Polar window estimate)**.** *Fix positive integers m, n and $\eta > 0$, and let $\delta_{\mathrm{pol}} = 60n + 56m$. Every window above satisfying (19) obeys*

$$\begin{aligned} T \ll_{m,n,\eta} a^\eta (X^5 + X^4 H^{1/2} + X^{14/3} \max(H, S', Y)^{1/17} \\ + X^{5-1/(3\delta_{\mathrm{pol}})} H^{1/\delta_{\mathrm{pol}}} + X^{5-2m/(3\delta_{\mathrm{pol}})} (S')^{m/\delta_{\mathrm{pol}}} + X^5 Y^{n/\delta_{\mathrm{pol}}}). \end{aligned} \tag{37}$$

Proof. The earlier exceptional patterns, conics and bounded cuts contribute $X^5 + X^4 H^{1/2}$, as in Proposition 9.11. Among the remaining patterns, Lemma 10.5 and Proposition 10.6 bound those that are branch-bad or satisfy $\Theta = 0$ by $O_\eta(a^\eta X^{14/3})$. Here $X \ll a^7$, so the extra X^η loss is absorbed into a^η. For these patterns use the unpowered graph of Section 8 ($m = n = 1$, so $\delta_{\min} = 17$), whose box has size $\max(H, S', Y)$; this gives the third term of (37).

For each remaining retained component, Lemma 10.3 gives degree 24, and (28) gives graph degree at least δ_{pol}. Its upper degree is bounded in terms of m, n by Lemma 8.12. The coefficient-uniform curve bound and Lemma 9.9 therefore give, for a pattern with shortest lattice length λ, a contribution

$$\ll_{m,n,\eta} a^\eta ((H/\lambda)^{1/\delta_{\mathrm{pol}}} + (S'/\lambda^2)^{m/\delta_{\mathrm{pol}}} + Y^{n/\delta_{\mathrm{pol}}}).$$

Summing with Lemma 9.6, at exponents $1/\delta_{\mathrm{pol}}$ and $2m/\delta_{\mathrm{pol}}$, produces the last three terms. □

Theorem 10.8 (Polar bound)**.** *For every fixed pair of positive integers m, n and every $\varepsilon > 0$,*

$$P(a, b) \ll_{m,n,\varepsilon} a^{E_{\mathrm{pol}}(m,n)+\varepsilon} \quad (0 < b < a),$$

where, with $\delta_{\mathrm{pol}} = 60n + 56m$,

$$E_{\mathrm{pol}}(m, n) = \max\left\{\frac{5}{17}, \frac{45\delta_{\mathrm{pol}}}{153\delta_{\mathrm{pol}} - 1}, \frac{45\delta_{\mathrm{pol}} - 3m}{153\delta_{\mathrm{pol}} - 11m}, \frac{15\delta_{\mathrm{pol}} + 6n}{51\delta_{\mathrm{pol}} + 20n}\right\}.$$

For $(m, n) = (3, 2)$ this exponent is $4317/14677$. Optimizing over the fixed powers gives

$$P(a, b) \ll_\varepsilon a^{\beta_* + \varepsilon}, \qquad \beta_* = \frac{41301 - 3\sqrt{82849}}{137480} = 0.2941336553\ldots.$$

Proof. Apply the window argument of Section 9 to (37). The three terms from the exceptional contribution $X^{14/3} \max(H, S', Y)^{1/17}$ give exponents $717/2440$ (twice, since H and S' satisfy the same bound $\ll a^{1+o(1)}/T^3$) and $732/2491$, all below $5/17$. The other terms give the displayed maximum. The six-step and drop bounds handle windows outside (19); dyadic summation and the factor a^η absorb the remaining losses.

For $(m,n)=(3,2)$, $\delta_{\mathrm{pol}}=288$ and the last three fractions are $12960/44063$, $4317/14677$ and $1083/3682$. For optimization put $r=n/m$. The curvature and rank-five fractions become

$$E_S(r)=\frac{2700r+2517}{9180r+8557},\qquad E_Y(r)=\frac{906r+840}{3080r+2856}.$$

The first decreases and the second increases. They cross at $r_*=(\sqrt{82849}-47)/360$, with common value β_*. Approximate r_* by positive rationals and scale the corresponding numerator and denominator until $45\delta_{\mathrm{pol}}/(153\delta_{\mathrm{pol}}-1)<\beta_*$. The arbitrarily small exponent loss absorbs the approximation, proving Theorem 1.1. □

11 Rises in the quotient increments

Throughout this section $k_i=q_{i+1}-q_i\ge 1$, $d_i=b_i-b_{i+1}\ge 1$, and a *rise* is an index $0\le i\le P(a,b)-3$ with $k_i<k_{i+1}$.

Lemma 11.1 (Six quotients in arithmetic progression)**.** *Let $q,h\ge 1$ and $x_0>x_1>\cdots>x_6\ge 0$ satisfy $x_j(q+jh)+x_{j+1}=a$ for $0\le j\le 5$. Then $q^4\le 324\,h^3a$.*

Proof. Let $f_j=x_j-x_{j+1}$, and let $E=(f_1-f_2)h$ and $F=(f_2-f_3)h$ be the curvatures of the blocks starting at x_1 and x_2 (all their increments equal h). With $U=f_0-2f_1+f_2$, the division equations $\mathcal{E}_j\colon x_j(q+jh)+x_{j+1}-a=0$ give the exact identity

$$qhU=(3h-1)E+F,\qquad\text{namely}\qquad qhU-(3h-1)E-F=h(\mathcal{E}_0-3\mathcal{E}_1+3\mathcal{E}_2-\mathcal{E}_3).$$

This is the case $k_0=k_1=k_2=k_3=h$ of Lemma 5.1, where $L=2h^3U$. By Lemma 3.2, $E,F\ge 1$, so the integer U is at least 1.

Now bound E and F. Subtracting consecutive division equations gives $q_jf_j\le hx_{j+1}$, where $q_j=q+jh$. Since $x_{j+1}q_{j+1}\le a$, this yields $q^2f_j\le ha$ for $1\le j\le 4$. By Lemma 3.4, $Ex_3\le(h+1)(f_1+f_2+f_3)^2\le 9h^2(h+1)a^2/q^4$. Also $a<x_3(q+3h+1)$, so $Eq^4\le 9h^2(h+1)(q+3h+1)a$, and likewise $Fq^4\le 9h^2(h+1)(q+4h+1)a$. Hence

$$q^5h\le q^4((3h-1)E+F)\le 27h^3(h+1)(q+4h+1)a.$$

If $q\ge h$ this gives $q^4\le 324h^3a$. If $q<h$ then $q^4<h^4\le h^3a$, since $h\le q_1\le a$. □

Lemma 11.2 (Drops decrease between rises)**.** *Whenever $b_{i+2}>0$, $d_{i+1}k_i<d_ik_{i+1}$. In particular, if $k_{i+1}\le k_i$ then $d_{i+1}<d_i$.*

Proof. This is $D=d_ik_{i+1}-d_{i+1}k_i\ge 1$ (Lemma 3.2). If $k_{i+1}\le k_i$ and $d_{i+1}\ge d_i$, then $d_ik_{i+1}\le d_{i+1}k_i$, which is a contradiction. □

Lemma 11.3 (Bounded multiplicity)**.** *Let $c_j\ge 1$ be integers, $j\in J$, such that no value occurs more than M times. Then $|J|^2\le 2M\sum_{j\in J}c_j$.*

Proof. Induct on $|J|$. Let c_{j_0} be the largest value. The values lie in $[1,c_{j_0}]$, so $|J|\le Mc_{j_0}$. Removing j_0 and using $|J|^2=(|J|-1)^2+2|J|-1$ completes the induction. □

Proposition 11.4 (Rises in a window)**.** *Let $x_0,\dots,x_{T-1}$ be a window run in $[A,2A)$ with $T\ge 2$, and let ρ be the number of rises among its $T-1$ increments. Then:*

1. $(T-1)^2\le 2(\rho+1)(A-1)$;

2. $T-1 \le \lfloor a/A \rfloor$;

3. *if* $6A \le a$, *then* $T-1 = \ell + s$ *with* $\ell^2 \le 8(\rho+1)\lfloor a/A \rfloor$ *and* $s^3 \le c_{\mathrm{low}} A$, *where* $c_{\mathrm{low}} = 324 \cdot 3^4 = 26244$.

Proof. Split the increments into *runs*: maximal stretches containing no rise. There are at most $\rho + 1$ runs.

(1) By Lemma 11.2, the drops strictly decrease along a run, so each value of d_j occurs at most once per run, hence at most $\rho + 1$ times. Lemma 11.3 and $\sum_j d_j = x_0 - x_{T-1} \le A - 1$ give the claim.

(2) The increments are ≥ 1, and they sum to $q_{T-1} - q_0 \le \lfloor a/A \rfloor$.

(3) In $c_{\mathrm{low}} = 324 \cdot 3^4$, the factor 324 comes from Lemma 11.1 and 3^4 from the bound $a < 3Aq$ below. Call an increment h *low* if $c_{\mathrm{low}} h^3 A^4 < a^3$. Five consecutive equal increments h inside the run give six quotients in arithmetic progression. Lemma 11.1 then gives $q^4 \le 324 h^3 a$. On the other hand, $a < 3Aq$, which follows from $a = qx_0 + x_1$, $x_1 < x_0 < 2A$ and $6A \le a$. Together these give $a^3 \le c_{\mathrm{low}} h^3 A^4$, so h is not low. Within a run the increments are nonincreasing, so equal values are consecutive. Hence each low value occurs at most 4 times per run, and at most $4(\rho+1)$ times in all. Lemma 11.3 bounds the number ℓ of low increments by $\ell^2 \le 8(\rho+1)\sum_j k_j \le 8(\rho+1)\lfloor a/A \rfloor$. Let s be the number of high increments. If $s = 0$, its bound is immediate. Otherwise let h_* be their smallest value. Then $s h_* \le \lfloor a/A \rfloor$ and $a^3 \le c_{\mathrm{low}} h_*^3 A^4$, so $s^3 a^3 \le c_{\mathrm{low}} A^4 (a/A)^3 = c_{\mathrm{low}} A a^3$. □

Lemma 11.5 (Geometric sums)**.** *Let* $\alpha, \beta, s, t > 0$ *and* $K \ge 0$. *Then*

$$\sum_{k<K} \min\bigl(\alpha 2^{-sk},\ \beta 2^{tk}\bigr) \le \Bigl(\frac{1}{1-2^{-s}} + \frac{2^t}{2^t-1}\Bigr)\alpha^{t/(s+t)}\beta^{s/(s+t)}.$$

Proof. Let $M = \alpha^{t/(s+t)}\beta^{s/(s+t)}$. Then $(\alpha 2^{-sk})^t (\beta 2^{tk})^s = M^{s+t}$, so for each k either $\beta 2^{tk} \le M$ or $\alpha 2^{-sk} < M$. The terms $\beta 2^{tk} \le M$ grow with ratio 2^t, so they sum to at most $M 2^t/(2^t-1)$. The terms $\alpha 2^{-sk} < M$ decay with ratio 2^{-s}, so they sum to at most $M/(1-2^{-s})$. □

Proof of Theorem 1.5. The rises inside a window are rises of the orbit, so $\rho + 1 \le R$. Combining the three parts of Proposition 11.4 gives, for the window $[2^k, 2^{k+1})$,

$$T_k \le 1 + \sqrt{R}\,\min\bigl(\sqrt{8a}\,2^{-k/2},\ \sqrt{2}\,2^{k/2}\bigr) + \min\bigl(a\,2^{-k},\ 30 \cdot 2^{k/3}\bigr);$$

when $6A > a$, part (2) gives $T - 1 \le a/A < 6 \le 30A^{1/3}$ instead of part (3). Lemma 11.5 gives

$$\sum_k \min\bigl(\sqrt{8a}\,2^{-k/2}, \sqrt{2}\,2^{k/2}\bigr) \ll \bigl(\sqrt{8a}\sqrt{2}\bigr)^{1/2} \ll a^{1/4}.$$

It also gives $\sum_k \min(a2^{-k}, 30 \cdot 2^{k/3}) \ll a^{1/4}$. Since $\log_2 a + 1 \ll a^{1/4}$, summing over k gives the theorem. □

Remark 11.6. By Theorem 1.5, an orbit with $P(a,b) \asymp a^{3/10}$ must have $R \gg a^{1/10}$. Proposition 11.4(3) says more: inside the critical window $A \asymp a^{3/5}$, a run of $T \asymp a^{3/10}$ remainders must contain $\gg a^{1/5}$ rises.

12 The Lean formalization

The source repository is https://github.com/jasondavies/pierce-length-bound (private during manuscript preparation).

The proof baseline is commit `c72edbcd67494c14e3b03d9629c2a20aeef2f1c2`. The development uses Lean 4 `v4.35.0` and Mathlib `v4.35.0`. The unconditional results of this paper are proved without additional mathematical hypotheses.

The strongest bound is exported as

```
theorem P_le_polar_optimal :
    ∀ ε > (0 : ℝ), ∃ C > 0, ∀ a b : ℕ,
    0 < b → b < a → (P a b : ℝ) ≤
    C * (a : ℝ) ^ (((41301 : ℝ) - 3 * Real.sqrt 82849) / 137480 + ε)
```

The quantifier order makes uniformity explicit: C may depend on ε, but not on a or b. The function `P a b` is defined by well-founded recursion,

$$P(a,0) = 0, \qquad P(a,b) = 1 + P(a, a \bmod b) \quad (b > 0),$$

and counts the positive terms of the zero-indexed orbit, as in Section 1. The rational bound 4317/14677 is exported separately as `P_le_4317_14677`. Both results are assembled in `PolarProjectionMain` from proved arithmetic, geometric and counting estimates. The repository README gives the correspondence between the other results and their Lean declarations.

The affine-curve theorem (24) is imported from the separate `ccdn-linear-growth` library at revision `9ade620`. That development proves the plane estimate, degree-preserving projection and box rescaling, using the same affine Hilbert degree as this paper. The present repository proves the monic Hilbert irreducibility estimate (Theorem 10.4) and its application to the branch-label family, so neither counting input is assumed.

The repository also contains a conditional refinement of Theorem 1.4, `P_le_loglog`, which proves $P(a,b) \ll a^{3/10}\sqrt{\log\log 3a}$ from two explicit hypotheses: the short-sum estimate for the Erdős–Hooley function from [10], over intervals $(x, x+y]$ with $y \geq x^{1/4}$, and the mean-value estimate from [2]. Their statements include the constants and thresholds; their proofs are not formalized here. This result is weaker than Theorem 1.1 and is not used elsewhere.

The proof library contains no `sorry` and uses no `native_decide`. Its axiom audit permits only `propext`, `Classical.choice` and `Quot.sound`, the standard axioms used by Mathlib. Fifteen public statements are compared with independent Mathlib-only specifications, and their proofs are replayed by Lean, NanoDa and con-ron. The specification file `Challenge.lean` deliberately contains theorem holes; it is separate from the proved solution. Build instructions, the full theorem map, and the verification commands are recorded in the README.

The divisor estimate uses `DivisorPowerBound.lean` from `openai/math` [11] under Apache-2.0, with unchanged proof bodies and adapted imports and module visibility. The repository records the source revision and the provenance of its other dependencies.

13 Limits of the method

The polar refinement brings the exponent close to 5/17, but does not remove that threshold. As explained in Section 1, 5/17 is the exponent obtained if every pattern carries $O(1)$ starts, and the estimates here never do better than that: the bounded cuts and conics contribute $a^{o(1)}$ starts for each of the $X^{5+o(1)}$ patterns, and the exceptional patterns and the windows where the line-exclusion

condition (19) fails also lead to exponent $5/17$, independently of the coordinate powers m, n. The branch-bad patterns and those with $\Theta = 0$ give only the exponents $717/2440$ and $732/2491$, below this threshold.

The improved graph degree gives

$$\beta_* - \frac{5}{17} = 0.00001600826\ldots.$$

Increasing the graph degree further can reduce this gap. Going below $5/17$ requires an improvement in the other counting contributions, or a different way to combine the information from successive orbit states. This is a limitation of the present estimates, not a lower bound for the Pierce iteration.

The elementary $3/10$ argument has a similar barrier: even with no divisor loss, its two window bounds balance at $A \asymp a^{3/5}$ (Section 4), so better constants or short-interval divisor estimates alone do not change that exponent.

Finally, there is no obvious monotone quantity along an orbit that could replace the window-by-window counting; in particular the curvatures need not decrease. For $a = 13242$, the orbit of 255 begins

$$255,\ 237,\ 207,\ 201,\ 177,\ 144,\ 138,\ 132, \ldots$$

with quotients $51, 55, 63, 65, 74, 91, 95, 100, \ldots$ and first five curvatures $24, 12, 6, 111, 30$. All these remainders lie in the single window $[128, 256)$. The rise bound gives one constraint on the order of the increments, but the present argument does not capture all compatibility conditions along an entire orbit. Shallit's conjectured $O((\log a)^2)$ bound remains open.

A Conic components

In this appendix we prove Proposition 8.9. Let C be a nonexceptional component of degree two. A component that is not retained already contributes $O(1)$ starts, by Lemma 8.2. We may therefore assume that the component C is retained. If it contains at most two free-node triples of good starts, there is nothing to prove. In the remaining case we first construct a parameter whose value is an integer at every such triple.

Lemma A.1 (A bounded integral parameter)**.** *Let C be a retained nonexceptional degree-two component containing the free-node triples of at least three good starts. After permuting the three free-node coordinates to (x, y, z), there are an integer $\Delta_c \neq 0$ and polynomials $\xi_0, \xi_1, \xi_2 \in \mathbb{Z}[t]$ such that*

$$\deg \xi_j \le 2, \qquad |\Delta_c|,\ \max_{j,k} |[t^k]\xi_j| \le (aX)^{O(1)}.$$

The map

$$t \longmapsto (\xi_0(t), \xi_1(t), \xi_2(t))/\Delta_c$$

parametrizes C over $\mathbb{Q}$. Its inverse takes an integer value at every integer point of C, and is injective on those points.

Proof. Every free node of a good start has absolute value at most a. Three distinct such triples lie on an integer plane

$$\lambda_z z + \lambda_x x + \lambda_y y + \lambda_1 = 0$$

with coefficients of size $a^{O(1)}$: take a cross product of two differences as the normal, or, if it vanishes, a nonzero perpendicular vector of the same size. A plane not containing an integral curve of degree

two meets it in at most two distinct points. Thus this plane contains C. Choose the eliminated coordinate z with $\lambda_z \neq 0$.

The six rational pattern coordinates (10) have common denominator $N_0 N_1 \neq 0$ and integer numerators of size at most X^2, so the pattern quadratic f has rational coefficients with numerators and denominators of size $X^{O(1)}$. Substitute $z = -(\lambda_x x + \lambda_y y + \lambda_1)/\lambda_z$ and clear denominators. The resulting plane equation is

$$\alpha_c x^2 + \beta_c xy + \gamma_c y^2 + \ell_x x + \ell_y y + \ell_0 = 0, \tag{38}$$

with integer coefficients of size $(aX)^{O(1)}$. The restriction of f to the plane is nonzero: otherwise the plane would be a component of the irreducible pattern quadric. It has degree two because it contains C, and hence defines C in that plane.

The pullback of the hyperplane at infinity to the normalization of C has degree two. Corollary 8.4 says that each of its positive multiplicities is even. Counting residue degrees, there is therefore just one branch over infinity, of multiplicity two and residue degree one. Consequently the leading binary quadratic in (38) is a nonzero rational scalar times the square of a rational linear form. Indeed, the normalization map is finite and surjective, so the projective conic meets the line at infinity only at the image of this rational point. Its leading binary quadratic therefore has a single rational zero, of multiplicity two. Interchanging x and y if necessary gives

$$\alpha_c \neq 0, \qquad \beta_c^2 - 4\alpha_c\gamma_c = 0.$$

Set

$$\kappa_c = 2\beta_c\ell_x - 4\alpha_c\ell_y, \qquad \epsilon_c = \ell_x^2 - 4\alpha_c\ell_0, \qquad t = 2\alpha_c x + \beta_c y + \ell_x.$$

Completing the square in (38) gives

$$t^2 = \kappa_c y + \epsilon_c. \tag{39}$$

Here $\kappa_c \neq 0$. If it were zero, a rational point of C with parameter t_0 would give $\epsilon_c = t_0^2$, and the equation would factor as $(t - t_0)(t + t_0) = 0$. Since C is irreducible, it would lie in one of these two lines, contradicting its degree.

We can therefore solve explicitly:

$$y = \frac{t^2 - \epsilon_c}{\kappa_c}, \qquad x = \frac{t - \beta_c y - \ell_x}{2\alpha_c}, \qquad z = -\frac{\lambda_x x + \lambda_y y + \lambda_1}{\lambda_z}.$$

Substitution verifies the conic and plane equations for every rational t, and the displayed linear expression in x, y recovers t. In particular, t is an integer whenever (x, y, z) is an integer point. Distinct points have distinct parameters. Taking $\Delta_c = 2\alpha_c\kappa_c\lambda_z$ clears all three denominators, and all the quantities involved are polynomials of bounded degree in coefficients of size $(aX)^{O(1)}$, which gives the stated bounds. □

Lemma A.2 (Properness and coefficient height)**.** *In the parameter of Lemma A.1, write*

$$E_0(t) = \frac{F_E(t)}{G_E(t)}, \qquad F_E, G_E \in \mathbb{Z}[t], \qquad \gcd_{\mathbb{Q}[t]}(F_E, G_E) = 1.$$

The polynomials may be chosen so that

$$\deg G_E - \deg F_E = 3, \qquad \max(\deg F_E, \deg G_E) \leq D_0, \qquad \max_k(|[t^k]F_E|, |[t^k]G_E|) \leq C_0 a^{k_0},$$

for absolute constants D_0, C_0, k_0. In particular, G_E is nonconstant. At the parameter t of every good start, $G_E(t) \neq 0$ and $F_E(t)/G_E(t) = I_3(i)$ is an integer.

Proof. The explicit parametrization identifies C with the affine line and its smooth projective model with $\mathbb{P}^1$. Its point at infinity has hyperplane multiplicity two. Lemma 8.6 therefore gives

$$\operatorname{ord}_{t=\infty} E_0 = 3.$$

The function is nonzero, since it is positive at a good start. For a reduced rational function on $\mathbb{P}^1$, its order at infinity is $\deg G_E - \deg F_E$, proving properness.

We next bound the coefficients uniformly in the component, including after cancellation. The six rational pattern coordinates, the three coordinate polynomials ξ_j/Δ_c, and the dividend a have the common nonzero denominator $N_0N_1\Delta_c$, and their numerators are polynomials in t of degree at most two with coefficients of size $(aX)^{O(1)}$. The cofactor formula defining E_0 is a quotient of two fixed polynomials in these inputs. Substituting and clearing a fixed power of the common denominator gives nonzero integer polynomials $\tilde{n}, \tilde{d}$ of bounded degree with coefficients of size $(aX)^{O(1)}$.

Choose a primitive $\Pi \in \mathbb{Z}[t]$ representing $\gcd_{\mathbb{Q}[t]}(\tilde{n}, \tilde{d})$. By Gauss's lemma, Π divides both $\tilde{n}$ and $\tilde{d}$ in $\mathbb{Z}[t]$, so $F_E = \tilde{n}/\Pi$ and $G_E = \tilde{d}/\Pi$ are coprime integer polynomials with the same ratio. Every integer factor of a nonzero integer polynomial of degree at most D_0 and height $\mathcal{H}$ has height at most $2^{D_0}(D_0+1)\mathcal{H}$, by comparing Mahler measures; so cancellation preserves the bound $(aX)^{O(1)}$. Since $X \le a^{O(1)}$, this is the asserted $C_0a^{k_0}$. Finally, the reconstruction denominator is nonzero at every good start, hence so are $\tilde{d}$ and its factor G_E at the corresponding parameter, and the reconstruction identity gives $F_E(t)/G_E(t) = I_3(i) \in \mathbb{Z}$. □

Lemma A.3 (Integer values of a rational function)**.** *Let $F_E, G_E \in \mathbb{Z}[t]$ be coprime over $\mathbb{Q}$, of degrees at most D_0, with G_E nonconstant and coefficient heights at most $C_0a^{k_0}$. For every $\eta > 0$, the number of integers t such that $G_E(t) \neq 0$ and $F_E(t)/G_E(t) \in \mathbb{Z}$ is $O_\eta(a^\eta)$. The implied constant depends only on η, D_0, C_0, k_0.*

Proof. The integer resultant $R = \operatorname{Res}(F_E, G_E)$ is nonzero. The adjugate of the Sylvester matrix gives $U, V \in \mathbb{Z}[t]$ with $UF_E + VG_E = R$. At any parameter in question, $G_E(t) \mid F_E(t)$, and hence $G_E(t) \mid R$. There are $2\tau(|R|)$ possible signed values of $G_E(t)$. For each such value v, the nonzero polynomial $G_E - v$ has at most $\deg G_E \le D_0$ roots. Thus the number of parameters is at most

$$2D_0\tau(|R|).$$

The Sylvester matrix has size at most $2D_0$, and its entries are bounded by $C_0a^{k_0}$. Expanding its determinant gives

$$1 \le |R| \le (2D_0)!\left(C_0a^{k_0}\right)^{2D_0}.$$

The divisor estimate $\tau(n) \ll_\varepsilon n^\varepsilon$, with ε chosen in terms of η and the fixed degree and height exponents, proves the claim. □

Proof of Proposition 8.9. Apply the last lemma to the reconstructed invariant E_0. The parameter is injective on free-node triples, and reconstruction of the strictly increasing q_i makes those triples injective in the good starts. Each retained nonexceptional conic therefore contributes $O_\eta(a^\eta)$ starts. The components with at most two free-node triples of good starts and the nonretained components were already bounded absolutely. Apply the conic estimate with $\eta/2$ and the pattern estimate $O_{\eta/2}(a^{\eta/2}X^5)$. Since there are at most twenty-four components per pattern, all nonexceptional conics together contribute $O_\eta(a^\eta X^5)$. □

B The upper bound for graph degrees

Proof of the upper bound in Lemma 8.12. Only total polynomial degrees in the nodes matter here, not the heights of the pattern-dependent coefficients. The cofactors c_j of Section 8.2 have bounded degree, hence so do the numerator and denominator of $E_0 = E_{\mathrm{n}}/E_{\mathrm{d}}$. Since H_3, H_3' have degree at most three and the gaps have degree one, the three curvature coordinates have the common denominator $\Omega = E_{\mathrm{d}} H_3 k_1$, and Ω and the numerators $\mathcal{N}_0, \mathcal{N}_1, \mathcal{N}_2$ have degree at most an absolute constant D_W. The numerator for J with this denominator is $\mathcal{N}_3 = J\Omega$, of degree at most $D_W + 3$. The denominator is nonzero on C, since it is nonzero at a good start.

Put $\bar{m} = m + n$, and represent the powered coordinates with denominator $\Omega^{\bar{m}}$ and numerators

$$\mathcal{N}_j^m \Omega^{\bar{m}-m} \quad (0 \le j \le 2), \qquad \mathcal{N}_3^n \Omega^{\bar{m}-n}.$$

All these numerators, and those of the node coordinates written with the same denominator, have degree at most $(D_W + 3)\bar{m} + 1$. So multiplying a polynomial of degree at most t in the graph coordinates by $\Omega^{\bar{m}t}$ gives a polynomial in the original nodes of degree at most $((D_W + 3)\bar{m} + 1)t$. This operation is injective on the corresponding spaces of functions on C, because Ω is nonzero in $\mathbb{Q}(C)$. Comparing affine Hilbert functions gives

$$\delta t + O(1) \le d((D_W + 3)\bar{m} + 1)t + O(1),$$

and hence the asserted bound with $D_{\mathrm{gr}} = D_W + 4$. □

C A reduced basis in dimension three

The constant $64/27 = (4/3)(16/9)$ in Lemma 9.3 comes from two successive size reductions. We give the elementary argument for completeness.

Proof of Lemma 9.3. Let V be the least nonzero $|\det|$ of a triple in Λ; the triple $(\kappa\mathbf{e}_1, \kappa\mathbf{e}_2, \kappa\mathbf{e}_3)$ gives $V \le \kappa^3$. A triple $(\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3)$ with $|\det| = V$ is a basis: if not, some $x \in \Lambda$ has a Cramer coordinate that is not an integer, say the first, $\det(x, \mathbf{f}_2, \mathbf{f}_3)/\det(\mathbf{f}_1, \mathbf{f}_2, \mathbf{f}_3)$, and subtracting the integer part of that coordinate times $\mathbf{f}_1$ from x gives a triple with smaller nonzero determinant. Among all bases (all have $|\det| = V$) minimize $g_1 = |\mathbf{f}_1|^2$, and then the Gram determinant $g_2 = g_1|\mathbf{f}_2|^2 - \langle \mathbf{f}_1, \mathbf{f}_2\rangle^2$. Put $g_3 = V^2$.

Choose an integer z_1 with $|2(\langle \mathbf{f}_1, \mathbf{f}_2\rangle + z_1 g_1)| \le g_1$. Then $(\mathbf{f}_2 + z_1\mathbf{f}_1, \mathbf{f}_1, \mathbf{f}_3)$ is a basis, so $g_1 \le |\mathbf{f}_2 + z_1\mathbf{f}_1|^2$, and $g_1|\mathbf{f}_2 + z_1\mathbf{f}_1|^2 = g_2 + (\langle \mathbf{f}_1, \mathbf{f}_2\rangle + z_1 g_1)^2$. Hence $3g_1^2 \le 4g_2$.

Let $\langle x, x'\rangle_1 = g_1\langle x, x'\rangle - \langle \mathbf{f}_1, x\rangle\langle \mathbf{f}_1, x'\rangle$, so that $\langle \mathbf{f}_2, \mathbf{f}_2\rangle_1 = g_2$, and Sylvester's identity gives $\langle \mathbf{f}_2, \mathbf{f}_2\rangle_1\langle \mathbf{f}_3, \mathbf{f}_3\rangle_1 - \langle \mathbf{f}_2, \mathbf{f}_3\rangle_1^2 = g_1 g_3$. Choose an integer z_2 with $|2(\langle \mathbf{f}_2, \mathbf{f}_3\rangle_1 + z_2 g_2)| \le g_2$ and put $y = \mathbf{f}_3 + z_2\mathbf{f}_2$. Then $(\mathbf{f}_1, y, \mathbf{f}_2)$ is a basis with first vector $\mathbf{f}_1$, so $g_2 \le g_1|y|^2 - \langle \mathbf{f}_1, y\rangle^2 = \langle y, y\rangle_1$, while $g_2\langle y, y\rangle_1 = (\langle \mathbf{f}_2, \mathbf{f}_3\rangle_1 + z_2 g_2)^2 + g_1 g_3$. Hence $3g_2^2 \le 4g_1 g_3$.

Finally choose an integer z_3 with $|2(\langle \mathbf{f}_1, y\rangle + z_3 g_1)| \le g_1$, and use the basis $\mathbf{f}_1$, $\mathbf{f}_2' = \mathbf{f}_2 + z_1\mathbf{f}_1$, $\mathbf{f}_3' = y + z_3\mathbf{f}_1$. The identities above give $3g_1|\mathbf{f}_2'|^2 \le 4g_2$ and $4g_1 g_2|\mathbf{f}_3'|^2 \le g_2^2 + 4g_1 g_3 + g_1^2 g_2$. With the two inequalities this gives $9g_2|\mathbf{f}_3'|^2 \le 16 g_3$. Since $|\mathbf{f}_1|^2 = g_1$, the product of the bounds $|\mathbf{f}_2'|^2 \le \frac{4}{3} g_2/g_1$ and $|\mathbf{f}_3'|^2 \le \frac{16}{9} g_3/g_2$ is the claim for the basis $\mathbf{f}_1, \mathbf{f}_2', \mathbf{f}_3'$; rename $\mathbf{f}_2', \mathbf{f}_3'$ as $\mathbf{f}_2, \mathbf{f}_3$. Replacing $\mathbf{f}_1$ by $-\mathbf{f}_1$ if necessary makes the determinant positive. For the coordinates, $x_1 V = \det(x, \mathbf{f}_2, \mathbf{f}_3)$, and Hadamard's inequality gives $\det(x, \mathbf{f}_2, \mathbf{f}_3)^2 \le |x|^2|\mathbf{f}_2|^2|\mathbf{f}_3|^2$, so $27|\mathbf{f}_1|^2 x_1^2 V^2 \le 64 V^2 |x|^2$; likewise for x_2 and x_3. □

D Certificates for the polar bound

D.1 The quartic initial form at the base point

At $(\rho, \varphi, \chi, \rho', \varphi', \chi') = (2, 5, -3, 3, 6, -5)$ the polar base point is $\mathbf{k}_* = (-1/7, -3/28, -1/2)$ and $\nabla f(\mathbf{k}_*) = (-4, 9, -5)$. Use tangent coordinates (s, t) with first-order displacement

$$\delta\mathbf{k} = ((9s - 5t)/4, s, t).$$

Thus the two axes used in Lemma 10.2 have tangent vectors $(9/4, 1, 0)$ and $(-5/4, 0, 1)$. Direct substitution in the Newton quartic, its remainder and the norm polynomial gives

$$\begin{aligned} P_{4,*}(x) &= x^4 - \frac{20}{7}x^3 + \frac{757}{112}x^2 - \frac{669}{112}x + 2, \\ R_1(x; s, t) &= \frac{77s + 7t}{14} + \frac{-56s - 124t}{14}x + \frac{-45s + 157t}{14}x^2 + \frac{28s - 28t}{14}x^3, \\ \Psi_1(s, t) &= \frac{3s + 85t}{28}. \end{aligned}$$

The resultant is

$$N_4(s, t) = \frac{32233977s^4 - 95047956s^3t + 147873750s^2t^2 - 111199572st^3 + 31702169t^4}{6272}.$$

In particular,

$$N_4(1, 0)\Psi_1(0, 1)^4 - N_4(0, 1)\Psi_1(1, 0)^4 = \frac{26291112152499}{60236288}.$$

The realizing matrix in Lemma 10.2 shows that Θ is nonzero on the Plücker cone; no orbit with these parameters is required.

D.2 The irreducible sextic

The specialization $E_\gamma = 24E_\star$ at $(\gamma_1, \gamma_2, \gamma_3) = (-7, 8, -4)$ reduces modulo 71 to a nonzero scalar multiple of

$$q(T) = T^6 + 20T^5 - 14T^4 + 15T^3 - 28T^2 + 5T - 4.$$

More precisely, $E_\star = 57q$ in $\mathbb{F}_{71}[T]$. The following certificates prove that q is irreducible. Repeated squaring modulo q gives

$$\begin{aligned} T^{71^2} &\equiv r_2(T) := 36T^5 + 48T^4 + 49T^3 + 53T^2 + 36T + 42, \\ T^{71^3} &\equiv r_3(T) := 37T^5 + 22T^4 + 60T^3 + 13T^2 + 66T + 35. \end{aligned}$$

For checking the iteration one may first use

$$T^{71} \equiv 14T^5 + 33T^4 + 31T^3 + 34T^2 + 48T \pmod{q}$$

and raise each remainder to the 71st power. Define

$$\begin{aligned} U_2 &= 20T^4 + 34T^3 + 31T^2 + 13T + 24, \\ V_2 &= 31T^5 + 61T^4 + 42T^3 + 61T^2 + 39T + 4, \\ U_3 &= 25T^4 + 11T^3 + 53T^2 + 59T + 48, \\ V_3 &= 7T^5 + 30T^4 + 65T^3 + 64T^2 + 25T + 40. \end{aligned}$$

Direct multiplication in $\mathbb{F}_{71}[T]$ verifies

$$U_2 q + V_2(r_2 - T) = 1, \qquad U_3 q + V_3(r_3 - T) = 1.$$

Thus q has no irreducible factor of degree one or two, since such a factor would divide $T^{71^2} - T$, and none of degree three, since it would divide $T^{71^3} - T$. Every reducible sextic has an irreducible factor of degree at most three, so q is irreducible. Reduction modulo 71 preserves degree, and Gauss's lemma proves $E_\star$ irreducible over $\mathbb{Q}$.

Acknowledgements

The argument was developed with the help of GPT-6-Astra. This includes the use of orthogonal polynomials and adjacent labels behind Section 7, the first proof of a bound $a^{3/10-\delta}$ through resultants, the use of Lemma 7.3 to bound the number of states per pattern, the sign identities of Lemma 7.5 and the slice count behind the 11/37 bound, the observation that the pattern integers form a Plücker vector, and the proposal to weight minor vectors by the shortest vectors of their lattices, behind the 45/152 bound. Subsequent AI-assisted work developed and formalized the joint-invariant graphs, their normalization and valuation arguments, the powered-coordinate optimization, and the external affine-curve counting theorem. OpenAI Codex assisted with this formalization and with automated statement, dependency and submission checks. These checks are not independent human peer review. The unconditional results, including the polar refinement and its arithmetic sieve, and the stated conditional implications were verified in Lean 4 against Mathlib. We thank the Mathlib community, and the authors of `openai/math` for the formalized divisor bound.

References

[1] O. Baraskar and I. Vukusic, *Bounds for sets of remainders*, arXiv:2508.20853 (2025).

[2] R. de la Bretèche and G. Tenenbaum, *Note on the mean value of the Erdős–Hooley Delta-function*, Acta Arith. **219** (2025), no. 4, 379–394, doi:10.4064/aa250107-13-2.

[3] W. Castryck, R. Cluckers, P. Dittmann and K. H. Nguyen, *The dimension growth conjecture, polynomial in the degree and without logarithmic factors*, Algebra Number Theory **14** (2020), no. 8, 2261–2294, doi:10.2140/ant.2020.14.2261.

[4] R. Cluckers, P. Dèbes, Y. I. Hendel, K. H. Nguyen and F. Vermeulen, *Improvements on dimension growth results and effective Hilbert's irreducibility theorem*, Forum Math. Sigma **13** (2025), e153, doi:10.1017/fms.2025.10096.

[5] Z. Chase and M. Pandey, *On the length of Pierce expansions*, arXiv:2211.08374 (2022).

[6] P. Erdős and J. O. Shallit, *New bounds on the length of finite Pierce and Engel series*, J. Théor. Nombres Bordeaux **3** (1991), no. 1, 43–53, doi:10.5802/jtnb.41.

[7] J. Franke, Yu. I. Manin and Yu. Tschinkel, *Rational points of bounded height on Fano varieties*, Invent. Math. **95** (1989), 421–435, doi:10.1007/BF01393904.

[8] L. de Moura and S. Ullrich, *The Lean 4 theorem prover and programming language*, in: Automated Deduction – CADE 28, Lecture Notes in Comput. Sci. **12699**, Springer, 2021, 625–635, doi:10.1007/978-3-030-79876-5_37.

[9] The mathlib Community, *The Lean mathematical library*, in: Proceedings of the 9th ACM SIGPLAN International Conference on Certified Programs and Proofs (CPP 2020), 367–381, doi:10.1145/3372885.3373824.

[10] M. Nair and G. Tenenbaum, *Short sums of certain arithmetic functions*, Acta Math. **180** (1998), 119–144, doi:10.1007/BF02392880.

[11] OpenAI, `openai/math`, https://github.com/openai/math.

[12] J. O. Shallit, *A discrete iteration related to Pierce expansions*, Open Problem Garden, https://www.openproblemgarden.org/op/a_discrete_iteration_related_to_pierce_expansions.

[13] T. A. Pierce, *On an algorithm and its use in approximating roots of algebraic equations*, Amer. Math. Monthly **36** (1929), no. 10, 523–525, doi:10.1080/00029890.1929.11987017.

[14] J. O. Shallit, *Metric theory of Pierce expansions*, Fibonacci Quart. **24** (1986), no. 1, 22–40, doi:10.1080/00150517.1986.12429786.

[15] J. O. Shallit, *Talks*, https://cs.uwaterloo.ca/~shallit/talks.html (accessed October 2026).