

# Relations and Functions

CMSC250

# Relations

# Definition

- Let  $A, B$  be sets. A **relation**  $R$  from  $A$  to  $B$  is **any subset** of  $A \times B$ .

# Examples

- $(<, \mathbb{R} \times \mathbb{R})$ 
  - $\{ \dots, (-1.5, -1.2), (-1.4, -1.2), (\sqrt{2}, \sqrt{3}), (\sqrt{2}, \sqrt{5}), \dots \}$

# Examples

- $(<, \mathbb{R} \times \mathbb{R})$ 
  - $\{ \dots, (-1.5, -1.2), (-1.4, -1.2), (\sqrt{2}, \sqrt{3}), (\sqrt{2}, \sqrt{5}), \dots \}$
- $(\leq, \mathbb{R} \times \mathbb{R})$ 
  - $\{ \dots, (2, 2), (2, 2.1), (\sqrt{2}, \sqrt{3}), (\sqrt{2}, \sqrt{5}), \dots \}$

# Examples

- $(<, \mathbb{R} \times \mathbb{R})$ 
  - $\{ \dots, (-1.5, -1.2), (-1.4, -1.2), (\sqrt{2}, \sqrt{3}), (\sqrt{2}, \sqrt{5}), \dots \}$
- $(\leq, \mathbb{R} \times \mathbb{R})$ 
  - $\{ \dots, (2, 2), (2, 2.1), (\sqrt{2}, \sqrt{3}), (\sqrt{2}, \sqrt{5}), \dots \}$
- $(R, \mathbb{R} \times \mathbb{N})$ 
  - $\{(r, n) \mid n \text{ appears in the decimal expansion of } r \}$
  - E.g:  $\{ \dots, (\pi, 1), (e, 7), (1/3, 3), \dots \}$
  - We would formally say that all of the above are elements of the relation  $R$

# Reflexivity

- A relation  $X \subseteq A \times A$  is **reflexive** if

$$(\forall a \in A)[(a, a) \in X]$$

# Reflexivity

- A relation  $X \subseteq A \times A$  is **reflexive** if

$$(\forall a \in A)[(a, a) \in X]$$

- Examples:
  - $(\leq, \mathbb{N} \times \mathbb{N})$  is reflexive, since  $(\forall n \in \mathbb{N})[n \leq n]$
  - $(<, \mathbb{N} \times \mathbb{N})$  is **not** reflexive, since  $\sim (\forall n \in \mathbb{N})[n < n]$  (in fact, there is no such  $n$ )
  - $(R, \mathbb{N} \times \mathbb{N})$  defined as  $\{(x, y) \mid x + y \geq 100\}$  is **not** reflexive (e.g  $10 \in \mathbb{N}$ , but  $(10, 10) \notin R$ )



# Symmetry

- A relation  $X \subseteq A \times A$  is **symmetric** if

$$(\forall a_1, a_2 \in A) [ ((a_1, a_2) \in X) \Rightarrow ((a_2, a_1) \in X) ]$$

# Symmetry

- A relation  $X \subseteq A \times A$  is **symmetric** if

$$(\forall a_1, a_2 \in A) [ ((a_1, a_2) \in X) \Rightarrow ((a_2, a_1) \in X) ]$$

- Examples:
  - $(\leq, \mathbb{N} \times \mathbb{N})$  is **not** symmetric since  $4 \leq 5$  but  $\neg (5 \leq 4)$
  - $(<, \mathbb{N} \times \mathbb{N})$  is **not** symmetric (see above)
  - $(R, \mathbb{N} \times \mathbb{N})$  defined as  $\{(x, y) \mid x + y \geq 100\}$  is **symmetric** since

$$(x + y \geq 100) \Rightarrow (y + x \geq 100)$$

# Transitivity

- A relation  $X \subseteq A \times A$  is **transitive** if

$$(\forall a_1, a_2, a_3 \in A) [ ((a_1, a_2) \in X) \wedge ((a_2, a_3) \in X) \Rightarrow (a_1, a_3) \in X ]$$

# Transitivity

- A relation  $X \subseteq A \times A$  is **transitive** if

$$(\forall a_1, a_2, a_3 \in A) [ ((a_1, a_2) \in X) \wedge ((a_2, a_3) \in X) \Rightarrow (a_1, a_3) \in X ]$$

- $(\leq, \mathbb{N} \times \mathbb{N})$  is **transitive** since  $((x \leq y) \wedge (y \leq z)) \Rightarrow (x \leq z)$
- $(<, \mathbb{N} \times \mathbb{N})$  is **transitive** (see above)
- $(R, \mathbb{N} \times \mathbb{N})$  defined as  $\{(x, y) \mid x + y \geq 100\}$  ???

# Transitivity

- A relation  $X \subseteq A \times A$  is **transitive** if

$$(\forall a_1, a_2, a_3 \in A) [ ((a_1, a_2) \in X) \wedge ((a_2, a_3) \in X) \Rightarrow (a_1, a_3) \in X ]$$


- $(\leq, \mathbb{N} \times \mathbb{N})$  is **transitive** since  $((x \leq y) \wedge (y \leq z)) \Rightarrow (x \leq z)$
- $(<, \mathbb{N} \times \mathbb{N})$  is **transitive** (see above)
- $(R, \mathbb{N} \times \mathbb{N})$  defined as  $\{(x, y) \mid x + y \geq 100\}$  is **not transitive** since (counter-example):

$$((1, 100) \in R) \wedge ((100, 5) \in R), \text{ but } (1, 5) \notin R$$

# Functions


# Functions

- Most basic representation: **Arrow Diagrams**



# Functions

- Most basic representation: **Arrow Diagrams**




A is called the **domain** and B is called the **co-domain** and we say that “ $f$  is such that it maps elements from A to B” ( $f: A \mapsto B$ )



# Functions

- Most basic representation: **Arrow Diagrams**



A is called the **domain** and B is called the **co-domain** and we say that “ $f$  is such that it maps elements from A to B”  
( $f: A \mapsto B$ )


$\backslash\text{mapsto}$  in LaTeX

# Example 1

- Is this a function?

Yes

No




# Example 1

- Is this a function?

Yes

No



- Domain:  $\{0, 1, 2, 3, 4\}$
- Co-domain:  $\{0, 1, 4\}$
- Formula (that we came up with):  $f(x) = x^2 \bmod 5$


Just because two 'x's map to the same 'y' doesn't make this a non-function... it just makes it a **non-injective** (not "1-1") function

## Example 2

- Is this a function?

Yes

No




## Example 2

- Is this a function?

Yes

No




- Every element of the domain should map to some co-domain element!

## Example 2

- Is this a function?

Yes

No




## Example 2

- Is this a function?

Yes

No



Fails the  
“vertical line”  
test (2 different  
`y`'s mapped to  
by the same `x`)

## Example 3

- Is this a function?

Yes

No

$$f: \mathbb{N} \mapsto \mathbb{N}, \text{ and } f(x) = x/2$$



## Example 3

- Is this a function?

Yes

No

$$f: \mathbb{N} \mapsto \mathbb{N}, \text{ and } f(x) = x/2$$

- For any **odd** selection of  $x \in \mathbb{N}$ , there is no  $x/2 \in \mathbb{N}$ !
- $f(4) = 2 \in \mathbb{N}$ , but  $f(5) = 2.5 \notin \mathbb{N}$

## Example 3

Yes

No

- Is this a function?

$$f: \mathbb{N} \mapsto \mathbb{N}, \text{ and } f(x) = x/2$$

- For any **odd** selection of  $x \in \mathbb{N}$ , there is no  $x/2 \in \mathbb{N}$ !
- $f(4) = 2 \in \mathbb{N}$ , but  $f(5) = 2.5 \notin \mathbb{N}$
- **What about this?**

$$f: \mathbb{N} \mapsto \mathbb{Q}, \text{ and } f(x) = x/2$$

## Example 3

- Is this a function?

Yes

No

$$f: \mathbb{N} \mapsto \mathbb{N}, \text{ and } f(x) = x/2$$

- For any **odd** selection of  $x \in \mathbb{N}$ , there is no  $x/2 \in \mathbb{N}$ !
- $f(4) = 2 \in \mathbb{N}$ , but  $f(5) = 2.5 \notin \mathbb{N}$
- **What about this?**

$$f: \mathbb{N} \mapsto \mathbb{Q}, \text{ and } f(x) = x/2$$

## Example 4

- Are the following **valid functions?**

Yes


No

## Example 4

- Are the following **valid functions**?

Yes

No




## Example 4

Yes

No

(As long as the  
domain is  $\mathbb{R}^{>0}$ !!)

Log function




## Example 5

- Are the following **valid functions**?

Yes

No




## Example 5

- Are the following **valid functions**?

Yes

No



Fails the  
“vertical line”  
test




# Example 5

- Are the following **valid functions**?

Yes

No




## Example 5

- Are the following **valid functions**?

Yes

No



It just happens to  
not be “1-1”

# Surjective functions

- A function  $f: X \mapsto Y$  is called **surjective (or onto)** iff

$$(\forall y \in Y, \exists x \in X)[f(x) = y]$$

# Surjective functions

- A function  $f: X \mapsto Y$  is called **surjective (or onto)** iff

$$(\forall y \in Y, \exists x \in X)[f(x) = y]$$

- **Intuitively:**  $f$ 's co-domain is “full” with pointer heads

# Surjective functions

- A function  $f: X \mapsto Y$  is called **surjective (or onto)** iff

$$(\forall y \in Y, \exists x \in X)[f(x) = y]$$


- **Intuitively:**  $f$ 's co-domain is “full” with pointer heads
- Mnemonic rule to remember the name “surjective”: in French, “sur” means “above”, “on”, or “**onto**”.

# Surjective functions

- A function  $f: X \mapsto Y$  is called **surjective (or onto)** iff

$$(\forall y \in Y, \exists x \in X)[f(x) = y]$$

- **Intuitively:**  $f$ 's co-domain is “full” with pointer heads
- Mnemonic rule to remember the name “surjective”: in French, “sur” means “above”, “on”, or “**onto**”.



# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**,  
given the following  
domain / co-domain pairs?

Yes

No

# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**,  
given the following  
domain / co-domain pairs?

a)  $D = \mathbb{R}, C = \mathbb{R}$

Yes

No



# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**,  
given the following  
domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$  No (e.g -1 is not mapped to)

# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**, given the following domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$  No (e.g -1 is not mapped to)

*b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$

# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**,  
given the following  
domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$  No (e.g -1 is not mapped to)

*b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$  Yes

# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**,  
given the following  
domain / co-domain pairs?

Yes

No

- a)*  $D = \mathbb{R}, C = \mathbb{R}$  No (e.g -1 is not mapped to)
- b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$  Yes
- c)*  $D = \mathbb{N}, C = \text{SQUARES}$

# Quiz on surjectivity

- Is  $f(x) = x^2$  **surjective**,  
given the following  
domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$  No (e.g -1 is not mapped to)

*b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$  Yes

*c)*  $D = \mathbb{N}, C = \text{SQUARES}$  Yes

# Injective functions

- A function  $f: X \mapsto Y$  is called **injective (or 1-1)** iff

$$(\forall x_1, x_2 \in X) [(f(x_1) = f(x_2)) \Rightarrow (x_1 = x_2)]$$

- **Intuitively:** Every element of the co-domain is mapped to by **at most one** element of the domain.

# Injective functions

- A function  $f: X \mapsto Y$  is called **injective (or 1-1)** iff

$$(\forall x_1, x_2 \in X) [(f(x_1) = f(x_2)) \Rightarrow (x_1 = x_2)]$$

- **Intuitively:** Every element of the co-domain is mapped to by **at most one** element of the domain.
- *Why “at most one” and not exactly one?*





# Injective functions

- A function  $f: X \mapsto Y$  is called **injective (or 1-1)** iff

$$(\forall x_1, x_2 \in X) [(f(x_1) = f(x_2)) \Rightarrow (x_1 = x_2)]$$

- **Intuitively:** Every element of the co-domain is mapped to by **at most one** element of the domain.
- *Why at most one and not exactly one?*
- *Because 1-1 but **not onto** functions are possible!*





# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**,  
given the following  
domain / co-domain pairs?

Yes

No

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**, given the following domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$ :

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**,  
given the following  
domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**, given the following domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )

*b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$ :

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**,  
given the following  
domain / co-domain pairs?

Yes

No

*a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )

*b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$ : **Non-Injective!** (same example works)

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**,  
given the following  
domain / co-domain pairs?

Yes

No

- a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )
- b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$ : **Non-Injective!** (same example works)
- c)*  $D = \mathbb{Z}, C = \text{SQUARES}$ :

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**,  
given the following  
domain / co-domain pairs?

Yes

No

- a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )
- b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$ : **Non-Injective!** (same example works)
- c)*  $D = \mathbb{Z}, C = \text{SQUARES}$ : **Non-Injective!** (same example works)

# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**,  
given the following  
domain / co-domain pairs?

Yes

No

- a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )
- b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$ : **Non-Injective!** (same example works)
- c)*  $D = \mathbb{Z}, C = \text{SQUARES}$ : **Non-Injective!** (same example works)

- Can this function **ever** be injective?



# Quiz on injectivity

- Is  $f(x) = x^2$  **injective**, given the following domain / co-domain pairs?

Yes

No

- a)*  $D = \mathbb{R}, C = \mathbb{R}$ : **Non-Injective!** (e.g  $(-3)^2 = 3^2 = 9$ )
- b)*  $D = \mathbb{R}, C = \mathbb{R}^{\geq 0}$ : **Non-Injective!** (same example works)
- c)*  $D = \mathbb{Z}, C = SQUARES$ : **Non-Injective!** (same example works)

- Can this function **ever** be injective?
  - Yes. Pick  $D = \mathbb{N}, C = SQUARES$

# Making functions onto or 1-1

- To make a function **onto**, we need to make the **co-domain smaller**.
- To make a function **1-1**, we need to make the **domain smaller**.

# Bijjective functions

- A function  $f: X \mapsto Y$  is called **bijjective (or a bijection, or a 1-1 correspondence)** iff it is **both** **surjective** and **injective**.
  - We will try to avoid using the term “**1-1 correspondence**” (your book uses it) since it can confuse us with the notion of an injective (or 1-1) **function**.

# Quiz on bijections

# Quiz on bijections

- Are the following functions **bijections**?  
*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

1.  $f(x) = |x|, x \in \mathbb{R}$

Yes

No

# Quiz on bijections


- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

- $f(x) = |x|, x \in \mathbb{R}$  **No**



**Non-injective!**

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

1.  $f(x) = |x|, x \in \mathbb{R}$

2.  $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$

Yes

No



# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

- $f(x) = |x|, x \in \mathbb{R}$  **No**
- $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**

Yes

No

!!!!

For  $a = 0$ , the graph of the function fails the “horizontal line test”!

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

1.  $f(x) = |x|, x \in \mathbb{R}$  **No**
2.  $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**
3.  $g(x) = a \cdot x^2, a, x \in \mathbb{R}, a > 0$

# Quiz on bijections


- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

- $f(x) = |x|, x \in \mathbb{R}$  **No**
- $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**
- $g(x) = a \cdot x^2, a, x \in \mathbb{R}, a > 0$  **No**



**Non-injective!**

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

1.  $f(x) = |x|, x \in \mathbb{R}$  **No**
2.  $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**
3.  $g(x) = a \cdot x^2, a, x \in \mathbb{R}, a > 0$  **No**
4.  $h(n) = 4n - 1, n \in \mathbb{Z}$

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

1.  $f(x) = |x|, x \in \mathbb{R}$  **No**
2.  $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**
3.  $g(x) = a \cdot x^2, a, x \in \mathbb{R}, a > 0$  **No**
4.  $h(n) = 4n - 1, n \in \mathbb{Z}$  **No**

**Non-surjective!** Set  $h(n) = y$  and solve for  $n$ :

$$4n - 1 = y \Rightarrow n = \frac{y + 1}{4}$$

There are infinitely many choices of  $y$  for which  $n \notin \mathbb{Z}$ !

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

1.  $f(x) = |x|, x \in \mathbb{R}$  **No**
2.  $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**
3.  $g(x) = a \cdot x^2, a, x \in \mathbb{R}, a > 0$  **No**
4.  $h(n) = 4n - 1, n \in \mathbb{Z}$  **No**
5.  $h(x) = 4x - 1, x \in \mathbb{R}$

# Quiz on bijections

- Are the following functions **bijections**?

*(In all examples,  $C = \mathbb{R}$ )*

Yes

No

1.  $f(x) = |x|, x \in \mathbb{R}$  **No**
2.  $f(x) = a \cdot x + b, (\forall a, x, b \in \mathbb{R})$  **No**
3.  $g(x) = a \cdot x^2, a, x \in \mathbb{R}, a > 0$  **No**
4.  $h(n) = 4n - 1, n \in \mathbb{Z}$  **No**
5.  $h(x) = 4x - 1, x \in \mathbb{R}$  **Yes**

Surjective and injective! Surjective, since, if we set  $h(n) = y$  and solve for  $n$ :

$$4n - 1 = y \Rightarrow n = \frac{y + 1}{4}$$

For every real  $y$ , there's always a **real** solution  $n$ . **Injective**, since it's of the form of (2) with  $a \neq 0$ .

# Some special functions

- Are the following Java functions injective, surjective, bijective?
  - `Comparable.compareTo()`
  - `Object.hashCode()`