

Statistics of Particle Concentration in Free-Surface Turbulence

Jason Larkin

Department Mechanical Engineering and Materials Science, University of Pittsburgh.

Mahesh M. Bandi

Center for Nonlinear Studies and Condensed Matter & Thermal Physics Division,
Los Alamos National Lab.

Walter Goldberg

Department of Physics & Astronomy, University of Pittsburgh



Alain Pumir

Institut Nonline'aire de Nice, CNRS, France

Outline

- I. Experiment Description/Overview
- II. Coarse-Graining Procedure
- III. Scale-Free Behavior
- IV. Probability Distribution
- V. Review
- VI. Questions

I. Experiment Overview



- Tank: 1 m x 1 m x 0.3 m driven by 8 hp pump through “sprinkler” system.
- Real particles 50 micron size and highly bouyant (sg 0.25), low Stokes number $St \simeq 0.1$
- Want to study particle coagulation due to surface effect, not inertia.
- Instantaneous velocity fields measured by high-speed camera (9 cm field of view) used to evolve virtual particles.

$$C = \frac{\langle (\nabla \cdot \mathbf{v})^2 \rangle}{\langle (\partial_x v_x)^2 + (\partial_x v_y)^2 + (\partial_y v_x)^2 + (\partial_y v_y)^2 \rangle} \simeq 0.5$$

II. Coarse-Graining Procedure

Coarse-Grained Concentration:


All analysis done in steady-state!

$$n_r = \frac{N_r}{\langle N_r \rangle} \quad \eta = \left(\frac{\nu^3}{\varepsilon} \right)^{1/4} = 0.02 cm$$


$$r = s / \eta \quad \langle n_r \rangle = \left\langle \frac{N_r}{\langle N_r \rangle} \right\rangle = 1$$

S


x




y




III. Scale-Free Distribution



$$\langle n_r^2 \rangle = 1$$



$$\langle n_r^2 \rangle = 4$$



$$\langle n_r^2 \rangle = 16$$

$$\langle n_r^2 \rangle \propto r^{-\alpha_2}$$

$$\text{Re}_\lambda = 169$$


Point: $\alpha_2 = 2$

Line: $\alpha_2 = 1$


Surface: $\alpha_2 = 0$

-For fractal distributions:

$$\langle n_r^m \rangle \propto r^{-\alpha_m}$$



IV. Concentration PDF




-For power-law distributions:

$$P(n_r) \propto n_r^{-\beta}$$

-Cumulative PDF:

$$\Pi_c(n_r) = \int_{n_r}^{\infty} P(n_r) dn_r \propto n_r^{-\beta+1}$$

Can this be explained by theory of compressible clustering?



V. Review

- Floaters coagulate into strings.
- Effect not due to surface waves or inertia.
- Distribution is scale dependent.

VI. Question?

Funding was provided by the US National Science Foundation grant # DMR-0604477. This work was partially carried out under the auspices of the National Nuclear Security Administration of the U.S. Department of Energy at Los Alamos National Laboratory under Contract No. DE-AC52-06NA25396.