



Review article

Multidisciplinary treatment of lung cancer in older patients: A review



Marco Radovic^a, Ravindran Kanesvaran^b, Achim Rittmeyer^c, Martin Früh^{a,d}, Fabrizio Minervini^{a,e}, Markus Glatzer^a, Paul Martin Putora^{a,*}

^a Dept of Radiation Oncology, Kantonsspital St Gallen, St Gallen, Switzerland

^b National Cancer Centre Singapore, Singapore, Singapore

^c Dept of Thoracic Oncology, Lungenfachklinik Immenhausen, Immenhausen, Germany

^d Dept of Oncology, Haematology, Kantonsspital St Gallen, St Gallen, Switzerland

^e Dept of Thoracic Surgery, Kantonsspital St Gallen, St Gallen, Switzerland

ARTICLE INFO

Article history:

Received 2 May 2018

Received in revised form 9 August 2018

Accepted 10 September 2018

Available online 3 October 2018

ABSTRACT

Lung cancer is the leading cause of cancer death worldwide. Older patients represent approximately half of the patient population and optimal management of these patients is challenging. In early-stage non-small cell lung cancer (NSCLC), lobectomy should be considered in fit older patients. For unfit patients, stereotactic body radiotherapy (SBRT) represents a good alternative. While data on the benefit and risk of concurrent chemo-radiotherapy (cCRT) in older patients with locally advanced NSCLC is conflicting, age alone should not preclude cCRT. Multidisciplinary collaboration is essential for appropriate patient selection. In limited disease small cell lung cancer (SCLC), older patients appear to benefit similarly from standard treatment compared to their younger counterparts, however, with a higher risk of toxicity. Appropriately selected older patients with lung cancer seem to derive as much benefit from active oncological treatment as their younger counterparts. Geriatric screening tests and comprehensive geriatric assessments (CGA) can be helpful when choosing between treatment strategies. Older patients are at risk for under-treatment; this should be avoided by proper selection and multidisciplinary management. This review outlines the management of lung cancer in older patients.

© 2018 Elsevier Ltd. All rights reserved.

Contents

1. Introduction	405
2. Non-Small Cell Lung Cancer	406
2.1. Early-Stage NSCLC: AJCC Stage I and II	406
2.2. Locally Advanced NSCLC (AJCC Stage III)	406
2.3. Adjuvant Systemic Treatment	406
2.4. Treatment for Metastatic NSCLC	407
2.5. NSCLC Geriatric Assessment	407
3. Small Cell Lung Cancer (SCLC)	407
3.1. SCLC Limited Disease	407
3.2. LD SCLC- Prophylactic Cranial Irradiation (PCI)	408
3.3. Extensive Disease SCLC	408
4. Summary/Conclusion	408
Author Contributions	408
References	408

1. Introduction

Lung cancer is the leading cause of cancer death worldwide. Each year, more people die of lung cancer than of colon, breast, and prostate

* Corresponding author at: Dept. of Radiation Oncology, Kantonsspital St Gallen, 9007 St. Gallen, Switzerland.
E-mail address: paul.putora@kssg.ch (P.M. Putora).

cancers combined [1]. Non-small cell lung cancer (NSCLC) is the most common histology (85%) [2]. Depending on tumour stage, surgery, radiotherapy, and systemic therapy alone or in combination represent standard treatment options [3,4]. While older patients (>70 years of age) represent approximately half of the patient population [5], this subgroup is under-represented in clinical trials where the median age is typically around 60 years [6].

Older patients do not tolerate treatments as well as their younger counterparts [7,8], which may be caused by higher rates of comorbidities as well as decreased regenerative capacity and organ function [9]. Providing optimal treatment for older patients is challenging. Geriatric screening tests, comprehensive geriatric assessments (CGA), and guideline-concordant treatment are tools to optimize lung cancer management in older patients [10–14]. Less than half of older patients receive guideline-concordant treatment according to a pooled analysis of SEER-data. Yet, survival was increased in the patients who received guideline concordant care [15]. For the purpose of this review we have followed the International Society of Geriatric Oncology's (SIOG's) definition of older patients as patients 70 years of age and older.

2. Non-Small Cell Lung Cancer

2.1. Early-Stage NSCLC: AJCC Stage I and II

According to a meta-analysis, standard treatment of stage IA NSCLC consists of a lobectomy [16]. It is associated with the best long-term outcomes in fit older patients in an unadjusted survival analysis [17]. However, a retrospective analysis suggested that sub-lobar resections achieve similar survival rates when compared with lobectomy in older patients with stage I NSCLC with tumours of <2 cm diameter and with low Forced Expiratory Pressure in one Second (FEV1) [18]. However, this could not be confirmed in a larger SEER database analysis [17].

When surgery is considered too risky in older patients, e.g. due to poor pulmonary function, stereotactic body radiotherapy (SBRT) represents a good alternative. SBRT should be considered in patients with advanced age with multiple comorbidities which preclude surgical resection, as the tolerability appears to be better compared to surgery [17,19–21]. According to cohort studies (propensity score matched/matched pairs), SBRT has demonstrated favourable results, particularly when compared to conventional fractionated radiotherapy, observation only or sub-lobar resection [17,19–21]. However, these results were not confirmed by a systematic review and meta-analysis of sixteen studies containing 11,540 patients where a sub-lobar resection was associated with significantly improved overall survival compared to irradiated patients ($p < 0.05$) [22,23]. While newer techniques are being investigated, such as protons and carbon-ions [24], these have not found their way into clinical practice yet. Three randomized trials of SBRT vs. resection in patients with operable stage I NSCLC (STARS [NCT00840749], ROSEL [NCT00687986], and ACOSOG Z4099 trial [NCT01336894] [22]) closed early due to poor accrual. While a pooled analysis [25] of the first two studies demonstrated a favourable trend in survival towards SBRT, results should be interpreted with caution due to the small sample sizes of the trials. Retrospectively, frailty is associated with lower overall survival in older patients treated with SBRT but should not be viewed as a contraindication as disease free survival and overall survival were better compared to no active treatment [26]. According to a registry-based study, SBRT could be applied safely even in patients over the age of 80 years old [27]. Recent retrospective data comparing 30- and 90-day mortality of surgery and SBRT showed differences in favour of SBRT, especially in patients older than 71 years [28].

Although, active treatment confers similar survival advantages in both younger and older patients, older patients are still at risk of under-treatment [29]. Survival after radical treatment for early-stage NSCLC depends on factors such as tumour stage, performance status, gender, and radiotherapy (RT) dose, but not on age [30]. SBRT introduction and implementation has led to a higher proportion of

older patients receiving potentially curative treatment, leading to improved overall survival [31]. The introduction of multi-disciplinary teams and tumour boards can increase the proportion of actively treated older patients [32].

When feasible, surgery followed by chemotherapy can be considered standard treatment in stage II NSCLC. However, there is no universally accepted standard for older patients with stage II disease, especially when surgery is not reasonably possible. While data for this subset is limited, treatment would typically involve radiotherapy (conventionally fractionated, hypofractionated, or stereotactic radiotherapy), often in combination with chemotherapy (for N+) when possible. Adjuvant immunotherapy may also be considered for selected patients.

2.2. Locally Advanced NSCLC (AJCC Stage III)

Concurrent chemo-radiotherapy (cCRT) is the standard treatment for many patients with locally advanced NSCLC [4]; its use is however not universally implemented for older patients.

Patients included in randomized controlled trials and represented in meta-analyses establishing cCRT were mostly under 70 years old [33–35].

A randomized phase III study was conducted to assess whether cCRT with carboplatin resulted in longer survival than radiotherapy alone in patients older than 70 years [36]. The median overall survival (OS) for cCRT and RT alone was 22.4 and 16.9 months ($p = 0.0179$), progression-free survival (PFS) was significantly longer with cCRT (8.9 vs. 6.8 months, $p = 0.009$). These results were confirmed by a recently published meta-analysis [37]. Studies addressing cCRT in older patients concluded that grade 3–4 haematological toxicity and grade 3 infections were increased with cCRT, but lung toxicity was not increased [36,37].

While the above-mentioned studies show the benefit of cCRT in older patients, pooled analysis of individual patient data on cCRT for Stage III NSCLC in older vs. younger patients (cut-off 65 years) demonstrated that older patients experienced shorter OS, more toxicity, more treatment discontinuation, and a higher rate of death during treatment [7]. Recent comparative effectiveness [38] and phase-II-data suggest that simultaneous treatment could cause shorter overall survival in older patients [NCT01166204] [39]. Therefore, older patients should be selected carefully for a cCRT approach [40]. Sequential chemoradiation (CRT) could be a viable alternative to cCRT, although overall survival has shown to be inferior in a meta-analysis [35].

While data on the benefit and risk of cCRT in older patients are conflicting, age alone should not preclude cCRT, and a multidisciplinary team could help to select appropriate patients. Evidence shows that treatment, as it is applied to younger patients, could be used for fit older patients. However, there is insufficient evidence that this is an appropriate approach for unfit older patients. Recently, the Pacific trial showed a longer progression free survival in patients with NSCLC stage III treated with durvalumab after cCRT compared to placebo. There was no age restriction in this study, and the median age was 64 years (range 23–90 years) but age-related results showed that the experimental arm may not be as efficacious in older adults. Older patients were not analysed separately [41].

2.3. Adjuvant Systemic Treatment

Adjuvant systemic therapy significantly improves survival in patients with NSCLC [42]. Age alone should not disqualify an older patient from adjuvant chemotherapy in NSCLC, although data demonstrating a survival benefit is missing for patients >75 years of age [43]. The decision to treat should include the expected benefit and the increased risk of serious adverse events [44,45]. In the absence of specific trials for older patients, we must rely on retrospective data and subgroup analyses. Post hoc subgroup analyses inherently lack power due to small numbers of older patients included. One such small subgroup analysis of a randomized trial even suggested that

adjuvant cisplatin-containing chemotherapy could be harmful for patients 75 years or older [46].

Another retrospective analysis was performed on populations which included individual patient data of 4584 patients randomized in cisplatin based adjuvant trials [43]. Impact on overall survival, treatment delivery, and toxicity were investigated according to age. Older patients (≥ 70 years old) made up 9% of the patients, and were compared to both the young group (< 65 years old, 71%) and an intermediate group (65–69 years old, 20%). The result of this analysis showed that selected older patients had similar survival benefits and toxicity rates including toxic death rates from chemotherapy compared to their younger counterparts. Separate SEER-Medicare database analyses revealed that up to 83% of older patients were actually treated with carboplatin instead of cisplatin. These analyses did not show significant survival differences between cisplatin and carboplatin but slightly less side effects in carboplatin treated patients [47,48].

Potentially, better tolerated new systemic treatment options such as immunotherapy are of particular interest in older patients [49]. Various studies are currently testing whether checkpoint-inhibitors or other targeted therapies could improve clinical outcomes in the adjuvant as well as in the neoadjuvant setting [50]. One recent study showed a DFS benefit for EGFR mutant patients who were given gefitinib targeted therapy compared to chemotherapy [51]. Like in the metastatic setting, these drugs have a better toxicity profile and are better tolerated in older patients.

2.4. Treatment for Metastatic NSCLC

As in younger patients, platinum-based doublet chemotherapy is associated with a survival benefit compared with vinorelbine or gemcitabine monotherapy in older patients with advanced NSCLC [52]. However, a recent Asian study of docetaxel plus cisplatin versus docetaxel monotherapy failed to demonstrate any survival advantage as first-line chemotherapy for advanced NSCLC in older patients [53]. Further information about treatment outcomes of cisplatin-based chemotherapy in older patients with lung cancer are available from subgroup analyses of patients with metastatic disease who participated in large cooperative group studies, but again, likely represent a highly selected group of older patients [54–56]. These studies mostly showed similar treatment outcomes for older patients with similar or slightly increased toxicity rates in selected older patients who fulfilled the stringent inclusion criteria of clinical trials.

Responses to palliative radiotherapy in older patients are similar to those in younger patients [57]. However, the QUARTZ-trial comparing best supportive care with or without whole brain radiotherapy (WBRT) in patients with brain metastases from NSCLC only showed a survival benefit for irradiated patients younger than 60 years and Karnofsky Performance Status (KPS) of $> 70\%$ in a subgroup analysis [58]. For patients at risk for neurocognitive disorders methods of minimizing WBRT associated neurotoxicity like the use of NMDA (*N*-methyl-D-aspartate) receptor antagonist memantine and/or hippocampal-sparing radiation techniques may become standard options in the future [59].

Trials evaluating targeted therapy in patients with EGFR mutations or ALK translocations have constantly demonstrated better tolerability and quality of life compared to platinum-based chemotherapy. Therefore, these treatments should also be suitable for a more vulnerable population such as patients with poor performance status or older patients [60].

As immunotherapy with checkpoint inhibitors represents a relatively novel treatment option for patients with metastatic NSCLC, limited data is available on the safety and efficacy of these agents in older patients. While a recent trial suggested good tolerability [61], another study demonstrated increased toxicity and worse outcomes [62]. Specific trials for older patients are needed to further explore the role of immunotherapy in this population. Recent phase III trials have

demonstrated an overall survival advantage of immunotherapy as second line treatment compared to docetaxel or, most recently, as first line treatment in patients whose tumours express high levels of PDL-1 or if added to standard platinum-based chemotherapy. All these trials enrolled relatively young (median age ranging from 63 to 65 years) as well as exclusively fit patients (PS 0–1) [63–70]. Due to the underrepresentation of older patients in these trials, only limited conclusions can be drawn because the majority of these studies did not analyse older patients separately. In the IMPOWER 150 trial, only 9% ($n = 64$) were older than 75 years of age, which does not allow meaningful conclusions to be drawn [66]. In this study, the addition of immunotherapy to first line carboplatin/paclitaxel and bevacizumab appeared to be beneficial with regards to PFS benefit, both in younger and also in patients aged 75 or older (hazard ratio 0.78). In the absence of sufficient data on the older population in these trials, it has to be noted that with the addition of immunotherapy to chemotherapy, toxicity rates were higher and tolerability of such an approach in older patients needs further evaluation. On the other hand, in a European single-arm, phase 2 trial (CheckMate 171) testing nivolumab in previously treated patients with metastatic squamous NSCLC, the subgroup of patients aged 70 years or older ($n = 155$) had a 56% rate of treatment related adverse events (TRAEs) and only 16 (6%) discontinued due to TRAEs [61]. These results indicate that anti-PD1 monotherapy tolerability in patients aged 70 or older may be comparable to the overall population.

2.5. NSCLC Geriatric Assessment

Retrospective analyses have demonstrated that comprehensive geriatric assessments (CGA) in older patients in addition to neuropsychological evaluations and blood tests help in predicting survival [10] and chemotherapy associated toxicity [11]. CGAs can better predict the risks of chemotherapy toxicity than KPS [71]. In a prospective study, treatment allocation based on CGAs in older patients with advanced NSCLC failed to improve the treatment failure free survival (primary endpoint) or OS, but reduced treatment toxicity [72]. This can be an important surrogate marker of quality of life.

Although the use of CGA for treatment allocation is currently not standard, their use may facilitate an individualized approach. As CGAs are time consuming, screening tools such as the G8 or Vulnerable Elders Survey (VES)-13 may be suitable to select patients for a CGA [12]. These short geriatric screenings include multiple tools to evaluate the need for a full comprehensive assessment, an estimation of the ability for self-care, risk assessment regarding excess treatment related toxicity, and early mortality [13]. All older patients with lung cancer should have a CGA done if possible, and if resources are an issue, they should at least get a CGA screening test to select those requiring a full CGA [14].

3. Small Cell Lung Cancer (SCLC)

The incidence of SCLC is gradually decreasing as the prevalence of smoking is declining [73]. Almost 40% of all SCLC patients are over 70 years old and 10% are older than 80 years [74]. Without therapy, the overall survival is limited from weeks to a few months [3]. Five year overall survival (5y-OS) for lung cancer independent of histology is 18.1% [5], whereas 5y-OS for SCLC is only 6.6%. For limited disease SCLC (LD-SCLC), 5y-OS is 12.1%; and only 1.6% for extensive disease (ED-SCLC) [3]. About 70% will present with ED-SCLC at time of diagnosis [75].

3.1. SCLC Limited Disease

The standard treatment for LD-SCLC is cCRT [76,77]. Older patients appear to benefit similarly from treatment compared to their younger counterparts, but with increased toxicity [56]. According to a meta-analysis including older trials [76], no survival benefit was observed for patients over 65 years, but this subgroup analysis should be

interpreted with caution. There are no randomized phase III studies available comparing combined chemo-radiotherapy to chemotherapy alone in the older population with LD-SCLC [78], but retrospective data suggest that age does not have an impact on the efficacy of combined treatment in contrast to the mentioned meta-analysis [79]. For example, Corso et al. showed that CRT has better results compared to chemotherapy alone in older patients after stratification for comorbidities, nodal status, radiation once/twice daily or concomitant/sequential treatment [78]. Subset analyses of CRT treatment sequence showed a survival benefit with cCRT over sequential CRT. The subset analysis restricted the chemotherapy cohort to patients for whom CRT was explicitly recommended but not delivered in an effort to reduce selection bias. The cohort was reduced to 8.9% of the original cohort (335 patients), but a survival benefit for CRT persisted.

In the CONVERT trial, radiation once daily up to 66 Gy did not provide a survival benefit or more severe adverse events compared to radiation twice daily up to 45 Gy [80]. 15% of patients were >70 years old with a median age of 62 years, but there were fewer older patients in the twice-daily-group. A separate report on outcome in the older population from this study showed similar results for older patients [81]. A phase II trial demonstrated similar outcomes with less esophagitis in patients treated once daily to 70 Gy with 2 Gy daily compared with radiation twice daily [82], however with no specific focus on older populations.

A large retrospective cohort study (SEER-Medicare Database) of 10,428 patients aged 65 years and older who were diagnosed with SCLC between 1992 and 2001, showed that 67.1% of patients received chemotherapy, and 41.6% received etoposide with carboplatin or cisplatin with or without other agents. One third of older patients with SCLC never received chemotherapy, and one sixth were never referred to a medical oncologist. These results indicate that the use of chemotherapy is limited in this vulnerable population. Definitive conclusions regarding survival benefit are difficult due to the retrospective nature of this analysis [83]. Evidence from a retrospective analysis of data from two randomized trials suggests that response rates to chemotherapy in the older patients are comparable to younger patients [56]. Chemotherapy regimens used are the same as in younger patients [84]. Therefore, while standard treatment for limited disease small cell lung cancer consists of cCRT once or twice daily, older patients should be selected carefully for combined treatment.

3.2. LD SCLC- Prophylactic Cranial Irradiation (PCI)

PCI should be part of standard care in LD SCLC in case of complete or good partial remission after primary treatment. According to a meta-analysis, there was an overall survival benefit in patients receiving PCI (25% of study participants were 65 years old or older) [85]. Based on additional data from a SEER-database analysis, older patients, who received PCI, had better survival than patients who did not receive PCI [86]. While age alone should not rule out PCI, it should be used with caution in older patients as they are at higher risk for neurocognitive deterioration [3].

3.3. Extensive Disease SCLC

ED SCLC is highly responsive to chemotherapy, but median survival is less than one year [87,88]. First line treatment in older patients consists of Carboplatin/Etoposide [73,89] or Cisplatin/Etoposide [90]. A trial including patients up to 75 years of age showed doubling of the 1y OS with PCI, as well as reduced rates of brain metastases at one year if no routine MRI of the brain was performed before the start of chemotherapy [91]. PCI in older patients with SCLC results in a longer survival, according to a pooled analysis of four prospective trials [92]. However, in a Japanese trial, PCI did not result in longer overall survival compared with observation in patients with ED SCLC (median age 69 years, maximum age 86 years) [93]. Regarding PCI, risk

of overtreatment should be avoided [94] and PCI is currently controversially discussed for ED SCLC even for younger patients. As mentioned previously, numerous approaches are being investigated to preserve neurocognitive function in patients receiving WBRT, these methods may prove most valuable in subsets of patients with relatively good prognoses – i.e. in prophylactic cranial irradiation.

The use of thoracic RT (TRT) in ED SCLC remains controversial and patients have to be chosen carefully for this approach which slightly increases survival rates [95–97]. The only two prospective studies included patients with a median age of 60 and 63 years, respectively, <10% were 75 years old or older. Given the overall poor prognosis, the older population with ED-SCLC require an individualized approach [98] and multidisciplinary team in the decision making in order to avoid both, over- and under-treatment.

4. Summary/Conclusion

Older patients with lung cancer should be discussed in a multidisciplinary setting. Selected older patients with lung cancer seem to derive as much benefit from active oncological treatment as their younger counterparts. Under-treatment is a risk in this patient population and should be avoided. Patient selection is of great importance as many older patients have other medical issues that need to be taken into account before selecting the appropriate treatment. This remains a challenging task as data for this specific population is scarce. While newer agents are being investigated in the neoadjuvant, concurrent or adjuvant setting, we should not lose sight of the fact that a major impact on survival and quality of life in potentially curative stages is still achieved by established local treatments (i.e. radiotherapy, surgery). While emerging treatments may provide improved outcomes with limited toxicity, patient preference should remain at the centre of our attention, especially in an older population where tolerability and logistics may significantly impact patient satisfaction. Having a multidisciplinary team and performing CGAs could help select appropriate treatments with improved outcomes (including quality of life) and reduce toxicities. Effective treatment should not be withheld based on age alone.

Author Contributions

M. Putora and M. Radovic contributed to the study concept. M. Glatzer, F. Minervini, M. Früh, A. Rittmeyer, R. Kanesvaran, M. Radovic and M. Putora contributed to manuscript preparation, editing, and review. All authors have no conflicts of interest to declare.

References

- [1] Society AC. Key Statistics for Lung Cancer. Available from: <https://www.cancer.org/cancer/non-small-cell-lung-cancer/about/key-statistics.html>; 2018, Accessed date: 18 June 2008.
- [2] Walters S, et al. Lung cancer survival and stage at diagnosis in Australia, Canada, Denmark, Norway, Sweden and the UK: A population-based study, 2004–2007. *Thorax* 2013;68(6):551–64.
- [3] Glatzer M, et al. Treatment of limited disease small cell lung cancer: The multidisciplinary team. *Eur Respir J* 2017;50(2).
- [4] Glatzer M, et al. Radio(chemo)therapy in locally advanced nonsmall cell lung cancer. *Eur Respir Rev* 2016;25(139):65–70.
- [5] Institute, N.C. Cancer stat facts: Lung and bronchus Cancer. Available from: <https://seer.cancer.gov/statfacts/html/lungb.html>; 2018, Accessed date: 18 June 2008.
- [6] Hutchins LF, et al. Underrepresentation of patients 65 years of age or older in cancer-treatment trials. *N Engl J Med* 1999;341(27):2061–7.
- [7] Stinchcombe TE, et al. Pooled Analysis of Individual Patient Data on Concurrent Chemoradiotherapy for Stage III Non-Small-Cell Lung Cancer in Elderly Patients Compared With Younger Patients Who Participated in US National Cancer Institute Cooperative Group Studies. *J Clin Oncol* 2017;35:2885–92.
- [8] Sud S, et al. Chemotherapy in the oldest old: The feasibility of delivering cytotoxic therapy to patients 80 years old and older. *J Geriatr Oncol* 2015;6(5):395–400.
- [9] Balducci L, Extermann M. Cancer and aging. An evolving panorama. *Hematol Oncol Clin North Am* 2000;14(1):1–16.
- [10] Kanesvaran R, et al. Analysis of prognostic factors of comprehensive geriatric assessment and development of a clinical scoring system in elderly Asian patients with cancer. *J Clin Oncol* 2011;29(27):3620–7.

- [11] Hurria A, et al. Validation of a prediction tool for chemotherapy toxicity in older adults with cancer. *J Clin Oncol* 2016;34(20):2366–71.
- [12] Decoster L, et al. Screening tools for multidimensional health problems warranting a geriatric assessment in older cancer patients: An update on SIOG recommendations. *Ann Oncol* 2015;26(2):288–300.
- [13] Honecker F. Screening tools versus a comprehensive geriatric assessment in oncology. *Schweizer Zeitschrift für Onkologie* 2015;6–9.
- [14] Kanesvaran R, Roy Chowdhury A, Krishna L. Practice pearls in the management of lung cancer in the elderly. *J Geriatr Oncol* 2016;7(5):362–7.
- [15] Nadpara PA, et al. Guideline-concordant lung cancer care and associated health outcomes among elderly patients in the United States. *J Geriatr Oncol* 2015;6(2):101–10.
- [16] Liu Y, et al. Sublobectomy versus lobectomy for stage IA (T1a) non-small-cell lung cancer: Ameta-analysis study. *World J Surg Oncol* 2014;12:138.
- [17] Shirvani SM, et al. Comparative effectiveness of 5 treatment strategies for early-stage non-small cell lung cancer in the elderly. *Int J Radiat Oncol Biol Phys* 2012;84(5):1060–70.
- [18] Qiu C, et al. Sublobectomy versus lobectomy for stage I non-small cell lung cancer in the elderly. *Int J Surg* 2017;37:1–7.
- [19] Shirvani SM, et al. Lobectomy, sublobar resection, and stereotactic ablative radiotherapy for early-stage non-small cell lung cancers in the elderly. *JAMA Surg* 2014;149(12):1244–53.
- [20] Palma D, et al. Treatment of stage I NSCLC in elderly patients: Apopulation-based matched-pair comparison of stereotactic radiotherapy versus surgery. *Radiother Oncol* 2011;101(2):240–4.
- [21] Ezer N, et al. Outcomes after stereotactic body radiotherapy versus limited resection in older patients with Early-stage lung cancer. *J Thorac Oncol* 2015;10(8):1201–6.
- [22] Wang HH, et al. Sublobar resection is associated with improved outcomes over radiotherapy in the management of high-risk elderly patients with Stage I non-small cell lung cancer: A systematic review and meta-analysis. *Oncotarget* 2017;8(4):6033–42.
- [23] Fernando HC, Timmerman R. American College of Surgeons Oncology Group Z4099/ Radiation Therapy Oncology Group 1021: A randomized study of sublobar resection compared with stereotactic body radiotherapy for high-risk stage I non-small cell lung cancer. *J Thorac Cardiovasc Surg* 2012;144(3):S35–8.
- [24] Grutters JP, et al. Comparison of the effectiveness of radiotherapy with photons, protons and carbon-ions for non-small cell lung cancer: Ameta-analysis. *Radiother Oncol* 2010;95(1):32–40.
- [25] Chang JY, et al. Stereotactic ablative radiotherapy versus lobectomy for operable stage I non-small-cell lung cancer: a pooled analysis of two randomised trials. *Lancet Oncol* 2015;16(6):630–7.
- [26] Franco I, et al. Use of frailty to predict survival in elderly patients with early stage non-small-cell lung cancer treated with stereotactic body radiation therapy. *J Geriatr Oncol* 2018;9(2):130–7.
- [27] Kreinbrink P, et al. Lung stereotactic body radiation therapy (SBRT) for early-stage non-small cell lung cancer in the very elderly (>=80 years old): Extremely safe and effective. *J Geriatr Oncol* 2017;8(5):351–5.
- [28] Stokes WA, et al. Post-treatment mortality after surgery and stereotactic body radiotherapy for early-stage non-small-cell lung cancer. *J Clin Oncol* 2018;36(7):642–51.
- [29] Higton AM, Monach J, Congleton J. Investigation and management of lung cancer in older adults. *Lung Cancer* 2010;69(2):209–12.
- [30] Palma DA, et al. Stage I non-small cell lung cancer (NSCLC) in patients aged 75 years and older: Does age determine survival after radical treatment? *J Thorac Oncol* 2010;5(6):818–24.
- [31] Palma D, et al. Impact of introducing stereotactic lung radiotherapy for elderly patients with stage I non-small-cell lung cancer: Apopulation-based time-trend analysis. *J Clin Oncol* 2010;28(35):5153–9.
- [32] Price A, Kerr Gillian, Gregor A, Ironside J. The impact of multidisciplinary teams and site specialisation on the use of radiotherapy in elderly people with non-small cell lung cancer (NSCLC). *Radiother Oncol* 2002;64(Suppl. 1):80.
- [33] Schaake-Koning C, et al. Effects of concomitant cisplatin and radiotherapy on inoperable non-small-cell lung cancer. *N Engl J Med* 1992;326(8):524–30.
- [34] Auperin A, et al. Concomitant radio-chemotherapy based on platinum compounds in patients with locally advanced non-small cell lung cancer (NSCLC): Ameta-analysis of individual data from 1764 patients. *Ann Oncol* 2006;17(3):473–83.
- [35] Auperin A, et al. Meta-analysis of concomitant versus sequential radiochemotherapy in locally advanced non-small-cell lung cancer. *J Clin Oncol* 2010;28(13):2181–90.
- [36] Atagi S, et al. Thoracic radiotherapy with or without daily low-dose carboplatin in elderly patients with non-small-cell lung cancer: A randomised, controlled, phase 3 trial by the Japan Clinical Oncology Group (JCOG0301). *Lancet Oncol* 2012;13(7):671–8.
- [37] Dawe DE, et al. Chemoradiotherapy versus radiotherapy alone in elderly patients with stage III non-small cell lung cancer: A systematic review and meta-analysis. *Lung Cancer* 2016;99:180–5.
- [38] Miller ED, et al. The addition of chemotherapy to radiation therapy improves survival in elderly patients with stage III non-small cell lung cancer. *J Thorac Oncol* 2018;13(3):426–35.
- [39] Van Loon J. Elderly Non-Small Cell Lung Cancer Patients are Less Likely to Benefit from Treatment with Simultaneous Radio- and Chemotherapy. Barcelona, Spain: ESTRO; 2018.
- [40] De Ruysscher D, van Loon J. Radical radiotherapy for locally advanced non-small cell lung cancer: When should concurrent chemoradiotherapy not be used? *Clin Oncol (R Coll Radiol)* 2016;28(11):708–11.
- [41] Antonia SJ, et al. Durvalumab after chemoradiotherapy in stage III non-small-cell lung cancer. *N Engl J Med* 2017;377:1919–29.
- [42] Pignon JP, et al. Lung adjuvant cisplatin evaluation: A pooled analysis by the LACE Collaborative Group. *J Clin Oncol* 2008;26(21):3552–9.
- [43] Fruh M, et al. Pooled analysis of the effect of age on adjuvant cisplatin-based chemotherapy for completely resected non-small-cell lung cancer. *J Clin Oncol* 2008;26(21):3573–81.
- [44] Wisnivesky JP, et al. Survival and risk of adverse events in older patients receiving postoperative adjuvant chemotherapy for resected stages II–IIIA lung cancer: Observational cohort study. *BMJ* 2011;343:d4013.
- [45] Pallis AG, et al. Management of elderly patients with NSCLC: updated expert's opinion paper: EORTC Elderly Task Force, Lung Cancer Group and International Society for Geriatric Oncology. *Ann Oncol* 2014;25(7):1270–83.
- [46] Pepe C, et al. Adjuvant vinorelbine and cisplatin in elderly patients: National Cancer Institute of Canada and Intergroup Study JBR.10. *J Clin Oncol* 2007;25(12):1553–61.
- [47] Gu F, Strauss GM, Wisnivesky JP. Platinum-based adjuvant chemotherapy (ACT) in elderly patients with non-small cell lung cancer (NSCLC) in the SEER-Medicare database: Comparison between carboplatin- and cisplatin-based regimens. *J Clin Oncol* 2011;29(15_suppl):7014.
- [48] Gu F, et al. Carboplatin versus cisplatin-based adjuvant chemotherapy in elderly patients with stages IB, II, and IIIA non-small cell lung cancer in the community setting. *J Clin Oncol* 2013;31(15_suppl):7533.
- [49] Meoni G, et al. Medical treatment of advanced non-small cell lung cancer in elderly patients: A review of the role of chemotherapy and targeted agents. *J Geriatr Oncol* 2013;4(3):282–90.
- [50] Simon GR, Somaiah N. A tabulated summary of targeted and biologic therapies for non-small-cell lung cancer. *Clin Lung Cancer* 2014;15(1):21–51.
- [51] Zhong WZ, et al. Gefitinib versus vinorelbine plus cisplatin as adjuvant treatment for stage II–IIIA (N1–N2) EGFR-mutant NSCLC (ADJUVANT/CTONG1104): A randomised, open-label, phase 3 study. *Lancet Oncol* 2018;19(1):139–48.
- [52] Quoix E, et al. Carboplatin and weekly paclitaxel doublet chemotherapy compared with monotherapy in elderly patients with advanced non-small-cell lung cancer: IFCT-0501 randomised, phase 3 trial. *Lancet* 2011;378(9796):1079–88.
- [53] Abe T, et al. Randomized phase III trial comparing weekly docetaxel plus cisplatin versus docetaxel monotherapy every 3 weeks in elderly patients with advanced non-small-cell lung cancer: The intergroup trial JCOG0803/WJOG4307L. *J Clin Oncol* 2015;33(6):575–81.
- [54] Langer CJ, et al. Cisplatin-based therapy for elderly patients with advanced non-small-cell lung cancer: Implications of Eastern Cooperative Oncology Group 5592, a randomized trial. *J Natl Cancer Inst* 2002;94(3):173–81.
- [55] Rocha Lima CM, et al. Therapy choices among older patients with lung carcinoma: An evaluation of two trials of the Cancer and Leukemia Group B. *Cancer* 2002;94(1):181–7.
- [56] Siu LL, et al. Influence of age on the treatment of limited-stage small-cell lung cancer. *J Clin Oncol* 1996;14(3):821–8.
- [57] Patterson CJ, et al. Retrospective study of radiotherapy for lung cancer in patients aged 75 years and over. *Age Ageing* 1998;27(4):515–8.
- [58] Mulvenna P, et al. Dexamethasone and supportive care with or without whole brain radiotherapy in treating patients with non-small cell lung cancer with brain metastases unsuitable for resection or stereotactic radiotherapy (QUARTZ): Results from a phase 3, non-inferiority, randomised trial. *Lancet* 2016;388(10055):2004–14.
- [59] Slade AN, Stanic S. The impact of RTOG 0614 and RTOG 0933 trials in routine clinical practice: The US Survey of Utilization of Memantine and IMRT planning for hippocampus sparing in patients receiving whole brain radiotherapy for brain metastases. *Contemp Clin Trials* 2016;47:74–7.
- [60] Inoue A, et al. First-line gefitinib for patients with advanced non-small-cell lung cancer harboring epidermal growth factor receptor mutations without indication for chemotherapy. *J Clin Oncol* 2009;27(9):1394–400.
- [61] Popat S. 1303PD - Nivolumab in previously treated patients with metastatic squamous NSCLC: Results of a European single-arm, phase 2 trial (CheckMate 171) including patients aged 70 years and with poor performance status in ESMO. *Ann Oncol* 2017;28 (mdx380.006 [Madrid, Spain]).
- [62] Gomes F. MA 10.07 - elderly lung cancer patients on immunotherapy: Preliminary results from the ELDERs study; 2017.
- [63] Lopes G. Pembrolizumab (pembro) versus platinum-based chemotherapy (chemo) as first-line therapy for advanced/metastatic NSCLC with a PD-L1 tumor proportion score (TPS) ≥ 1%: Open-label, phase 3 KEYNOTE-042 study. (abstract LBA4). 2018 ASCO Annual Meeting; 2018 [Chicago].
- [64] Brahmer J, et al. Nivolumab versus docetaxel in advanced squamous-cell non-small-cell lung cancer. *N Engl J Med* 2015;373(2):123–35.
- [65] Reck M, et al. Pembrolizumab versus chemotherapy for PD-L1-positive non-small-cell lung cancer. *N Engl J Med* 2016;375(19):1823–33.
- [66] Socinski MA, et al. Atezolizumab for first-line treatment of metastatic nonsquamous NSCLC. *N Engl J Med* 2018;378(24):2288–301.
- [67] Gandhi L, et al. Pembrolizumab plus chemotherapy in metastatic non-small-cell lung cancer. *N Engl J Med* 2018;378(22):2078–92.
- [68] Herbst RS, et al. Pembrolizumab versus docetaxel for previously treated, PD-L1-positive, advanced non-small-cell lung cancer (KEYNOTE-010): A randomised controlled trial. *Lancet* 2016;387(10027):1540–50.
- [69] Rittmeyer A, et al. Atezolizumab versus docetaxel in patients with previously treated non-small-cell lung cancer (OAK): A phase 3, open-label, multicentre randomised controlled trial. *Lancet* 2017;389(10066):255–65.
- [70] Borghaei H, et al. Nivolumab versus docetaxel in advanced nonsquamous non-small-cell lung cancer. *N Engl J Med* 2015;373(17):1627–39.
- [71] Nie X, et al. Predicting chemotherapy toxicity in older adults with lung cancer. *J Geriatr Oncol* 2013;4(4):334–9.

- [72] Corre R, et al. Use of a comprehensive geriatric assessment for the management of elderly patients with advanced non-small-cell lung cancer: The phase III randomized ESOGIA-GFPC-GECP08-02 study. *J Clin Oncol* 2016;34(13):1476–83.
- [73] Kalemkerian GP, et al. Small cell lung cancer. *J Natl Compr Canc Netw* 2013;11(1):78–98.
- [74] Owonikoko TK, et al. Lung cancer in elderly patients: An analysis of the surveillance, epidemiology, and end results database. *J Clin Oncol* 2007;25(35):5570–7.
- [75] Fruh M, et al. Small-cell lung cancer (SCLC): ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol* 2013;24(Suppl. 6):vi99–105.
- [76] Pignon JP, et al. A meta-analysis of thoracic radiotherapy for small-cell lung cancer. *N Engl J Med* 1992;327(23):1618–24.
- [77] Warde P, Payne D. Does thoracic irradiation improve survival and local control in limited-stage-small-cell carcinoma of the lung? A meta-analysis. *J Clin Oncol* 1992;10(6):890–5.
- [78] Corso CD, et al. Role of chemoradiotherapy in elderly patients with limited-stage-small-cell lung cancer. *J Clin Oncol* 2015;33(36):4240–6.
- [79] Gridelli C, et al. Treatment of limited-stage small cell lung cancer in the elderly, chemotherapy vs. sequential chemoradiotherapy vs. concurrent chemoradiotherapy: That's the question. *Transl Lung Cancer Res* 2016;5(2):150–4.
- [80] Faivre-Finn C, et al. Concurrent once-daily versus twice-daily chemoradiotherapy in patients with limited-stage-small-cell lung cancer (CONVERT): An open-label, phase 3, randomised, superiority trial. *Lancet Oncol* 2017;18(8):1116–25.
- [81] Christodoulou M. Compliance and outcome of elderly patients treated in the concurrent once-daily versus twice-daily Radiotherapy (CONVERT) Trial; 2016.
- [82] Bogart JA, et al. 70Gy thoracic radiotherapy is feasible concurrent with chemotherapy for limited-stage-small-cell lung cancer: Analysis of Cancer and Leukemia Group B study 39808. *Int J Radiat Oncol Biol Phys* 2004;59(2):460–8.
- [83] Caprario LC, Kent DM, Strauss GM. Effects of chemotherapy on survival of elderly patients with small-cell lung cancer: Analysis of the SEER-medicare database. *J Thorac Oncol* 2013;8(10):1272–81.
- [84] Rossi A, et al. Treatment of small cell lung cancer in the elderly. *Oncologist* 2005;10(6):399–411.
- [85] Auperin A, et al. Prophylactic cranial irradiation for patients with small-cell lung cancer in complete remission. Prophylactic Cranial Irradiation Overview Collaborative Group. *N Engl J Med* 1999;341(7):476–84.
- [86] Eaton BR, et al. Effect of prophylactic cranial irradiation on survival in elderly patients with limited-stage small cell lung cancer. *Cancer* 2013;119(21):3753–60.
- [87] Johnson BE, Janne PA. Basic treatment considerations using chemotherapy for patients with small cell lung cancer. *Hematol Oncol Clin North Am* 2004;18(2):309–22.
- [88] Demedts IK, Vermaelen KY, van Meerbeeck JP. Treatment of extensive-stage small cell lung carcinoma: Current status and future prospects. *Eur Respir J* 2010;35(1):202–15.
- [89] Okamoto H, et al. Randomised phase III trial of carboplatin plus etoposide vs split doses of cisplatin plus etoposide in elderly or poor-risk patients with extensive disease small-cell lung cancer: JCOG 9702. *Br J Cancer* 2007;97(2):162–9.
- [90] Hanna N, et al. Randomized phase III trial comparing irinotecan/cisplatin with etoposide/cisplatin in patients with previously untreated extensive-stage disease small-cell lung cancer. *J Clin Oncol* 2006;24(13):2038–43.
- [91] Slotman B, et al. Prophylactic cranial irradiation in extensive small-cell lung cancer. *N Engl J Med* 2007;357(7):664–72.
- [92] Rule WG, et al. Prophylactic cranial irradiation in elderly patients with small cell lung cancer: Findings from a North Central Cancer Treatment Group pooled analysis. *J Geriatr Oncol* 2015;6(2):119–26.
- [93] Takahashi T, et al. Prophylactic cranial irradiation versus observation in patients with extensive-disease-small-cell lung cancer: A multicentre, randomised, open-label, phase 3 trial. *Lancet Oncol* 2017;18(5):663–71.
- [94] Urbanic JJ. The tippy barstool of prophylactic cranial irradiation. *J Oncol Pract* 2017;13(11):742–3.
- [95] Slotman BJ, et al. Use of thoracic radiotherapy for extensive stage small-cell lung cancer: A phase 3 randomised controlled trial. *Lancet* 2015;385(9962):36–42.
- [96] Jeremic B, et al. Role of radiation therapy in the combined-modality treatment of patients with extensive disease small-cell lung cancer: A randomized study. *J Clin Oncol* 1999;17(7):2092–9.
- [97] Palma DA, et al. Thoracic Radiotherapy for Extensive Stage Small-Cell Lung Cancer: A Meta-Analysis. *Clin Lung Cancer* 2016;17(4):239–44.
- [98] Montella TC, et al. Evaluation of elderly patients with extended disease small-cell lung cancer. *J Clin Oncol* 2013;31(15_suppl):e18515.