



## MYELOYDYSPLASTIC SYNDROMES

# MDS overlap disorders and diagnostic boundaries

Tiffany N. Tanaka and Rafael Bejar

Division of Hematology and Oncology, Moores Cancer Center, University of California, San Diego, La Jolla, CA

**Myelodysplastic syndromes (MDS) are clonal diseases defined by clinical, morphologic, and genetic features often shared by related myeloid disorders. The diagnostic boundaries between these diseases can be arbitrary and not necessarily reflective of underlying disease biology or outcomes. In practice, measures that distinguish MDS from related disorders may be difficult to quantify and can vary as disease progression occurs. Patients may harbor findings that are not consistent with a single diagnostic category. Several overlap disorders have been formally described, such as the myelodysplastic/myeloproliferative neoplasms (MDS/MPNs). These disorders are characterized by hematopoietic dysplasia with increased proliferation of monocytes, neutrophils, or platelets. They may have mutational profiles that distinguish them from the disorders they resemble and reflect important differences in pathophysiology. MDS also shares diagnostic borders with other diseases. For**

**example, aplastic anemia and hypoplastic MDS can be difficult to distinguish in patients with pancytopenia and bone marrow hypocellularity. Genetic features may help in this regard, because they can identify differences in prognosis and risk of progression. The boundary between MDS and secondary acute myeloid leukemia (sAML) is arbitrarily defined and has been redefined over the years. Genetic studies have demonstrated that sAML clones can precede clinical progression from MDS by many months, suggesting that MDS with excess blasts could be viewed as an overlap between a dysplastic bone marrow failure syndrome and an oligoblastic leukemia. This review will describe the diagnostic boundaries between MDS, MDS/MPNs, sAML, clonal hematopoiesis of indeterminate potential, clonal cytopenia of undetermined significance, and aplastic anemia and how genetic approaches may help to better define them. (*Blood*. 2019;133(10):1086-1095)**

## Introduction

Myelodysplastic syndromes (MDS) are clonal hematopoietic disorders that typically present with features indicative of bone marrow failure, including inefficient hematopoiesis, morphologic dysplasia, and cytopenias of the peripheral blood. This clinical phenotype is nonspecific and can be a consequence of a variety of benign or malignant conditions. MDS and its mimics show an increased incidence with age, often making it challenging to arrive at the appropriate diagnosis. This is exacerbated by the fact that the apparently well-defined diagnostic boundaries between MDS and related conditions can, in practice, be much more vague and difficult to characterize. This can occur at initial presentation, when a patient with MDS-like features might also have evidence of hypoplasia or a myeloproliferative neoplasm (MPN), and over time, because MDS can evolve into another diagnosis, such as a secondary acute myeloid leukemia (sAML). Recent advances in our understanding about the genetics of MDS and its diagnostic neighbors may help to sharpen their boundaries or, ultimately, redefine them altogether.

Correctly diagnosing patients with MDS overlap syndromes can have important clinical implications. Often, the prognosis associated with an overlap syndrome is distinct from the individual disorders that they resemble. This is driven, in part, by differences

in their genetic profiles and pathobiology. Consequently, overlap disorders may be amenable to different therapeutic options and can harbor unique molecular vulnerabilities. Although genetics can aid in the diagnosis of overlap disorders, somatic mutations rarely define them independent of the clinical context in which they are found. Other factors, including patient characteristics, epigenetic alterations, and microenvironmental interactions, such as inflammation and adaptive immune responses, help to shape the disease phenotype. Together, these characteristics can help to establish an accurate diagnosis in cases with overlapping features.

This review will focus on those diagnostic categories that have features of MDS combined with elements of other disorders in the context of our greater understanding about their underlying molecular genetics. This includes the individual MDS/MPN overlap disorders recognized in the World Health Organization (WHO) classification of myeloid neoplasms. We will also examine the diagnostic boundaries between MDS and two related conditions, aplastic anemia (AA) and sAML, disorders that can lead to, or arise from, MDS respectively. Finally, we explore the diagnostic boundaries between lower-risk MDS, clonal hematopoiesis of indeterminate potential (CHIP), and idiopathic cytopenias of undetermined significance (ICUS), a substantial fraction of which harbor somatic mutations typical of MDS.

## The MDS/MPN overlap disorders

MDS/MPN overlap disorders are considered distinct from MDS and MPN, according to the WHO classification scheme. They include diagnoses with very different clinical manifestations, underlying genetics, and overall prognosis. Their shared features can include cellular dysplasia or cytopenias, in addition to an elevation in  $\geq 1$  blood cell count. At the molecular level, MDS/MPN disorders are more likely to carry gene mutations associated with the activation of growth factor signaling pathways in conjunction with mutations in epigenetic regulators or splicing factors associated with morphologic dysplasia.

Chronic myelomonocytic leukemia (CMML) is the most common of the MDS/MPN overlap diseases, even though its prevalence is estimated to be only  $\sim 10\%$  of that for MDS. CMML is defined by the presence of monocytosis in addition to  $\geq 1$  cytopenia (typically anemia) and bone marrow findings that typically meet MDS diagnostic criteria. The monocytosis in CMML has to be relative ( $\geq 10\%$  of white blood cells) and absolute ( $\geq 1 \times 10^9/L$ ) and must persist for  $\geq 3$  months.<sup>1,2</sup> Criteria indicative of other myeloid neoplasms and alternative causes of monocytosis should be absent. Historically, the French–American–British classification scheme considered CMML to be a subtype of MDS. The subsequent WHO classification divided CMML into a “proliferative type,” with a total white blood cell (WBC) count  $\geq 13 \times 10^9/L$ , and a “dysplastic type,” with a WBC count below this threshold, to reflect clinical and genetic distinctions between these 2 subtypes.<sup>3,4</sup> In the most recent WHO classification, CMML is considered a separate entity from MDS and is classified into subtypes based on blood and bone marrow blast proportions and not on total WBC count (Table 1).

Despite what seems like an arbitrary numerical distinction between MDS and CMML, there is evidence that the underlying pathobiology in these disorders is quite different. At the cellular level, patients with CMML have a high percentage of classical monocytes that are CD14<sup>+</sup> and CD16<sup>+</sup>.<sup>5,6</sup> These cells show a hypersensitivity to growth factor stimulation with granulocyte-macrophage colony-stimulating factor that is not observed in MDS. At the genetic level, CMML patients also have distinct mutational profiles (Figure 2). Somatic mutations in several genes, including *TET2*, *ASXL1*, *SRSF2*, *EZH2*, *NRAS*, *KRAS*, and *CBL*, are significantly more common in patients with CMML and, therefore, are more likely to co-occur.<sup>7–9</sup> In fact, the triad of *TET2*, *ASXL1*, and *SRSF2* mutations is highly specific for CMML.<sup>10–12</sup> In contrast, mutations of *SF3B1* and *TP53* are observed less often than in MDS.

Clinically, patients with proliferative type CMML are enriched for RAS signaling pathway mutations and appear to have slightly greater disease-related risk than MDS patients with similar blast proportions.<sup>10,13</sup> The original and revised International Prognostic Scoring Systems only included a small fraction of patients with dysplastic CMML (WBC  $< 13 \times 10^9/L$ ), whereas proliferative CMML was excluded.<sup>14,15</sup> This has led to the development of several CMML-specific prognostic tools.<sup>10,16–19</sup> Where these models consider molecular abnormalities, mutations of *ASXL1* are universally identified as independent adverse prognostic markers.

Therapeutic approaches for CMML aim to improve symptoms related to peripheral cytopenias or blood count proliferation.

Similar to patients with MDS, hypomethylating agents may be considered for patients with CMML if poor prognostic factors or excess blasts are present. Response rates and benefit from treatment with hypomethylating agents appear to be comparable between patients with CMML and MDS.<sup>20,21</sup> Surprisingly, responding CMML patients can revert to a normal monocyte profile with improved blood counts, without demonstrating changes in clonal burden.<sup>22</sup> And unlike in MDS, DNA-methylation profiles predictive of hypomethylating agent response have been identified in CMML.<sup>23</sup> Allogeneic hematopoietic stem cell transplantation (HSCT) remains the only curative treatment for CMML and should be considered in younger patients with higher-risk CMML, although the increased use of reduced-intensity conditioning and alternative donor sources has allowed increased implementation of HSCT in older patients. Expert opinion, including a recent international panel, suggests treatment before HSCT, particularly when marrow blasts are  $> 10\%$  or other higher-risk features are present.<sup>24,25</sup>

The next most common MDS/MPN overlap disorder is MDS/MPN with ring sideroblasts and thrombocytosis (MDS/MPN-RS-T). This entity has very little resemblance to CMML, despite being in the same diagnostic category. Patients with MDS/MPN-RS-T meet criteria for MDS with ring sideroblasts  $\geq 15\%$  and must have a persistently elevated platelet count ( $\geq 450 \times 10^9/L$ ). Classical hotspot mutations of *SF3B1* are found in  $> 80\%$  of cases, resembling the rate of *SF3B1* mutation observed in MDS-RS patients with single-lineage dysplasia (MDS-RS-SLD), and are often the likely founder mutation based on variant frequency and occurrence as the sole abnormality in some patients.<sup>26</sup> MDS/MPN-RS-T patients also have a high rate of mutations in *JAK2* (50–70%), *CALR* (10–20%), and *MPL* (2–5%), comparable to the mutational spectrum observed in essential thrombocythemia (ET).<sup>27,28</sup> Mutations in several other genes may be present, including *TET2*, *DNMT3A*, *ASXL1*, and *SETBP1*, with the latter 2 being considered prognostically adverse.<sup>29</sup> The prognosis of patients with MDS/MPN-RS-T lies between that of patients with MDS-RS-SLD and ET; the leukemic transformation rate per 100 years is similar in MDS/MPN-RS-T (1.8) and MDS-RS-SLD (2.4) and is higher in MDS/MPN-RS-T compared with ET (0.7).<sup>30</sup> Rates of thrombosis are similar in MDS/MPN-RS-T and ET but are higher than in MDS-RS.<sup>30,31</sup> In patients with anemia, treatment is usually supportive, with erythropoiesis-stimulating agents and transfusions following guidelines for lower-risk MDS. Low-dose aspirin may be prescribed for patients with *JAK2* mutations, older age, or cardiovascular risk factors. Although del(5q) is not common in MDS/MPN-RS-T, case reports have described activity of lenalidomide, a drug that typically causes thrombocytopenia.<sup>32</sup> Cytoreductive therapy is generally avoided because it can exacerbate anemia, but it may be implemented in the presence of multiple risk factors for thrombosis, vasomotor symptoms, or acquired von Willebrand syndrome.

Atypical chronic myeloid leukemia (aCML) is another WHO-recognized MDS/MPN overlap syndrome that is characterized by leukocytosis.<sup>33</sup> As its name suggests, it is distinct from classical chronic myeloid leukemia (CML) driven by the *BCR-ABL1* fusion gene. Specifically, aCML requires some degree of dysgranulopoiesis in the blood and bone marrow, minimal or no absolute basophilia (common in CML), minimal or no absolute monocytosis (common in CMML), and no gene rearrangements associated with other neoplasms (eg, *BCR-ABL1*, *PDGFRA*, *PDGFRB*,

**Table 1. Clinical features associated with different MDS/MPN overlap conditions and disorders at the diagnostic boundary with MDS**

	Median age (y)	Female/male ratio	Laboratory features	Physical features	Bone marrow features		
					Dysplasia	Cellularity	Other
MDS	70-71	1:1.5	Anemia is the most common cytopenia seen, 25-40% have thrombocytopenia	Hepatosplenomegaly or extramedullary involvement not seen	Present	Hypercellular (10-20% hypocellular)	Dysplasia in $\geq 1$ cell lineage; <20% blasts; 10% may have fibrosis
CMML	65-75	1:1.5-3	Absolute monocyte count $\geq 1.0 \times 10^9/L$ , accounting for $\geq 10\%$ of total WBCs for $\geq 3$ mo	Splenomegaly present in up to half of patients; hepatomegaly and extramedullary involvement (skin, LNs) may be seen	Typically present but not required for diagnosis	Hypercellular	Dysplasia typical in $\geq 1$ cell lineage, but may be absent; <20% blasts
MDS/MPN-RS-T	72-73	1:1	Anemia and platelet count $\geq 450 \times 10^9/L$	Thromboembolism may occur	Present	Hypercellular	$\geq 15\%$ erythroid precursors with ring sideroblasts; megakaryocytic atypia; <5% blasts
aCML	69-72	1:1.5	WBCs $> 13 \times 10^9/L$ , increased dysplastic neutrophils, no or minimal monocytosis and basophilia	Splenomegaly may be present	Present	Hypercellular	Dysplasia in $\geq 1$ cell lineage; <20% blasts
JMML	1-2	1:2-3	Absolute monocyte count $\geq 1.0 \times 10^9/L$ accounting for $\geq 10\%$ of total WBCs for $\geq 3$ mo	Splenomegaly common; monocytic and granulocytic infiltration of LNs, liver, skin, GI tract, and lungs also seen	Present	Hypercellular	<20% blasts
sAML	70-73	1:1.5	Variety of peripheral blood cytopenias may be seen, with or without leukocytosis	Hepatosplenomegaly may be present; infiltration of skin, gingiva, and CNS common in monocytic subtypes	Present	Hypercellular	$\geq 20\%$ blasts; Auer rods
SAA	50% <50 y old	1:1	Pancytopenia	Hepatosplenomegaly is not common; congenital anomalies may suggest an inherited marrow failure syndrome	Absent	Hypocellular	Profoundly hypocellular with all myeloid cell lineages diminished; marrow primarily composed of fat/stroma
CCUS	65-75	1:1.5	Anemia most common; other cytopenias can occur, often isolated	Hepatosplenomegaly or extramedullary involvement not seen	Absent or minimal	Normocellular or hypercellular	Does not meet MDS diagnostic criteria; fewer mutations in MDS genes but with comparable VAF

CNS, central nervous system; GI, gastrointestinal; LN, lymph node.

*FGFR1*, or *PCM1-JAK2*). Mutations typical of *BCR-ABL1*<sup>+</sup> MPN, like those in *JAK2*, *CALR*, and *MPL*, make a diagnosis of aCML less likely. The same is true for mutations of *CSF3R*, which are found in <10% of patients with aCML compared with 80% to 90% of patients with chronic neutrophilic leukemia, a clinically similar disorder that also presents with leukocytosis but no dysgranulopoiesis.<sup>33-35</sup> No single molecular abnormality specific for aCML has been described, although *SETBP1* mutations occur more frequently in aCML (25%) compared with CMML (6-15%) and JMML (3%).<sup>36</sup> Recurrent mutations in several other CMML-like genes have also been detected in patients with aCML (Figure 2).<sup>37</sup> In general, aCML patients tend to have a more aggressive disease course compared with patients with MDS/MPN, unclassifiable.<sup>33</sup> Although there is no consensus on the role of HSCT, long-term remissions have been reported with this strategy.<sup>38</sup> Other commonly used treatments include hypomethylating agent therapy and cytooreduction with hydroxyurea. The investigational use of JAK inhibitors has also been implemented in aCML and chronic neutrophilic leukemia based on the knowledge that some *CSF3R* mutations, most commonly *CSF3R*<sup>T618I</sup>, may activate the JAK/STAT pathway.<sup>39</sup>

Juvenile myelomonocytic leukemia (JMML) is an uncommon MDS/MPN overlap syndrome that occurs in early childhood, with a median age of 2 years. Clinical outcomes vary in JMML, with a minority of patients experiencing spontaneous remission, particularly those with germline diseases, such as Noonan or CBL syndrome, and some patients relapsing despite HSCT. Although there are shared clinical features with CMML, such as monocytosis and marked hepatosplenomegaly, the genetic landscape in JMML is distinct from adult myeloid neoplasms by the near absence of mutations in epigenetic and splicing modifiers. Up to 95% of children with JMML will possess a somatic or germline mutation in a Ras pathway gene (*PTPN11*, *NF1*, *NRAS*, *KRAS*, *CBL*).<sup>40-42</sup> Despite some patients having identical genetic mutation profiles, differing clinical outcomes are observed. Recently, DNA-methylation patterns were shown to improve the prediction of outcomes, distinguishing JMML patients who experienced spontaneous remission from those who experienced an aggressive disease course.<sup>43</sup>

## Other myeloid neoplasms with overlapping dysplastic and proliferative features

WHO-defined MDS/MPNs are considered distinct diagnoses, separate from the overlapping syndromes that they resemble. Yet myeloid malignancies can co-occur or have such nebulous boundaries that there exists an area of apparent diagnostic overlap. This can be challenging clinically, because treatment recommendations may differ across what may be rather arbitrary diagnostic borders. Consideration of clinical and molecular features may help to determine which condition should take precedence.

Due to its unique clinical and pathologic features, systemic mastocytosis (SM) is now recognized as its own disease category by the WHO. SM is divided into indolent SM (ISM), smoldering SM, SM with an associated clonal hematologic non-mast-cell lineage disease that was renamed as systemic mastocytosis with associated hematologic neoplasm (SM-AHN) in the WHO 2016 update, aggressive SM, and mast cell leukemia.<sup>1</sup> In addition to activating mutations in *KIT*, mutations in *TET2*, *SRSF2*, *ASXL1*, *CBL*, *RUNX1*,


and *RAS* have been identified in patients with SM-AHN, aggressive SM, and mast cell leukemia.<sup>44</sup> Additionally, mutations in *ETNK1* are frequently seen in patients with SM with eosinophilia.<sup>45</sup> Among patients with SM-AHN, these mutations may be coexpressed with *KIT* D816V in the same cells or expressed by other non-mast-cell myeloid cells.<sup>46,47</sup> Colony assay studies found that *KIT* D816V mutations are often late events that are frequently preceded by mutations of *TET2*, *SRSF2*, and *ASXL1*, indicating that SM-AHN is a multimitigated malignancy with diverging molecular evolution in subclones that have distinct differentiation potential.

Myeloid neoplasms account for 90% of all SM-AHN patients, including SM MPN (45%), SM CMML (29%), and SM MDS (23%).<sup>48</sup> The largest study to date comparing patients with SM CMML (n = 50) vs CMML alone (n = 501) evaluated differences in clinical, cytogenetic, and genetic features, as well as clinical outcomes.<sup>49</sup> Both groups had similar mutation profiles, with the exception of *KIT* and *CBL* mutations in the SM CMML cohort, suggesting that late *KIT* mutations may alter an initial CMML phenotype into one consistent with SM CMML.

Of note, *KIT*D816V may be viewed as a differentiation inducer in neoplastic cells rather than a dominant driver of oncogenesis, because patients with ISM express *KIT* D816V and do not typically have limited survival.<sup>50,51</sup> Additional pathways mediated by oncogenic lesions preceding *KIT* mutations are likely responsible for a more aggressive disease phenotype, treatment resistance, and shortened survival. To this end, treatment of SM-AHN is focused on which disease component requires more immediate intervention. For example, a patient with an associated higher-risk CMML and resultant peripheral cytopenias may be treated with a hypomethylating agent, whereas mast cell-directed therapy may be appropriate for a patient with a lower-risk non-mast-cell malignancy and symptoms or organ dysfunction ("C findings") related to the mast-cell component of the disease.<sup>52</sup> Midostaurin is an approved tyrosine kinase inhibitor with activity against *KIT* D816V that demonstrated an overall response rate of 60% among patients with advanced SM.<sup>53</sup> Additional studies are ongoing to evaluate alternative more selective *KIT* inhibitors. Future treatment strategies that extend beyond *KIT* are under investigation and include targeting pathways involving RAS, PI3K, mTOR, STAT5, and members of the BCL2 family.<sup>54,55</sup>

## AA and hypoplastic MDS

Another area of diagnostic overlap occurs between AA and hypoplastic MDS (hMDS). Although the etiology of AA is typically considered distinct from that of MDS, with AA driven by immune-mediated destruction of hematopoietic stem/progenitor cells and MDS driven by a selective growth advantage of somatically mutated clonal hematopoietic stem/progenitor cells, in practice, these mechanisms may co-occur (Figure 1). First, among a subset of patients with lower-risk MDS, immune activation and inflammation drive the selection of somatically mutated clones, potentiating the response to immunosuppressive therapies (ISTs).<sup>56,57</sup> Second, up to 15% of patients with severe AA (SAA) will have their disease evolve into MDS and/or acute myeloid leukemia (AML).<sup>58,59</sup> Distinguishing AA from hMDS may be challenging, because patients with these diseases share many clinical features, such as bone marrow hypocellularity that



**Figure 1.** Diagram depicting myeloid disorders with clinical and genetic features shared with MDS and the degree to which they are driven by proliferative and immunologic mechanisms.

hinders accurate evaluation of morphologic dysplasia, clonal cytogenetic and/or genetic abnormalities, and clinically meaningful responses to ISTs. Additionally, a subset of patients with AA harbor somatically mutated clones defined by mutations recurrently found in patients with MDS.<sup>60</sup> A recent study evaluated

somatic mutations in bone marrow samples from 150 patients with AA and no morphologic dysplasia.<sup>61</sup> Excluding *PIGA* mutations, 29 of 150 (19%) patients harbored mutations, predominantly in *ASXL1*, *DNMT3A*, and *BCOR* (Figure 2). A total of 17 (11%) patients experienced progression to MDS, with 11 of these patients belonging to the group of 29 patients who possessed mutations. Somatic mutations were significantly associated with longer disease duration, shorter telomere lengths, and a greater likelihood of progressing to MDS or AML compared with patients without mutations. A similar study of 439 patients with AA found clonal hematopoiesis in 47% of patients, with inferior survival outcomes seen among patients with *DNMT3A* and *ASXL1* mutations and higher IST response rates seen among patients with *BCOR* and *PIGA* mutations.<sup>62</sup> SAA patients with MDS-like mutations were more likely to have these clones expand over time, particularly after IST. Other patients can harbor somatic copy number–neutral loss of heterozygosity at the HLA locus on chromosome arm 6p<sup>63</sup> or mutations in HLAs and related pathways.<sup>64,65</sup> These abnormalities appear to provide escape from HLA-restricted T-cell immunity driving SAA, occur more often in younger patients, and are associated with lower rates of neoplastic progression.<sup>66,67</sup>

Approximately 15% to 20% of MDS bone marrows are hypocellular for age. These cases have differences in genetic profiles that include a lower rate of mutations and a lower frequency of splicing factor gene mutations compared with hyperplastic patients.<sup>68</sup> This pattern is more similar to that observed in SAA. Because bone marrow cellularity is limited, morphologic dysplasia

Mutated Gene	MDS	CMML	MDS/MPN-RS-T	aCML	JMML	sAML	SAA	CCUS	CHIP	Mutation Frequency
TET2										unknown
DNMT3A										rare or absent
ASXL1										< 2%
EZH2										2-5%
SETBP1										6-15%
SF3B1										16-25%
SRSF2										> 25%
U2AF1										
RUNX1										
TP53										
NF1										
NRAS										
KRAS										
CBL										
JAK2										
CALR										
MPL										
FLT3										
CSF3R										
IDH1										
IDH2										
NPM1										
BCOR/-L1										
PIGA										
ETNK1										

**Figure 2.** Differences in gene mutation frequency across different MDS/MPN overlap conditions and disorders at the diagnostic boundary with MDS.



is difficult to evaluate when considering hMDS vs SAA or non-severe AA. Other morphologic features outside of dysplasia that support a diagnosis of hMDS, or an MDS/MPN overlap syndrome, over AA include excess bone marrow blasts ( $\geq 2\%$ ), ring sideroblasts, extensive fibrosis, and circulating pseudo-Pelger-Huet cells. Certain cytogenetic abnormalities, such as *del(5q)*, monosomy 7, and inversion 3, are considered presumptive evidence of MDS.<sup>69,70</sup> The paucity of splicing factor and cohesin mutations in AA suggests that these lesions may also help to define the distinction between these disorders in the future. A lack of common MDS mutations or the presence of abnormalities of *BCOR*, *PIGA*, or the HLA loci correlate with more favorable outcomes in SAA and may be surrogate molecular markers of this disorder absent MDS defining features. In the meantime, a practical approach would be to minimize the distinction between hMDS and AA and simply consider patients at this boundary to be potentially responsive to immune suppression, reserving molecular studies to identify patients at risk for evolution to higher-risk disease.

One important caveat to this approach involves patients with inherited bone marrow failure syndromes, many of which can evolve into MDS or AML. For example, individuals with germline mutations of *GATA2*, *DDX41*, Fanconi anemia genes, or telomerase complex genes can have hypoplastic marrow findings well before the development of a clonal myeloid disorder that, in some cases, might never occur. Identifying these individuals is critical because their marrow failure does not respond to immune suppression. There are also important implications involving the health of family members, related stem cell donor candidates, and increased toxicity of IST or cytotoxic therapy. To make matters worse, some germline predisposition mutations, such as those in *RUNX1* and *ANKRD26*, may cause thrombocytopenia and megakaryocyte dysplasia that could be mistaken for MDS-defining criteria.<sup>61,71</sup> This diagnosis should not be made in the absence of other diagnostic elements.<sup>72</sup> In this context, however, the presence of somatic mutations may indicate a greater risk for neoplastic progression.<sup>73,74</sup> Mutation testing of presumed *de novo* MDS patients may also detect germline variants, because many of the genes tested are included in these panels.<sup>75</sup> These variants can occur even in patients without a family history, young age of onset, or associated physical findings typical of germline predisposition syndromes.<sup>76</sup> Dedicated testing of nonhematopoietic tissue is recommended in cases in which such a germline variant is suspected.<sup>77,78</sup>

## Clonal hematopoiesis, unexplained cytopenias, and lower-risk MDS

Another diagnostic boundary with MDS involves patients with unexplained cytopenias often described as ICUS. These patients lack MDS-defining bone marrow criteria that include an increased blast proportion, specific cytogenetic abnormalities, or morphologic dysplasia in at least 10% of cells of a given lineage (Figure 3).<sup>79</sup> Sequencing studies have identified somatic abnormalities indicative of clonal hematopoiesis in nearly 40% of ICUS patients, and closer to 70% in those who have some degree of dysplasia.<sup>80,81</sup> These individuals are described as having a clonal cytopenia of undetermined significance (CCUS). Patients with CCUS can have many of the same mutated genes observed in lower-risk MDS and have comparable variant allele

frequencies although mutations of *SF3B1* appear to be more specific for MDS. Patients with a CCUS have a high rate of progression to MDS or other myeloid malignancies, particularly if they carry higher-risk features. These include somatic mutations in *JAK2*, *RUNX1*, any of the commonly mutated splicing factors (*SF3B1*, *SRSF2*, *U2AF1*, *ZRSR2*), or  $\geq 2$  somatically mutated myeloid malignancy genes.<sup>82</sup> This risk may be as high as 90% at 5 years. For single mutations of *DNMT3A*, *TET2*, or *ASXL1*, the risk of progression is lower:  $\sim 50\%$  at 5 years. An absence of mutations on a broad panel of the 40 most frequently mutated MDS genes has a very low rate of progression approaching 1% per year of follow-up. Because MDS-defining bone marrow dysplasia can be hard to quantify, future revisions to MDS diagnostic criteria may include more clearly defined higher-risk CCUS patients, just as *SF3B1* mutations are currently accepted as evidence of MDS-RS in patients with as few as 5% ring sideroblasts.<sup>1</sup>

An important caveat is that somatic mutations typical of MDS can also occur in the blood cells of hematologically normal persons, with a prevalence that increases markedly with age.<sup>83,84</sup> These individuals are said to have clonal hematopoiesis of indeterminate potential (CHIP) and, in the absence of cytopenias (or another concerning clinical context, such as a germline predisposition), are believed to have a very low risk for neoplastic progression ( $\sim 1\%$  per year). CHIP mutations are most often found in *DNMT3A*, *TET2*, or *ASXL1* (Figure 2) as isolated lesions with a low variant allele frequency (VAF) ( $<10\%$ ) and should not be considered diagnostic of MDS or any myeloid neoplasm. CHIP should also not be equated with CCUS, in which mutations are more frequent, of greater abundance, and associated with a much higher probability of malignant progression.<sup>80-82,85,86</sup>

## MDS progression to sAML

At the other end of the prognostic spectrum for MDS lies the boundary with sAML. The border between these disorders has shifted over time, with the WHO classification for sAML currently defined as  $\geq 20\%$  bone marrow and/or peripheral blood blasts. Under the earlier French-American-British schema, MDS patients with 20% to 29% blasts were considered to have refractory anemia with excess blasts in transformation. The poor outcome of this latter group prompted the lower blast threshold set by the WHO; however, in retrospect, it is not clear that MDS patients with 10% to 19% bone marrow blasts have meaningfully different outcomes. In practice, these 2 groups straddling the divide between MDS and AML are treated in a similar fashion, receiving hypomethylating agents and considered for HSCT when appropriate. For these reasons, there have been calls to do away with the concepts of MDS with excess blasts and low-blast-count sAML, unifying them under the term "oligoblastic leukemia."<sup>87</sup> However, arbitrarily redefining the boundary between MDS and AML may not be enough. The challenge will be to identify those MDS patients who are headed toward leukemic progression and those who may have excess blasts but largely fail to progress.

The existence of the latter group can be inferred from the population of prognostically higher risk patients who live longer than the median for their revised International Prognostic Scoring System risk group.<sup>88</sup> These individuals have a time-dependent risk

MDS by WHO 2016						
	Non-Clonal ICUS	CHIP	CCUS	Low Blast MDS	High Blast MDS	sAML/AML-MRC
VAf	N/A	~9%	~10-50%	~30-50%	~40-50%	~40-50%
Dysplasia	-	-	-	+	+	+
Cytopenias	+	-	+	+	+	+
BM Blast %	< 2%	< 2%	< 2%	< 2%	2-19%	20+%
Overall Risk	Very Low	Very Low	Low	Low/Int	High	Very High
Treatments	Observation	None	Obs/BSC/GF	Obs/BSC/GF IMiD/IST	HMA/HST	HMA/IC/HST
			Clonal Cytopenias		Oligoblastic Leukemia	

**Figure 3. Comparison of features between cytopenic and clonal hematopoietic states that border MDS.** BSC, best supportive care; GF, growth factors; HMA, hypomethylating agent; HST, hematopoietic stem cell transplant; IC, induction chemotherapy; IMiD, immunomodulatory imide drugs; Obs, observation.

that more closely resembles that of MDS patients with lower-risk disease. Because this determination is not made at diagnosis, several studies have attempted to risk-stratify MDS patients based on their leukemic potential earlier in the course of their disease. For example, Makishima et al examined tumor samples from >2000 patients for mutated genes enriched in higher-risk MDS and sAML.<sup>89</sup> Mutations of *NPM1*, *IDH1*, *IDH2*, *WT1*, *NRAS*, *PTPN11*, and *FLT3* were found significantly more often in the sAML cohort. Mutations of these genes were typically subclonal to a more abundant mutation (suggesting that they were acquired later) and were associated with significantly shorter progression-free survival. In MDS, acquisition of these gene mutations may define leukemic clones that might not lead to a clinical definition of sAML for many months. Such patients could be said to harbor an overlap disorder between MDS and sAML. One implication of this hypothesis is that therapies targeted at these subclones (eg, *IDH* or *FLT3* inhibitors) may not lead to traditionally defined hematologic responses but, nonetheless, may delay leukemic transformation. This prediction will have to be tested in prospective clinical trials.

Patterns of gene expression have also been used to identify MDS patients at greatest risk of leukemic progression. Shiozawa et al examined the transcriptomes of CD34<sup>+</sup> bone marrow cells from patients with MDS.<sup>90</sup> Unsupervised clustering identified 2 major subgroups: 1 enriched for the expression of genes associated with erythroid and megakaryocytic differentiation and another defined by transcripts associated with immature progenitors (IMPs). The erythroid and megakaryocytic differentiation subgroup had longer overall survival and was associated with *SF3B1* mutations, ring sideroblasts, and a strong erythroid signature. In contrast, the IMP subgroup had lower platelet counts, increased marrow blasts, and higher-risk mutations. Strikingly, only patients in the IMP subgroup had disease that transformed into sAML, suggesting that, even in the absence of "leukemic" mutations, some forms of MDS have greater leukemic potential that can be recognized well before progression takes place.

The second area of overlap between MDS and AML involves patients who may not have had a recognized antecedent MDS

but are diagnosed with AML with myelodysplasia-related changes (AML-MRC), suggesting a pathogenic link with MDS.<sup>91,92</sup> Molecular profiling may help to segregate those with MDS and AML-MRC from those with de novo AML, providing prognostic information for the patient and clinician. These patients will frequently harbor MDS-associated cytogenetic abnormalities and are often classified as having higher-risk disease. Not surprisingly, patients with AML-MRC are more likely to harbor mutated genes typical of MDS and sAML, including splicing factors (*SRSF2*, *SF3B1*, and *U2AF1*), chromatin modifiers (*EZH2* and *ASXL1*), and *STAG2* and *BCOR* (Figure 2).<sup>93,94</sup> Older individuals with AML are more likely to carry somatic mutations in these genes, even if they are not described as having AML-MRC. Importantly, older de novo AML patients without these mutations have a more favorable response to therapy and duration of remission, making it important to identify them at diagnosis.

From another perspective, one could consider MDS with excess blasts to be an overlap syndrome between lower-risk MDS defined by clonal cytopenias with bone marrow failure and oligoblastic myeloid leukemia (Figure 3).<sup>86,95</sup> In practice, patients with higher-risk MDS or low-blast-count AML have a similar prognosis and are often treated with hypomethylating agents or, if appropriate, considered for stem cell transplantation.<sup>96</sup> Altering our diagnostic boundaries between MDS and AML based on underlying mutations and clinical phenotypes may more accurately classify patients with these conditions.

## Conclusions

MDS overlap syndromes are genetically and clinically heterogeneous disorders that can represent distinct biological entities or areas of diagnostic ambiguity. Although WHO-defined disease classifications rely largely on morphologic criteria, molecular markers of disease are increasingly able to identify differences in clinical phenotypes when considered in the appropriate clinical context. There are no specific mutations that stringently diagnose MDS overlap syndromes or unequivocally define diagnostic boundaries with MDS;

however, future classification schemes are sure to incorporate our growing understanding of the molecular basis of these disorders.

## Authorship

Contribution: T.N.T. and R.B. cowrote this review and edited the submitted manuscript.

Conflict-of-interest disclosure: R.B. has served as a consultant for Celgene and Genoptix, has received research funding from Celgene and Takeda, has served on a data safety monitoring board for Celgene-sponsored clinical trials, and has received honoraria for speaking at

education conferences sponsored by Celgene and Xian Janssen. T.N.T. declares no competing financial interests.

ORCID profiles: T.N.T., 0000-0002-7656-5249; R.B., 0000-0002-5603-4598.

Correspondence: Rafael Bejar, University of California San Diego Moores Cancer Center, 3855 Health Sciences Dr, MC 0820, La Jolla, CA 92093-0820; e-mail: rabejar@ucsd.edu.

## Footnote

Submitted 15 October 2018; accepted 11 December 2018. Prepublished online as *Blood* First Edition paper, 22 January 2019; DOI 10.1182/blood-2018-10-844670.

## REFERENCES

- Arber DA, Orazi A, Hasserjian R, et al. The 2016 revision to the World Health Organization classification of myeloid neoplasms and acute leukemia. *Blood*. 2016;127(20):2391-2405.
- Solary E, Itzykson R. How I treat chronic myelomonocytic leukemia. *Blood*. 2017;130(2):126-136.
- Ricci C, Fermo E, Corti S, et al. RAS mutations contribute to evolution of chronic myelomonocytic leukemia to the proliferative variant. *Clin Cancer Res*. 2010;16(8):2246-2256.
- Cervera N, Itzykson R, Coppin E, et al. Gene mutations differently impact the prognosis of the myelodysplastic and myeloproliferative classes of chronic myelomonocytic leukemia. *Am J Hematol*. 2014;89(6):604-609.
- Selimoglu-Buet D, Badaoui B, Benayoun E, et al; Groupe Francophone des Myélodysplasies. Accumulation of classical monocytes defines a subgroup of MDS that frequently evolves into CMML. *Blood*. 2017;130(6):832-835.
- Talati C, Zhang L, Shaheen G, et al. Monocyte subset analysis accurately distinguishes CMML from MDS and is associated with a favorable MDS prognosis. *Blood*. 2017;129(13):1881-1883.
- Meggendorfer M, Roller A, Haferlach T, et al. SRSF2 mutations in 275 cases with chronic myelomonocytic leukemia (CMML). *Blood*. 2012;120(15):3080-3088.
- Patnaik MM, Itzykson R, Lasho TL, et al. ASXL1 and SETBP1 mutations and their prognostic contribution in chronic myelomonocytic leukemia: a two-center study of 466 patients. *Leukemia*. 2014;28(11):2206-2212.
- Papaemmanuil E, Gerstung M, Malcovati L, et al; Chronic Myeloid Disorders Working Group of the International Cancer Genome Consortium. Clinical and biological implications of driver mutations in myelodysplastic syndromes. *Blood*. 2013;122(22):3616-3627, quiz 3699.
- Itzykson R, Kosmider O, Renneville A, et al. Prognostic score including gene mutations in chronic myelomonocytic leukemia. *J Clin Oncol*. 2013;31(19):2428-2436.
- Padron E, Garcia-Manero G, Patnaik MM, et al. An international data set for CMML validates prognostic scoring systems and demonstrates a need for novel prognostication strategies. *Blood Cancer J*. 2015;5(7):e333.
- Mughal TI, Cross NC, Padron E, et al. An International MDS/MPN Working Group's perspective and recommendations on molecular pathogenesis, diagnosis and clinical characterization of myelodysplastic/myeloproliferative neoplasms. *Haematologica*. 2015;100(9):1117-1130.
- Kantarjian H, O'Brien S, Ravandi F, et al. Proposal for a new risk model in myelodysplastic syndrome that accounts for events not considered in the original International Prognostic Scoring System. *Cancer*. 2008;113(6):1351-1361.
- Greenberg P, Cox C, LeBeau MM, et al. International scoring system for evaluating prognosis in myelodysplastic syndromes. *Blood*. 1997;89(6):2079-2088.
- Greenberg PL, Tuechler H, Schanz J, et al. Revised international prognostic scoring system for myelodysplastic syndromes. *Blood*. 2012;120(12):2454-2465.
- Nazha A, Patnaik MM. Making sense of prognostic models in chronic myelomonocytic leukemia. *Curr Hematol Malig Rep*. 2018;13(5):341-347.
- Wassie EA, Itzykson R, Lasho TL, et al. Molecular and prognostic correlates of cytogenetic abnormalities in chronic myelomonocytic leukemia: a Mayo Clinic-French Consortium Study. *Am J Hematol*. 2014;89(12):1111-1115.
- Beran M, Wen S, Shen Y, et al. Prognostic factors and risk assessment in chronic myelomonocytic leukemia: validation study of the M.D. Anderson Prognostic Scoring System. *Leuk Lymphoma*. 2007;48(6):1150-1160.
- Elena C, Galli A, Such E, et al. Integrating clinical features and genetic lesions in the risk assessment of patients with chronic myelomonocytic leukemia. *Blood*. 2016;128(10):1408-1417.
- Drummond MW, Pocock C, Boissinot M, et al. A multi-centre phase 2 study of azacitidine in chronic myelomonocytic leukaemia. *Leukemia*. 2014;28(7):1570-1572.
- Tantravahi SK, Szankasi P, Khorashad JS, et al. A phase II study of the efficacy, safety, and determinants of response to 5-azacitidine (Vidaza) in patients with chronic myelomonocytic leukemia. *Leuk Lymphoma*. 2016;57(10):2441-2444.
- Merlevede J, Droin N, Qin T, et al. Mutation allele burden remains unchanged in chronic myelomonocytic leukaemia responding to hypomethylating agents. *Nat Commun*. 2016;7(1):10767.
- Meldi K, Qin T, Buchi F, et al. Specific molecular signatures predict decitabine response in chronic myelomonocytic leukemia. *J Clin Invest*. 2015;125(5):1857-1872.
- de Witte T, Bowen D, Robin M, et al. Allogeneic hematopoietic stem cell transplantation for MDS and CMML: recommendations from an international expert panel. *Blood*. 2017;129(13):1753-1762.
- Robin M, Fenaux P. Hypomethylating agents as bridging therapy before allogeneic hematopoietic stem cell transplantation in patients with chronic myelomonocytic leukemia? *Biol Blood Marrow Transplant*. 2016;22(1):1-2.
- Jeromin S, Haferlach T, Weissmann S, et al. Refractory anemia with ring sideroblasts and marked thrombocytosis cases harbor mutations in SF3B1 or other spliceosome genes accompanied by JAK2V617F and ASXL1 mutations. *Haematologica*. 2015;100(4):e125-e127.
- Patnaik MM, Tefferi A. Refractory anemia with ring sideroblasts (RARS) and RARS with thrombocytosis (RARS-T): 2017 update on diagnosis, risk-stratification, and management. *Am J Hematol*. 2017;92(3):297-310.
- Meggendorfer M, Jeromin S, Haferlach C, Kern W, Haferlach T. The mutational landscape of 18 investigated genes clearly separates four subtypes of myelodysplastic/myeloproliferative neoplasms. *Haematologica*. 2018;103(5):e192-e195.
- Patnaik MM, Lasho TL, Finke CM, et al. Predictors of survival in refractory anemia with ring sideroblasts and thrombocytosis (RARS-T) and the role of next-generation sequencing. *Am J Hematol*. 2016;91(5):492-498.
- Broseus J, Florensa L, Zipperer E, et al. Clinical features and course of refractory anemia with ring sideroblasts associated with marked thrombocytosis. *Haematologica*. 2012;97(7):1036-1041.
- Patnaik MM, Lasho TL, Finke CM, et al. Vascular events and risk factors for thrombosis in refractory anemia with ring sideroblasts and thrombocytosis. *Leukemia*. 2016;30(11):2273-2275.
- Huls G, Mulder AB, Rosati S, van de Loosdrecht AA, Vellenga E, de Wolf JT. Efficacy of



- single-agent lenalidomide in patients with JAK2 (V617F) mutated refractory anemia with ring sideroblasts and thrombocytosis. *Blood*. 2010;116(2):180-182.
33. Wang SA, Hasserjian RP, Fox PS, et al. Atypical chronic myeloid leukemia is clinically distinct from unclassifiable myelodysplastic/myeloproliferative neoplasms. *Blood*. 2014;123(17):2645-2651.
  34. Maxson JE, Gotlib J, Pollyea DA, et al. Oncogenic CSF3R mutations in chronic neutrophilic leukemia and atypical CML. *N Engl J Med*. 2013;368(19):1781-1790.
  35. Pardanani A, Lasho TL, Laborde RR, et al. CSF3R T618I is a highly prevalent and specific mutation in chronic neutrophilic leukemia. *Leukemia*. 2013;27(9):1870-1873.
  36. Makishima H, Yoshida K, Nguyen N, et al. Somatic SETBP1 mutations in myeloid malignancies. *Nat Genet*. 2013;45(8):942-946.
  37. Gotlib J, Maxson JE, George TI, Tyner JW. The new genetics of chronic neutrophilic leukemia and atypical CML: implications for diagnosis and treatment. *Blood*. 2013;122(10):1707-1711.
  38. Langabeer SE, McCarron SL, Haslam K, O'Donovan MT, Conneally E. The CSF3R T618I mutation as a disease-specific marker of atypical CML post allo-SCT. *Bone Marrow Transplant*. 2014;49(6):843-844.
  39. Dao KH, Solti MB, Maxson JE, et al. Significant clinical response to JAK1/2 inhibition in a patient with CSF3R-T618I-positive atypical chronic myeloid leukemia. *Leuk Res Rep*. 2014;3(2):67-69.
  40. Stieglitz E, Taylor-Weiner AN, Chang TY, et al. The genomic landscape of juvenile myelomonocytic leukemia [published correction appears in *Nat Genet*. 2015;47(11):1333]. *Nat Genet*. 2015;47(11):1326-1333.
  41. Sakaguchi H, Okuno Y, Muramatsu H, et al. Exome sequencing identifies secondary mutations of SETBP1 and JAK3 in juvenile myelomonocytic leukemia. *Nat Genet*. 2013;45(8):937-941.
  42. Caye A, Strullu M, Guidez F, et al. Juvenile myelomonocytic leukemia displays mutations in components of the RAS pathway and the PRC2 network. *Nat Genet*. 2015;47(11):1334-1340.
  43. Stieglitz E, Mazor T, Olshen AB, et al. Genome-wide DNA methylation is predictive of outcome in juvenile myelomonocytic leukemia. *Nat Commun*. 2017;8(1):2127.
  44. Schwaab J, Schnittger S, Sotlar K, et al. Comprehensive mutational profiling in advanced systemic mastocytosis. *Blood*. 2013;122(14):2460-2466.
  45. Lasho TL, Finke CM, Zblewski D, et al. Novel recurrent mutations in ethanolamine kinase 1 (ETNK1) gene in systemic mastocytosis with eosinophilia and chronic myelomonocytic leukemia. *Blood Cancer J*. 2015;5(1):e275.
  46. Jawhar M, Schwaab J, Schnittger S, et al. Additional mutations in SRSF2, ASXL1 and/or RUNX1 identify a high-risk group of patients with KIT D816V(+) advanced systemic mastocytosis. *Leukemia*. 2016;30(1):136-143.
  47. Jawhar M, Schwaab J, Schnittger S, et al. Molecular profiling of myeloid progenitor cells in multi-mutated advanced systemic mastocytosis identifies KIT D816V as a distinct and late event. *Leukemia*. 2015;29(5):1115-1122.
  48. Pardanani A, Lim KH, Lasho TL, et al. Prognostically relevant breakdown of 123 patients with systemic mastocytosis associated with other myeloid malignancies. *Blood*. 2009;114(18):3769-3772.
  49. Patnaik MM, Rangit Vallapureddy, Lasho TL, et al. A comparison of clinical and molecular characteristics of patients with systemic mastocytosis with chronic myelomonocytic leukemia to CMML alone. *Leukemia*. 2018;32(8):1850-1856.
  50. Mayerhofer M, Gleixner KV, Hoelbl A, et al. Unique effects of KIT D816V in BaF3 cells: induction of cluster formation, histamine synthesis, and early mast cell differentiation antigens. *J Immunol*. 2008;180(8):5466-5476.
  51. Lim KH, Tefferi A, Lasho TL, et al. Systemic mastocytosis in 342 consecutive adults: survival studies and prognostic factors. *Blood*. 2009;113(23):5727-5736.
  52. Scherber RM, Borate U. How we diagnose and treat systemic mastocytosis in adults. *Br J Haematol*. 2018;180(1):11-23.
  53. Gotlib J, Kluin-Nelemans HC, George TI, et al. Efficacy and safety of midostaurin in advanced systemic mastocytosis. *N Engl J Med*. 2016;374(26):2530-2541.
  54. Gleixner KV, Mayerhofer M, Cerny-Reiterer S, et al. KIT-D816V-independent oncogenic signaling in neoplastic cells in systemic mastocytosis: role of Lyn and Btk activation and disruption by dasatinib and bosutinib. *Blood*. 2011;118(7):1885-1898.
  55. Peter B, Cerny-Reiterer S, Hadzijušević E, et al. The pan-Bcl-2 blocker obatoclax promotes the expression of Puma, Noxa, and Bim mRNA and induces apoptosis in neoplastic mast cells. *J Leukoc Biol*. 2014;95(1):95-104.
  56. Stahl M, DeVeaux M, de Witte T, et al. The use of immunosuppressive therapy in MDS: clinical outcomes and their predictors in a large international patient cohort. *Blood Adv*. 2018;2(14):1765-1772.
  57. Passweg JR, Giagounidis AA, Simcock M, et al. Immunosuppressive therapy for patients with myelodysplastic syndrome: a prospective randomized multicenter phase III trial comparing antithymocyte globulin plus cyclosporine with best supportive care--SAKK 33/99. *J Clin Oncol*. 2011;29(3):303-309.
  58. Socié G, Henry-Amar M, Bacigalupo A, et al. European Bone Marrow Transplantation-Severe Aplastic Anaemia Working Party. Malignant tumors occurring after treatment of aplastic anemia. *N Engl J Med*. 1993;329(16):1152-1157.
  59. Socié G, Rosenfeld S, Frickhofen N, Gluckman E, Tichelli A. Late clonal diseases of treated aplastic anemia. *Semin Hematol*. 2000;37(1):91-101.
  60. Stanley N, Olson TS, Babushok DV. Recent advances in understanding clonal haematopoiesis in aplastic anaemia. *Br J Haematol*. 2017;177(4):509-525.
  61. Kulasekararaj AG, Jiang J, Smith AE, et al. Somatic mutations identify a subgroup of aplastic anemia patients who progress to myelodysplastic syndrome. *Blood*. 2014;124(17):2698-2704.
  62. Yoshizato T, Dumitriu B, Hosokawa K, et al. Somatic mutations and clonal Hematopoiesis in aplastic anemia. *N Engl J Med*. 2015;373(1):35-47.
  63. Katagiri T, Sato-Otsubo A, Kashiwase K, et al; Japan Marrow Donor Program. Frequent loss of HLA alleles associated with copy number-neutral 6pLOH in acquired aplastic anemia. *Blood*. 2011;118(25):6601-6609.
  64. Maciejewski JP, Follmann D, Nakamura R, et al. Increased frequency of HLA-DR2 in patients with paroxysmal nocturnal hemoglobinuria and the PNH/aplastic anemia syndrome. *Blood*. 2001;98(13):3513-3519.
  65. Sauntharajah Y, Nakamura R, Nam JM, et al. HLA-DR15 (DR2) is overrepresented in myelodysplastic syndrome and aplastic anemia and predicts a response to immunosuppression in myelodysplastic syndrome. *Blood*. 2002;100(5):1570-1574.
  66. Führer M, Durner J, Brännler G, et al. HLA association is different in children and adults with severe acquired aplastic anemia. *Pediatr Blood Cancer*. 2007;48(2):186-191.
  67. Babushok DV, Duke JL, Xie HM, et al. Somatic HLA mutations expose the role of class I-mediated autoimmunity in aplastic anemia and its clonal complications. *Blood Adv*. 2017;1(22):1900-1910.
  68. Nazha A, Seastone D, Radivoyevitch T, et al. Genomic patterns associated with hypoplastic compared to hyperplastic myelodysplastic syndromes. *Haematologica*. 2015;100(11):e434-e437.
  69. Afable MG II, Wlodarski M, Makishima H, et al. SNP array-based karyotyping: differences and similarities between aplastic anemia and hypocellular myelodysplastic syndromes. *Blood*. 2011;117(25):6876-6884.
  70. Mikhailova N, Sessarego M, Fugazza G, et al. Cytogenetic abnormalities in patients with severe aplastic anemia. *Haematologica*. 1996;81(5):418-422.
  71. Noris P, Favier R, Alessi MC, et al. ANKRD26-related thrombocytopenia and myeloid malignancies. *Blood*. 2013;122(11):1987-1989.
  72. Niemeyer CM, Mecucci C. Practical considerations for diagnosis and management of patients and carriers. *Semin Hematol*. 2017;54(2):69-74.
  73. Schaefer EJ, Lindsley RC. Significance of clonal mutations in bone marrow failure and inherited myelodysplastic syndrome/acute myeloid leukemia predisposition syndromes. *Hematol Oncol Clin North Am*. 2018;32(4):643-655.
  74. Churpek JE, Pyrtel K, Kanchi K-L, et al. Genomic analysis of germ line and somatic variants in familial myelodysplasia/acute myeloid leukemia. *Blood*. 2015;126(22):2484-2490.
  75. Drazer MW, Kadri S, Sukhanova M, et al. Prognostic tumor sequencing panels frequently identify germ line variants associated

- with hereditary hematopoietic malignancies. *Blood Adv*. 2018;2(2):146-150.
76. Brown AL, Churpek JE, Malcovati L, Döhner H, Godley LA. Recognition of familial myeloid neoplasia in adults. *Semin Hematol*. 2017; 54(2):60-68.
77. National Comprehensive Cancer Network. Myelodysplastic syndromes (Version 2.2019). [https://www.nccn.org/professionals/physician\\_gls/pdf/mds.pdf](https://www.nccn.org/professionals/physician_gls/pdf/mds.pdf). Accessed 26 November 2018.
78. Guidugli L, Johnson AK, Alkorta-Aranburu G, et al. Clinical utility of gene panel-based testing for hereditary myelodysplastic syndrome/acute leukemia predisposition syndromes. *Leukemia*. 2017;31(5):1226-1229.
79. Valent P, Orazi A, Steensma DP, et al. Proposed minimal diagnostic criteria for myelodysplastic syndromes (MDS) and potential pre-MDS conditions. *Oncotarget*. 2017;8(43):73483-73500.
80. Kwok B, Hall JM, Witte JS, et al. MDS-associated somatic mutations and clonal hematopoiesis are common in idiopathic cytopenias of undetermined significance. *Blood*. 2015;126(21):2355-2361.
81. Cargo CA, Rowbotham N, Evans PA, et al. Targeted sequencing identifies patients with preclinical MDS at high risk of disease progression. *Blood*. 2015;126(21):2362-2365.
82. Malcovati L, Galli A, Travaglini E, et al. Clinical significance of somatic mutation in unexplained blood cytopenia. *Blood*. 2017; 129(25):3371-3378.
83. Jaiswal S, Fontanillas P, Flannick J, et al. Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med*. 2014; 371(26):2488-2498.
84. Genovese G, Kähler AK, Handsaker RE, et al. Clonal hematopoiesis and blood-cancer risk inferred from blood DNA sequence. *N Engl J Med*. 2014;371(26):2477-2487.
85. Bejar R. CHIP, ICUS, CCUS and other four-letter words. *Leukemia*. 2017;31(9):1869-1871.
86. Valent P. ICUS, IDUS, CHIP and CCUS: diagnostic criteria, separation from MDS and clinical implications. *Pathobiology*. 2019;86(1):30-38.
87. Lichtman MA. Does a diagnosis of myelogenous leukemia require 20% marrow myeloblasts, and does <5% marrow myeloblasts represent a remission? The history and ambiguity of arbitrary diagnostic boundaries in the understanding of myelodysplasia. *Oncologist*. 2013;18(9):973-980.
88. Pfeilstöcker M, Tuechler H, Sanz G, et al. Time-dependent changes in mortality and transformation risk in MDS. *Blood*. 2016;128(7): 902-910.
89. Makishima H, Yoshizato T, Yoshida K, et al. Dynamics of clonal evolution in myelodysplastic syndromes. *Nat Genet*. 2017;49(2):204-212.
90. Shiozawa Y, Malcovati L, Galli A, et al. Gene expression and risk of leukemic transformation in myelodysplasia. *Blood*. 2017;130(24): 2642-2653.
91. Sperling AS, Gibson CJ, Ebert BL. The genetics of myelodysplastic syndrome: from clonal haematopoiesis to secondary leukemias. *Nat Rev Cancer*. 2017;17(1):5-19.
92. Corces-Zimmerman MR, Hong WJ, Weissman IL, Medeiros BC, Majeti R. Preleukemic mutations in human acute myeloid leukemia affect epigenetic regulators and persist in remission. *Proc Natl Acad Sci USA*. 2014; 111(7):2548-2553.
93. Lindsley RC, Mar BG, Mazzola E, et al. Acute myeloid leukemia ontogeny is defined by distinct somatic mutations. *Blood*. 2015; 125(9):1367-1376.
94. Yokoyama K, Shimizu E, Yokoyama N, et al. Cell-lineage level-targeted sequencing to identify acute myeloid leukemia with myelodysplasia-related changes. *Blood Adv*. 2018;2(19):2513-2521.
95. Bejar R. What biologic factors predict for transformation to AML? *Best Pract Res Clin Haematol*. 2018;31(4):341-345.
96. DiNardo CD, Garcia-Manero G, Pierce S, et al. Interactions and relevance of blast percentage and treatment strategy among younger and older patients with acute myeloid leukemia (AML) and myelodysplastic syndrome (MDS). *Am J Hematol*. 2016;91(2): 227-232.



**blood**®

2019 133: 1086-1095

doi:10.1182/blood-2018-10-844670 originally published  
online January 22, 2019

## **MDS overlap disorders and diagnostic boundaries**

Tiffany N. Tanaka and Rafael Bejar

---

Updated information and services can be found at:

<http://www.bloodjournal.org/content/133/10/1086.full.html>

Articles on similar topics can be found in the following Blood collections

[Clinical Trials and Observations](#) (4981 articles)

[Hematopoiesis and Stem Cells](#) (3581 articles)

[Myeloid Neoplasia](#) (1978 articles)

[Review Articles](#) (839 articles)

[Review Series](#) (244 articles)

---

Information about reproducing this article in parts or in its entirety may be found online at:

[http://www.bloodjournal.org/site/misc/rights.xhtml#repub\\_requests](http://www.bloodjournal.org/site/misc/rights.xhtml#repub_requests)

Information about ordering reprints may be found online at:

<http://www.bloodjournal.org/site/misc/rights.xhtml#reprints>

Information about subscriptions and ASH membership may be found online at:

<http://www.bloodjournal.org/site/subscriptions/index.xhtml>