




Patient-reported outcome measures for facial skin cancer: a systematic review and evaluation of the quality of their measurement properties

T.D. Dobbs ^{1,2} H. Samarendra,³ S. Hughes,^{4,5} H.A. Hutchings ⁴ and I. Whitaker ^{1,2}

¹Reconstructive Surgery and Regenerative Medicine Research Group (ReconRegen), ⁴Patient and Population Health and Informatics, Institute of Life Science 2, Swansea University Medical School, Swansea, U.K.

²The Welsh Centre for Burns and Plastic Surgery, Morriston Hospital, Swansea, U.K.

³Oxford University Medical School, Oxford, U.K.

⁵Abertawe Bro Morgannwg University Health Board, Princess of Wales Hospital, Bridgend, U.K.

Linked Comment: Rodrigues. *Br J Dermatol* 2019; **180**:976.

Summary

Correspondence

Thomas D. Dobbs.

E-mail: tomddobbs@doctors.org.uk

Accepted for publication

5 October 2018

Funding sources

This work has received no specific funding.

T.D.D. is funded by the Welsh Clinical Academic Training Fellowship.

Conflicts of interest

None to declare.

DOI 10.1111/bjd.17342

Background Skin cancer is the commonest malignancy worldwide, often occurring on the face. Both the condition and treatment can lead to scarring and facial disfigurement, affecting a patient's health-related quality of life (HRQoL), which can be measured using patient-reported outcome measures (PROMs).

Objectives This systematic review identifies PROMs for facial skin cancer and appraises their methodological quality and psychometric properties using up-to-date methods. **Methods** MEDLINE, Embase, PsycINFO, Cochrane and CINAHL were searched systematically in accordance with PRISMA guidelines, identifying all PROMs designed for or validated in facial skin cancer. Methodological quality and evidence of psychometric properties were assessed using the CONsensus-based Standards for the Selection of Health Measurement INstruments (COSMIN) checklist and criteria proposed by Terwee and colleagues. A best-evidence synthesis and assessment of instrument focus on post-resection reconstruction was also performed.

Results We included 24 studies on 11 PROMs. Methodological quality and psychometric evidence was variable, with the Patient Outcome of Surgery – Head/Neck (POS-H/N), Skin Cancer Index (SCI), Skin Cancer Quality of Life Impact Tool (SCQOLIT) and Essers and colleagues demonstrating the greatest level of validation. None scored well in their relevance to post-skin cancer reconstruction of the face.

Conclusions This systematic review critically appraises PROMs for facial skin cancer using internationally accepted criteria. The identified PROMs demonstrate a variation in the quality of validation performed, with a need to improve this across all PROMs in the field. Only through improving the quality of available PROMs and their focus on the post-treatment aesthetic and functional outcome will we be able to truly appreciate the concerns of our patients and improve the management of facial skin cancer.

What's already known about this topic?

- Patient-reported outcome measures (PROMs) are important in both research and daily clinical practice.
- This is especially true in facial skin cancer, where both the condition and the resulting aesthetic outcome of treatment are important.
- PROMs for facial skin cancer exist; however, their validity against the contemporary international consensus have yet to be reported.
- The relevance of these PROMs to patients' views of treatment outcomes is yet to be investigated.

What does this study add?

- This systematic review provides a comprehensive assessment of the validity of those PROMs used for facial skin cancer using current best-practice assessment tools, helping clinicians and researchers to select the most appropriate PROM to use.
- Each PROM is also assessed for relevance to the post-treatment aesthetic outcome, with a recommendation that further validated items are required to adequately assess this important area of skin cancer treatment.

Skin cancer is the commonest malignancy worldwide,¹ affecting one in five Americans during their lifetime.² The incidence of nonmelanoma skin cancers (NMSC) in England is 98.85 per 100 000 person-years and predominantly affects the face.^{3,4} There were 70 000 new diagnoses of NMSC made in the U.K. in 2013,⁵ presenting a significant and growing health burden. Although skin cancer mortality is low, particularly for NMSC,^{5,6} the diagnosis is often psychologically damaging, including anxiety over the cancer diagnosis⁷ and concerns over visible scarring, especially on the face,⁸ affecting health-related quality of life (HRQoL).

HRQoL has been given a number of definitions,⁹ but broadly represents an individual's perception of the effects of an illness and/or treatment on physical, psychological and social aspects of their life.¹⁰ One method for assessing HRQoL is the use of patient-reported outcome measures (PROMs). PROMs are standardized, validated questionnaires that are completed by patients and capture one or more aspects of their health and well-being.^{11,12} They are considered by the U.K. Department of Health to be the current best method for quantifying a patient's clinical experience. Currently only four conditions have routine PROM data collected at a national level in the U.K.,¹² although PROM data collection in many different cancer registries and dermatological trials is now commonplace.^{13,14}

Previous reviews have demonstrated a number of PROMs used in the assessment of patients with both skin cancer generally^{15,16} and facial skin cancer.¹⁷ However, none have used current 'gold-standard' methodology for assessing the methodological quality of included studies, or the quality of those PROMs' measurement properties. Furthermore, given the burden associated with cosmetic outcomes in post-skin cancer facial reconstruction, no review has yet assessed available PROMs for their focus on this. In an era of core outcome sets (COS),^{18,19} where agreed upon minimum sets of outcomes when reporting research are expected, it is important that PROMs are appraised for their validity. If validation, or relevant items for the condition of interest are lacking, it is important that this is identified and rectified before inclusion in a COS.

The objectives of this systematic review are therefore to: (i) identify PROMs that have been designed for and/or validated in patients with facial skin cancer; (ii) assess the methodological quality of the included studies; (iii) assess the psychometric

properties of those identified PROMs; (iv) make an assessment of the focus of each PROM on the reconstructive aspect of patient care; and (v) make recommendations that could lead to the development of a facial skin cancer COS.

Methods

Search strategy and selection criteria

A systematic review protocol was developed in accordance with the Preferred Reporting for Items for Systematic Reviews and Meta-Analyses-Protocols (PRISMA-P)^{20,21} and registered with PROSPERO (CRD42016043181).

The search strategy was constructed in line with PRISMA guidelines,²² the Cochrane handbook²³ and guidance from Terwee *et al.*²⁴ To identify all papers that discussed some aspect of PROM development or validation for facial skin cancer, three separate constructs were explored: target condition, target body area and measurement instrument (e.g. PROM). Keywords and MeSH terms were selected where available and searches were performed in: MEDLINE (Ovid), Embase (Ovid), PsycINFO (Ovid), Cochrane and CINAHL (EBSCO). An example search strategy can be seen in Supplementary Figure S1 (see Supporting Information). Grey literature and reference lists were also searched using Google, Google Scholar and known PROMs-based websites. Searches were performed

Table 1 Inclusion and exclusion criteria used when screening identified studies

Inclusion criteria	1	Head and neck skin cancer population
	2	Papers discussing some aspect of PROM development or validation
	3	English-only articles
Exclusion criteria	1	Questionnaires not developed or validated in patients with head and neck skin cancer
	2	Oropharyngeal head and neck cancer population
	3	Questionnaires developed to assess nodal or distant metastatic disease
	4	General oncology questionnaires unless specifically validated in a head and neck skin cancer population
PROM, patient-reported outcome measure		

by two independent researchers (T.D.D. and H.S.) on the same day in August 2016, with results uploaded to the reference management software package, EndNote® Version X7 (Clarivate Analytics, Philadelphia, PA, U.S.A.). The search strategy was re-run prior to submission in January 2018 to identify any further studies that matched the inclusion criteria. Duplicates were removed using the functionality in EndNote®, with all references transferred to the online programme Covidence (www.covidence.org) for title and abstract screening. References were screened by two independent reviewers (T.D.D. and H.S.) according to the inclusion and exclusion criteria (Table 1), with all remaining articles downloaded in full-text format and re-screened. Discrepancies were discussed between the two reviewers with a third reviewer (H.A.H.) consulted if required.

Assessment of the methodological quality of included studies

The methodological quality of included studies was assessed using the CONsensus-based Standards for the Selection of Health Measurement INstruments (COSMIN).^{25,26} The COSMIN checklist contains nine main sections, each assessing a different measurement property: internal consistency, reliability, measurement error, content validity, construct validity (structural validity and hypothesis testing), cross-cultural validity, criterion validity and responsiveness. An updated checklist with a four-category rating scale (4, excellent; 3, good; 2, fair; 1, poor) was used.²⁷ Each paper included in the review was compared against the 98 items in the checklist, and for those where evidence was presented in the paper, a score on the four-category scale was given. One is able to assess criterion validity only where the PROM in question was compared with a longer version. Any paper describing criterion validity but not actually assessing against a 'gold standard' or long version was not assessed in the criterion validity category of COSMIN. The final rating for methodological quality in any given area of assessment is considered to be the lowest score (i.e. if a property such as internal consistency is scored 'excellent' in one question, but 'poor' in another, the methodological quality for that property is considered to be 'poor').

The COSMIN checklist has good interrater agreement and reliability;²⁸ however, to account for bias and subjectivity when rating studies it is considered good practice to compare results between two independent reviewers. A randomly selected sample of 30% of the included studies were assessed by two reviewers (T.D.D. and S.H.) and compared using intra-class coefficient (ICC),²⁹ Cohen's κ ³⁰ and percentage agreement. If agreement was low in this sample, all included studies would be doubly assessed.

Assessment of psychometric properties

The psychometric quality of each PROM was assessed using criteria developed by Terwee *et al.*³¹ and updated in 2016.³² Supplementary Figure S2 (see Supporting Information)

describes the measurement properties that are assessed according to these criteria. Each criterion is rated as criteria met (+), criteria not met (−) or not all information present (?).

Data analysis and best-evidence synthesis

Data were collated in Excel for Mac (V14.5.7) and presented as tables and narrative synthesis. Interrater reliability statistics were calculated for the COSMIN analysis using the Statistical Package for the Social Sciences (SPSS) software V.22 (IBM Corp., New York, NY, U.S.A.).

A best-evidence synthesis was performed by applying the levels of evidence summary as described by Furlan *et al.*³³ to the combined results of the COSMIN and Terwee *et al.*³¹ assessments. The Outcome Measures in Rheumatology (OMERACT) criteria were then used to categorize each instrument into (A) instrument meets all requirements and is recommended for use; (B) instrument meets two or more required items and therefore has potential for use; (C) instrument has low quality in at least one area and is not recommended for use; and (D) instrument has almost no validation.³⁴ This method has previously been used by Gerbens *et al.* in the dermatology literature.³⁵

Assessment of reconstructive relevance

The focus of each PROM on reconstruction post-skin cancer has never been assessed before and therefore there is no framework to work from. We thus performed a subjective assessment of the included questions based on specialist knowledge of the topic area by the authors. As a reconstructive PROM was not the aim of the original scale developers, we have performed this assessment separately and did not let this influence the COSMIN analysis when judging content validity.

Results

Two reviewers independently reviewed 4886 articles. With the addition of articles identified during reference searching a total of 24 studies were finally included (Fig. 1).^{7,36–58} Of those articles included, 11 different PROMs were identified: two generic PROMs (SF-36 and FACT-G) and nine skin cancer-specific (FACT-M, POS-H/N, SCI, SCQoL, aBCCdex, SCQOLIT, FACE-Q, DLQI, Essers *et al.*). As per the inclusion criteria, all included PROMs demonstrated some aspect of validation in the facial skin cancer population. A summary of identified PROMs and included papers describing aspects of design or validation are presented in Table 2.^{7,36–58} A more detailed assessment of each instrument is presented in Supplementary Results S1 (see Supporting Information).

Methodological quality of those included studies

Raw individual category scores for each PROM are presented in Table 3.^{36–38,40–58} Of the 11 PROMs included, there was a range of methodological quality, with only one paper scoring in all eight of the COSMIN categories (FACT-M). The spread of




Fig 1. PRISMA (Preferred Reporting for Items for Systematic Reviews and Meta-Analyses) flow diagram demonstrating the identification and screening of studies for inclusion.

Table 2 Summary of patient-reported outcome measures (PROMs) and corresponding papers identified using the inclusion and exclusion criteria of this systematic review. The number of items in each questionnaire and domains assessed are documented

PROM	Papers included	Generic or condition-specific	Number of items	Domains
SF-36	Rhee 2003 ⁴³	Generic	36	<ul style="list-style-type: none"> • Vitality • Physical functioning • Bodily pain • General health perception • Physical role functioning • Emotional role functioning • Social role functioning • Mental health
FACT-G	Rhee 2003 ⁴³	Generic	27	<ul style="list-style-type: none"> • Physical • Social/family • Emotional • Functional well-being
FACT-M	Cormier 2005 ³⁷ Cormier 2008 ³⁸ Askew 2009 ³⁹ Swartz 2012 ⁴⁰ Winstanley 2013 ⁴¹	Condition-specific	24 (in FACT-M subscale) 18 in reduced version	<ul style="list-style-type: none"> • Physical well-being • Emotional well-being • Social well-being
POS-H/N	Cano 2006 ⁴²	Condition-specific	15 (6 preoperatively and 9 postoperatively)	<ul style="list-style-type: none"> • Psychological functioning and cosmetic appearance • Satisfaction
SCI	Rhee 2005 ³⁶ Matthews 2006 ⁴⁴ Rhee 2006 ⁴⁵ Rhee 2007 ⁴⁶ de Troya-Martín 2015 ⁴⁷ Körner 2016 ⁷	Condition-specific	15	<ul style="list-style-type: none"> • Emotion • Social • Appearance
SCQoL	Vinding 2013 ⁴⁸ Vinding 2014 ⁴⁹	Condition-specific	9	<ul style="list-style-type: none"> • Function • Emotions • Control
aBCCdex	Mathias 2014 ⁵⁰ Mathias 2015 ⁵¹	Condition-specific	26	<ul style="list-style-type: none"> • Worry about future lesions • Mental health • Social/Relationships • Lesion symptoms • Life impact
SCQOLIT	Burdon-Jones 2010 ⁵² Burdon-Jones 2013 ⁵³	Condition-specific	10	<ul style="list-style-type: none"> • Psychosocial • Physical
FACE-Q	Lee 2015 ⁵⁴	Condition-specific	N/A	N/A
DLQI	Finlay 1994 ⁵⁵ Blackford 1996 ⁵⁶	Generic skin PROM	10	<ul style="list-style-type: none"> • Symptoms and feelings • Daily activities • Leisure • Work and school • Personal relationships • Treatment
Essers <i>et al.</i>	Essers 2006 ⁵⁷ Essers 2007 ⁵⁸	Condition-specific	22	<ul style="list-style-type: none"> • Worrying about facial health • Susceptibility for facial BCC • Fear of developing a new BCC

aBCCdex, Advanced Basal Cell Carcinoma Index; BCC, basal cell carcinoma; DLQI, Dermatology Life Quality Index; FACE-Q, evaluates outcomes from facial cosmetic procedures; FACT-G, Functional Assessment of Cancer Therapy – General; FACT-M, Functional Assessment of Cancer Therapy – Melanoma; POS-H/N, Patient Outcome of Surgery – Head/Neck; SCI, Skin Cancer Index; SCQoL, Skin Cancer Quality of Life; SCQOLIT, Skin Cancer Quality of Life Impact Tool; SF-36, 36-Item Short Form Health Survey

Table 3 Individual category scores for each study for all included patient-reported outcome measures (PROMs) as assessed by the COSMIN four-point scale. Each domain is made up of a number of questions as part of the COSMIN checklist, with the lowest-scoring category representing the overall methodological quality for that domain in the paper assessed

PROM	Paper	Internal consistency	Reliability	Measurement error	Content validity	Structural validity	Hypotheses testing	Criterion validity	Responsiveness
SF-36	Rhee 2003 ⁴³	Poor							
FACT-G	Rhee 2003 ⁴³	Poor							
FACT-M	Cormier 2005 ³⁷				Excellent				
	Cormier 2008 ³⁸	Poor	Poor	Poor	Poor	Poor	Good	Poor	Fair
	Winstanley 2013 ⁴¹	Excellent				Excellent			
	Swartz 2012 ⁴⁰	Excellent	Poor			Excellent			
POS-H/N	Cano 2006 ⁴²	Excellent	Poor	Excellent	Excellent		Poor		Poor
FSCI/SCI	Rhee 2005 ³⁶	Poor			Excellent				
	Matthews 2006 ⁴⁴	Poor	Poor		Excellent	Poor			
	Rhee 2006 ⁴⁵	Good					Good		
	Rhee 2007 ⁴⁶								Fair
	de Troya-Martín 2015 ⁴⁷	Good	Good		Excellent	Good	Poor		
SCQoL	Vinding 2013 ⁴⁸ (IRT ^a)	Poor			Excellent	Excellent	Fair		
	Vinding 2014 ⁴⁹						Fair		Fair
aBCCdex	Mathias 2014 ⁵⁰				Excellent				
	Mathias 2015 ⁵¹	Good	Good			Good	Fair		Poor
SCQOLIT	Burdon-Jones 2010 ⁵²				Excellent				
	Burdon-Jones 2013 ⁵³	Good	Good	Poor	Excellent	Poor	Good		Poor
FACE-Q (SCM)	Lee 2015 ⁵⁴								
DLQI	Finlay, 1994 ⁵⁵		Fair		Excellent	Poor			
	Blackford 1996 ⁵⁶						Poor		
Essers <i>et al.</i>	Essers 2006 ⁵⁷	Good			Fair	Fair			
	Essers 2007 ⁵⁸								

All domains are scored according to the COSMIN checklist with four-point scale;²⁷ potential categories include: excellent, good, fair and poor. (Blank) indicates domains not measured in a study. ^aRefers to the use of Item Response Theory, rather than Classical Test Theory. aBCCdex, Advanced Basal Cell Carcinoma Index; COSMIN, COnsensus-based Standards for the Selection of Health Measurement INstruments; DLQI, Dermatology Life Quality Index; FACE-Q, evaluates outcomes from facial cosmetic procedures; FACT-G, Functional Assessment of Cancer Therapy – General; FACT-M, Functional Assessment of Cancer Therapy – Melanoma; FSCI, Facial Skin Cancer Index; POS-H/N, Patient Outcome of Surgery – Head/Neck; SCI, Skin Cancer Index; SCM, skin cancer module; SCQoL, Skin Cancer Quality of Life; SCQOLIT, Skin Cancer Quality of Life Impact Tool; SF-36, 36-Item Short Form Health Survey

ratings between the four categories (excellent, good, fair and poor) was relatively even, with 28% being 'excellent', 18% 'good', 14% 'fair' and 40% 'poor'. The content validity for all but one condition-specific PROMs (Essers *et al.*^{57,58}) demonstrated 'excellent' methodology. Of the other categories, internal consistency and structural validity are the next two most commonly reported on and appropriately investigated areas of PROM development and validation in the identified studies.

An ICC of 0.844 (0.796–0.88), κ of 0.648 ($P < 0.005$) and a percentage agreement of 97.84% were observed between the two reviewers, demonstrating good agreement.

Psychometric properties of included patient-reported outcome measures

The results of the psychometric evaluation are shown in Table 4. Of the 11 PROMs assessed, none scored positively in all domains. The PROMs with the lowest-scoring psychometric measurement properties as assessed using criteria produced by

Terwee *et al.*³¹ were SF-36, FACT-G and FACE-Q skin cancer module. The FACE-Q skin cancer module was described in outline only in one paper;⁵⁴ hence the scores noted in Table 4.

Content validity and internal consistency are the two most commonly reported on and well-designed aspects of PROMs validation papers. Seven out of nine condition-specific PROMs showed 'appropriate assessment of content validity', demonstrating appropriate use of commonly used methods to generate items specific to the patient group.⁵⁹ Good internal consistency, as demonstrated by having a Cronbach's α of 0.70–0.95, was shown in eight of the nine condition-specific PROMs.

However, the presentation of data in the included studies required to assess the other criteria of Terwee *et al.*³¹ was more sporadic.

Best-evidence synthesis

A summary of the best-evidence synthesis using the method outlined can be seen in Table 5. Using the OMERACT filter no

Table 4 Individual category scores assessing psychometric properties for each study for all included patient-reported outcome measures (PROMs) as developed by Terwee *et al.*^{31,32}

PROM	Paper	Internal consistency	Reliability	Measurement error	Content validity	Structural validity	Hypotheses testing	Criterion validity	Responsiveness
SF-36	Rhee 2003 ⁴³	—							
FACT-G	Rhee 2003 ⁴³	—							
FACT-M	Cormier 2005 ³⁷				+				
	Cormier 2008 ³⁸	+	?	?	+	?	+	?	—
	Winstanley 2013 ⁴¹	+				+			
	Swartz 2012 ⁴⁰	—	?			+			
POS-H/N	Cano 2006 ⁴²	+	+	?	+		+		+
FSCI/SCI	Rhee, 2005 ³⁶	?			+				
	Matthews 2006 ⁴⁴	+	?		+	—			
	Rhee 2006 ⁴⁵	+					+		
	Rhee 2007 ⁴⁶								+
	de Troya-Martín 2015 ⁴⁷	—	+		+	?	+		
SCQoL	Vinding 2013 ⁴⁸ (IRT) ^a	+			+	+	—		
	Vinding 2014 ⁴⁹						—		—
aBCCdex	Mathias 2014 ⁵⁰				+				
	Mathias 2015 ⁵¹	+	—			+	+		+
SCQOLIT	Burdon-Jones 2010 ⁵²				+				
	Burdon-Jones 2013 ⁵³	+	+	?	+	?	+		?
FACE-Q (SCM)	Lee 2015 ⁵⁴								
DLQI	Finlay 1994 ⁵⁵		?		+	?			
	Blackford 1996 ⁵⁶						+		
Essers <i>et al.</i>	Essers 2006 ⁵⁷	+			+	?			
	Essers 2007 ⁵⁸								

Each criterion is assessed as either: positive rating (+), negative rating (–) or indeterminate rating (?). (Blank) indicates domains not measured or where no evidence is presented. ^aRefers to the use of Item Response Theory, rather than Classical Test Theory. aBCCdex, Advanced Basal Cell Carcinoma Index; DLQI, Dermatology Life Quality Index; FACE-Q, evaluates outcomes from facial cosmetic procedures; FACT-G, Functional Assessment of Cancer Therapy – General; FACT-M, Functional Assessment of Cancer Therapy – Melanoma; FSCI, Facial Skin Cancer Index; POS-H/N, Patient Outcome of Surgery – Head/Neck; SCI, Skin Cancer Index; SCM, skin cancer module; SCQoL, Skin Cancer Quality of Life; SCQOLIT, Skin Cancer Quality of Life Impact Tool; SF-36, 36-Item Short Form Health Survey

PROMs met the criteria for an 'A'-graded PROM, four PROMs were considered to be a 'B'-graded PROM, four were 'C'-grade PROMs, two were 'D'-grade PROMs and one was not gradeable.

Focus on reconstructive aspects in each questionnaire

An assessment of the questions included in each questionnaire was made for their relevance to and focus on the reconstructive aspects and cosmesis of facial skin cancer. A summary of the questions that hold some relevance to reconstruction for each PROM is shown in Table 6.

Discussion

This systematic review has been designed to identify all PROMs that are validated for use in patients with facial skin cancer. At a time when the use of PROMs is being encouraged in both research and clinical use, it is important that only those PROMs that show evidence of validation are used. In the ideal world these would be validated in the exact population in which they were being implemented; however, in practice

this is often too time-consuming and expensive. Previous systematic reviews on this topic^{16,17} have demonstrated many similar PROMs to this review; however, we have assessed the methodological quality of these studies using internationally accepted criteria to minimize the risk of bias. This was performed using the COSMIN checklist,²⁶ which is now routinely accepted across the systematic review literature and has been used extensively in orthopaedics,⁶⁰ paediatrics,⁶¹ neurology⁶² and dermatology.³⁵ A further update to the COSMIN methodology has been published, although this was after this review was performed.⁶³ We also assessed the quality of the psychometric properties of the included PROMs using the criteria of Terwee *et al.*³¹ and performed a best-evidence synthesis.

Of the two generic instruments identified, SF-36 and FACT-G, only rudimentary validation was provided in one paper.⁴³ Both instruments are well established in the literature for their general use; however, due to poor evidence of validation in the facial skin cancer population their use in this setting is difficult to recommend. This is mainly due to the instruments initially being designed for a different population to the one studied here and therefore they lack face and content validity. For example, the issues affecting a facial skin cancer population are likely

Table 5 Best-evidence synthesis and grading according to the OMERACT filter

PROM	Internal consistency	Reliability	Measurement error	Content validity	Structural validity	Hypothesis testing	Criterion validity	Responsiveness	OMERACT recommendation
SF-36	?								D
FACT-G	?								D
FACT-M	±	?	?	±	±	++	?	–	C
POS-H/N	+++	?	±	+++		?		?	B
SCI	±	±		+++	±	±		+	B
SCQoL	?			+++	+++	–		–	C
aBCCdex	++	– –		+++	++	+		?	C
SCQOLIT	++	++	?	+++	?	++		?	B
FACE-Q									N/A
DLQI		–		+++	?	?			C
Essers <i>et al.</i>	++			+	±				B

Positive rating for measurement property (+++, consistent findings in multiple studies of good methodological quality or one study of excellent quality; ++, consistent findings in multiple studies of fair methodological quality or one good study; +, one study of fair methodological quality). Negative rating for measurement property (– –, consistent findings in multiple studies of good methodological quality or one study of excellent quality; –, consistent findings in multiple studies of fair methodological quality or one good study; –, one study of fair methodological quality). ?, indeterminate due to poor quality study; ±, conflicting evidence. OMERACT filter using categories of A, B, C, D as discussed in the Methods. aBCCdex, Advanced Basal Cell Carcinoma Index; DLQI, Dermatology Life Quality Index; FACE-Q, evaluates outcomes from facial cosmetic procedures; FACT-G, Functional Assessment of Cancer Therapy – General; FACT-M, Functional Assessment of Cancer Therapy – Melanoma; OMERACT, Outcome Measures in Rheumatology; POS-H/N, Patient Outcome of Surgery – Head/Neck; SCI, Skin Cancer Index; SCQoL, Skin Cancer Quality of Life; SCQOLIT, Skin Cancer Quality of Life Impact Tool; SF-36, 36-Item Short Form Health Survey

to be very different from those affecting the population groups used to design the SF-36. There was a range of quality with respect to design and validation across the nine condition-specific PROMs identified. Internal consistency was the most commonly reported area of validation across all PROMs, with measurement error and criterion validity least commonly reported on. Unfortunately, measurement error is an important concept required to design high-quality prospective studies using these instruments. Evidence for content validity was excellent in all but one condition-specific PROM (Essers *et al.*^{57,58}), with all condition-specific PROMs attempting to include representative patients in their design and validation.

Combining the results of the COSMIN and Terwee *et al.*³¹ analysis into a best-evidence synthesis identified four PROMs that are currently the most appropriate for inclusion in a COS for facial skin cancer: POS-H/N, SCI, SCQOLIT and Essers *et al.*^{57,58} However, all of these still have deficiencies in their validation (see Table 5) and further studies are advised. Furthermore, the FACE-Q skin cancer module has the potential to be a well-designed and validated instrument, but further studies are awaited.

This is the first systematic review on the subject to assess each PROM for their focus on the post-resection reconstruction of facial skin cancer. The results show that this is poorly addressed, even in PROMs designed specifically for facial skin cancer. Questions relating to the degree of scarring, how noticeable it is, physical symptoms such as pain and itch and psychological concerns all featured, but no single instrument adequately addressed this area. This is an important finding. In an era where skin cancer is treatable the long-term sequelae of the treatment given is important, especially where this results

in visible and potentially disfiguring scarring on the face. The only way in which the medical community will be able to improve the treatment offered is by asking patients what they think, through the medium of PROMs. Therefore, it is important that PROMs exist which include relevant and valid items relating to issues such as the reconstruction if they are to be included in a facial skin cancer COS. A COS for basal cell carcinomas is already in creation⁶⁴ and the CSG-COUSIN (Cochrane Skin Group Core Outcome Set Initiative) group⁶⁵ plan many more in the dermatology world. We therefore hope and implore that these take into account areas such as aesthetic and functional outcomes of reconstructive surgery.

The use of the COSMIN checklist is a strength of this study; however, despite being validated and well accepted in the literature, there are limitations associated with it. Firstly, scoring of each item in the checklist is reliant on author judgment and therefore can be subjective. Secondly, the checklist is extensive and while this means it is considered to be the 'gold-standard' it is potentially difficult for the non-health outcome specialist to use.

In this systematic review we tried to control for interrater reliability issues by two independent reviewers assessing a randomly selected selection of papers. An ICC score of 0.844 (considered 'good' by Koo and Li⁶⁶), κ statistic of 0.648 ($P < 0.005$) ('moderate agreement'⁶⁷) and percentage agreement of 97.84% validated our interrater reliability and therefore COSMIN scores. While this assessment provides some reassurance when using COSMIN, we appreciate that it is feasible that another review team may score items differently.

Another strength of this systematic review is the use of a validated and highly sensitive search strategy, using guidance

Table 6 Assessment of each questionnaire for a focus on questions relating to reconstruction and post-treatment aesthetics

PROM	Questions with a focus relevant to reconstruction	Global summary of focus on reconstruction
SF-36	No questions relevant to reconstruction	Absent
FACT-G	No questions relevant to reconstruction	Absent
FACT-M	Four items show some relevance: <ul style="list-style-type: none"> • I feel numbness at my surgical site • I have pain at my melanoma site or surgical scar • I worry about the appearance of surgical scars • I have swelling as a result of surgery 	Poor
POS-H/N	Postsurgical questionnaire attempts to address aspects of the operation and outcomes: <ul style="list-style-type: none"> • Are the results of the operation on your head/neck skin growths better/about the same/worse than expected? • If a friend has a similar head/neck skin growth(s) that you had before your operation would you recommend the same operation you had? 	Average
SCI	Two items relating to scarring: <ul style="list-style-type: none"> • Worried about how large the scar will be? • Thought about how noticeable the scar will be to others? 	Average
SCQoL	No focus on the treatment or reconstructive aspect. One question with a vague reference to aesthetics: <ul style="list-style-type: none"> • During the past week, I have used such things as make-up or clothing to hide my skin cancer from others 	Poor
aBCCdex	Items relevant to appearance: <ul style="list-style-type: none"> • Your appearance changing due to surgery or procedures Three items relating to the lesion: <ul style="list-style-type: none"> • Bleeding from lesion(s) • Oozing or pus from lesions(s) • Sensitive/tender skin around lesion(s) However, no questions with a focus on the reconstruction	Poor
SCQOLIT	One item relating to disfigurement and one relating to discomfort following the treatment: <ul style="list-style-type: none"> • Over the last week, how much have you been bothered about any disfigurement or scarring, in respect to your skin cancer or its treatment? • Over the last week, how much skin discomfort or inconvenience have you experienced, in respect to your skin cancer or its treatment? 	Poor/Average
FACE-Q	No specific questionnaire items have yet to be published but one of the aims of the new skin cancer module is to address areas around facial aesthetics	Absent
DLQI	No questions relevant to reconstruction	Absent
Essers <i>et al.</i>	No questions relevant to reconstruction	Absent

aBCCdex, Advanced Basal Cell Carcinoma Index; DLQI, Dermatology Life Quality Index; FACE-Q, evaluates outcomes from facial cosmetic procedures; FACT-G, Functional Assessment of Cancer Therapy – General; FACT-M, Functional Assessment of Cancer Therapy – Melanoma; POS-H/N, Patient Outcome of Surgery – Head/Neck; PROM, patient-reported outcome measure; SCI, Skin Cancer Index; SCQoL, Skin Cancer Quality of Life; SCQOLIT, Skin Cancer Quality of Life Impact Tool; SF-36, 36-Item Short Form Health Survey

from the Cochrane group²³ and Terwee *et al.*²⁴ We used a broad search strategy to identify all relevant studies demonstrating some aspect of design or validation of a PROM for facial skin cancer. However, this could also be a limitation in that we included only those studies that demonstrated aspects of design or validation. Studies that used a PROM in the facial skin cancer population but did not assess validation were excluded, potentially missing PROMs, which if they were validated, might be useful in this population group.

This systematic review has identified a number of different PROMs relevant to the facial skin cancer population. The identified PROMs demonstrated variable psychometric validation and all poorly addressed the reconstructive aspects of facial skin cancer. While POS-H/N, SCI, SCQOLIT and Essers *et al.*^{57,58} all show potential, further validation work is required before they could be confidently included in a COS.

In order to move forward and improve our understanding of patients' views on facial skin cancer and the difference

between treatment options, it is important that these deficiencies in validation studies are addressed. Furthermore, additional items, either as an addition to a current PROM or included in an entirely new PROM, are required to specifically address the reconstruction and aesthetic outcomes of facial skin cancer. It is hoped that in time the tools will exist to confidently assess our patients' views on their facial skin cancer and treatment outcomes, reducing the psychological and social burden associated with this disease.

References

- Geller AC, Annas GD. Epidemiology of melanoma and non-melanoma skin cancer. *Semin Oncol Nurs* 2003; **19**:2–11.
- The Skin Cancer Foundation. Understanding skin cancer. Available at: www.skincancer.org (last accessed 23 November 2017).
- Lomas A, Leonardi-Bee J, Bath-Hextall F. A systematic review of worldwide incidence of nonmelanoma skin cancer. *Br J Dermatol* 2012; **166**:1069–80.
- Franceschi S, Levi F, Randimbison L, La Vecchia C. Site distribution of different types of skin cancer: new aetiological clues. *Int J Cancer* 1996; **67**:24–8.
- Cancer Research UK statistics. 2016 Apr 7; 1–1. Available at: <http://www.cancerresearchuk.org/health-professional/cancer-statistics/statistics-by-cancer-type/skin-cancer/incidence> (last accessed 15 October 2017).
- Weinstock MA, Bogaars HA, Ashley M *et al.* Nonmelanoma skin cancer mortality. A population-based study. *Arch Dermatol* 1991; **127**:1194–7.
- Körner A, Garland R, Czajkowska Z *et al.* Supportive care needs and distress in patients with non-melanoma skin cancer: nothing to worry about? *Eur J Oncol Nurs* 2016; **20**:150–5.
- Sobanko JF, Sarwer DB, Zvargulis Z, Miller CJ. Importance of physical appearance in patients with skin cancer. *Dermatol Surg* 2015; **41**:183.
- Karimi M, Brazier J. Health, health-related quality of life, and quality of life: what is the difference? *Pharmacoeconomics* 2016; **34**:645–9.
- U.S. Department of Health and Human Services FDA Center for Drug Evaluation and Research; U.S. Department of Health and Human Services FDA Center for Biologics Evaluation and Research; U.S. Department of Health and Human Services FDA Center for Devices and Radiological Health. Guidance for industry: patient-reported outcome measures: use in medical product development to support labeling claims: draft guidance. *Health Qual Life Outcomes* 2006; **4**:79.
- McGrail K, Bryan S, Davis J. Let's all go to the PROM: the case for routine patient-reported outcome measurement in Canadian healthcare. *Healthc Pap* 2011; **11**:8–18.
- Devlin NJ, Appleby J. Getting the most out of PROMS. Putting health outcomes at the heart of NHS decision-making. *Health Economics* 2010. Available at: <http://www.kingsfund.org.uk/publications/proms.html> (last accessed 23 November 2017).
- Kotronoulas G, Kearney N, Maguire R *et al.* What is the value of the routine use of patient-reported outcome measures toward improvement of patient outcomes, processes of care, and health service outcomes in cancer care? A systematic review of controlled trials. *J Clin Oncol* 2014; **32**:1480–501.
- Brass D, Fouweather T, Stocken DD *et al.* An observer-blinded randomized controlled pilot trial comparing localized immersion psoralen–ultraviolet A with localized narrowband ultraviolet V for the treatment of palmar hand eczema. *Br J Dermatol* 2018; **179**:63–71.
- Lee EH, Klassen AF, Nehal KS *et al.* A systematic review of patient-reported outcome instruments of nonmelanoma skin cancer in the dermatologic population. *J Am Acad Dermatol* 2013; **69**:e59–67.
- Gibbons E, Casañas i Comabella C, Fitzpatrick R. A structured review of patient-reported outcome measures for patients with skin cancer, 2013. *Br J Dermatol* 2013; **168**:1176–86.
- Bates AS, Davis CR, Takwale A, Knevil GJ. Patient-reported outcome measures in nonmelanoma skin cancer of the face: a systematic review. *Br J Dermatol* 2013; **168**:1187–94.
- Williamson P, Altman D, Blazeby J *et al.* Driving up the quality and relevance of research through the use of agreed core outcomes. *J Health Serv Res Policy* 2012; **17**:1–2.
- Porter ME, Larsson S, Lee TH. Standardizing patient outcomes measurement. *N Engl J Med* 2016; **374**:504–6.
- Moher D, Shamseer L, Clarke M *et al.*; PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015 statement. *Syst Rev* 2015; **4**:1.
- Shamseer L, Moher D, Clarke M *et al.*; PRISMA-P Group. Preferred reporting items for systematic review and meta-analysis protocols (PRISMA-P) 2015: elaboration and explanation. *BMJ* 2015; **350**:g7647.
- Moher D, Liberati A, Tetzlaff J, Altman DG. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. *Ann Intern Med* 2009; **151**:264–9.
- Higgins JPT, Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. Version 5-1. 0 [updated March 2011]. The Cochrane Collaboration; 2011. Available at: <http://handbook-5-1.cochrane.org/> (last accessed 14 November 2018).
- Terwee CB, Jansma EP, Riphagen II, de Vet HC. Development of a methodological PubMed search filter for finding studies on measurement properties of measurement instruments. *Qual Life Res* 2009; **18**:1115–23.
- Mokkink LB, Terwee CB, Patrick DL *et al.* The COSMIN checklist for assessing the methodological quality of studies on measurement properties of health status measurement instruments: an international Delphi study. *Qual Life Res* 2010; **19**:539–49.
- Mokkink LB, Terwee CB, Patrick DL *et al.* The COSMIN study reached international consensus on taxonomy, terminology, and definitions of measurement properties for health-related patient-reported outcomes. *J Clin Epidemiol* 2010; **63**:737–45.
- Terwee CB, Mokkink LB, Knol DL *et al.* Rating the methodological quality in systematic reviews of studies on measurement properties: a scoring system for the COSMIN checklist. *Qual Life Res* 2012; **21**:651–7.
- Mokkink LB, Terwee CB, Gibbons E *et al.* Inter-rater agreement and reliability of the COSMIN (Consensus-based Standards for the selection of health status Measurement INstruments) checklist. *BMC Med Res Methodol* 2010; **10**:82.
- Shrout PE, Fleiss JL. Intraclass correlations: uses in assessing rater reliability. *Psychol Bull* 1979; **86**:420–8.
- Cohen J. A coefficient of agreement for nominal scales. *Educ Psychol Meas* 1960; **20**:37–46.
- Terwee CB, Bot SDM, de Boer MR *et al.* Quality criteria were proposed for measurement properties of health status questionnaires. *J Clin Epidemiol* 2007; **60**:34–42.
- Prinsen CAC, Vohra S, Rose MR *et al.* How to select outcome measurement instruments for outcomes included in a 'Core Outcome Set' – a practical guideline. *Trials* 2016; **17**:449.
- Furlan AD, Pennick V, Bombardier C *et al.* Updated method guidelines for systematic reviews in the Cochrane Back Review Group. *Spine* 2009; **34**:1929–41.
- Boers M, Brooks P, Strand CV, Tugwell P. The OMERACT filter for outcome measures in rheumatology. *J Rheumatol* 1998; **25**:198–99.

- 35 Gerbens LAA, Prinsen CAC, Chalmers JR *et al.* Evaluation of the measurement properties of symptom measurement instruments for atopic eczema: a systematic review. *Allergy* 2017; **72**:146–63.
- 36 Rhee JS, Matthews BA, Neuburg M *et al.* Creation of a quality of life instrument for nonmelanoma skin cancer patients. *Laryngoscope* 2005; **115**:1178–85.
- 37 Cormier JN, Davidson L, Xing Y *et al.* Measuring quality of life in patients with melanoma: development of the FACT-melanoma subscale. *J Support Oncol* 2005; **3**:139–45.
- 38 Cormier JN, Ross MI, Gershenwald JE *et al.* Prospective assessment of the reliability, validity, and sensitivity to change of the Functional Assessment of Cancer Therapy-Melanoma questionnaire. *Cancer* 2008; **112**:2249–57.
- 39 Askew RL, Xing Y, Palmer JL *et al.* Evaluating minimal important differences for the FACT-Melanoma quality of life questionnaire. *Value Health* 2009; **12**:1144–50.
- 40 Swartz RJ, Baum GP, Askew RL *et al.* Reducing patient burden to the FACT-Melanoma quality-of-life questionnaire. *Melanoma Res* 2012; **22**:158–63.
- 41 Winstanley JB, Saw R, Boyle F, Thompson J. The FACT-Melanoma Quality-of-Life instrument: comparison of a five-point and four-point response scale using the Rasch measurement model. *Melanoma Res* 2013; **23**:61–9.
- 42 Cano SJ, Browne JP, Lamping DL *et al.* The Patient Outcomes of Surgery-Head/Neck (POS-head/neck): a new patient-based outcome measure. *J Plast Reconstr Aesthet Surg* 2006; **59**:65–73.
- 43 Rhee JS, Loberiza FR, Matthews BA *et al.* Quality of life assessment in nonmelanoma cervicofacial skin cancer. *Laryngoscope* 2003; **113**:215–20.
- 44 Matthews BA, Rhee JS, Neuburg M *et al.* Development of the Facial Skin Care Index: a health-related outcomes index for skin cancer patients. *Dermatol Surg* 2006; **32**:924–34.
- 45 Rhee JS, Matthews BA, Neuburg M *et al.* Validation of a quality-of-life instrument for patients with nonmelanoma skin cancer. *Arch Facial Plast Surg* 2006; **8**:314–18.
- 46 Rhee JS, Matthews BA, Neuburg M *et al.* The Skin Cancer Index: clinical responsiveness and predictors of quality of life. *Laryngoscope* 2007; **117**:399–405.
- 47 de Troya-Martín M, Rivas-Ruiz F, Blázquez-Sánchez N *et al.* A Spanish version of the Skin Cancer Index: a questionnaire for measuring quality of life in patients with cervicofacial nonmelanoma skin cancer. *Br J Dermatol* 2015; **172**:160–8.
- 48 Vinding GR, Christensen KB, Esmann S *et al.* Quality of life in non-melanoma skin cancer – the Skin Cancer Quality of Life (SCQoL) questionnaire. *Dermatol Surg* 2013; **39**:1784–93.
- 49 Vinding GR, Esmann S, Olesen AB *et al.* Interpretation of the Skin Cancer Quality of Life score: a validated quality of life questionnaire for non-melanoma skin cancer. *Dermatology* 2014; **229**:123–9.
- 50 Mathias SD, Chren M-M, Colwell HH *et al.* Assessing health-related quality of life for advanced basal cell carcinoma and basal cell carcinoma nevus syndrome. *JAMA Dermatol* 2014; **150**:169–8.
- 51 Mathias SD, Chren MM, Crosby RD *et al.* Reliability and validity of the Advanced Basal Cell Carcinoma Index (aBCCdex). *Br J Dermatol* 2015; **173**:713–19.
- 52 Burdon-Jones D, Thomas P, Baker R. Quality of life issues in non-metastatic skin cancer. *Br J Dermatol* 2010; **162**:147–51.
- 53 Burdon-Jones D, Gibbons K. The Skin Cancer Quality of Life Impact Tool (SCQOLIT): a validated health-related quality of life questionnaire for non-metastatic skin cancers. *J Eur Acad Dermatol Venerol* 2013; **27**:1109–13.
- 54 Lee EH, Klassen AF, Lawson JL *et al.* Patient experiences and outcomes following facial skin cancer surgery: a qualitative study. *Australas J Dermatol* 2015; **57**:e100–4.
- 55 Finlay AY, Khan GK. Dermatology Life Quality Index (DLQI) – a simple practical measure for routine clinical use. *Clin Exp Dermatol* 1994; **19**:210–16.
- 56 Blackford S, Roberts D, Salek MS, Finlay A. Basal cell carcinomas cause little handicap. *Qual Life Res* 1996; **5**:191–4.
- 57 Essers BAB, Nieman FHM, Prins MH *et al.* Determinants of satisfaction with the health state of the facial skin in patients undergoing surgery for facial basal cell carcinoma. *Patient Educ Couns* 2006; **60**:179–86.
- 58 Essers B, Nieman F, Prins M *et al.* Perceptions of facial aesthetics in surgical patients with basal cell carcinoma. *J Eur Acad Dermatol Venerol* 2007; **21**:1209–14.
- 59 Streiner DL, Norman GR, Cairney J. *Health Measurement Scales*. Fifth edition. Oxford: Oxford University Press, 2015.
- 60 Gagnier JJ, Mullins M, Huang H *et al.* A systematic review of measurement properties of patient-reported outcome measures used in patients undergoing total knee arthroplasty. *J Arthroplasty* 2017; **32**:1688–97. e7.
- 61 Speyer R, Cordier R, Parsons L *et al.* Psychometric characteristics of non-instrumental swallowing and feeding assessments in pediatrics: a systematic review using COSMIN. *Dysphagia* 2017; **31**:1–14.
- 62 Haywood KL, Mars TS, Potter R *et al.* Assessing the impact of headaches and the outcomes of treatment: a systematic review of patient-reported outcome measures (PROMs). *Cephalalgia* 2018; **38**:1374–86.
- 63 Prinsen CAC, Mokkink LB, Bouter LM *et al.* COSMIN guideline for systematic reviews of patient-reported outcome measures. *Qual Life Res* 2018; **27**:1147–57.
- 64 Schlessinger DI, Iyengar S, Yanes AF *et al.* Development of a core outcome set for clinical trials in basal cell carcinoma: study protocol for a systematic review of the literature and identification of a core outcome set using a Delphi survey. *Trials* 2017; **18**:490.
- 65 Schmitt J, Deckert S, Alam M *et al.* Report from the kick-off meeting of the Cochrane Skin Group Core Outcome Set Initiative (CSG-COUSIN). *Br J Dermatol* 2016; **174**:287–95.
- 66 Koo TK, Li MY. A guideline of selecting and reporting intraclass correlation coefficients for reliability research. *J Chiropr Med* 2016; **15**:155–63.
- 67 McHugh ML. Interrater reliability: the kappa statistic. *Biochemia Medica* 2012; **22**:276–82.
- 68 Ware JE, Sherbourne CD. The MOS 36-item short-form health survey (SF-36). I. Conceptual framework and item selection. *Med Care* 1992; **30**:473–83.
- 69 Brazier JE, Harper R, Jones NM *et al.* Validating the SF-36 health survey questionnaire: new outcome measure for primary care. *BMJ* 1992; **305**:160–4.
- 70 McHorney CA, Ware JE, Raczek AE. The MOS 36-Item Short-Form Health Survey (SF-36): II. Psychometric and clinical tests of validity in measuring physical and mental health constructs. *Med Care* 1993; **31**:247–63.
- 71 Cella DF, Tulsky DS, Gray G *et al.* The Functional Assessment of Cancer Therapy scale: development and validation of the general measure. *J Clin Oncol* 1993; **11**:570–9.
- 72 Rhee JS, Matthews BA, Neuburg M *et al.* Skin cancer and quality of life: assessment with the Dermatology Life Quality Index. *Dermatol Surg* 2004; **30**:525–9.
- 73 Nunes DH, Fröde TS. Quality of life in basal cell carcinoma patients in Brazil: a pilot cross sectional study. *Dermatol Surg* 2013; **39**:620–6.
- 74 Janz NK, Becker MH. The health belief model: a decade later. *Health Educ Q* 1984; **11**:1–47.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's website:

Fig S1. Example search strategy used for MEDLINE (Ovid) between 1946 and August 2016.

Fig S2. Summary of psychometric properties assessed in quality criteria proposed by Terwee *et al.*³¹ and updated by Prinsen *et al.*³²

Results S1. Generic patient-reported outcome measures. Skin cancer-specific patient-reported outcome measures.^{7,36–58,68–74}

Powerpoint S1 Journal Club Slide Set.