

Cross-study synthetic-guided transcriptomics identifies thymic proliferative suppression and soleus metabolic remodeling in spaceflown mice

Organ-specific flight responses across NASA OSDR mouse transcriptomes

Jason Trinh

Space Life Sciences Training Program, NASA Ames Research Center, Moffett Field, California, USA

Correspondence: jasontrinh@berkeley.edu

Research manuscript draft for author review. Author list, acknowledgments, repository release URL, and archival DOI require final review before submission.

Abstract

Background: Mouse spaceflight studies provide access to tissues that cannot be sampled extensively from astronauts, but individual experiments are small and differ in design. We asked whether synthetic gene-expression models could help identify reproducible, organ-specific flight responses without treating generated profiles as new animals.

Methods: We assembled 1,610 mouse flight and ground-control bulk RNA-seq profiles through the NASA Open Science Data Repository API. A conditional diffusion model was pretrained on 17,244 tissue-diverse ARCHS4 mouse profiles after excluding every GEO series linked to OSDR, then adapted with tissue, flight status, study, and material-type conditioning. Evaluation proceeded from a pooled synthetic-data benchmark to tissue-specific selection among five downstream uses, real-only cross-study association testing, and complete-study transfer. Generated profiles could affect prediction or feature selection, but flight effects and false-discovery rates were estimated exclusively from real samples.

Results: One pooled augmentation policy was not useful: balanced accuracy was 0.754 with real-only training and 0.737 with real-plus-synthetic training. Tissue-specific selection nevertheless improved balanced accuracy and AUROC during development in thymus, spleen, skin, kidney, pooled skeletal muscle, soleus, and other tissues. Across canonical tissues and anatomical muscle groups, 49 real-data BH-FDR associations also entered stable synthetic-informed selection: 26 promoted and 23 reinforced. Whole-study transfer sharply narrowed this evidence. In OSD-457 thymus, synthetic-guided balanced accuracy increased from 0.500 to 0.833 and AUROC from 0.840 to 0.979; lung failed fixed confirmation, and an initial pooled-muscle augmentation gain did not generalize across 11 held-out accessions. Flight thymus showed lower expression of eight mitotic genes, including *Cdk1*, *Ccnb1*, *Ccnb2*, *Birc5*, and *Ube2c*, together with lower APC/C, G2/M-checkpoint, and DNA-replication programs. Anatomical separation of skeletal muscle exposed a soleus response involving lower *Bdh1*, *Ech1*, *Bnip3*, and *Decr1* with higher *Tpm1*. Kidney supplied a secondary promoted *Inpp4b* and reinforced *Slc37a4* result; spleen supplied a developmental adhesion and cytoskeletal hypothesis.

Conclusions: Synthetic-guided analysis added the most information when it prioritized genes for testing in real samples rather than increasing the apparent sample size. The findings support reduced thymic proliferative renewal, reinforce a soleus mitochondrial and contractile response, and nominate focused renal and spleen candidates for prospective testing.

Keywords: spaceflight; bulk RNA-seq; synthetic data; thymus; soleus; kidney; NASA OSDR; ARCHS4

Introduction

Spaceflight affects immune, musculoskeletal, metabolic, and barrier tissues through a combination of microgravity, radiation, confinement, altered nutrition, stress, and mission-specific procedures. Mouse flight experiments provide tissue access that is unavailable in astronauts, but their transcriptomic interpretation is difficult. Individual studies are small, missions differ in strain and duration, and condition labels can be entangled with study, material, genotype, or collection protocol. Pooling samples without preserving these design variables can convert study effects into apparent flight biology.

The NASA Open Science Data Repository (OSDR) now exposes sample metadata and processed assay data through a queryable biological API [1]. This makes it possible to assemble a cross-study cohort without relying on a precombined raw HDF5 object and to retain accession-level provenance. Public reference resources offer a second opportunity. ARCHS4 uniformly processes a large fraction of public human and mouse RNA-seq data [2], providing tissue-diverse reference profiles for pretraining models that would be underdetermined on OSDR alone.

Deep generative models can learn high-dimensional expression distributions. Conditional WGAN-GP models have reproduced tissue and cancer properties in GTEx and TCGA [3]. More recently, Lacan and colleagues adapted denoising diffusion probabilistic and implicit models to bulk transcriptomics and


reported strong gene-correlation, neighborhood, adversarial, and downstream classification metrics [4]. GeneJEPA instead learns masked-gene representations without reconstructing expression [5]. These approaches solve different problems: a generator can sample expression, whereas a representation learner needs an additional decoder or generative objective before it can do so.

Synthetic expression is commonly presented as a remedy for small sample size. Generated profiles, however, are not new biological replicates. We therefore separated four questions: whether one pooled augmentation strategy helped at all, whether different tissues benefited from different synthetic-data uses, whether prioritized genes were associated with flight in real samples, and whether a frozen policy transferred to an entirely excluded study.


Our primary biological questions were whether this approach could clarify the tissue response to spaceflight, whether anatomical separation would expose muscle-specific responses hidden by pooling, and whether the resulting signals complemented pathway-level findings from expiMap.

The clearest findings arose in thymus and soleus. Thymus supplied a held-out test in a study excluded from ARCHS4 pretraining, OSDR adaptation, and feature-policy development. Soleus supplied a cross-study metabolic signal that is biologically coherent but still needs confirmation in a newly collected or fully unseen study. Kidney, pooled skeletal muscle, spleen, skin, and adrenal gland supplied secondary developmental candidates. Results from the remaining tissues define the exploratory boundary of the approach.

A Data and conditional expression model



B Evaluation funnel



Synthetic use is selected per tissue; biological effects and FDR are always estimated from real OSDR samples.

Figure 1. Evaluation funnel. (A) A tissue-conditioned mouse expression model was pretrained with ARCHS4 and adapted to API-derived OSDR flight and ground-control profiles. (B) Evaluation moved from a pooled benchmark to tissue-specific synthetic-use selection, real-only association testing, and complete-study transfer. Biological effects and false-discovery rates always came from real OSDR samples.

Materials and methods

Data sources

The OSDR Biological Data API was used to identify *Mus musculus* bulk RNA-seq assays with flight or ground-control labels [1]. Tissue and material names were harmonized while preserving study provenance. The resulting cohort contained 1,610 biological profiles from 75 accessions: 835 flight and

775 ground control. Full-transcriptome expression was converted to transcripts per million before selecting a 974-gene mouse landmark panel. No raw integrated OSD R H5 file was used.

The local ARCHS4 mouse resource contained 997,515 public RNA-seq profiles [2]. All nine GEO series linked from the API-derived OSD R metadata were excluded before reference selection, removing 108 otherwise eligible profiles and preventing cross-resource sample overlap. A healthy-preferred, tissue-balanced subset of 17,244 profiles spanning 20 tissue classes was then used for model pretraining. Complete GEO series were assigned to one reference role, producing 10,150 training, 2,466 validation, and 4,628 test profiles with no series overlap. This backbone was used for both the held-out test and the broad development screens.

Table 1. Data scope.

Source	Profiles used	Biological scope	Role
ARCHS4 mouse v2.5	17,244	20 tissue classes	Mouse tissue pretraining
NASA OSD R API	1,610	75 accessions; 835 flight and 775 ground control	Spaceflight analysis

Conditional expression model

We reproduced the bulk-expression diffusion architecture of Lacan et al. [4], based on denoising diffusion and implicit sampling [13,14]. The ARCHS4-pretrained model was adapted to OSD R while representing tissue, flight status, study, and material type as separate conditions. This allowed the same model lineage to generate flight or ground-control profiles for represented biological and study contexts. A designated accession-excluded adaptation was used for the fixed lung/thymus test, while a separate broad adaptation supplied the repeated all-tissue development screens.

Generated profiles were checked for preservation of tissue identity, expression structure, diversity, and flight-related differences on withheld real profiles. The evaluation included correlation, neighborhood overlap, adversarial separability, and distributional distance [15,16]. The exact architecture, transformations, split construction, calibration, thresholds, and comparator-model results are reported in the supplementary methods.

Evaluation funnel and synthetic-guided analysis

Evaluation used four stages. First, a pooled locked benchmark compared real-only, generated-only, and real-plus-generated training. This asked whether one synthetic-data policy generalized across tissues. Second, each tissue was evaluated separately with five candidate uses: real-only training, generated-only training, real-plus-generated training, synthetic-guided feature ranking with a real-only classifier, and guided training with generated profiles assigned 0.05 total weight. Nested development splits withheld profiles but retained representation from the same accessions. These are within-study development results, not evidence of transfer to a new study. Synthetic attribution was retained only when the selected arm was nonworse than real-only training across balanced accuracy, AUROC, and average precision under the frozen eligibility rule.

Third, stable features from the selected tissue arm were compared with stable real-only features. Genes stable under both approaches were interpreted as reinforced. Genes stable only under the eligible synthetic-informed arm were interpreted as synthetic-promoted. "Synthetic-promoted" describes repeated feature selection; it does not mean that a gene was absent from real expression or biologically novel.

Fourth, whole-study transfer was tested by excluding complete OSD R accessions from generator adaptation and classifier development. This is stricter than holding out profiles from represented studies because the test study contributes no training context. An earlier accession-held-out augmentation screen and its expanded skeletal-muscle evaluation were retained as context, while the fixed lung and thymus experiment supplied the primary transfer test.

Flight-minus-ground effects were estimated within each OSD R study and then summarized with a random-effects model [17]. Within each tissue, the 974 gene-level meta-analysis P values were adjusted

with the Benjamini-Hochberg procedure, and BH FDR below 0.05 defined the primary statistical inclusion rule [6]. Benjamini-Hochberg correction controls the expected proportion of false discoveries among the reported genes while retaining more power than family-wise error correction. Synthetic selection status was then recorded separately as reinforced, synthetic-promoted, real-only, or not stably selected. Accession-direction agreement, between-study heterogeneity, and leave-one-accession-out results were retained as interpretation and sensitivity measures rather than inclusion requirements. The complete BH-FDR inventory is provided in Supplementary Table S17, its synthetic-guided subset in Table S16, and tissue-level counts in Table S18. Reactome was used to group selected genes into biological processes [7].

Interpretation of the four stages

The stages answer different questions and should not be collapsed into one performance claim. Development screens identify where and how synthetic data may be useful. Real-only random-effects tests establish association in the observed studies. Complete-study transfer asks whether a frozen policy applies outside its development studies and therefore provides the strongest validation.

Table 2. Evaluation stages and permitted interpretation.

Stage	Analysis	Data separation	Question answered	Permitted interpretation
1	Pooled utility benchmark	Locked profiles from represented studies	Does one synthetic-data policy help across tissues?	Method-level positive or negative result; no biological claim
2	Tissue-specific five-arm screen	Held-out profiles within represented accessions	Which synthetic use, if any, helps each tissue?	Developmental utility and stable feature nomination
3	Real-data random-effects BH FDR	FLT-GC effects estimated separately within each accession	Are nominated genes associated with flight in observed studies?	Real biological association; synthetic status reported separately
4	Whole-study transfer	Complete OSDR accessions absent from adaptation and policy development	Does a frozen synthetic-informed policy transfer to a new study context?	Strongest synthetic-utility evidence; still requires prospective replication

Signals supported by Stages 2 and 3 but lacking Stage 4 evidence are described as developmental or exploratory. Generative findings were also compared with the separate expiMap pathway analysis [12] to identify convergent and complementary tissue responses.

For a secondary spleen comparison, study-level *Igfbp3* effects were checked against full-transcriptome TPM values. Healthy sorted-cell and single-cell references, GSE156162 and E-MTAB-7703, were then examined to identify the normal spleen populations expressing *Igfbp3* [18,19]. The model did not select *Igfbp3*, so this analysis is reported as real-data context rather than a synthetic-guided finding.

Results

Pooled augmentation failed, but tissue-specific synthetic use helped during development

The generator preserved broad tissue structure well enough to support downstream testing. Real-trained and generated-trained classifiers both achieved tissue balanced accuracy 0.781 on the complete-series ARCHS4 test split, and the adapted OSDR model passed its locked finite-sample fidelity checks. The stricter ARCHS4 gene-correlation gate was not met, however, so downstream value was evaluated directly rather than inferred from generative similarity alone. Full fidelity metrics and comparator-model results are reported in Supplementary Figures S1-S4 and Supplementary Tables S1-S3.

A single pooled augmentation policy was not useful. On the locked OSDR test, balanced accuracy was 0.754 with real-only training, 0.695 with generated-only training, and 0.737 with real-plus-generated training. This negative result did not imply that every tissue responded identically. When synthetic use was selected separately by tissue, several arms improved balanced accuracy, AUROC, and average

precision within represented studies (Table 3). Different tissues selected different uses: spleen, skin, and soleus selected real-plus-generated training; pooled skeletal muscle selected feature guidance with a real-only classifier; kidney and thymus selected low-weight guided training; and lung and adrenal gland selected generated-only training. These are development-screen results because profiles, rather than complete studies, were withheld.

Table 3. Selected tissue-specific development results. Every displayed synthetic-informed arm met the balanced-accuracy, AUROC, and average-precision eligibility rule. Full results are in Supplementary Table S20.

Tissue	Selected synthetic use	Balanced accuracy, real to selected	AUROC, real to selected
Thymus	Low-weight guided training	0.733 to 0.844	0.818 to 0.887
Skeletal muscle, pooled	Feature-guided real-only classifier	0.861 to 0.932	0.924 to 0.961
Soleus	Real plus generated	0.925 to 0.963	0.980 to 0.980
Kidney	Low-weight guided training	0.654 to 0.707	0.680 to 0.771
Spleen	Real plus generated	0.517 to 0.648	0.540 to 0.704
Skin	Real plus generated	0.671 to 0.756	0.691 to 0.768
Lung	Generated only	0.664 to 0.742	0.680 to 0.830
Adrenal gland	Generated only	0.781 to 0.922	0.906 to 0.984

Synthetic-informed selection identified real-data associations

The real-data screen yielded 202 BH-FDR tissue-gene associations across 10 of 22 canonical tissue analyses and 257 across the five anatomical muscle groups. These are tissue-gene results rather than 459 unique or independent discoveries: a gene can be significant in more than one tissue, and pooled skeletal muscle overlaps its anatomical subgroups. Forty-nine associations also entered stable synthetic-informed selection: 26 were synthetic-promoted and 23 were reinforced by both real-only and synthetic-informed selection.

All 459 P values and FDR values came from real profiles. Synthetic data affected only whether a gene was repeatedly prioritized and, for eligible augmentation arms, classifier fitting. Synthetic labels were suppressed where the selected arm failed its metric gate; quadriceps, EDL, and liver therefore contribute real-data associations but no synthetic-informed claims. The complete inventory, including associations with mixed study directions, is provided in Supplementary Tables S16-S18.

Whole-study transfer narrowed the evidence to thymus

The development gains did not uniformly transfer to excluded studies. In an initial six-tissue feature-guidance transfer screen, only lung and thymus met the advancement rule. Liver and kidney gained balanced accuracy but lost AUROC, while retina and skin did not improve. In the fixed follow-up, thymus improved on all three metrics, whereas lung balanced accuracy declined from 0.400 to 0.350 and the result was rejected.

Separate frozen augmentation experiments reached the same general conclusion. Heart balanced accuracy improved in a five-profile test, but AUROC and average precision were unchanged, making this a small exploratory result. Pooled skeletal muscle improved in the first two-accession test, but extension of the same recipe to all 11 eligible held-out accessions changed balanced accuracy only from 0.655 to 0.658 while AUROC fell from 0.718 to 0.690. Spleen also failed its initial held-out criterion. Thus tissue-specific development gains identify useful hypotheses, but only thymus supplied a strong retained whole-study result.

Table 4. Selected complete-accession tests. Experiments used separately frozen policies and are shown to distinguish initial gains from broader transfer.

Tissue and experiment	Accessions / profiles	Synthetic use	Balanced accuracy, real to informed	AUROC, real to informed	Interpretation
Thymus, fixed test	1 / 24	Feature guidance	0.500 to 0.833	0.840 to 0.979	Retained primary transfer result
Lung, fixed test	1 / 20	Low-weight guidance	0.400 to 0.350	0.450 to 0.470	Failed confirmation
Heart, early screen	1 / 5	Real plus generated	0.333 to 0.583	0.667 to 0.667	Exploratory because of test size
Skeletal muscle, initial	2 / 24	Real plus generated	0.875 to 0.917	0.958 to 0.972	Initial gain
Skeletal muscle, full extension	11 / 159	Real plus generated	0.655 to 0.658	0.718 to 0.690	Initial gain did not generalize
Spleen, initial	2 / 29	Real plus generated	0.750 to 0.725	0.766 to 0.794	Failed criterion

Thymus shows lower proliferative renewal during spaceflight

OSD-457 provided the strongest result. Its GEO series and every other OSDR-linked GEO series were excluded from ARCHS4 pretraining, and OSD-457 was excluded from OSDR adaptation and feature-policy development. In this held-out test, synthetic-guided gene selection improved separation of flight and ground-control thymus samples from balanced accuracy 0.500 to 0.833 and AUROC 0.840 to 0.979. The result held in both wild-type and Nrf2-knockout mice. Flight effects were closely aligned between the two genotypes.

The core genes *Birc5*, *Ccne2*, *Gmnn*, *Ube2c*, *Cdk1*, *Nusap1*, *Ccnb1*, and *Ccnb2* were lower in flight in both strata (Fig. 2A). These were not genes absent from the real-data analysis: all eight were among its 26 highest-ranked candidates, including four among the top ten. Synthetic guidance instead assembled a broader, coherent cell-cycle panel that transferred more effectively to the study-excluded test. Together, these genes regulate DNA replication, chromosome progression, mitotic entry, and completion of cell division. Reactome analysis connected them to APC/C-mediated cyclin degradation, G2/M checkpoints, DNA synthesis, and broader cell-cycle control (Fig. 2B).

The study-held-out result is therefore interpreted as panel-level reinforcement and organization, not de novo gene discovery. In the repeated development screen, *Birc5*, *Cdk1*, *Nusap1*, *Ccne2*, and *Ccnb2* crossed the stable-selection threshold only with synthetic guidance; *Gmnn* and *Ube2c* were reinforced by both arms; and *Ccnb1* remained BH-significant but did not cross either stability threshold. These labels describe repeated selection behavior and do not override the stronger held-out result. The complete cross-analysis mapping is provided in Supplementary Table S19.

This result is aligned with, but more specific than, prior thymus studies. STS-135 mouse thymus showed changes in cell-cycle and DNA-damage programs, including lower checkpoint-related expression [8]. A later ISS experiment reported marked thymus mass loss and partial artificial-gravity rescue of cell-cycle expression [9]. The current signature emphasizes mitotic completion and replication rather than acute apoptosis alone. Agreement between genotype strata suggests that the predictive signature is not confined to one Nrf2 background, but it does not prove Nrf2 independence.

Bulk thymus expression cannot distinguish lower transcription within proliferating thymocytes from loss or redistribution of proliferating cell populations. The defensible biological conclusion is lower abundance of a mitotic transcript program in flight, consistent with reduced proliferative renewal. Cell-resolved or histological confirmation is required to assign the effect to a cell-intrinsic mechanism.

Held-out thymus test prioritizes a flight-lower mitotic program in both genotypes




Figure 2. Thymus response to spaceflight. (A) Flight-minus-ground effects in real OSD-457 profiles for eight cell-cycle genes that were lower in both wild-type and Nrf2-knockout mice. (B) The genes converge on a mitotic and DNA-replication process family.

Anatomical separation exposes a soleus-specific metabolic program

Aggregate skeletal muscle concealed substantial anatomical heterogeneity. We therefore examined extensor digitorum longus, gastrocnemius, quadriceps, soleus, and tibialis anterior separately. Soleus produced the clearest biological pattern: its selected genes showed consistent flight effects across three accessions and converged on related metabolic processes.

The soleus screen selected real-plus-generated training. Balanced accuracy increased from 0.925 to 0.963, AUROC remained 0.980, and average precision increased from 0.980 to 0.986. Five BH-FDR genes were stable in both real-only and synthetic-guided selection: *Bdh1*, *Ech1*, *Bnip3*, and *Decr1* were lower in flight in all three accessions, while *Tpm1* was higher (Fig. 3B). Four genes passed leave-one-accession-out FDR; *Decr1* did not.

The model did not promote a soleus gene absent from stable real-only selection. Its contribution was reinforcement of a coherent existing pattern. Reactome connected the retained genes to mitochondrial protein turnover, mitochondrial fatty-acid beta-oxidation, and lipid metabolism (Fig. 3C). *Bdh1* and *Ech1* support oxidative substrate handling, *Bnip3* supports mitochondrial quality control, *Decr1* contributes to unsaturated fatty-acid oxidation, and *Tpm1* suggests contractile remodeling.

Prior 30-day spaceflight profiling of mouse soleus reported a slow-to-fast shift and broad changes in oxidative metabolism, PPAR signaling, and contractile genes [10]. Unloading studies have also reported reduced soleus fatty-acid oxidation [11]. The result is therefore literature-aligned panel reinforcement rather than de novo gene discovery.

Unlike thymus, soleus was represented during model development. Its cross-study consistency makes it a focused biological hypothesis, but an entirely unseen soleus study is still needed for independent confirmation.

Anatomical separation reveals a soleus-specific oxidative-metabolism hypothesis




Figure 3. Skeletal-muscle and soleus response. (A) Number of synthetic-informed genes that also pass real leave-one-accession-out FDR in each anatomical muscle group. (B) Five reinforced soleus BH-FDR genes with consistent real flight effects. (C) Their strongest shared biological processes center on mitochondrial turnover and lipid metabolism.

Other muscle groups provide narrower hypotheses

The pooled skeletal-muscle screen selected synthetic-guided feature ranking and improved balanced accuracy, AUROC, and average precision by 0.071, 0.036, and 0.037. Twelve BH-FDR genes were synthetic-informed, nine of which passed leave-one-accession-out sensitivity. The associated gene sets included interferon signaling and sialic-acid metabolism. Because pooled muscle combines anatomical groups with distinct physiology, this result is interpreted alongside, rather than instead of, the soleus analysis.

Among the remaining anatomical groups, tibialis anterior retained a real-plus-generated arm and gastrocnemius retained a low-weight guided arm. Tibialis anterior reinforced *Cdkn1a*, *St3gal5*, and *Bnip3* and promoted *Cebpd*; gastrocnemius promoted *Fhl2* and *Nfkbia*. None of these genes passed leave-one-accession-out FDR. EDL and quadriceps selected real-only arms, so their BH-FDR genes, including quadriceps *Rbm6*, are retained in the complete real-data inventory but are not attributed to the synthetic workflow.

Spleen screen nominates adhesion and cytoskeletal genes

The spleen screen selected real-plus-generated training. Balanced accuracy increased from 0.517 to 0.648, AUROC from 0.540 to 0.704, and average precision from 0.589 to 0.749. Synthetic guidance promoted flight-higher *Rai14*, *Ptprk*, and *My19* and reinforced flight-higher *Loxl1*. *Rai14*, *Ptprk*, and *Loxl1* had the same direction in all six studies; *My19* agreed in five of six. None passed leave-one-accession-out FDR, and the set did not produce significant Reactome enrichment.

The four genes provide a tentative structural hypothesis rather than a pathway claim. RAI14 links actin-associated mechanosensing to Hippo signaling, PTPRK regulates cell-cell junctions, MYL9 contributes to actomyosin contractility, and LOXL1 supports elastic extracellular-matrix maintenance [25-27]. Their shared direction is compatible with altered adhesion, mechanics, or tissue architecture, but the present bulk data cannot identify a common source cell or establish one mechanism.

Igf1bp3 remains a notable but separate real-data association. It was higher in flight in all six eligible spleen studies, with a random-effects FDR of $1.76e-9$; the largest FDR after removing one study at a time was 0.00385 . A full-transcriptome TPM sensitivity calculation showed 9% to 63% higher group means in flight. Healthy references localized baseline *Igf1bp3* mainly to fibroblastic reticular, collagen-producing, and perivascular stromal populations [18,19], and spaceflight proteomics has reported

increased IGFBP3 in serum and femur under microgravity relative to onboard artificial gravity [20]. However, the synthetic workflow did not select *Igfbp3*. It is therefore retained as a real-data stromal hypothesis, not evidence that synthetic guidance discovered or reinforced the gene.

Other organ responses were heterogeneous

Lung produced large repeated development-screen gains and cell-cycle, senescence, and PI3K/AKT-related enrichment, but no selected gene passed the real-data BH-FDR screen. More importantly, the fixed OSD-900 test lost balanced accuracy and failed its validation gate. Lung therefore demonstrates why within-study separation and pathway enrichment cannot substitute for study-held-out transfer.

Skin selected real-plus-generated training and promoted flight-higher *Plscr1*. *Plscr1* passed BH FDR and had the same direction in all six studies, although it did not pass leave-one-accession-out FDR. Broader selected sets were enriched for G1/S and DNA-repair processes, consistent with prior OSDR skin analyses reporting strong mission, strain, recovery-interval, and anatomical-site dependence [21]. This is a literature-aligned developmental candidate rather than a confirmed transferable signal.

Kidney produced the clearest secondary gene-level result. *Slc37a4* was reinforced by both selection arms and was higher in flight in all six kidney accessions. Synthetic guidance additionally promoted flight-higher *Inpp4b*; it passed BH FDR and remained significant in leave-one-accession-out analysis, although one of six accession effects was negative. The selected pair did not yield significant Reactome enrichment. *Slc37a4* supports renal glucose handling, while *Inpp4b* nominates phosphoinositide signaling. Both require prospective study-level confirmation and should be interpreted alongside prior reports of strain-dependent lipid, ECM, TGF-beta, Wnt, and nephron-remodeling responses [22-24].

Adrenal gland produced flight-lower synthetic-promoted *Psmb8* and reinforced *Tspan4*, each with the same direction in all three studies but neither passing leave-one-accession-out FDR. Retina showed repeated predictive gains and selected-set enrichment but no synthetic-informed BH-FDR gene. Liver selected the real-only arm: its 19 real-data BH-FDR genes remain in the complete inventory, but none support a synthetic-guided claim. Quadriceps and EDL likewise selected real-only arms.

Comparison with expiMap shows convergence and complementarity

The evidence distribution therefore differs from the separate expiMap analysis. Both approaches prioritize thymus. expiMap produced its broadest pathway evidence in thymus, skin, spleen, and kidney, whereas the generative workflow added its clearest complementary result in soleus and a focused kidney pair. The methods examine different biological representations: expiMap tests predefined pathway activity, while synthetic guidance helps prioritize individual genes and their shared processes.

Evidence is strongest in thymus and soleus, with secondary tissue candidates



Figure 4. Tissue evidence. (A) Repeated development-screen changes relative to real-only models for selected tissues. (B) BH-FDR genes promoted or reinforced by the synthetic workflow; synthetic labels are suppressed where the generated arm failed its metric gate. (C) Thymus remains the strongest result, soleus and pooled muscle provide the clearest process-level complement, kidney supplies a focused secondary pair, and spleen, skin, and adrenal gland remain developmental.

Table 5. Selected synthetic-guided biological interpretations by tissue. Complete real-data BH-FDR results are in Supplementary Tables S17-S18.

Tissue	Main signal	Interpretation
Thymus	Lower mitotic genes; APC/C, G2/M, and DNA replication	Strongest result; supports reduced proliferative renewal
Soleus	Reinforced FLT-lower <i>Bdh1</i> , <i>Ech1</i> , <i>Bnip3</i> , and <i>Decr1</i> ; FLT-higher <i>Tpm1</i>	Coherent mitochondrial and lipid-metabolism program; no promoted gene
Skeletal muscle, pooled	12 synthetic-informed BH-FDR genes; interferon and sialic-acid terms	Developmental complement to the anatomically specific soleus result
Kidney	Promoted <i>Inpp4b</i> and reinforced <i>Slc37a4</i> , both FLT-higher	Focused renal metabolic-signaling hypothesis
Spleen	Promoted <i>Rai14</i> , <i>Ptprk</i> , and <i>Myl9</i> ; reinforced <i>Loxl1</i>	Developmental adhesion/cytoskeletal hypothesis
Skin	Promoted FLT-higher <i>Plscr1</i> plus cell-cycle/DNA-repair enrichment	Literature-aligned developmental candidate
Adrenal gland	Promoted <i>Psmb8</i> and reinforced <i>Tspan4</i> , both FLT-lower	Three-study developmental candidate
Tibialis anterior	Reinforced <i>Cdkn1a</i> , <i>St3gal5</i> , and <i>Bnip3</i> ; promoted <i>Cebpd</i>	Exploratory stress and metabolic response
Gastrocnemius	Promoted <i>Fhl2</i> and <i>Nfkb1a</i>	Exploratory two-gene result without a coherent pathway
Lung	Within-study pathway candidates but failed OSD-900 transfer	No retained study-held-out synthetic-guided result
Retina	Predictive and pathway gains without a synthetic-informed BH-FDR gene	Exploratory gene/pathway mismatch
Quadriceps, EDL, liver	Screen retained real-only arms	Real-data BH-FDR associations are not synthetic-guided findings

Discussion

What synthetic data added

The pooled benchmark and tissue-specific screens answer different questions. Adding the same generated cohort across tissues did not improve classification. In tissue-specific development, however, direct augmentation, low-weight augmentation, generated-only training, and feature guidance each helped at least one tissue. There was therefore no universally best use of synthetic expression.

Whole-study transfer was much more selective. The initial pooled-muscle gain disappeared when the same frozen augmentation recipe was extended to 11 accessions, and lung failed fixed confirmation despite a strong development screen. Thymus was the exception: synthetic-guided feature selection transferred to OSD-457 and organized real-supported genes into a coherent cell-cycle panel. This pattern supports synthetic data as a tissue-specific regularizer or feature prior, not as independent biological sample size.

Thymus demonstrates panel-level organization: its central genes already ranked strongly in real data, but the synthetic-guided panel transferred after the study was removed from both reference pretraining and OSDR adaptation and converged on one interpretable cell-cycle process. The development-screen labels describe which genes crossed repeated selection thresholds and are not claims of biological novelty.

Soleus demonstrates reinforcement rather than discovery. The synthetic arm retained the same five BH-FDR genes as stable real-only selection and sharpened balanced accuracy without improving AUROC. The value is the convergence of independently selected genes on mitochondrial turnover and fatty-acid metabolism, not nomination of a new soleus gene.

Kidney provides the clearest promotion result outside thymus. Synthetic guidance added *Inpp4b* to reinforced *Slc37a4*, and the real-data association for *Inpp4b* survived leave-one-accession-out analysis. Spleen provides a weaker multi-gene example: *Rai14*, *Ptprk*, and *Myl9* were promoted and *Lox11* reinforced, but none passed LOO FDR and no pathway was significant. The previously emphasized *Igfbp3* association remains biologically interesting but does not demonstrate synthetic contribution. Quadriceps and EDL selected real-only arms and therefore no longer support synthetic-guided claims.

Thymus and soleus define the strongest organ responses

The thymus result refines established spaceflight immune biology. Prior work documents thymic involution and altered cell-cycle expression [8,9]. The current signature concentrates on cyclins, CDK1, UBE2C, BIRC5, NUSAP1, geminin, APC/C-mediated protein turnover, and G2/M control. Together these support lower proliferative renewal or a lower proportion of cycling thymocytes. Because the data are bulk, composition and cell-intrinsic regulation remain inseparable.

The soleus result addresses a different physiological axis. Weight-bearing slow muscle is especially sensitive to unloading. Lower *Bdh1*, *Ech1*, *Bnip3*, and *Decr1* together with higher *Tpm1* describe reduced oxidative substrate handling, altered mitochondrial quality control, and contractile remodeling. This is compatible with known soleus atrophy, altered lipid metabolism, and slow-to-fast transition [10,11]. It is complementary to expiMap because synthetic guidance reinforced a compact fatty-acid-oxidation panel observed consistently across soleus studies.

Why the other tissues remain useful

The other tissues constrain the method's scope. Kidney combines one promoted and one reinforced gene but lacks pathway enrichment. Spleen combines strong within-study predictive gains with four BH-FDR genes, yet none survives the stricter LOO criterion. Skin promotes one directionally consistent gene while broader pathway enrichment remains heterogeneous. Adrenal gland has only three studies. Retina has predictive and pathway gains without a synthetic-informed BH-FDR gene. Lung fails the study-held-out test despite strong development-screen separation. Liver, quadriceps, and EDL select real-only arms.

These results define direct follow-up priorities. A new thymus study should test the fixed eight-gene panel prospectively. Soleus experiments should quantify the five-gene mitochondrial and contractile panel together with fatty-acid oxidation. Kidney should test *Inpp4b* and *Slc37a4* jointly in an unseen mission. Spleen should test *Rai14*, *Ptprk*, *Myl9*, and *Lox11* in stroma-preserving assays, while *Igfbp3* should be evaluated separately as a real-data-derived stromal hypothesis. Lung requires a prospectively genotype-stratified study rather than further tuning on OSD-900.

Limitations

No tissue currently has a completely prospective independent biological test. OSD-457 was excluded from ARCHS4 pretraining, OSDR adaptation, and feature-policy development. Because its outcome had been examined during earlier model development, it still requires prospective replication. All other BH-FDR findings were identified within the broader model-development domain and still lack equivalent study-excluded testing. Agreement across represented real studies supports focused follow-up but does not establish transfer to a new mission or study.

The analysis used a 974-gene landmark panel, so relevant genes outside that panel could not be discovered. BH FDR was controlled separately within each declared tissue family, not once across every tissue-gene combination in the project. Direction disagreement and high heterogeneity do not remove a BH-significant association, but they reduce confidence that its pooled effect represents a common response across missions. Bulk tissue also cannot distinguish a transcriptional change within a cell type from a change in cell composition. The thymus result could reflect lower expression in proliferating thymocytes, fewer proliferating thymocytes, or both. The spleen structural candidates likewise cannot distinguish altered expression from altered stromal or contractile-cell abundance.

Finally, the generator represents tissues and study contexts available during training. The ARCHS4 reference failed its strict correlation-structure gate. After OSDR adaptation, the locked within-study test

passed its finite-sample fidelity rule, but the preceding validation screen missed its correlation floor and muscle accession-effect gate. The broad tissue screens measure interpolation within represented studies and remain hypothesis-generating. The model should not be assumed to reproduce a new mission, strain, or sample-processing protocol without additional testing. Exact model limitations, sensitivity analyses, and statistical safeguards are documented in the supplementary methods.

Conclusions

Synthetic expression was useful only when its role was selected for the tissue and evaluated against real-only training. A pooled augmentation policy failed, and most development gains did not establish whole-study transfer. The strongest retained use was thymus feature guidance; soleus supplied a coherent development-stage reinforcement result. The analysis supports a flight-lower thymus mitotic program, reinforces soleus mitochondrial lipid metabolism and contractile remodeling, and identifies kidney *Inpp4b* and *Slc37a4* as focused secondary candidates.

Thymus remains the strongest result because it survived complete OSDR-series removal from reference pretraining and accession exclusion from adaptation. Soleus supplies the clearest complementary process-level result. Pooled skeletal muscle, kidney, spleen, skin, adrenal gland, gastrocnemius, and tibialis anterior provide developmental hypotheses requiring unseen-study validation. Quadriceps, EDL, and liver do not support synthetic-guided claims because their real-only arms were retained. Lung failed transfer to OSD-900, and pooled-muscle augmentation did not improve the full 11-accession benchmark. Together, these results provide a reproducible framework for deciding when synthetic expression can guide biological analysis and when it should be rejected.

Data and code availability

OSDR data were accessed through the public Biological Data API [1]. ARCHS4 and Reactome are public resources [2,7]. Repository scripts are under `src/nasa_mouse_rna_diffusion/`; frozen run outputs are under `outputs/generative_benchmark/`; this paper package is under `paper/synthetic_guided_spaceflight/`. The figure and table builder is `nasa_mouse_rna_diffusion.build_synthetic_guided_paper`. SHA-256 hashes of every frozen analysis input are included in `source_data/frozen_input_manifest.tsv`. Public repository URL and archival DOI will be added after author review.

Ethics statement

This study is a secondary computational analysis of publicly available animal-study data. No new animals were used.

Competing interests

The author declares no competing interests.

Acknowledgments

The author acknowledges the NASA Open Science Data Repository, GeneLab data-processing teams, original flight-study investigators, ARCHS4, Reactome, and the developers of the evaluated generative-model implementations. Program and mentor acknowledgments require final author review.

References

1. NASA Open Science Data Repository. OSDR Biological Data API. <https://visualization.osdr.nasa.gov/biodata/api/>. Accessed July 28, 2026.
2. Lachmann A, Torre D, Keenan AB, et al. Massive mining of publicly available RNA-seq data from human and mouse. *Nature Communications*. 2018;9:1366. <https://doi.org/10.1038/s41467-018-03751-6>.
3. Viñas R, Andrés-Terré H, Liò P, Bryson K. Adversarial generation of gene expression data. *Bioinformatics*. 2022;38:730-737. <https://doi.org/10.1093/bioinformatics/btab035>.
4. Lacan A, André R, Sebag M, Hanczar B. In silico generation of gene expression profiles using diffusion models. *BMC Bioinformatics*. 2026. <https://doi.org/10.1186/s12859-026-06470-8>.
5. Litman E, Myers T, Agarwal V, et al. GeneJEPa: A predictive world model of the transcriptome. *bioRxiv*. 2025. <https://doi.org/10.1101/2025.10.14.682378>.
6. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal Statistical Society Series B*. 1995;57:289-300. <https://doi.org/10.1111/j.2517-6161.1995.tb02031.x>.
7. Milacic M, Beavers D, Conley P, et al. The Reactome Pathway Knowledgebase 2024. *Nucleic Acids Research*. 2024;52:D672-D678. <https://doi.org/10.1093/nar/gkad1025>.
8. Gridley DS, Mao XW, Stodieck LS, et al. Changes in mouse thymus and spleen after return from the STS-135 mission in space. *PLoS ONE*. 2013;8:e75097. <https://doi.org/10.1371/journal.pone.0075097>.
9. Horie K, Kato T, Kudo T, et al. Impact of spaceflight on the murine thymus and mitigation by exposure to artificial gravity during spaceflight. *Scientific Reports*. 2019;9:19866. <https://doi.org/10.1038/s41598-019-56432-9>.
10. Gambaro G, Salanova M, Ciciliot S, et al. Gene expression profiling in slow-type calf soleus muscle of 30 days space-flown mice. *PLoS ONE*. 2017;12:e0169314. <https://doi.org/10.1371/journal.pone.0169314>.
11. Stein TP, Schluter MD, Grindeland RE, Moran MM, Baer LA, Wade CE. Rate controlling steps in fatty acid oxidation by unloaded rodent soleus muscle. *Journal of Gravitational Physiology*. 2002;9:P165-P166. PMID: 15002531.
12. Lotfollahi M, Rybakov S, Hrovatin K, et al. Biologically informed deep learning to query gene programs in single-cell atlases. *Nature Cell Biology*. 2023;25:337-350. <https://doi.org/10.1038/s41556-022-01072-x>.
13. Ho J, Jain A, Abbeel P. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*. 2020;33:6840-6851.
14. Song J, Meng C, Ermon S. Denoising diffusion implicit models. *International Conference on Learning Representations*. 2021. <https://arxiv.org/abs/2010.02502>.
15. Heusel M, Ramsauer H, Unterthiner T, Nessler B, Hochreiter S. GANs trained by a two time-scale update rule converge to a local Nash equilibrium. *Advances in Neural Information Processing Systems*. 2017;30.
16. Kynkäänniemi T, Karras T, Laine S, Lehtinen J, Aila T. Improved precision and recall metric for assessing generative models. *Advances in Neural Information Processing Systems*. 2019;32.
17. DerSimonian R, Laird N. Meta-analysis in clinical trials. *Controlled Clinical Trials*. 1986;7:177-188. [https://doi.org/10.1016/0197-2456\(86\)90046-2](https://doi.org/10.1016/0197-2456(86)90046-2).
18. Pezoldt J, Wiechers C, Erhard F, et al. Single-cell transcriptional profiling of splenic fibroblasts reveals subset-specific innate immune signatures in homeostasis and during viral infection. *Communications Biology*. 2021;4:1355. <https://doi.org/10.1038/s42003-021-02882-9>.
19. Cheng HW, Onder L, Novkovic M, et al. Origin and differentiation trajectories of fibroblastic reticular cells in the splenic white pulp. *Nature Communications*. 2019;10:1739. <https://doi.org/10.1038/s41467-019-09728-3>.

20. Kimura Y, Nakai Y, Ino Y, Akiyama T, Ryo A, Hirano H. Identification of gravity-responsive serum proteins in spaceflight mice using a quantitative proteomic approach with data-independent acquisition mass spectrometry. *Proteomics*. 2024;24:2300214. <https://doi.org/10.1002/pmic.202300214>.
21. Cope H, Elsborg J, Demharter S, et al. Transcriptomics analysis reveals molecular alterations underpinning spaceflight dermatology. *Communications Medicine*. 2024;4:106. <https://doi.org/10.1038/s43856-024-00532-9>.
22. Finch RH, Vitry G, Siew K, et al. Spaceflight causes strain-dependent gene expression changes in the kidneys of mice. *npj Microgravity*. 2025;11:11. <https://doi.org/10.1038/s41526-025-00465-0>.
23. Siew K, et al. Cosmic kidney disease: an integrated pan-omic, physiological and morphological study into spaceflight-induced renal dysfunction. *Nature Communications*. 2024;15:4923. <https://doi.org/10.1038/s41467-024-49212-1>.
24. Suzuki N, Iwamura Y, Nakai T, et al. Gene expression changes related to bone mineralization, blood pressure and lipid metabolism in mouse kidneys after space travel. *Kidney International*. 2022;101:92-105. <https://doi.org/10.1016/j.kint.2021.09.031>.
25. Jeong W, Kwon H, Park SK, Lee IS, Jho EH. Retinoic acid-induced protein 14 links mechanical forces to Hippo signaling. *EMBO Reports*. 2024;25:4033-4061. <https://doi.org/10.1038/s44319-024-00228-0>.
26. Fearnley GW, Young KA, Edgar JR, et al. The homophilic receptor PTPRK selectively dephosphorylates multiple junctional regulators to promote cell-cell adhesion. *eLife*. 2019;8:e44597. <https://doi.org/10.7554/eLife.44597>.
27. Liu X, Zhao Y, Gao J, et al. Elastic fiber homeostasis requires lysyl oxidase-like 1 protein. *Nature Genetics*. 2004;36:178-182. <https://doi.org/10.1038/ng1297>.
28. Machour FE, Abu-Zhayia ER, Awwad SW, et al. RBM6 splicing factor promotes homologous recombination repair of double-strand breaks and modulates sensitivity to chemotherapeutic drugs. *Nucleic Acids Research*. 2021;49:11708-11727. <https://doi.org/10.1093/nar/gkab976>.
29. Benbarche S, Bello Pineda JM, Baquero Galvis L, et al. GPATCH8 modulates mutant SF3B1 mis-splicing and pathogenicity in hematologic malignancies. *Molecular Cell*. 2024;84:1886-1903.e10. <https://doi.org/10.1016/j.molcel.2024.04.006>.
30. Wiltshire C, Matsushita M, Tsukada S, Gillespie DAF, May GHW. A new c-Jun N-terminal kinase (JNK)-interacting protein, Sab (SH3BP5), associates with mitochondria. *Biochemical Journal*. 2002;367:577-585. <https://doi.org/10.1042/BJ20020553>.