


Logic and Sets 1

Introduction to Propositional Logic



Organization - Lectures and exercise classes

Lectures on **logic** on *Tuesdays*

- ▶ lecturer: Patrick Koopmann
`p.k.koopmann@vu.nl`

Lectures on **sets** on *Thursdays*

- ▶ lecturer: Robert van der Vorst

Organization - Lectures and exercise classes

Lectures on **logic** on *Tuesdays*

- ▶ lecturer: Patrick Koopmann
`p.k.koopmann@vu.nl`

Lectures on **sets** on *Thursdays*

- ▶ lecturer: Robert van der Vorst

Enroll *in Canvas* for both **logic** and **sets** exercise classes

Organization - Examination

Midterm exam (E_1): Monday March 3

Final exam (E_2): Tuesday March 25

Organization - Examination

Midterm exam (E_1): Monday March 3

Final exam (E_2): Tuesday March 25

Overall grade: $0.4 E_1 + 0.6 E_2$

At least a 5 must be achieved for both parts (and a 5.5 overall)

Organization - Examination

Midterm exam (E_1): Monday March 3

Final exam (E_2): Tuesday March 25


Overall grade: $0.4 E_1 + 0.6 E_2$

At least a 5 must be achieved for both parts (and a 5.5 overall)

The resit covers both parts at once


Material logic lectures

- ▶ M. Huth, M. Ryan, *Logic in Computer Science*, 2nd edition, Cambridge University Press, 2004
 - ▶ we treat only some parts (Chapters 1, 2, 6)
 - ▶ some topics in logic lectures are not covered
 - ▶ copies of relevant parts of the book on Canvas




Material logic lectures

- ▶ M. Huth, M. Ryan, *Logic in Computer Science*, 2nd edition, Cambridge University Press, 2004
 - ▶ we treat only some parts (Chapters 1, 2, 6)
 - ▶ some topics in logic lectures are not covered
 - ▶ copies of relevant parts of the book on Canvas
- ▶ Lecture slides available on Canvas
- ▶ Exercises for working groups also available on Canvas



Question




Which of the three ladies want a beer ?

Importance of logic

Hardware circuits are based on (Boolean) logic

Software is basically a piece of logic

Artificial intelligence is founded on logic




Importance of logic

Hardware circuits are based on (Boolean) logic

Software is basically a piece of logic


Artificial intelligence is founded on logic



Reasoning about a computer program or AI application requires

- ▶ a formal logical framework, and
- ▶ a trained mind in logical thought

Propositional logic: Syntax



A line of argument (in natural language)

We want to understand why arguments such as these hold

Example

If the train arrives late and there are no taxis at the station,
then Jane is late for her meeting

A line of argument (in natural language)

We want to understand why arguments such as these hold

Example

If the train arrives late and there are no taxis at the station,
then Jane is late for her meeting

Jane is not late for the meeting

A line of argument (in natural language)

We want to understand why arguments such as these hold

Example

If the train arrives late and there are no taxis at the station,
then Jane is late for her meeting

Jane is not late for the meeting

The train does arrive late

A line of argument (in natural language)

We want to understand why arguments such as these hold

Example

If the train arrives late and there are no taxis at the station,
then Jane is late for her meeting


Jane is not late for the meeting

The train does arrive late

Therefore, there are taxis at the station


Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement



Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement




Examples

► $5 > 3$

Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement




Examples

► $5 > 3$

► $5 < 3$

Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement




Examples

- ▶ $5 > 3$
- ▶ $5 < 3$
- ▶ grass is green

Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement




Examples

- ▶ $5 > 3$
- ▶ $5 < 3$
- ▶ grass is green
- ▶ grass is green *and* roses are blue

Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement




Examples

- ▶ $5 > 3$
- ▶ $5 < 3$
- ▶ grass is green
- ▶ grass is green *and* roses are blue
- ▶ grass is green *or* roses are blue

Declarative sentences

A *declarative sentence* (or *proposition*) is a *true* or *false* statement



Examples

- ▶ $5 > 3$
- ▶ $5 < 3$
- ▶ grass is green
- ▶ grass is green *and* roses are blue
- ▶ grass is green *or* roses are blue
- ▶ *if* $x > 1$, *then* $x^2 \neq x$

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza
- ▶ Every even number > 2 is the sum of two prime numbers
Goldbach conjecture, 1742

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza
- ▶ Every even number > 2 is the sum of two prime numbers
Goldbach conjecture, 1742
- ▶ It is possible to derive $0 = 1$ in the natural numbers
Gödel's 1st incompleteness theorem, 1931

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza
- ▶ Every even number > 2 is the sum of two prime numbers
Goldbach conjecture, 1742
- ▶ It is possible to derive $0 = 1$ in the natural numbers
Gödel's 1st incompleteness theorem, 1931
- ▶ You are enjoying this lecture

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza
- ▶ Every even number > 2 is the sum of two prime numbers
Goldbach conjecture, 1742
- ▶ It is possible to derive $0 = 1$ in the natural numbers
Gödel's 1st incompleteness theorem, 1931
- ▶ You are enjoying this lecture

Non-examples

- ▶ Could you please be quiet during my lecture?

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza
- ▶ Every even number > 2 is the sum of two prime numbers
Goldbach conjecture, 1742
- ▶ It is possible to derive $0 = 1$ in the natural numbers
Gödel's 1st incompleteness theorem, 1931
- ▶ You are enjoying this lecture

Non-examples

- ▶ Could you please be quiet during my lecture?
- ▶ Prepare for the exercise classes!

Declarative sentences

More examples

- ▶ All Martians like pepperoni on their pizza
- ▶ Every even number > 2 is the sum of two prime numbers
Goldbach conjecture, 1742
- ▶ It is possible to derive $0 = 1$ in the natural numbers
Gödel's 1st incompleteness theorem, 1931
- ▶ You are enjoying this lecture

Non-examples

- ▶ Could you please be quiet during my lecture?
- ▶ Prepare for the exercise classes!
- ▶ May you do well at the exam

Argument abstraction

Example (Jane)

If the train arrives late, and there are no taxis at the station,
then Jane is late for her meeting

Jane is not late for her meeting

The train does arrive late

Therefore, there are taxis at the station

Argument abstraction

Example (Jane)

If the train arrives late, **and** there are **no** taxis at the station,
then Jane is late for her meeting

Jane is **not** late for her meeting

The train does arrive late

Therefore, there are taxis at the station

Argument abstraction

Example (Jane)

If the train arrives late, **and** there are **no** taxis at the station,
then Jane is late for her meeting

Jane is **not** late for her meeting

The train does arrive late

Therefore, there are taxis at the station

Key of translation

- p* the train arrives late
- q* there are taxis at the station
- r* Jane is late for her meeting

Argument abstraction

Example (Jane)

If the train arrives late, and there are no taxis at the station,
then Jane is late for her meeting

Jane is not late for her meeting

The train does arrive late

Therefore, there are taxis at the station

Key of translation

- p the train arrives late
- q there are taxis at the station
- r Jane is late for her meeting

Abstraction:

If p and not q , then r Not r p *Therefore*, q

Argument abstraction (another example)

Example (John)

If it is raining, and John did not take his umbrella with him,
then he will get wet

John is not getting wet

It is raining

Therefore,

Argument abstraction (another example)

Example (John)

If it is raining, and John did not take his umbrella with him,
then he will get wet

John is not getting wet

It is raining

Therefore, John took his umbrella with him

Argument abstraction (another example)

Example (John)

If it is raining, and John did not take his umbrella with him,
then he will get wet

John is not getting wet

It is raining

Therefore, John took his umbrella with him

Key of translation

- p it is raining
- q John takes his umbrella with him
- r John is getting wet

Argument abstraction (another example)

Example (John)

If it is raining, and John did not take his umbrella with him,
then he will get wet

John is not getting wet

It is raining

Therefore, John took his umbrella with him

Key of translation

p it is raining

q John takes his umbrella with him

r John is getting wet

Abstraction:

If p and not q , then r Not r p *Therefore*, q

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$\neg q$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$p \wedge \neg q$$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$(p \wedge \neg q) \rightarrow r$$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$(p \wedge \neg q) \rightarrow r \qquad \neg r \quad p$$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$(p \wedge \neg q) \rightarrow r \qquad \neg r \wedge p$$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$((p \wedge \neg q) \rightarrow r) \wedge (\neg r \wedge p)$$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$(((p \wedge \neg q) \rightarrow r) \wedge (\neg r \wedge p)) \rightarrow q$$

Argument formalization

	Example (Jane)	Example (John)
p	the train is late	it is raining
q	there are taxis at the station	John has his umbrella with him
r	Jane is late for her meeting	John is getting wet

Both arguments have the same abstraction:

If p and not q , then r Not r p Therefore, q

with as logical formalization:

$$(((p \wedge \neg q) \rightarrow r) \wedge (\neg r \wedge p)) \rightarrow q$$

That the two arguments **hold** is due to their *logical form*

It does not depend on the actual content of propositions p , q and r

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$

‘and’

conjunction

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$

‘and’

conjunction

$\vee$

‘or’

disjunction

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$

‘and’

conjunction

$\vee$

‘or’

disjunction

$\oplus$

‘either ... or ...’

exclusive or

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$	'and'	<i>conjunction</i>
$\vee$	'or'	<i>disjunction</i>
$\oplus$	'either ... or ...'	<i>exclusive or</i>
$\neg$	'not'	<i>negation</i>

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$	'and'	<i>conjunction</i>
$\vee$	'or'	<i>disjunction</i>
$\oplus$	'either ... or ...'	<i>exclusive or</i>
$\neg$	'not'	<i>negation</i>
$\rightarrow$	'if ... then ...'	<i>implication</i>

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$	'and'	<i>conjunction</i>
$\vee$	'or'	<i>disjunction</i>
$\oplus$	'either ... or ...'	<i>exclusive or</i>
$\neg$	'not'	<i>negation</i>
$\rightarrow$	'if ... then ...'	<i>implication</i>
$\leftrightarrow$	'if and only if'	<i>bi-implication</i>

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$	'and'	<i>conjunction</i>
$\vee$	'or'	<i>disjunction</i>
$\oplus$	'either ... or ...'	<i>exclusive or</i>
$\neg$	'not'	<i>negation</i>
$\rightarrow$	'if ... then ...'	<i>implication</i>
$\leftrightarrow$	'if and only if'	<i>bi-implication</i>

Not in the scope of propositional logic are constructs like:

- ▶ *for all*, *there exists* (will be treated in lecture 6)

Symbols of propositional logic

*We want to study logic without being distracted by
the concrete contents of propositions*

Propositional variables:

$p, q, r, \dots$

Connectives:

$\wedge$	'and'	<i>conjunction</i>
$\vee$	'or'	<i>disjunction</i>
$\oplus$	'either ... or ...'	<i>exclusive or</i>
$\neg$	'not'	<i>negation</i>
$\rightarrow$	'if ... then ...'	<i>implication</i>
$\leftrightarrow$	'if and only if'	<i>bi-implication</i>

Not in the scope of propositional logic are constructs like:

- ▶ *for all*, *there exists* (will be treated in lecture 6)
- ▶ *eventually* (temporal logic), *I know that* (epistemic logic)

Sentences and formulas

Propositional structure of sentences

► $5 > 3$

p

Sentences and formulas

Propositional structure of sentences

- ▶ $5 > 3$ p
- ▶ grass is green p

Sentences and formulas

Propositional structure of sentences

- ▶ $5 > 3$ p
- ▶ grass is green p
- ▶ grass is green and roses are blue $p \wedge q$

Sentences and formulas

Propositional structure of sentences

- ▶ $5 > 3$ p
- ▶ grass is green p
- ▶ grass is green and roses are blue $p \wedge q$
- ▶ grass is green or roses are blue $p \vee q$

Sentences and formulas

Propositional structure of sentences

- ▶ $5 > 3$ p
- ▶ grass is green p
- ▶ grass is green and roses are blue $p \wedge q$
- ▶ grass is green or roses are blue $p \vee q$
- ▶ if $x > 1$, then $x^2 \neq x$ $p \rightarrow \neg q$

Question

What are p and q in the last example ?

Sentences and formulas

Propositional structure of sentences

- ▶ $5 > 3$ p
- ▶ grass is green p
- ▶ grass is green and roses are blue $p \wedge q$
- ▶ grass is green or roses are blue $p \vee q$
- ▶ if $x > 1$, then $x^2 \neq x$ $p \rightarrow \neg q$

Question

What are p and q in the last example ?

Example

The sentence “I don’t wear glasses” can be expressed by

- ▶ $\neg p$ where p represents “I wear glasses”

Sentences and formulas

Propositional structure of sentences

- ▶ $5 > 3$ p
- ▶ grass is green p
- ▶ grass is green and roses are blue $p \wedge q$
- ▶ grass is green or roses are blue $p \vee q$
- ▶ if $x > 1$, then $x^2 \neq x$ $p \rightarrow \neg q$

Question

What are p and q in the last example ?

Example

The sentence “I don’t wear glasses” can be expressed by

- ▶ $\neg p$ where p represents “I wear glasses”
- ▶ q where q represents “I don’t wear glasses”

It's all Greek to me

Following Huth and Ryan, we use *Greek letters* for propositional formulas

ϕ phi

ψ psi

χ chi



Formulas of propositional logic

Building blocks:

- ▶ $p, q, r, \dots$ are propositional variables

Formulas of propositional logic

Building blocks:

- ▶ $p, q, r, \dots$ are propositional variables
- ▶ $\neg$ is a unary connective

Formulas of propositional logic

Building blocks:

- ▶ $p, q, r, \dots$ are propositional variables
- ▶ $\neg$ is a unary connective
- ▶ $\wedge, \vee, \oplus, \rightarrow, \leftrightarrow$ are binary connectives

Formulas of propositional logic

Building blocks:

- ▶ $p, q, r, \dots$ are propositional variables
- ▶ $\neg$ is a unary connective
- ▶ $\wedge, \vee, \oplus, \rightarrow, \leftrightarrow$ are binary connectives

The construction of formulas:

Inductive definition

1. (BASE STEP) every propositional variable is a formula

Formulas of propositional logic

Building blocks:

- ▶ $p, q, r, \dots$ are **propositional variables**
- ▶ $\neg$ is a **unary connective**
- ▶ $\wedge, \vee, \oplus, \rightarrow, \leftrightarrow$ are **binary connectives**

The construction of **formulas**:

Inductive definition

1. (BASE STEP) every propositional variable is a **formula**
2. (CONSTRUCTION STEPS)
 - 2.1 if ϕ is a **formula**, then so is $\neg(\phi)$
 - 2.2 if ϕ and ψ are **formulas**, then so are $(\phi \wedge \psi)$, $(\phi \vee \psi)$, $(\phi \oplus \psi)$, $(\phi \rightarrow \psi)$ and $(\phi \leftrightarrow \psi)$

Formulas of propositional logic

Building blocks:

- ▶ $p, q, r, \dots$ are **propositional variables**
- ▶ $\neg$ is a **unary connective**
- ▶ $\wedge, \vee, \oplus, \rightarrow, \leftrightarrow$ are **binary connectives**

The construction of **formulas**:

Inductive definition

1. (BASE STEP) every propositional variable is a **formula**
2. (CONSTRUCTION STEPS)
 - 2.1 if ϕ is a **formula**, then so is $\neg(\phi)$
 - 2.2 if ϕ and ψ are **formulas**, then so are $(\phi \wedge \psi)$, $(\phi \vee \psi)$, $(\phi \oplus \psi)$, $(\phi \rightarrow \psi)$ and $(\phi \leftrightarrow \psi)$

We take the liberty to omit the outermost pair of parentheses

Exercise: Construct a formal language


Arithmetic:

- ▶ Think of simple arithmetic terms with natural numbers
- ▶ E.g. $7 + 3$, $12 * (3 + 7)$, -5 , etc

Parsing a formula

$$(\neg(p) \wedge q) \rightarrow (p \wedge (q \vee \neg(r)))$$

Parsing a formula



Exercise: Parse the following formula

Exercise: Parse the following formula

$\neg q$

Exercise: Parse the following formula

$$p \wedge \neg q$$

Exercise: Parse the following formula

$$(p \wedge \neg q) \rightarrow r$$

Exercise: Parse the following formula

$$(p \wedge \neg q) \rightarrow r \quad \neg r \quad p$$

Exercise: Parse the following formula

$$(p \wedge \neg q) \rightarrow r \qquad \neg r \wedge p$$

Exercise: Parse the following formula


$$((p \wedge \neg q) \rightarrow r) \wedge (\neg r \wedge p)$$

Exercise: Parse the following formula

$$(((p \wedge \neg q) \rightarrow r) \wedge (\neg r \wedge p)) \rightarrow q$$


Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:




Reconstructing a formula

A formula can be reconstructed from its parse tree:



Reconstructing a formula

A formula can be reconstructed from its parse tree:



Omitting parentheses

Question

How do you read the arithmetic expression $3 \times 5 + 2$?

Is it 17 or 21 ?

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:



Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$\neg$		
$\wedge$	$\vee$	$\oplus$
$\rightarrow$	$\leftrightarrow$	

Examples

- ▶ $\neg(p) \vee q$
- ▶ $\neg(p \vee q)$
- ▶ $\neg(r) \rightarrow (p \vee q)$
- ▶ $r \wedge (p \vee q)$
- ▶ $\neg(\neg(p))$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$\neg$		
$\wedge$	$\vee$	$\oplus$
$\rightarrow$	$\leftrightarrow$	

Examples

- ▶ $\neg(p) \vee q \rightsquigarrow \neg p \vee q$
- ▶ $\neg(p \vee q)$
- ▶ $\neg(r) \rightarrow (p \vee q)$
- ▶ $r \wedge (p \vee q)$
- ▶ $\neg(\neg(p))$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$$\frac{\neg}{\frac{\wedge \vee \oplus}{\rightarrow \leftrightarrow}}$$

Examples

► $\neg(p) \vee q \rightsquigarrow \neg p \vee q$

► $\neg(p \vee q)$

► $\neg(r) \rightarrow (p \vee q)$

► $r \wedge (p \vee q)$

► $\neg(\neg(p))$

Note: $\nrightarrow \neg p \vee q$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$$\frac{\neg}{\frac{\wedge \vee \oplus}{\rightarrow \leftrightarrow}}$$

Examples

► $\neg(p) \vee q \rightsquigarrow \neg p \vee q$

► $\neg(p \vee q)$

Note: $\not\rightsquigarrow \neg p \vee q$

► $\neg(r) \rightarrow (p \vee q) \rightsquigarrow \neg r \rightarrow p \vee q$

► $r \wedge (p \vee q)$

► $\neg(\neg(p))$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$$\frac{\neg}{\frac{\wedge \vee \oplus}{\rightarrow \leftrightarrow}}$$

Examples

► $\neg(p) \vee q \rightsquigarrow \neg p \vee q$

► $\neg(p \vee q)$

Note: $\not\rightsquigarrow \neg p \vee q$

► $\neg(r) \rightarrow (p \vee q) \rightsquigarrow \neg r \rightarrow p \vee q$

► $r \wedge (p \vee q)$

Note: $\not\rightsquigarrow r \wedge p \vee q$

► $\neg(\neg(p))$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$$\frac{\neg}{\frac{\wedge \vee \oplus}{\rightarrow \leftrightarrow}}$$

Examples

► $\neg(p) \vee q \rightsquigarrow \neg p \vee q$

► $\neg(p \vee q)$

Note: $\not\rightsquigarrow \neg p \vee q$

► $\neg(r) \rightarrow (p \vee q) \rightsquigarrow \neg r \rightarrow p \vee q$

► $r \wedge (p \vee q)$

Note: $\not\rightsquigarrow r \wedge p \vee q$

► $\neg(\neg(p)) \rightsquigarrow \neg\neg p$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$\neg$		
$\wedge$	$\vee$	$\oplus$
$\rightarrow$	$\leftrightarrow$	

Examples

► $\neg(p) \vee q \rightsquigarrow \neg p \vee q$

► $\neg(p \vee q)$

Note: $\not\rightsquigarrow \neg p \vee q$

► $\neg(r) \rightarrow (p \vee q) \rightsquigarrow \neg r \rightarrow p \vee q$

► $r \wedge (p \vee q)$

Note: $\not\rightsquigarrow r \wedge p \vee q$

► $\neg(\neg(p)) \rightsquigarrow \neg\neg p$

Question

Which two pairs of parentheses cannot be omitted from

$$(((p \wedge \neg(q)) \rightarrow r) \wedge (\neg(r) \wedge p)) \rightarrow q \text{ ?}$$

Omitting parentheses

To omit parentheses from formulas, *without causing ambiguity*, we use the priority schema:

$$\frac{\neg}{\frac{\wedge \vee \oplus}{\rightarrow \leftrightarrow}}$$

Examples

► $\neg(p) \vee q \rightsquigarrow \neg p \vee q$

► $\neg(p \vee q)$

Note: $\nrightarrow \neg p \vee q$

► $\neg(r) \rightarrow (p \vee q) \rightsquigarrow \neg r \rightarrow p \vee q$

► $r \wedge (p \vee q)$

Note: $\nrightarrow r \wedge p \vee q$

► $\neg(\neg(p)) \rightsquigarrow \neg\neg p$

Question

$$(p \wedge \neg q \rightarrow r) \wedge (\neg r \wedge p) \rightarrow q$$

Exercise: Simplify and parse


$$((((\neg(\neg q)) \vee (\neg q)) \wedge q) \rightarrow (p \wedge r))$$

Remember:

$$\frac{\neg}{\frac{\wedge \vee \oplus}{\rightarrow \leftrightarrow}}$$


Parse tree

By the conventions regarding omitted parentheses,
the parse tree of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ is:



Parse tree

By the conventions regarding omitted parentheses, the parse tree of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ is:



Remark: Feel free to write all parentheses, if you prefer this

Syntax and semantics

The **syntax** is how a word (or formula) is *written*

The **semantics** is its *meaning*

Syntax and semantics

The **syntax** is how a word (or formula) is *written*


The **semantics** is its *meaning*



For example, the syntax for the animal shown above is *cow*

The semantics of *cow* is a female mammal, white with black spots, eats grass, gives milk, says moo

Propositional logic: Semantics



See? You forgot to close a parenthesis in its master control program.

Truth value semantics for propositional logic

Formulas of propositional logic are used to express declarative statements, which are either *true* or *false*

We introduce the **truth values** $\mathbf{T}$ and $\mathbf{F}$, corresponding to truth and falsehood, respectively

Truth value semantics for propositional logic

Formulas of propositional logic are used to express declarative statements, which are either *true* or *false*

We introduce the **truth values** $\mathbf{T}$ and $\mathbf{F}$, corresponding to truth and falsehood, respectively

The truth value of a formula such as $\phi \wedge \psi$ is determined by the truth values of its components, ϕ and ψ

For each connective, this is expressed by its **truth table**

Negation

A negation $\neg\phi$ (“not ϕ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is false} \\ \text{false} & \text{if } \phi \text{ is true} \end{cases}$$

Negation

A negation $\neg\phi$ (“not ϕ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is false} \\ \text{false} & \text{if } \phi \text{ is true} \end{cases}$$

Truth table for $\neg$:

ϕ	$\neg\phi$
T	
F	

T = true

F = false

Negation

A negation $\neg\phi$ (“not ϕ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is false} \\ \text{false} & \text{if } \phi \text{ is true} \end{cases}$$

Truth table for $\neg$:

ϕ	$\neg\phi$
T	F
F	T

T = true

F = false

Negation

A negation $\neg\phi$ (“not ϕ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is false} \\ \text{false} & \text{if } \phi \text{ is true} \end{cases}$$

Truth table for $\neg$:

ϕ	$\neg\phi$
T	F
F	T

T = true

F = false

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$$

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$
T	T	

We first list all possible combinations of assignments to ϕ and ψ

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$
T	T	
T	F	

We first list all possible combinations of assignments to ϕ and ψ

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$
T	T	
T	F	
F	T	
F	F	

We first list all possible combinations of assignments to ϕ and ψ

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$
T	T	
T	F	
F	T	
F	F	

We first list all possible combinations of assignments to ϕ and ψ

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$
T	T	T
T	F	
F	T	
F	F	

Conjunction

A conjunction $\phi \wedge \psi$ (“ ϕ and ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true and } \psi \text{ is true} \\ \text{false} & \text{in all other cases} \end{cases}$

Truth table for $\wedge$:

ϕ	ψ	$\phi \wedge \psi$
T	T	T
T	F	F
F	T	F
F	F	F

Disjunction

A disjunction $\phi \vee \psi$ (“ ϕ or ψ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is true or } \psi \text{ is true} \\ \text{false} & \text{otherwise} \end{cases}$$

Disjunction

A disjunction $\phi \vee \psi$ (“ ϕ or ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ is true or } \psi \text{ is true} \\ \text{false} & \text{otherwise} \end{cases}$

Truth table for $\vee$:

ϕ	ψ	$\phi \vee \psi$
T	T	
T	F	
F	T	
F	F	

Disjunction

A disjunction $\phi \vee \psi$ (“ ϕ or ψ ”) is

$$\begin{cases} \text{true} & \text{if } \phi \text{ is true or } \psi \text{ is true} \\ \text{false} & \text{otherwise} \end{cases}$$

Truth table for $\vee$:

ϕ	ψ	$\phi \vee \psi$
T	T	
T	F	T
F	T	T
F	F	

Disjunction

A disjunction $\phi \vee \psi$ (“ ϕ or ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if } \phi \text{ is true or } \psi \text{ is true} \\ \text{false} & \text{otherwise} \end{array} \right.$

Truth table for $\vee$:

ϕ	ψ	$\phi \vee \psi$
T	T	?
T	F	T
F	T	T
F	F	F

Disjunction

A disjunction $\phi \vee \psi$ (“ ϕ or ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if } \phi \text{ is true or } \psi \text{ is true (or both)} \\ \text{false} & \text{otherwise} \end{array} \right.$

Truth table for $\vee$:

ϕ	ψ	$\phi \vee \psi$
T	T	T
T	F	T
F	T	T
F	F	F

Examples in natural language:

- ▶ Inclusive:

Do you take sugar or cream in your coffee ?

Examples in natural language:

- ▶ Inclusive:

Do you take sugar or cream in your coffee ?

- ▶ Exclusive:

Do you want a cappuccino or an espresso ?

Exclusive or

An exclusive or $\phi \oplus \psi$ (“either ϕ or ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if either } \phi \text{ or } \psi \text{ is true (but not both)} \\ \text{false} & \text{otherwise} \end{array} \right.$

Exclusive or

An exclusive or $\phi \oplus \psi$ (“either ϕ or ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if either } \phi \text{ or } \psi \text{ is true (but not both)} \\ \text{false} & \text{otherwise} \end{array} \right.$

Truth table for $\oplus$:

ϕ	ψ	$\phi \oplus \psi$
T	T	
T	F	
F	T	
F	F	

Exclusive or

An exclusive or $\phi \oplus \psi$ (“either ϕ or ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if either } \phi \text{ or } \psi \text{ is true (but not both)} \\ \text{false} & \text{otherwise} \end{array} \right.$

Truth table for $\oplus$:

ϕ	ψ	$\phi \oplus \psi$
T	T	
T	F	T
F	T	T
F	F	

Exclusive or

An exclusive or $\phi \oplus \psi$ (“either ϕ or ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if either } \phi \text{ or } \psi \text{ is true (but not both)} \\ \text{false} & \text{otherwise} \end{array} \right.$

Truth table for $\oplus$:

ϕ	ψ	$\phi \oplus \psi$
T	T	F
T	F	T
F	T	T
F	F	F

Implication

$\phi \rightarrow \psi$ means: *if ϕ is true, then ψ is true*

Question

For which truth values of ϕ and ψ is $\phi \rightarrow \psi$ false ?

Implication

An implication $\phi \rightarrow \psi$ (“if ϕ , then ψ ”) is

$\begin{cases} \text{false} & \text{if } \phi \text{ is true and } \psi \text{ is false} \\ \text{true} & \text{otherwise} \end{cases}$

Truth table for $\rightarrow$:

ϕ	ψ	$\phi \rightarrow \psi$
T	T	
T	F	
F	T	
F	F	

Implication

An implication $\phi \rightarrow \psi$ (“if ϕ , then ψ ”) is

$\begin{cases} \text{false} & \text{if } \phi \text{ is true and } \psi \text{ is false} \\ \text{true} & \text{otherwise} \end{cases}$

Truth table for $\rightarrow$:

ϕ	ψ	$\phi \rightarrow \psi$
T	T	F
T	F	
F	T	
F	F	

Implication

An implication $\phi \rightarrow \psi$ (“if ϕ , then ψ ”) is

$\left\{ \begin{array}{ll} \text{false} & \text{if } \phi \text{ is true and } \psi \text{ is false} \\ \text{true} & \text{otherwise} \end{array} \right.$

Truth table for $\rightarrow$:

ϕ	ψ	$\phi \rightarrow \psi$
T	T	T
T	F	F
F	T	T
F	F	T

Implication

Truth table for $\rightarrow$:

Consider $(n > 2) \rightarrow (n + 1 > 2)$

Clearly, this implication is *true* for each n

ϕ	ψ	$\phi \rightarrow \psi$
T	T	
T	F	
F	T	
F	F	

Implication

Truth table for $\rightarrow$:

Consider $(n > 2) \rightarrow (n + 1 > 2)$

Clearly, this implication is *true* for each n

$n = 1$: both $n > 2$ and $n + 1 > 2$ are *false*

ϕ	ψ	$\phi \rightarrow \psi$
T	T	
T	F	
F	T	
F	F	T

Implication

Truth table for $\rightarrow$:

Consider $(n > 2) \rightarrow (n + 1 > 2)$

Clearly, this implication is *true* for each n

ϕ	ψ	$\phi \rightarrow \psi$
T	T	
T	F	
F	T	T
F	F	T

$n = 1$: both $n > 2$ and $n + 1 > 2$ are *false*

$n = 2$: $n > 2$ is *false* while $n + 1 > 2$ is *true*

Implication

Truth table for $\rightarrow$:

Consider $(n > 2) \rightarrow (n + 1 > 2)$

Clearly, this implication is *true* for each n

ϕ	ψ	$\phi \rightarrow \psi$
T	T	T
T	F	
F	T	T
F	F	T

$n = 1$: both $n > 2$ and $n + 1 > 2$ are *false*

$n = 2$: $n > 2$ is *false* while $n + 1 > 2$ is *true*

$n = 3$: both $n > 2$ and $n + 1 > 2$ are *true*

Implication

Truth table for $\rightarrow$:

ϕ	ψ	$\phi \rightarrow \psi$
T	T	T
T	F	
F	T	T
F	F	T

Consider $(n > 2) \rightarrow (n + 1 > 2)$

Clearly, this implication is *true* for each n

$n = 1$: both $n > 2$ and $n + 1 > 2$ are *false*

$n = 2$: $n > 2$ is *false* while $n + 1 > 2$ is *true*

$n = 3$: both $n > 2$ and $n + 1 > 2$ are *true*

Finally, consider $(n = 2) \rightarrow (n = 1)$

Implication

Truth table for $\rightarrow$:

ϕ	ψ	$\phi \rightarrow \psi$
T	T	T
T	F	F
F	T	T
F	F	T

Consider $(n > 2) \rightarrow (n + 1 > 2)$

Clearly, this implication is *true* for each n

$n = 1$: both $n > 2$ and $n + 1 > 2$ are *false*

$n = 2$: $n > 2$ is *false* while $n + 1 > 2$ is *true*

$n = 3$: both $n > 2$ and $n + 1 > 2$ are *true*

Finally, consider $(n = 2) \rightarrow (n = 1)$

This implication is *false* for $n = 2$ (but *true* for any other value of n)

Principle of explosion


In natural language,

if I'm a movie star, then I'm rich

makes sense, unlike

if I'm a movie star, then the moon is made of cheese

But they are both true (because I'm not a movie star)



Bi-implication

A bi-implication $\phi \leftrightarrow \psi$ (“ ϕ if and only if ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ and } \psi \text{ have the same truth value} \\ \text{false} & \text{otherwise} \end{cases}$

Truth table for $\leftrightarrow$:

ϕ	ψ	$\phi \leftrightarrow \psi$
T	T	
T	F	
F	T	
F	F	

Bi-implication

A bi-implication $\phi \leftrightarrow \psi$ (“ ϕ if and only if ψ ”) is

$\begin{cases} \text{true} & \text{if } \phi \text{ and } \psi \text{ have the same truth value} \\ \text{false} & \text{otherwise} \end{cases}$

Truth table for $\leftrightarrow$:

ϕ	ψ	$\phi \leftrightarrow \psi$
T	T	T
T	F	
F	T	
F	F	T

Bi-implication

A bi-implication $\phi \leftrightarrow \psi$ (“ ϕ if and only if ψ ”) is

$\left\{ \begin{array}{ll} \text{true} & \text{if } \phi \text{ and } \psi \text{ have the same truth value} \\ \text{false} & \text{otherwise} \end{array} \right.$

Truth table for $\leftrightarrow$:

ϕ	ψ	$\phi \leftrightarrow \psi$
T	T	T
T	F	F
F	T	F
F	F	T

Exercise: What are we missing?

- ▶ How many other logical operators are there?
- ▶ Let me introduce Sheffer Stroke " $|$ "

Truth values and valuations

The truth value of a formula is determined by an *assignment of truth values* to its propositional variables

Such an assignment is called a *valuation*

Truth values and valuations

The truth value of a formula is determined by an *assignment of truth values* to its propositional variables

Such an assignment is called a *valuation*

Example

Given the valuation: p and q are $\mathbb{T}$, and r is $\mathbb{F}$

What is the truth value of the formula $p \vee \neg q \rightarrow r$?

Truth values and valuations

The truth value of a formula is determined by an *assignment of truth values* to its propositional variables

Such an assignment is called a *valuation*

Example

Given the valuation: p and q are T , and r is F

What is the truth value of the formula $p \vee \neg q \rightarrow r$?

We reason:

- ▶ $p \vee \neg q$ is T because p is T

Truth values and valuations

The truth value of a formula is determined by an *assignment of truth values* to its propositional variables

Such an assignment is called a *valuation*

Example

Given the valuation: p and q are T , and r is F

What is the truth value of the formula $p \vee \neg q \rightarrow r$?

We reason:

- ▶ $p \vee \neg q$ is T because p is T
- ▶ So, since r is F , it follows that $p \vee \neg q \rightarrow r$ is F

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T			
T	T	F			
T	F	T			
T	F	F			
F	T	T			
F	T	F			
F	F	T			
F	F	F			

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T			
T	T	F			
T	F	T			
T	F	F			
F	T	T			
F	T	F			
F	F	T			
F	F	F			

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T	F		
T	T	F			
T	F	T			
T	F	F			
F	T	T			
F	T	F			
F	F	T			
F	F	F			

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T			
T	T	F	F	T	
T	F	T			
T	F	F			
F	T	T			
F	T	F			
F	F	T			
F	F	F			

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T			
T	T	F	F	T	F
T	F	T			
T	F	F			
F	T	T			
F	T	F			
F	F	T			
F	F	F			

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T	F	T	T
T	T	F	F	T	F
T	F	T	T	T	T
T	F	F	T	T	F
F	T	T	F	F	T
F	T	F	F	F	T
F	F	T	T	T	T
F	F	F	T	T	F

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T	F	T	T
T	T	F	F	T	F
T	F	T	T	T	T
T	F	F	T	T	F
F	T	T	F	F	T
F	T	F	F	F	T
F	F	T	T	T	T
F	F	F	T	T	F

For $p \vee \neg q \rightarrow r$ there are 8 ($= 2^3$) rows in the truth table

Valuations of a formula in a truth table

A *valuation* corresponds to one row in the truth table of a formula

Example: Truth table of $p \vee \neg q \rightarrow r$

p	q	r	$\neg q$	$p \vee \neg q$	$p \vee \neg q \rightarrow r$
T	T	T	F	T	T
T	T	F	F	T	F
T	F	T	T	T	T
T	F	F	T	T	F
F	T	T	F	F	T
F	T	F	F	F	T
F	F	T	T	T	T
F	F	F	T	T	F

For $p \vee \neg q \rightarrow r$ there are 8 ($= 2^3$) rows in the truth table

Question

How many rows has a truth table for a formula with n variables ?

Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

▶ $p = \text{T}$

▶ $q = \text{F}$

▶ $r = \text{T}$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

► $p = \text{T}$

► $q = \text{F}$

► $r = \text{T}$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

► $p = \text{T}$

► $q = \text{F}$

► $r = \text{T}$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

▶ $p = \text{T}$

▶ $q = \text{F}$

▶ $r = \text{T}$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

► $p = \text{T}$

► $q = \text{F}$

► $r = \text{T}$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

► $p = \text{T}$

► $q = \text{F}$

► $r = \text{T}$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

► $p = T$

► $q = F$

► $r = T$



Bottom-up truth assignment via a parse tree


The truth value of a formula can be computed via its *parse tree*

Bottom-up evaluation of $\neg p \wedge q \rightarrow p \wedge (q \vee \neg r)$ with:

► $p = T$

► $q = F$

► $r = T$



Semantic equivalence $\equiv$

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T			
T	F			
F	T			
F	F			

Semantic equivalence $\equiv$

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T	T		
T	F	F		
F	T	T		
F	F	T		

Semantic equivalence $\equiv$

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T	T	F	
T	F	F	F	
F	T	T	T	
F	F	T	T	

Semantic equivalence $\equiv$

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T	T	F	T
T	F	F	F	F
F	T	T	T	T
F	F	T	T	T

Semantic equivalence $\equiv$

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T	T	F	T
T	F	F	F	F
F	T	T	T	T
F	F	T	T	T

The columns for $p \rightarrow q$ and $\neg p \vee q$ are identical

Semantic equivalence $\equiv$

Definition

Formulas ϕ and ψ are *semantically equivalent*, notation $\phi \equiv \psi$, if they have identical columns in their truth tables

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T	T	F	T
T	F	F	F	F
F	T	T	T	T
F	F	T	T	T

The columns for $p \rightarrow q$ and $\neg p \vee q$ are identical

Semantic equivalence $\equiv$

Definition

Formulas ϕ and ψ are *semantically equivalent*, notation $\phi \equiv \psi$, if they have identical columns in their truth tables

Example:

Truth tables for $p \rightarrow q$ and $\neg p \vee q$:

p	q	$p \rightarrow q$	$\neg p$	$\neg p \vee q$
T	T	T	F	T
T	F	F	F	F
F	T	T	T	T
F	F	T	T	T

The columns for $p \rightarrow q$ and $\neg p \vee q$ are identical

Hence: $p \rightarrow q \equiv \neg p \vee q$

Expressing connectives

$\leftrightarrow$ can be expressed using $\rightarrow$ and $\wedge$:

$$\phi \leftrightarrow \psi \quad \equiv \quad (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi)$$

Expressing connectives

$\leftrightarrow$ can be expressed using $\rightarrow$ and $\wedge$:

$$\phi \leftrightarrow \psi \quad \equiv \quad (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi)$$

Questions

Express $\phi \oplus \psi$ using $\vee$, $\wedge$ and $\neg$

Expressing connectives

$\leftrightarrow$ can be expressed using $\rightarrow$ and $\wedge$:

$$\phi \leftrightarrow \psi \quad \equiv \quad (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi)$$

Questions

Express $\phi \oplus \psi$ using $\vee$, $\wedge$ and $\neg$

Express $\phi \wedge \psi$ using $\vee$ and $\neg$

Expressing connectives

$\leftrightarrow$ can be expressed using $\rightarrow$ and $\wedge$:

$$\phi \leftrightarrow \psi \quad \equiv \quad (\phi \rightarrow \psi) \wedge (\psi \rightarrow \phi)$$

Questions

Express $\phi \oplus \psi$ using $\vee$, $\wedge$ and $\neg$

Express $\phi \wedge \psi$ using $\vee$ and $\neg$

Express $\phi \vee \psi$ using $\wedge$ and $\neg$

Semantic equivalences

$$\neg\neg\phi \equiv \phi$$

Semantic equivalences

$$\neg\neg\phi \equiv \phi$$

$$\phi \wedge \psi \equiv \neg(\neg\phi \vee \neg\psi)$$

Semantic equivalences

$$\neg\neg\phi \equiv \phi$$

$$\phi \wedge \psi \equiv \neg(\neg\phi \vee \neg\psi)$$

So

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$$

Semantic equivalences

$$\neg\neg\phi \equiv \phi$$

$$\phi \wedge \psi \equiv \neg(\neg\phi \vee \neg\psi)$$

So

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$$

$$\phi \vee \psi \equiv \neg(\neg\phi \wedge \neg\psi)$$

So

$$\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$$

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi$$

$$\phi \vee \psi \equiv \psi \vee \phi$$

commutativity

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi$$

commutativity

$$\phi \vee \psi \equiv \psi \vee \phi$$

$$\phi \wedge \phi \equiv \phi$$

idempotence

$$\phi \vee \phi \equiv \phi$$

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi$$

commutativity

$$\phi \vee \psi \equiv \psi \vee \phi$$

$$\phi \wedge \phi \equiv \phi$$

idempotence

$$\phi \vee \phi \equiv \phi$$

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

associativity

$$(\phi \vee \psi) \vee \chi \equiv \phi \vee (\psi \vee \chi)$$

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi$$

commutativity

$$\phi \vee \psi \equiv \psi \vee \phi$$

$$\phi \wedge \phi \equiv \phi$$

idempotence

$$\phi \vee \phi \equiv \phi$$

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

associativity

$$(\phi \vee \psi) \vee \chi \equiv \phi \vee (\psi \vee \chi)$$

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$$

De Morgan

$$\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$$

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi$$

commutativity

$$\phi \vee \psi \equiv \psi \vee \phi$$

$$\phi \wedge \phi \equiv \phi$$

idempotence

$$\phi \vee \phi \equiv \phi$$

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

associativity

$$(\phi \vee \psi) \vee \chi \equiv \phi \vee (\psi \vee \chi)$$

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$$

De Morgan

$$\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$$

$$\neg\neg\phi \equiv \phi$$

involution

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi$$

commutativity

$$\phi \vee \psi \equiv \psi \vee \phi$$

$$\phi \wedge \phi \equiv \phi$$

idempotence

$$\phi \vee \phi \equiv \phi$$

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

associativity

$$(\phi \vee \psi) \vee \chi \equiv \phi \vee (\psi \vee \chi)$$

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$$

De Morgan

$$\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$$

$$\neg\neg\phi \equiv \phi$$

involution

$$\phi \rightarrow \psi \equiv \neg\phi \vee \psi$$

Important semantic equivalences

$$\phi \wedge \psi \equiv \psi \wedge \phi \quad \text{commutativity}$$

$$\phi \vee \psi \equiv \psi \vee \phi$$

$$\phi \wedge \phi \equiv \phi \quad \text{idempotence}$$

$$\phi \vee \phi \equiv \phi$$

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi) \quad \text{associativity}$$

$$(\phi \vee \psi) \vee \chi \equiv \phi \vee (\psi \vee \chi)$$

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi \quad \text{De Morgan}$$

$$\neg(\phi \vee \psi) \equiv \neg\phi \wedge \neg\psi$$

$$\neg\neg\phi \equiv \phi \quad \text{involution}$$

$$\phi \rightarrow \psi \equiv \neg\phi \vee \psi$$

Try to develop **intuition** for semantic equivalences by practicing
(like learning arithmetic as a child)

Exercise: Let us prove de Morgan

$$\neg(\phi \wedge \psi) \equiv \neg\phi \vee \neg\psi$$

Associativity of conjunction and disjunction

Conjunction is **associative**

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

Associativity of conjunction and disjunction

Conjunction is **associative**

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

So we can skip parentheses: $\phi \wedge \psi \wedge \chi$

Associativity of conjunction and disjunction

Conjunction is **associative**

$$(\phi \wedge \psi) \wedge \chi \equiv \phi \wedge (\psi \wedge \chi)$$

So we can skip parentheses: $\phi \wedge \psi \wedge \chi$

Disjunction is also associative

$$(\phi \vee \psi) \vee \chi \equiv \phi \vee (\psi \vee \chi)$$

So we can skip parentheses: $\phi \vee \psi \vee \chi$

Is the exclusive or associative ?

Question

Does $(\phi \oplus \psi) \oplus \chi \equiv \phi \oplus (\psi \oplus \chi)$ hold ?

1. Yes

2. No

Is the exclusive or associative ?

Question

Does $(\phi \oplus \psi) \oplus \chi \equiv \phi \oplus (\psi \oplus \chi)$ hold ?

1. Yes

2. No

Feel free to discuss with your neighbors



Associativity of exclusive or

$\phi \oplus \psi \oplus \chi$ is true *if and only if* an odd number of the arguments ϕ , ψ and χ is true

ϕ	ψ	χ	$(\phi \oplus \psi) \oplus \chi$	$\phi \oplus (\psi \oplus \chi)$
T	T	T	T	T
T	T	F	F	F
T	F	T	F	F
T	F	F	T	T
F	T	T	F	F
F	T	F	T	T
F	F	T	T	T
F	F	F	F	F

Is implication associative ?

Question

Does $(\phi \rightarrow \psi) \rightarrow \chi \equiv \phi \rightarrow (\psi \rightarrow \chi)$ hold ?

1. Yes
2. No

Is implication associative ?

Question

Does $(\phi \rightarrow \psi) \rightarrow \chi \equiv \phi \rightarrow (\psi \rightarrow \chi)$ hold ?

1. Yes

2. No

ϕ	ψ	χ	$(\phi \rightarrow \psi) \rightarrow \chi$	$\phi \rightarrow (\psi \rightarrow \chi)$
T	T	T	T	T
T	T	F	F	F
T	F	T	T	T
T	F	F	T	T
F	T	T	T	T
F	T	F	F	T
F	F	T	T	T
F	F	F	F	T

Final quiz question

Question

Does $\phi \rightarrow (\psi \rightarrow \chi) \equiv \phi \wedge \psi \rightarrow \chi$ hold?

1. Yes

2. No

Final quiz question

Question

Does $\phi \rightarrow (\psi \rightarrow \chi) \equiv \phi \wedge \psi \rightarrow \chi$ hold?

1. Yes

2. No

ϕ	ψ	χ	$\psi \rightarrow \chi$	$\phi \rightarrow (\psi \rightarrow \chi)$	$\phi \wedge \psi$	$\phi \wedge \psi \rightarrow \chi$
T	T	T	T	T	T	T
T	T	F	F	F	T	F
T	F	T	T	T	F	T
T	F	F	T	T	F	T
F	T	T	T	T	F	T
F	T	F	F	T	F	T
F	F	T	T	T	F	T
F	F	F	T	T	F	T

Tautologies, contradictions and contingent formulas

Tautology = always true

$$p \vee \neg p$$

Tautologies, contradictions and contingent formulas

Tautology = always true

$$p \vee \neg p$$

Contradiction = always false

$$p \wedge \neg p$$

Tautologies, contradictions and contingent formulas

Tautology = always true

$$p \vee \neg p$$

Contradiction = always false

$$p \wedge \neg p$$

Contingent = sometimes true and sometimes false

$$p$$
$$p \wedge \neg q$$

Tautologies

Definition

A formula is a **tautology** (also called *valid*) if:
its column in a truth table has **T** on every row

So: a tautology is true for every valuation

Tautologies

Definition

A formula is a **tautology** (also called *valid*) if:
its column in a truth table has **T** on every row

So: a tautology is true for every valuation

Examples:

▶ $p \vee \neg p$

▶ $p \rightarrow p$

▶ $p \rightarrow (q \rightarrow p)$



Tautologies

Definition

A formula is a **tautology** (also called *valid*) if:
its column in a truth table has **T** on every row

So: a tautology is true for every valuation

Examples:

- ▶ $p \vee \neg p$
- ▶ $p \rightarrow p$
- ▶ $p \rightarrow (q \rightarrow p)$
- ▶

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$
T	T		
T	F		
F	T		
F	F		

Tautologies

Definition

A formula is a **tautology** (also called *valid*) if:
its column in a truth table has **T** on every row

So: a tautology is true for every valuation

Examples:

- ▶ $p \vee \neg p$
- ▶ $p \rightarrow p$
- ▶ $p \rightarrow (q \rightarrow p)$
- ▶

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$
T	T	T	
T	F	T	
F	T	F	
F	F	T	

Tautologies

Definition

A formula is a **tautology** (also called *valid*) if:
its column in a truth table has **T** on every row

So: a tautology is true for every valuation

Examples:

- ▶ $p \vee \neg p$
- ▶ $p \rightarrow p$
- ▶ $p \rightarrow (q \rightarrow p)$
- ▶

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$
T	T	T	T
T	F	T	T
F	T	F	T
F	F	T	T

Tautologies

Definition

A formula is a **tautology** (also called *valid*) if:
its column in a truth table has **T** on every row

So: a tautology is true for every valuation

Examples:

► $p \vee \neg p$

► $p \rightarrow p$

► $p \rightarrow (q \rightarrow p)$

► $(p \wedge \neg q \rightarrow r) \wedge \neg r \wedge p \rightarrow q$

p	q	$q \rightarrow p$	$p \rightarrow (q \rightarrow p)$
T	T	T	T
T	F	T	T
F	T	F	T
F	F	T	T

(The example of Jane)

Contradictions

Definition

A formula is a **contradiction** if:
its column in a truth table has **F** on every row

So: a contradiction is false for every valuation

Contradictions

Definition

A formula is a **contradiction** if:
its column in a truth table has **F** on every row

So: a contradiction is false for every valuation

Examples:

- ▶ $p \wedge \neg p$
- ▶ $(p \rightarrow q) \wedge p \wedge \neg q$

Contradictions

Definition

A formula is a **contradiction** if:

its column in a truth table has **F** on every row

So: a contradiction is false for every valuation

Examples:

► $p \wedge \neg p$

► $(p \rightarrow q) \wedge p \wedge \neg q$

p	q	$p \rightarrow q$	$\neg q$	$(p \rightarrow q) \wedge p \wedge \neg q$
T	T	T	F	F
T	F	F	T	F
F	T	T	F	F
F	F	T	T	F

More contradictions, and contingent formulas

Questions

Are the following formulas contradictions ?

► $p \rightarrow \neg p$

More contradictions, and contingent formulas

Questions

Are the following formulas contradictions ?

► $p \rightarrow \neg p$

► $((p \rightarrow q) \rightarrow p) \wedge \neg p$

More contradictions, and contingent formulas

Questions

Are the following formulas contradictions ?

▶ $p \rightarrow \neg p$

▶ $((p \rightarrow q) \rightarrow p) \wedge \neg p$

▶ $\neg \phi$ with ϕ a tautology

More contradictions, and contingent formulas

Questions

Are the following formulas contradictions ?

- ▶ $p \rightarrow \neg p$
- ▶ $((p \rightarrow q) \rightarrow p) \wedge \neg p$
- ▶ $\neg \phi$ with ϕ a tautology

Definition

A propositional formula is **contingent** if:
it is neither a tautology, nor a contradiction

Introduction to propositional logic

- ▶ from natural language to formal notation
- ▶ syntax of propositional logic
 - ▶ from formula to parse tree, and back
- ▶ truth value semantics
 - ▶ truth tables, valuations
- ▶ semantic equivalence
- ▶ tautologies and contradictions

Introduction to propositional logic

- ▶ from natural language to formal notation
- ▶ syntax of propositional logic
 - ▶ from formula to parse tree, and back
- ▶ truth value semantics
 - ▶ truth tables, valuations
- ▶ semantic equivalence
- ▶ tautologies and contradictions

Exercise classes today and tomorrow

- ▶ *Enroll for working groups in Canvas*
- ▶ *Try your hand at the exercises !*