ECS 122A Lecture 9 Jasper Lee

Shortest Path Algorithms

Setting: Given directed weighted graph (each edge e has weight w(e)) may be negative

compute shortest paths b/w vertices.

Variations:

- Single source: given vertex u, compute shortest path from u to all other vertices v
- All pairs: for all pairs u, v, compute shortest u to v path.

Single-source shortest path

Goal: Derive the Bellman-Ford alg.

Different intuition from basic "either or"

DP table?

D for distance → D[v] = shortest path len from u to v

Just rephrased original problem...

Q: DP needs subproblems. What are the subproblems here?

Any path is a seq of hops through graph.

⇒ Limit the # of hops for subproblems

DP table

$D[v, k] \doteq$ shortest path from u to v using $\leq k$ hops.

Equivalently, insert self loops of weight 0 into graph

$D[v, k] \doteq$ shortest path from u to v using exactly k hops.

Recurrence

$$D[v,0] \doteq \begin{cases} \infty & \text{if } v \neq u \\ 0 & \text{if } v = u \end{cases}$$

$$D[v,k] = \min\left(D[v,k-1] + w_{v \to v},\ \min_{x \neq v} D[x,k-1] + w_{x \to v} \right)$$

($w_{v \to v} = 0$; non-existent edges have weight ∞)

How large can k be? $n-1$

Implementation:

Initialize $D \leftarrow \infty$, $D[u,0] = 0$.

for $k = 1$ to $n-1$

  for each edge $a \to v$

    $D[v,k] \leftarrow \min(D[v,k], w_{a \to v} + D[a,k-1])$

$O(|V||E|)$ time

only enumerate edges, instead of all vertex pairs (v, x)

(without constraint on #hops)

Issue: Shortest path undefined if there is negative cycle reachable from u...

Claim: There is a negative cycle reachable from u if and only if

$\exists v$ s.t. $D[v,n] < D[v,n-1]$.

Proof

$\exists v \Rightarrow$ neg cycle

Let $u \to q_1 \to q_2 \to \cdots \to q_{n-1} \to v$ (= q_n) be a path of min weight for $D[v,n]$.

Since there are only n vertices, by pigeonhole, $\exists\, a_i = a_j$ for $i < j$

Now, the $n-1$ hop path (padded to $n-1$ hops)

$u \to q_1 \to \cdots \to a_i \to a_{j+1} \to v\ (\to \cdots \to v)$

has weight $\geq D[v, n-1] > D[v,n]$.

So $a_i \to \cdots \to a_j = a_i$ is a negative cycle.

Neg cycle $\Rightarrow \exists v$

$(\forall v, D[v,n] \geq D[v,n-1]) \Rightarrow$ No neg cycle

Consider any cycle $a_1 \to \cdots \to a_1$

$D[a_{i+1}, n-1] \leq D[a_{i+1}, n] \leq D[a_i, n-1] + w_{a_i a_{i+1}}$

Summing over cycle $\Rightarrow \sum w_{a_i a_{i+1}} \geq 0$.

Alg: Do one more inner loop to check if there are any updates.
If yes ⇒ neg cycle
no ⇒ no neg cycle.

Optimizing implementation:

- Only need to keep index $k-1$ to compute index k.

- In fact, can even ignore index and keep updating

Initialize $D[v] \leftarrow \infty$ for all $v \neq v$, $D[v] \leftarrow 0$

for $k = 1$ to $n-1$
  for each edge $u \to v$
    $D[v] \leftarrow \min(D[v], w_{u \to v} + D[u])$.

Do another inner loop to check neg cycle

Correctness proof slightly messier.

All-pairs shortest path

If we run Bellman-Ford on all source vertices, we can get all-pairs shortest path in $O(|V|^2|E|)$ time.

Floyd-Warshall improves to $O(|V|^3)$ time.

Assumes no negative cycles

$|E|$ can be as big as $O(|V|^2)$

$|E| \geq |V|$ so no worse than repeating Bellman-Ford.

Subproblem structure not necessarily obv

$D[i,j,k] =$ shortest path from i to j where intermediate vertices are all in $\{1..k\}$.

Recurrence

$$D[i,j,k] = \min(D[i,j,k-1], D[i,k,k-1] + D[k,j,k-1])$$

$D[i,j,0] = w_{i \to j}$ (which may be ∞)

Correctness:
Either use vertex k or not.
If using, must reach k using $\{1..k-1\}$ and then reach j using $\{1..k-1\}$.

```
for k = 1 to n
    for i = 1 to n
        for j = 1 to n
            D[i,j,k] ← min(D[i,j,k-1], D[i,k,k-1]+D[k,j,k-1])
```

Same optimization as before

$D[i,j] \leftarrow \min(D[i,j], D[i,k] + D[k,j])$