## Age Distribution by Wearable Clusters

$W_{\text{Mann-Whitney}} = 18.50$, $p = 0.30$, $\hat{r}_{\text{biserial}}^{\text{rank}} = -0.34$, $\text{CI}_{95\%}$ [−0.74, 0.25], $n_{\text{obs}} = 15$

$\hat{\mu}_{\text{median}} = 52.00$

$\hat{\mu}_{\text{median}} = 57.00$

Wearable Cluster 1 (n = 7)

Wearable Cluster 2 (n = 8)

Wearable Clusters

Age (years)

Age Distribution by Gender (Wearable Cohort)

$W_{\text{Mann-Whitney}} = 32.00$, $p = 0.68$, $\hat{r}_{\text{biserial}}^{\text{rank}} = 0.14$, $\text{CI}_{95\%}$ [−0.43, 0.63], $n_{\text{obs}} = 15$

$\hat{\mu}_{\text{median}} = 53.00$

$\hat{\mu}_{\text{median}} = 55.00$

Female (n = 7)

Male (n = 8)

Gender

Age (years)

Gender Distribution (Wearable Cohort)

PPV Patients (N = 15)

7 (46.7%)

8 (53.3%)

Total: 15

(r0 + r1)/2

Freq

group

Female

Male

Wearable Cluster Distribution

PPV Patients (N = 15)

7 (46.7%)

8 (53.3%)

Total: 15

(r0 + r1)/2

Freq

group

Wearable Cluster 1

Wearable Cluster 2