Assignment 5

Jatin Tarachandani-CS20BTECH11021

May 2021

Download latex codes from

https://github.com/jatin-tarachandani/AI1103/blob/main/Assignment_5/main.tex

1 Problem Statement

Let X, Y, have the joint discrete distribution such that $X|Y = y \sim \text{Binomial}(y, 0.5)$ and $Y \sim \text{Poisson}(\lambda)$, $\lambda > 0$, where λ is an unknown parameter. Let T = T(X, Y) be any unbiased estimator of λ . Then

- 1. $Var(T) \leq Var(Y)$ for all λ
- 2. $Var(T) \ge Var(Y)$ for all λ
- 3. $Var(T) \ge \lambda$ for all λ
- 4. Var(T) = Var(Y) for all λ

2 Solution

Definition 2.1. Suppose we have an estimator T(X, Y), acting on a set of 2 RVs X, Y; which estimates a parameter θ , then if

$$E(T(X,Y)) = \theta \tag{2.0.1}$$

the estimator is unbiased.

Lemma 2.1. Since Y has a Poisson distribution, we know that:

$$Var(Y) = \lambda \tag{2.0.2}$$

From Bayes Theorem,

$$Pr(X = x, Y = y) = Pr(Y = y) Pr(X = x | Y = y)$$
 (2.0.3)

$$= {}^{y}C_{x}\frac{1}{2^{y}}\frac{\lambda^{y}}{y!}e^{-\lambda}$$
 (2.0.4)

Let us represent Pr(X = x, Y = y) as a function of x, y, and λ : $f(x, y; \lambda)$.

Definition 2.2. If T(X, Y) is an unbiased estimator of a parameter λ , the Cramer-Rao bound states that:

$$Var(T(X,Y)) \ge -\frac{1}{E\left(\frac{\partial^2 \ln(f(x,y;\lambda))}{\partial \lambda^2}\right)}$$
 (2.0.5)

From using (2.0.5) on T(X, Y),

$$Var(T(X,Y)) \ge -\frac{1}{E\left(\frac{\partial^2 \ln(f(x,y;\lambda))}{\partial \lambda^2}\right)}$$
 (2.0.6)

$$\geq -\frac{1}{E\left(-\frac{y}{\lambda^2}\right)}\tag{2.0.7}$$

$$\geq \frac{\lambda^2}{E(y)} \tag{2.0.8}$$

$$\geq \lambda$$
 (2.0.9)

because the expectation value of a Poisson distribution with parameter λ is λ .

The correct options are options (2) and (3), since by (2.0.9), we see that

$$Var(T) \ge \lambda = Var(Y)$$

(from (2.0.2)). (1) and (4) do not hold for all λ .