ARJUNA JEE 2027

Lecture -05 | Mathematics

TRIGONOMETRIC FUNCTIONS

By- Sachin Jakhar Sir



Topics *to be covered*

A **Value at Larger Angles**

B **Compound Angle Formula**

C **Question Practice**

D





LAST CLASS RECAP

Dealing with Negative Angle:

$$\cos(-\theta) = \cos\theta$$

$$\sec(-\theta) = \sec\theta$$

$$\sin(-\theta) = -\sin\theta$$

$$\tan(-\theta) = -\tan\theta$$

$$\operatorname{cosec}(-\theta) = -\operatorname{cosec}\theta$$

$$\cot(-\theta) = -\cot\theta$$

Question Practice

Question

If θ is an acute angle such that $5\cot\theta = 3$, then find the value of $\frac{(5\sin\theta - 3\cos\theta)}{(4\sin\theta + 3\cos\theta)}$

A $\frac{11}{18}$

B $\frac{16}{29}$


C $\frac{14}{27}$

D None of these

$0^\circ < A.A. < 90^\circ$

I quad.

$\cot\theta = \frac{3}{5} = \frac{B}{P}$



$\sin\theta = +\frac{5}{\sqrt{34}}$

$\cos\theta = +\frac{3}{\sqrt{34}}$

$\theta \rightarrow$ acute

II. $\leftarrow$ $\frac{\pi}{2} + \theta$

I. $\leftarrow$ $\frac{\pi}{2} - \theta$

III. $\leftarrow$ $\pi + \theta$


II. $\leftarrow$ $\pi - \theta$

IV. $\leftarrow$ $\frac{3\pi}{2} + \theta$


III. $\leftarrow$ $\frac{3\pi}{2} - \theta$

I. $\leftarrow$ $2\pi + \theta$

IV. $\leftarrow$ $2\pi - \theta$



REDUCTION FORMULA



$2\pi - \theta = 4\frac{\pi}{2} - \theta$

(1) $\sin(2\pi - \theta) = -\sin\theta$ (IV)

(2) $\tan\left(\frac{3\pi}{2} + \theta\right) = -\cot\theta$ (IV)

(3) $\text{cosec}(5\pi - \theta) = +\text{cosec}\,\theta$ (II)

(4) $\sec(2\pi + \theta) = +\sec\theta$ (I)

$4\frac{\pi}{2}$

(5) $\cos\left(\frac{5\pi}{2}-\theta\right) = +\sin\theta$ (I)

(6) $\sin\left(\theta-\frac{3\pi}{2}\right) = \sin\left(-\left(\frac{3\pi}{2}-\theta\right)\right) = -\sin\left(\frac{3\pi}{2}-\theta\right) = -\left(-\cos\theta\right) = +\cos\theta$. (III)

(7) $\cos\left(\theta-\frac{\pi}{2}\right) = \cos\left(-\left(\frac{\pi}{2}-\theta\right)\right) = \cos\left(\frac{\pi}{2}-\theta\right) = +\sin\theta$. (I)

RAPID FIRE ROUND

1. $\sin(131\pi - \theta) =$ _____.
2. $\cos\left(57\frac{\pi}{2} - \theta\right) =$ _____.
3. $\cos(82\pi - \theta) =$ _____.
4. $\cos\left(33\frac{\pi}{2} + \theta\right) =$ _____.
5. $\tan(183 \times 90^\circ + \theta) =$ _____.
6. $\tan(181 \times 90^\circ - \theta) =$ _____.
7. $\cot(57 \times 90^\circ + \theta) =$ _____.
8. $\sec\left(323\frac{\pi}{2} - \theta\right) =$ _____.
9. $\operatorname{cosec}(62\pi + \theta) =$ _____.
10. $\operatorname{cosec}\left(\theta - 71\frac{\pi}{2}\right) =$ _____.



REDUCTION FORMULA


$\pi - \theta = (2)\frac{\pi}{2} - \theta$

$\pi + \theta = (2)\frac{\pi}{2} + \theta$

I	II
$\sin\left(\frac{\pi}{2}-\theta\right) = +\cos\theta$	$\sin\left(\frac{\pi}{2}+\theta\right) = +\cos\theta$
$\cos\left(\frac{\pi}{2}-\theta\right) = +\sin\theta$	$\cos\left(\frac{\pi}{2}+\theta\right) = -\sin\theta$
$\tan\left(\frac{\pi}{2}-\theta\right) = +\cot\theta$	$\tan\left(\frac{\pi}{2}+\theta\right) = -\cot\theta$
$\cot\left(\frac{\pi}{2}-\theta\right) = +\tan\theta$	$\cot\left(\frac{\pi}{2}+\theta\right) = -\tan\theta$
$\operatorname{cosec}\left(\frac{\pi}{2}-\theta\right) = +\sec\theta$	$\operatorname{cosec}\left(\frac{\pi}{2}+\theta\right) = +\sec\theta$
$\sec\left(\frac{\pi}{2}-\theta\right) = +\operatorname{cosec}\theta$	$\sec\left(\frac{\pi}{2}+\theta\right) = -\operatorname{cosec}\theta$

II	III
$\sin(\pi-\theta) = +\sin\theta$	$\sin(\pi+\theta) = -\sin\theta$
$\cos(\pi-\theta) = -\cos\theta$	$\cos(\pi+\theta) = -\cos\theta$
$\tan(\pi-\theta) = -\tan\theta$	$\tan(\pi+\theta) = +\tan\theta$
$\cot(\pi-\theta) = -\cot\theta$	$\cot(\pi+\theta) = +\cot\theta$
$\operatorname{cosec}(\pi-\theta) = +\operatorname{cosec}\theta$	$\operatorname{cosec}(\pi+\theta) = -\operatorname{cosec}\theta$
$\sec(\pi-\theta) = -\sec\theta$	$\sec(\pi+\theta) = -\sec\theta$

Class-X.



$\sin\theta = \frac{a}{c}$


$\cos\theta = \frac{b}{c}$

$\sin\left(\frac{\pi}{2}-\theta\right) = \cos\theta$

$\sin\left(\frac{\pi}{2}-\theta\right) = \frac{b}{c}$


$\cos\left(\frac{\pi}{2}-\theta\right) = \frac{a}{c}$

$\sin\theta = \cos\left(\frac{\pi}{2}-\theta\right)$



$\sin\theta = \frac{b}{r}$


$\cos\theta = \frac{a}{r}$



$\frac{\pi}{2}+\theta$

$\sin(90^\circ+\theta) = \frac{a}{r} = \cos\theta$

$\cos(90^\circ+\theta) = -\frac{b}{r} = -\sin\theta$



Congurent ✓

$\Delta ODC \cong \Delta BOA$

$\frac{OD}{BO} = \frac{DC}{OA} = \frac{OC}{BA}$

$1 = \frac{DC}{a} = \frac{OC}{b}$

Module Question

The value of $\cos(270^\circ + \theta) \cdot \cos(90^\circ - \theta) - \sin(270^\circ - \theta)\cos\theta$ is

A 0

B −1

C 1/2

D 1

$\cos(3\times 90^\circ + \theta)$ — IV

$(+\sin\theta)(+\sin\theta)$ — $\sin(3\times 90^\circ - \theta)$ — III

$(-\cos\theta)\cos\theta$

$\sin^2\theta + \cos^2\theta = 1$

VALUE AT LARGER ANGLES

Process

↳ Same as Reduc^{n} formula.

$240^\circ = 2\times 90^\circ + 60^\circ$
$= 3\times 90^\circ - 30^\circ$

Find value of following:

(1) $\cos(240^\circ)$ (III) $= \cos(2\times 90^\circ + 60^\circ) = -\cos 60^\circ = -\frac{1}{2}$ ✓ (OR $\cos(3\times 90^\circ - 30^\circ) = -\sin 30^\circ = -\frac{1}{2}$ ✓)

(2) $\tan(-300^\circ)$ $= -\tan(300^\circ) = -\tan(3\times 90^\circ + 30^\circ)$ (IV) $= +(+\cot 30^\circ) = \cot 30^\circ = \sqrt{3}$
$= -\tan(4\times 90^\circ - 60^\circ)$ (IV) $= +(+\tan 60^\circ) = \tan 60^\circ = \sqrt{3}$.

(3) $\sin\left(-\frac{11\pi}{3}\right)$

(4) $\tan\left(\frac{19\pi}{3}\right)$

(5) $\cot(-1470^\circ)$

Find value of following:

(1) $\cos(240^\circ)$

(2) $\tan(-300^\circ)$

(3) $\sin\left(-\frac{11\pi}{3}\right)$

$-\sin\left(\frac{11\pi}{3}\right) = -\sin\left(4\pi - \frac{\pi}{3}\right) = +\left(+\sin\frac{\pi}{3}\right) = +\frac{\sqrt{3}}{2}$ ✓

IV, 60°

$8\frac{\pi}{2} - \frac{\pi}{3}$

$-\sin\left(3\pi + \frac{2\pi}{3}\right) = -\left(-\sin\frac{2\pi}{3}\right) = \sin\left(\frac{2\pi}{3}\right)$

IV, cos

$= \sin\left(\pi - \frac{\pi}{3}\right) = +\sin\frac{\pi}{3}$

$= \frac{\sqrt{3}}{2}$ ✓

(4) $\tan\left(\frac{19\pi}{3}\right) = \tan\left(6\pi + \frac{\pi}{3}\right) = +\tan\frac{\pi}{3} = +\sqrt{3}$.

I

(5) $\cot(-1470^\circ)$

Find value of following:

(1) $\cos(240^\circ)$

(2) $\tan(-300^\circ)$

(3) $\sin\left(-\frac{11\pi}{3}\right)$

(4) $\tan\left(\frac{19\pi}{3}\right)$

(5) $\cot(-1470^\circ)$

$-\cot(1470^\circ) = -\cot(16\times 90^\circ + 30^\circ)$ (I)

$= -(+\cot 30^\circ) = -\sqrt{3}$



VALUE OF TRIGONOMETRIC FUNCTIONS

Radian	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	$\frac{5\pi}{6}$	π	$\frac{7\pi}{6}$	$\frac{5\pi}{4}$	$\frac{4\pi}{3}$	$\frac{3\pi}{2}$	$\frac{5\pi}{3}$	$\frac{7\pi}{4}$	$\frac{11\pi}{6}$	2π
Degree	0	30°	45°	60°	90°	120°	135°	150°	180°	210°	225°	240°	270°	300°	315°	330°	360°
sin																	
cos																	
tan																	
cot																	

!! FIBY !! for more practice.

Question

$\sin 420^\circ \cos 390^\circ + \cos(-300^\circ)\sin(-330^\circ)$ equals

A 0

B 1

C −1

D 2

$\frac{\sqrt{3}}{2} \times \frac{\sqrt{3}}{2} + \frac{1}{2} \cdot \frac{1}{2} = \frac{3}{4} + \frac{1}{4} = 1$

$420^\circ = 4 \times 90^\circ + 60^\circ$

$\sin 420^\circ = \sin(\underbrace{4 \times 90^\circ + 60^\circ}_{\text{I}})$

$= +\sin 60^\circ$

$= \frac{\sqrt{3}}{2}$

$\sin(-330^\circ) = -\sin(330^\circ)_{\text{IV}}$

$= -\sin(4 \times 90^\circ - 30^\circ)$

$= +(+\sin 30^\circ)$

$= \frac{1}{2}$

$\cos 390^\circ = \cos(\underbrace{4 \times 90^\circ + 30^\circ}_{\text{I}})$

$= +\cos 30^\circ$

$= \frac{\sqrt{3}}{2}$

$\cos(-300^\circ) = \cos(300^\circ)_{\text{IV}}$

$= \cos(4 \times 90^\circ - 60^\circ)$

$= +\cos 60^\circ$

$= +\frac{1}{2}$

Blue बादल

"ajeeb se angle"

(ex :- $\frac{\pi}{7}, \frac{3\pi}{11}, \frac{7\pi}{13}$ .....)

- First and last ko add.
- $\pi\ (180^\circ)$ में से add or subtract करो
- $\frac{\pi}{2}\ (90^\circ)$ में से add or subtract करो

} Reduc^{n} formula.

Module Question

$$\cos\frac{\pi}{7}+\cos\frac{2\pi}{7}+\cos\frac{3\pi}{7}+\cos\frac{4\pi}{7}+\cos\frac{5\pi}{7}+\cos\frac{6\pi}{7}=$$

A 1/2

B −1/2

C 0

D 1

$$\frac{\pi}{7}+\frac{6\pi}{7}=\frac{7\pi}{7}=\pi$$

$$\cos\frac{\pi}{7}=\cos\left(\pi-\underbrace{\frac{6\pi}{7}}_{\theta}\right)$$

$$=-\cos\frac{6\pi}{7}$$

$$\cos\frac{\pi}{7}+\cos\frac{6\pi}{7}=0$$

$$\cos\frac{5\pi}{7}=\cos\left(\pi-\frac{2\pi}{7}\right)$$

$$=-\cos\frac{2\pi}{7}$$

If

$A+B=\pi$

$\cos A=-\cos B$

$\cos(\pi-B)$

$\cos A+\cos B=0$

Question

$\tan\frac{\pi}{11}+\tan\frac{2\pi}{11}+\tan\frac{4\pi}{11}+\tan\frac{7\pi}{11}+\tan\frac{9\pi}{11}+\tan\frac{10\pi}{11}$ equals

A 0

B 1

C −1

D 2

$$\tan\left(\frac{10\pi}{11}\right)=\tan\left(\pi-\overset{\theta}{\left(\frac{\pi}{11}\right)}\right)=-\tan\frac{\pi}{11}$$

$$\#\quad \frac{\pi}{11}+\frac{10\pi}{11}=\frac{11\pi}{11}=\pi$$

Acidic Question

Evaluate :

$$\frac{\sin\frac{11\pi}{17}\cos\frac{10\pi}{13}\tan\frac{\pi}{7}}{\cos\frac{3\pi}{13}\sin\frac{6\pi}{17}\tan\frac{6\pi}{7}} = +1$$

$\frac{11\pi}{17} + \frac{6\pi}{17} = \pi$.

$$\sin\left(\frac{11\pi}{17}\right) = \sin\left(\overset{\pi-\theta}{\pi - \frac{6\pi}{17}}\right)$$

$$= +\sin\left(\frac{6\pi}{17}\right)$$

$\frac{10\pi}{13} + \frac{3\pi}{13} = \pi$.

$$\cos\frac{10\pi}{13} = \cos\left(\overset{\pi-\theta}{\pi - \frac{10\pi}{13}}\right)$$

$$= -\cos\frac{10\pi}{13}$$

$$\tan\left(\frac{6\pi}{7}\right) = \tan\left(\overset{\pi-\theta}{\pi - \frac{\pi}{7}}\right)$$

$$= -\tan\frac{\pi}{7}$$

Acidic Question

HW.

Evaluate :

(i) $$\frac{\sin\frac{\pi}{5}\cos\frac{7\pi}{9}\tan\frac{6\pi}{11}}{\cos\frac{2\pi}{9}\sin\frac{4\pi}{5}\tan\frac{5\pi}{11}}$$

(ii) $$\frac{\sin\frac{\pi}{7}\cos\frac{5\pi}{11}\tan\frac{3\pi}{7}}{\cos\frac{5\pi}{14}\sin\frac{\pi}{22}\tan\frac{4\pi}{7}}$$

COMPOUND ANGLE FORMULA's

For Sine & Cosine:

$$\sin(A+B) = \sin A \cdot \cos B + \cos A \sin B$$

$$\sin(A-B) = \sin A \cos B - \cos A \sin B$$

$$\cos(A+B) = \cos A \cdot \cos B - \sin A \sin B$$

$$\cos(A-B) = \cos A \cos B + \sin A \cdot \sin B$$

RAR

COMPOUND ANGLE FORMULA's

For Tan & Cot:

$$\tan(A \pm B) = \frac{\sin(A+B)}{\cos(A+B)} = \frac{\frac{\sin A\cos B}{\cos A\cos B} + \frac{\cos A\sin B}{\cos A\cos B}}{\frac{\cos A\cos B}{\cos A\cos B} - \frac{\sin A}{\cos A}\frac{\sin B}{\cos B}} = \frac{\tan A \pm \tan B}{1 \mp \tan A\cdot\tan B}$$

$$\cot(A-B) = \frac{\cos(A-B)}{\sin(A-B)} = \frac{\frac{\cos A\cos B}{\sin A\sin B} + \frac{\sin A\sin B}{\sin A\sin B}}{\frac{\sin A}{\sin A}\frac{\cos B}{\sin B} - \frac{\cos A}{\sin A}\frac{\sin B}{\sin B}} = \frac{\cot A\cdot\cot B + 1}{\cot B - \cot A}$$

Sachin Sir Special Summary (S^4):

RAR

$$\sin(A+B) = \sin A\cos B + \cos A\sin B$$

$$\sin(A-B) = \sin A\cos B - \cos A\sin B$$

$$\cos(A+B) = \cos A\cos B - \sin A\sin B$$

$$\cos(A-B) = \cos A\cos B + \sin A\sin B$$

(1) $\tan(A+B) = \left(\frac{\tan(A)+\tan(B)}{1-\tan A\tan B}\right)$

(2) $\tan(A-B) = \left(\frac{\tan A-\tan B}{1+\tan A\tan B}\right)$

(3) $\cot(A+B) = \left(\frac{\cot A\cot B-1}{\cot A+\cot B}\right)$

(4) $\cot(A-B) = \left(\frac{\cot A\cot B+1}{\cot B-\cot A}\right)$

$$\tan\left(\frac{\pi}{4}+\theta\right) = \left(\frac{1+\tan\theta}{1-\tan\theta}\right)$$

$$\tan\left(\frac{\pi}{4}-\theta\right) = \left(\frac{1-\tan\theta}{1+\tan\theta}\right)$$

$$\tan\left(\underbrace{\frac{\pi}{4}}_{A} + \underbrace{\theta}_{B}\right) = \frac{\tan\left(\pi/4\right) + \tan\theta}{1 - \tan\pi/4 \cdot \tan\theta} = \frac{1 + \tan\theta}{1 - \tan\theta}$$

($\frac{\pi}{4} = 45°$)

RAPID FIRE

Prove that : $\tan A \cot\frac{A}{2} - 1 = \sec A$

$$LHS = \frac{\sin A}{\cos A} \cdot \frac{\cos\frac{A}{2}}{\sin\frac{A}{2}} - 1 = \frac{\sin A \cdot \cos\frac{A}{2} - \cos A \cdot \sin\frac{A}{2}}{\cos A \cdot \sin\frac{A}{2}} = \sin\left(A - \frac{A}{2}\right)$$

$$= \sec A = RHS$$

Hope.

RAPID FIRE

Prove that $\cos(45° - A)\cos(45° - B) - \sin(45° - A)\sin(45° - B) = \sin(A + B)$

($45° - A$ = X, $45° - B$ = Y, $45° - A$ = X, $45° - B$ = Y)

$$LHS = \cos X \cdot \cos Y - \sin X \cdot \sin Y$$

$$= \cos(X + Y)$$

$$= \cos(45° - A + 45° - B)$$

$$= \cos(90° - \underbrace{(A+B)}_{\theta}) = \sin\theta$$

$$= \sin(A+B) = RHS.$$

Question

Prove that: $\frac{\sin(A-B)}{\cos A\cos B}+\frac{\sin(B-C)}{\cos B\cos C}+\frac{\sin(C-A)}{\cos C\cos A}=0$



Question

Value of $\left(\frac{\tan 20^\circ}{\operatorname{cosec} 70^\circ}\right)^2 + \left(\frac{\cot 20^\circ}{\sec 70^\circ}\right)^2 + 2\tan 15^\circ \tan 37^\circ \tan 53^\circ \tan 60^\circ \tan 75^\circ$ is



DO IT BY YOURSELF (DIBY)

DIBY-06

Prove that: $\sin^8 A - \cos^8 A = (\sin^2 A - \cos^2 A)(1 - 2\sin^2 A \cos^2 A)$

$a^8 - b^8 = (a^4)^2 - (b^4)^2$

$= (a^4 - b^4)(a^4 + b^4) = (a^2 - b^2)(a^2 + b^2)(a^4 + b^4)$

$(\sin^2 A - \cos^2 A)(\sin^2 A + \cos^2 A)(\sin^4 A + \cos^4 A)$ → $\sin^2 A + \cos^2 A = 1$

DIBY-07

$(1 + \tan\alpha \tan\beta)^2 + (\tan\alpha - \tan\beta)^2 =$

(A) $\tan^2\alpha \tan^2\beta$

(B) $\sec^2\alpha \sec^2\beta$

(C) $\tan^2\alpha \cot^2\beta$

(D) $\sec^2\alpha \cos^2\beta$

[Ans. : (B)]

$1 + \underbrace{\tan^2\alpha}_{x}\,\underbrace{\tan^2\beta}_{y} + 2\tan\alpha\tan\beta + \tan^2\alpha + \tan^2\beta - 2\tan\alpha\tan\beta.$

$1 + xy + x + y = (1+y) + x(1+y) = (1+y)(1+x) = (1+\tan^2\beta)(1+\tan^2\alpha)$

$\sec^2\beta \quad \sec^2\alpha.$

DIBY-08

Prove that : $\frac{\sin A}{1+\cos A}+\frac{1+\cos A}{\sin A}=2\operatorname{cosec} A$

DIBY-09

Prove that : $\sqrt{\frac{1-\sin A}{1+\sin A}}=\sec A-\tan A$

DIBY-10

Prove that : $\frac{\operatorname{cosec} A}{\operatorname{cosec} A-1}+\frac{\operatorname{cosec} A}{\operatorname{cosec} A+1}=2\sec^2 A$

DIBY-11

Prove that : $\frac{\operatorname{cosec} A}{\cot A+\tan A}=\cos A$

DIBY-12

Find value of following

(i) $\sin^2\frac{\pi}{6}+\cos^2\frac{\pi}{3}-\tan^2\frac{\pi}{4}=$

(ii) $2\sin^2\frac{\pi}{6}+\operatorname{cosec}^2\frac{7\pi}{6}\cos^2\frac{\pi}{3}=$

(iii) $\cot^2\frac{\pi}{6}+\operatorname{cosec}\frac{5\pi}{6}+3\tan^2\frac{\pi}{6}=$

(iv) $2\sin^2\frac{3\pi}{4}+2\cos^2\frac{\pi}{4}+2\sec^2\frac{\pi}{3}=$

[Ans. : (i) $\frac{-1}{2}$, (ii) $\frac{3}{2}$, (iii) 6, (iv) 10]

DIBY-13

$\frac{\tan(90°-\theta)\sec(180°-\theta)\sin(-\theta)}{\sin(180°+\theta)\cos(360°+\theta)\operatorname{cosec}(90°-\theta)}$ is equals

(A) $\operatorname{cosec}\theta$ (B) 1 (C) –1 (D) $-\operatorname{cosec}\theta$

[Ans. : (D)]

DIBY-14

$\sec(270^\circ - A)\sec(90^\circ - A) - \tan(270^\circ - A)\tan(90^\circ + A)$

[Ans. : –1]

DIBY-15

$\cot A + \tan(180^\circ + A) + \tan(90^\circ + A) + \tan(360^\circ - A)$

[Ans. : 0]

DIBY-16

$$\frac{\tan(180^\circ-\theta)\cos(180^\circ-\theta)\tan(90^\circ-\theta)}{\sin(90^\circ+\theta)\cot(90^\circ-\theta)\tan(90^\circ+\theta)}$$

[Ans. : –1]

DIBY-17

If $\tan A = 2$ and $A \in \left(\pi, \frac{3\pi}{2}\right)$ then the expression $\frac{\cos A}{\sin^3 A + \cos^3 A}$ is equal to

[Ans. : 5/9]

DIBY-18

$\sin^2\frac{\pi}{18}+\sin^2\frac{\pi}{9}+\sin^2\frac{7\pi}{18}+\sin^2\frac{4\pi}{9}$ equal

(A) 0 (B) 1 (C) –1 (D) 2

[Ans. : (D)]

DIBY-19

Prove that

1. $\tan 720^\circ-\cos 270^\circ-\sin 150^\circ\cos 120^\circ=\frac{1}{4}$

2. $\sin 780^\circ\sin 480^\circ+\cos 120^\circ\sin 150^\circ=\frac{1}{2}$

DIBY-20

Prove that $\sec\left(\frac{3\pi}{2}-\theta\right)\sec\left(\theta-\frac{5\pi}{2}\right)+\tan\left(\frac{5\pi}{2}+\theta\right)\tan\left(\theta-\frac{3\pi}{2}\right)=-1.$



Homework

Re-attempt all Class Examples

DPP & DIBY

Module: Only 2 Exercise

Topicwise & Bridge(Setu)- 4 sittings each of 1 HR

It is never about end result

It's always about Journey

Thank You

#futureIITians