



**UNIVERSIDAD NACIONAL AUTÓNOMA DE MÉXICO**  
**POSGRADO EN CIENCIAS FÍSICAS**

**OPTICAL RESPONSE OF PARTIALLY EMBEDDED  
NANOSPHERES**

**TESIS**

**QUE PARA OPTAR POR EL GRADO DE:  
MAESTRO EN CIENCIAS (FÍSICA)**

**PRESENTA:**

**JONATHAN ALEXIS URRUTIA ANGUIANO**

**TUTOR:**

**DR. ALEJANDRO REYES CORONADO**  
FACULTAD DE CIENCIAS, UNAM

**MIEMBROS DEL COMITÉ TUTOR**

**DRA. CITLALI SÁNCHEZ-AKÉ**  
INSTITUTO DE CIENCIAS APLICADAS Y TECNOLOGÍA, UNAM

**DR. GIUSEPPE PIRRUCCIO**  
INSTITUTO DE FÍSICA, UNAM

**CIUDAD DE MÉXICO, MÉXICO, 2021**







# Abstract/Resumen

---

Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

# Contents

<b>Abstract</b>	<b>iii</b>
<b>Introduction</b>	<b>1</b>
<b>1 Optical properties of single plasmonic nanoparticles</b>	<b>3</b>
1.1 The Optical Theorem: Amplitude Matrix and Cross Sections . . . . .	3
1.2 Scattering of spherical particles . . . . .	10
<b>2 Results and discussion</b>	<b>13</b>
2.1 Finite Element Method and Analytical Solutions . . . . .	13
2.2 Incrustation Degree of a Spherical Particle . . . . .	13
<b>3 Conclusions</b>	<b>15</b>
3.1 Future Work: Application on Metasurfaces . . . . .	15
<b>A The Finite Element Method</b>	<b>17</b>
<b>List of Figures</b>	<b>20</b>
<b>Index</b>	<b>21</b>

# Introduction

---

It is recommended to fill in this part of the document with the following information:

- Your field: Context about the field your are working  
**Plasmonics -> Metamaterials -> Biosensing**
- Motivation: Background about your thesis work and why did you choose this project and why is it important.  
**Fabrication -> Partially embedded NPs -> No analytical (approximated) method physically introduces the incrustation degree. There are numerical solutions and Effective Medium Theories approaching the problem but the later only as a fitting method.**
- Objectives: What question are you answering with your work.  
**Can optical non invasive tests (IR-Vis) retrieve the average incrustation degree for monolayers of small spherical particles?**
- Methodology: What are your secondary goals so you achieve your objective. Also, how are you answering your question: which method or model.  
**Bruggeman homogenization theories on bidimensional systems?  
Is the dipolar approximation is enough or do we need more multipolar terms?  
Do we need the depolarization factors?**
- Structure: How is this thesis divides and what is the content of each chapter.





# Optical properties of single plasmonic nanoparticles

The problem studied in this thesis corresponds to the theoretical analysis of the Localized Surface Plasmon Resonances (LSPR) excited on plasmonic spherical nanoparticles (NPs) when these are under realistic experimental conditions, such as those present on plasmonic biosensors, where the NPs are partially embedded into a substrate [1]. The theoretical analysis consists on the numerical calculation of the absorption, scattering and extinction cross sections of a partially embedded metal NP employing the Finite Element Method (FEM), nevertheless, to verify the validity of the obtained results, the problem of the absorption and scattering of light by an isolated particle must be addressed. In this chapter, we revisit the general solution of the light absorption and scattering by both an arbitrary particle and by a spherical particle, given by the Mie Theory [2].

## 1.1 The Optical Theorem: Amplitude Matrix and Cross Sections

Let  $\mathbf{E}^i = \mathbf{E}_0^i \exp(i\mathbf{k}^i \cdot \mathbf{r})$  be the electric field of an incident monochromatic plane wave with constant amplitude  $\mathbf{E}_0^i$  traveling through a non-dispersive medium with refractive index  $n_m$ , denominated matrix, in the direction  $\mathbf{k}^i = k_m \hat{\mathbf{k}}^i$ , with  $k_m$  the wave number of the plane wave into the matrix, and let  $\mathbf{E}^{\text{sca}}$  the electric far field of the scattered field due to a particle with arbitrary shape embedded into the matrix. In general, the scattered electric field propagates in all directions but for a given point  $\mathbf{r} = r\hat{\mathbf{e}}_r$  the traveling direction is defined by the vector  $\mathbf{k}^{\text{sca}} = k_m \hat{\mathbf{k}}^{\text{sca}} = k_m \hat{\mathbf{e}}_r$ . Due to the linearity of the Maxwell's equations, the incident and scattered electric fields are related in the far field by a linear relation [3], that is,

$$\mathbf{E}^{\text{sca}} = \frac{\exp(i\mathbf{k}^{\text{sca}} \cdot \mathbf{r})}{r} \mathbb{F}(\hat{\mathbf{k}}^{\text{sca}}, \hat{\mathbf{k}}^i) \mathbf{E}^i, \quad (1.1)$$

where  $\mathbb{F}(\hat{\mathbf{k}}^{\text{sca}}, \hat{\mathbf{k}}^i)$  is the scattering amplitude matrix from direction  $\hat{\mathbf{k}}^i$  into  $\hat{\mathbf{k}}^{\text{sca}}$ . Since only the far field is considered, both the incident and the scattered electric field can be decomposed into two linearly independent components perpendicular to  $\mathbf{k}^i$  and  $\mathbf{k}^{\text{sca}}$ , respectively, each forming a right-hand orthonormal system. If the particle acting as a scatterer has a symmetric shape, it is convenient to define the orthonormal systems relative to the scattering plane, which is the

plane containing  $\mathbf{k}^i$  and  $\mathbf{k}^{sca}$ , since the elements of  $\mathbb{F}(\hat{\mathbf{k}}^{sca}, \hat{\mathbf{k}}^i)$  simplify when represented in these bases [3]. By defining the directions perpendicular ( $\perp$ ) and parallel ( $\parallel$ ) to the scattering plane, the incident and scattered electric fields can be written as


$$\mathbf{E}^i = (E_{\parallel}^i \hat{\mathbf{e}}_{\parallel}^i + E_{\perp}^i \hat{\mathbf{e}}_{\perp}^i) \exp(i\mathbf{k}^i \cdot \mathbf{r}), \quad (1.2)$$

$$\mathbf{E}^{sca} = (E_{\parallel}^{sca} \hat{\mathbf{e}}_{\parallel}^{sca} + E_{\perp}^{sca} \hat{\mathbf{e}}_{\perp}^{sca}) \frac{\exp(i\mathbf{k}^{sca} \cdot \mathbf{r})}{r}, \quad (1.3)$$

where the harmonic time dependence  $\exp(-i\omega t)$  has been suppressed, and where it has been assumed that the scattered field is described by a spherical wave; the superindex “i” (“sca”) denotes the orthonormal system defined by the incident plane wave (scattered fields). Since  $\{\hat{\mathbf{e}}_{\perp}^i, \hat{\mathbf{e}}_{\parallel}^i, \hat{\mathbf{k}}^i\}$  and  $\{\hat{\mathbf{e}}_{\perp}^{sca}, \hat{\mathbf{e}}_{\parallel}^{sca}, \hat{\mathbf{k}}^{sca}\}$  are right-hand orthonormal systems, they are related by

$$\hat{\mathbf{e}}_{\perp}^i = \hat{\mathbf{e}}_{\perp}^{sca} = \hat{\mathbf{k}}^{sca} \times \hat{\mathbf{k}}^i, \quad \hat{\mathbf{e}}_{\parallel}^i = \hat{\mathbf{k}}^i \times \hat{\mathbf{e}}_{\perp}^i, \quad \hat{\mathbf{e}}_{\parallel}^{sca} = \hat{\mathbf{k}}^{sca} \times \hat{\mathbf{e}}_{\perp}^{sca}. \quad (1.4)$$

As the Eqs. (1.4) suggest, the unit vector bases of the orthonormal systems relative to the scattering plane depend on the scattering direction. For example, if the incident plane wave travels along the  $z$  axis, then  $\hat{\mathbf{k}}^i = \hat{\mathbf{e}}_z$  and  $\hat{\mathbf{k}}^{sca} = \hat{\mathbf{e}}_r$ . Thus, according to Eqs. (1.4), the unit vector bases of the systems relative to the scattering plane are  $\hat{\mathbf{e}}_{\parallel}^i = \cos \varphi \hat{\mathbf{e}}_x + \sin \varphi \hat{\mathbf{e}}_y$ ,  $\hat{\mathbf{e}}_{\parallel}^{sca} = \hat{\mathbf{e}}_{\theta}$  and  $\hat{\mathbf{e}}_{\perp}^i = \hat{\mathbf{e}}_{\perp}^{sca} = -\hat{\mathbf{e}}_{\varphi}$ , with  $\theta$  the polar angle and  $\varphi$  azimuthal angle. In Fig. 1.1 the unit vector systems (purple) based on the scattering plane (green) defined by the vectors  $\hat{\mathbf{k}}^i = \hat{\mathbf{e}}_z$  and  $\hat{\mathbf{k}}^{sca} = \hat{\mathbf{e}}_r$  are shown, along with the Cartesian (blue) and spherical (black) unit vector bases.



**Fig. 1.1:** The scattering plane (green) is defined by the vectors  $\hat{\mathbf{k}}^i$ , direction of the incident plane wave (red), and  $\hat{\mathbf{k}}^{sca}$ , direction of the scattered field in a given point  $\vec{r}$ . If the direction of the incident plane wave is chose to be  $\hat{\mathbf{e}}_z$ , the parallel and perpendicular components of the incident field relative to the scattering plane are  $\hat{\mathbf{e}}_{\parallel}^i = \cos \varphi \hat{\mathbf{e}}_x + \sin \varphi \hat{\mathbf{e}}_y$  and  $\hat{\mathbf{e}}_{\perp}^i = -\hat{\mathbf{e}}_{\varphi}$ , while the components of the scattering field relative to the scattering plane are  $\hat{\mathbf{e}}_{\parallel}^{sca} = \hat{\mathbf{e}}_{\theta}$ ,  $\hat{\mathbf{e}}_{\perp}^{sca} = -\hat{\mathbf{e}}_{\varphi}$ . The cartesian unit vector basis is shown in blue, the spherical unit vector basis in black, while the basis of the orthonormal systems relative to the scattering plane are shown in purple.

After a incident plane wave interacts with a particle with a possible complex refractive index  $n_p(\omega)$ , the total electric field outside the particle is given by the sum of the incident and the scattered fields. Therefore, the time averaged Poynting vector  $\langle \mathbf{S} \rangle_t$ , denoting the power flow per unit area, of the total field is given by

$$\langle \mathbf{S} \rangle_t = \underbrace{\frac{1}{2} \operatorname{Re} (\mathbf{E}^i \times \mathbf{H}^{i*})}_{\langle \mathbf{S}^i \rangle_t} + \underbrace{\frac{1}{2} \operatorname{Re} (\mathbf{E}^{\text{sca}} \times \mathbf{H}^{\text{sca}*})}_{\langle \mathbf{S}^{\text{sca}} \rangle_t} + \underbrace{\frac{1}{2} \operatorname{Re} (\mathbf{E}^i \times \mathbf{H}^{\text{sca}*} + \mathbf{E}^{\text{sca}} \times \mathbf{H}^{i*})}_{\langle \mathbf{S}^{\text{ext}} \rangle_t}, \quad (1.5)$$

where  $(*)$  denotes the complex conjugate operation and where the total Poynting vector is separated into the contribution from the incident field  $\langle \mathbf{S}^i \rangle_t$ , from the scattered field  $\langle \mathbf{S}^{\text{sca}} \rangle_t$  and from their cross product denoted by  $\langle \mathbf{S}^{\text{ext}} \rangle_t$ . By means of the Faraday-Lenz Law and Eq. (1.1), the contribution to the Poynting vector from the incident and the scattered fields can be rewritten as

$$\langle \mathbf{S}^i \rangle_t = \frac{\|\mathbf{E}_0^i\|^2}{2Z_m} \hat{\mathbf{k}}^i, \quad \text{and} \quad \langle \mathbf{S}^{\text{sca}} \rangle_t = \frac{\|\mathbf{E}^{\text{sca}}\|^2}{2Z_m} \hat{\mathbf{k}}^{\text{sca}} = \frac{\|\mathbb{F}(\hat{\mathbf{k}}^{\text{sca}}, \hat{\mathbf{k}}^i) \mathbf{E}^i\|^2}{2Z_m r^2} \hat{\mathbf{k}}^{\text{sca}}, \quad (1.6)$$

with  $Z_m = \sqrt{\mu_m/\varepsilon_m}$ , the impedance of the non-dispersive matrix, while the crossed contribution is given by

$$\begin{aligned} \langle \mathbf{S}^{\text{ext}} \rangle_t = \operatorname{Re} \left\{ \frac{\exp[-i(\mathbf{k}^{\text{sca}} - \mathbf{k}^i) \cdot \mathbf{r}]}{2Z_m r^2} \left[ \hat{\mathbf{k}}^{\text{sca}} (\mathbf{E}_0^i \cdot \mathbb{F}^* \mathbf{E}^{i*}) - \mathbb{F}^* \mathbf{E}^{i*} (\mathbf{E}_0^i \cdot \hat{\mathbf{k}}^{\text{sca}}) \right] \right. \\ \left. + \frac{\exp[i(\mathbf{k}^{\text{sca}} - \mathbf{k}^i) \cdot \mathbf{r}]}{2Z_m r^2} \left[ \hat{\mathbf{k}}^i (\mathbb{F} \mathbf{E}^i \cdot \mathbf{E}_0^{i*}) - \mathbf{E}_0^{i*} (\mathbb{F} \mathbf{E}^i \cdot \hat{\mathbf{k}}^i) \right] \right\}, \end{aligned} \quad (1.7)$$

where the scattering amplitude matrix is evaluated as  $\mathbb{F}(\hat{\mathbf{k}}^{\text{sca}}, \hat{\mathbf{k}}^i)$ .

The power scattered by the particle can be calculated by integrating  $\langle \mathbf{S}^{\text{sca}} \rangle_t$  in a closed surface surrounding the particle; if the scattered power is normalized by the irradiance of the incident field  $\|\langle \mathbf{S}^i \rangle_t\|$ , it is obtained a quantity with units of area known as the scattering cross section  $C_{\text{sca}}$ , given by

#### Scattering Cross Section

$$C_{\text{sca}} = \frac{2Z_m}{\|\mathbf{E}_0^i\|^2} \oint \langle \mathbf{S}^{\text{sca}} \rangle_t \cdot d\mathbf{a} = \oint \frac{\|\mathbb{F}(\hat{\mathbf{k}}^{\text{sca}}, \hat{\mathbf{k}}^i) \mathbf{E}^i\|^2}{\|\mathbf{E}_0^i\|^2} d\Omega, \quad (1.8)$$

where  $d\Omega$  is the solid angle differential. In a similar manner, an absorption cross section  $C_{\text{abs}}$  can be defined as well. On the one side, the absorption cross section is given by the integral on a closed surface of  $-\langle \mathbf{S} \rangle_t$  [Eq. (1.5)] divided by the irradiance of the incident field, where the minus sign is chosen so that  $C_{\text{abs}} > 0$  if the particle absorbs energy [2]. On the other side, if an Ohmic material for the particle with a conductivity  $\sigma(\omega) = i\omega n_p^2(\omega)$  [4] is assumed, through Joule's Heating Law [3] the absorption cross section can be computed as

## Ohmic Particle - Absorption Cross Section

$$C_{\text{abs}} = \frac{1}{2} \int \frac{\text{Re}(\mathbf{j} \cdot \mathbf{E}^{\text{int}*})}{\|\mathbf{E}_0^{\text{i}}\|^2 / 2Z_{\text{m}}} dV = \int \omega Z_{\text{m}} \text{Im}(n_p^2) \frac{\|\mathbf{E}^{\text{int}}\|^2}{\|\mathbf{E}_0^{\text{i}}\|^2} dV, \quad (1.9)$$

where integration is performed inside the particle, and  $\mathbf{j}$  and  $\mathbf{E}^{\text{int}}$ , are the volumetric electric current density and the total electric field in this region. Both the scattering and the absorption cross sections are quantities related to the optical signature of a particle [5], and their relation can be made explicit by performing the surface integral representation of  $C_{\text{abs}}$  and defining  $C_{\text{ext}}$ , that is,

$$\begin{aligned} C_{\text{abs}} &= -\frac{2Z_{\text{m}}}{\|\mathbf{E}_0^{\text{i}}\|^2} \int \left( \langle \mathbf{S}^{\text{i}} \rangle_t + \langle \mathbf{S}^{\text{sca}} \rangle_t + \langle \mathbf{S}^{\text{ext}} \rangle_t \right) \cdot d\mathbf{a} \\ &= -C_{\text{sca}} - \frac{2Z_{\text{m}}}{\|\mathbf{E}_0^{\text{i}}\|^2} \int \langle \mathbf{S}^{\text{ext}} \rangle_t \cdot \hat{\mathbf{e}}_r d\Omega \\ &= -C_{\text{sca}} + C_{\text{ext}}, \end{aligned} \quad (1.10)$$

where the contribution of  $\langle \mathbf{S}^{\text{i}} \rangle_t$  to the integral is zero since a non-dispersive matrix was assumed. From Eq.(1.10) it can be seen that  $C_{\text{ext}}$  takes into account both mechanisms for energy losses (scattering and absorption), thus it is called the extinction cross section. To solve the integral in Eq. (1.10) let us define  $\theta$  as the angle between  $\hat{\mathbf{k}}^{\text{sca}}$  and  $\hat{\mathbf{k}}^{\text{i}}$  as the polar angle and  $\varphi$  as the azimuthal angle as shown in Fig 1.1. With this election of coordinates, the extinction cross section can be computed as

$$\begin{aligned} C_{\text{ext}} &= -\text{Re} \left\{ \frac{\exp(-ik_m r)}{\|\mathbf{E}_0^{\text{i}}\|^2} \oint \exp(ik_m r \cos \theta) (1) (\mathbf{E}^{\text{i}} \cdot \mathbb{F}^* \mathbf{E}^{\text{i}*}) d\Omega \right. \\ &\quad + \frac{\exp(ik_m r)}{\|\mathbf{E}_0^{\text{i}}\|^2} \oint \exp(-ik_m r \cos \theta) \cos \theta (\mathbf{E}^{\text{i}*} \cdot \mathbb{F} \mathbf{E}^{\text{i}}) d\Omega \\ &\quad \left. + \frac{\exp(ik_m r)}{\|\mathbf{E}_0^{\text{i}}\|^2} \oint \exp(-ik_m r \cos \theta) \sin \theta (E_{0,x}^{\text{i}} \cos \varphi + E_{0,y}^{\text{i}} \sin \varphi) (\mathbb{F} \mathbf{E}^{\text{i}} \cdot \mathbf{k}^{\text{i}}) d\Omega \right\} \end{aligned} \quad (1.11)$$

where the relations  $\hat{\mathbf{k}}^{\text{sca}} \cdot \hat{\mathbf{e}}_r = 1$ ,  $\hat{\mathbf{k}}^{\text{i}} \cdot \hat{\mathbf{e}}_r = \cos \theta$  and  $\mathbf{E}^{\text{sca}} \cdot \hat{\mathbf{e}}_r = 0$  were employed. The integrals in Eq. (1.11) can be solved by a two fold integration by parts on the polar angle  $\theta$  and by depreciating the terms proportional to  $r^{-2}$ . This process leads to a zero contribution from the integrand proportional to  $\sin \theta$  of Eq. (1.11), and after arranging the other terms in their real and imaginary parts, it follows that  $C_{\text{ext}}$  depends only in the forward direction  $\hat{\mathbf{k}}^{\text{sca}} = \hat{\mathbf{k}}^{\text{i}}$  ( $\theta = 0$ ). This result is known as the Optical Theorem and whose mathematical expression is given by [3, 5, 6]

## Optical Theorem - Extinction Cross Section

$$C_{\text{ext}} = C_{\text{abs}} + C_{\text{sca}} = \frac{4\pi}{k_m \|\mathbf{E}_0^{\text{i}}\|^2} \text{Im} \left[ \mathbf{E}_0^{\text{i}} \cdot \mathbb{F}^* (\hat{\mathbf{k}}^{\text{i}}, \hat{\mathbf{k}}^{\text{i}}) \mathbf{E}^{\text{i}*} \right]. \quad (1.12)$$

From Eqs. (1.5) and (1.12) it can be seen that the extinction of light, the combined result of scattering and absorption as energy loss mechanisms, is also a manifestation of the interference between the incident and the scattered fields and that the over all effect of the light extinction can be fully understood by analyzing the amplitude of the scattering field in the forward direction. It is worth noting that Eq. (1.12) is an exact relation but its usefulness is bound to the correct evaluation of the scattering amplitude matrix  $\mathbb{F}$  [3]. Thus, in the following sections a scattering problem with spherical symmetry will be assumed, so that the exact solution to the scattering amplitude matrix can be developed; this solution is known as Mie Theory.




Fig. 1.2






Fig. 1.3



**Fig. 1.4:** The explanation of your figures. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.



**Fig. 1.5:** The explanation of your figures. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Etiam lobortis facilisis sem. Nullam nec mi et neque pharetra sollicitudin. Praesent imperdiet mi nec ante. Donec ullamcorper, felis non sodales commodo, lectus velit ultrices augue, a dignissim nibh lectus placerat pede. Vivamus nunc nunc, molestie ut, ultricies vel, semper in, velit. Ut porttitor. Praesent in sapien. Lorem ipsum dolor sit amet, consectetur adipiscing elit. Duis fringilla tristique neque. Sed interdum libero ut metus. Pellentesque placerat. Nam rutrum augue a leo. Morbi sed elit sit amet ante lobortis sollicitudin. Praesent blandit blandit mauris. Praesent lectus tellus, aliquet aliquam, luctus a, egestas a, turpis. Mauris lacinia lorem sit amet ipsum. Nunc quis urna dictum turpis accumsan semper.

## 1.2 Scattering of spherical particles

In the previous section, it was concluded that the extinction of light due to the interaction between a particle and a monochromatic plane wave can be determined through the amplitude of the scattered field in the forward direction. The past is stated in the Optical Theorem and it is an exact result but unaccuracies can arise when either the scattering amplitude matrix or extinction cross section is approximated<sup>1</sup>. A particular case in which the scattering amplitude matrix can be exactly calculated is when the scatterer has spherical symmetry. In order to address this special case it will be introduced a vectorial base with spherical symmetry, known as the Vectorial Spherical Harmonics (SVH).

The electric and magnetic field, denoted as  $\mathbf{E}$  and  $\mathbf{B}$ , respectively, are a solution to the homogeneous vectorial Helmholtz when an harmonic time dependence and a spacial domain with no external charge nor current densities is assumed, that is,

### Vectorial Helmholtz Equation

$$\nabla^2 \mathbf{E}(\mathbf{r}, \omega) + k_m^2 \mathbf{E}(\mathbf{r}, \omega) = \mathbf{0}, \quad (1.13a)$$

$$\nabla^2 \mathbf{B}(\mathbf{r}, \omega) + k_m^2 \mathbf{B}(\mathbf{r}, \omega) = \mathbf{0}. \quad (1.13b)$$

where the vectorial operator  $\nabla^2$  must be understood as  $\nabla^2 = \nabla(\nabla \cdot) - \nabla \times \nabla \times$ , and  $k_m$  is the wave number in the matrix. It is possible to build a basis set for the electric and magnetic fields as long as the elements of this basis are also solution to Eq. (1.21). One alternative is to employ the following set of vector functions

$$\mathbf{L} = \nabla \psi, \quad (1.14a)$$

$$\mathbf{M} = \nabla \times (\mathbf{r} \psi), \quad (1.14b)$$

$$\mathbf{N} = \frac{1}{k_m} \nabla \times \mathbf{M}, \quad (1.14c)$$

that are solution to the homogeneous vectorial Helmholtz equation as long as the scalar function  $\psi$  is solution to the scalar Helmholtz equation<sup>2</sup>

$$\nabla^2 \psi + k_m^2 \psi = 0. \quad (1.15)$$

The triad  $\{\mathbf{L}, \mathbf{M}, \mathbf{N}\}$  is a set of orthogonal vectors<sup>3</sup> that obey Helmholtz equation *i.e.*, they can be directly identify as electric or magnetic fields. The elements of the vector basis from Eq. (1.14) are known as the Vectorial Spherical Harmonics (VSH) as defined by Stratton [8], and Bohren and Huffman [2]. The scalar function  $\psi$  is known as the generating function of the VSH.

<sup>1</sup>See for example Section 2.4 from Ref. [3] on the Rayleigh Scattering and Section 21.7 from Ref. [7] on Thompson scattering.

<sup>2</sup>This result can be proven by considering the following: Let  $f$  be  $\mathbb{C}^3$  and  $\mathbf{F}$  a  $\mathbb{C}^2$ . Then, it is true that  $\nabla^2(\nabla f) = \nabla(\nabla^2 f)$ , and  $\nabla \times (\nabla^2 \mathbf{F}) = \nabla^2(\nabla \times \mathbf{F})$ .

<sup>3</sup>Employing the Einstein sum convention with  $\epsilon_{ijk}$  the Levi-Civita symbol, Eq. (1.14b) can be written as follows:  $M_i = [\nabla \times (\mathbf{r} \psi)]_i = \epsilon_{ijk} \partial_j (r_k \psi) = \psi \epsilon_{ijk} \partial_j (r_k) - \epsilon_{ijk} r_k \partial_j \psi = \psi [\nabla \times \mathbf{r}]_i - [\mathbf{r} \times \nabla \psi]_i = -[\mathbf{r} \times \nabla \psi]_i = [\mathbf{L} \times \mathbf{r}]_i$ , therefore  $\mathbf{M}$  is orthogonal to  $\mathbf{L}$  and  $\mathbf{r}$ . From Eq. (1.14c)  $\mathbf{M} \cdot \mathbf{N} = 0$ . **Falta probar  $\mathbf{L} \cdot \mathbf{N} = 0$ .**



If spherical coordinates are chosen, and it is assume that  $\psi(r, \theta, \varphi) = R(r)\Theta(\theta)\Phi(\varphi)$ , then Eq. (1.15) can be decouple into three ordinary differential equations:

$$\frac{1}{\Phi} \frac{\partial^2 \psi}{\partial \varphi^2} + m^2 \Phi = 0, \quad (1.16)$$

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{d\Theta}{d\theta} \right) + \left[ \ell(\ell + 1) - \frac{m^2}{\sin^2 \theta} \right] \Theta = 0, \quad (1.17)$$

$$\frac{d}{dr} \left( r^2 \frac{dR}{dr} \right) + [(k_m r)^2 - \ell(\ell + 1)] R = 0, \quad (1.18)$$

where  $\ell$  can takes natural values and zero, and  $|m| \leq \ell$  so  $\Phi$  and  $\Theta$  are univaluated and finite on a sphere. Eqs. (1.17) and (1.18) can be rewritten as

$$(1 - \mu^2) \frac{d^2 \Theta}{d\mu^2} - 2\mu \frac{d\Theta}{d\mu} + \left[ \ell(\ell + 1) - \frac{m^2}{1 - \mu^2} \right] \Theta = 0, \quad \text{with } \mu = \cos \theta, \quad (1.19)$$

$$\rho \frac{d}{d\rho} \left( \rho \frac{dZ}{d\rho} \right) + \left[ \rho^2 - \left( \ell + \frac{1}{2} \right)^2 \right] Z = 0, \quad \text{with } Z = R\sqrt{\rho} \text{ and } \rho = k_m r. \quad (1.20)$$

The solution to Eq. (1.19) are the Legendre Associated Functions  $P_\ell^m(\mu)$  and to Eq. (1.20) the solution is given by the Spherical Bessel Functions  $z_\ell = j_\ell, y_\ell$ . Following the convention from most literature on Mie Scattering [7], the solution to Eq. (1.16) will be decompose in an odd (*o*) and an even (*e*) solution, that is, as sine and cosine functions. After this procedure, it is determined that the generating function of the VSH is given by

**$\psi$ : Generating function of the vectorial spherical harmonics**

$$\psi_{e\ell m}(r, \theta, \varphi) = \cos(m\varphi) P_\ell^m(\cos \theta) z_\ell(k_m r), \quad (1.21a)$$

$$\psi_{o\ell m}(r, \theta, \varphi) = \sin(m\varphi) P_\ell^m(\cos \theta) z_\ell(k_m r). \quad (1.21b)$$



## Results and discussion

---

### **2.1 Finite Element Method and Analytical Solutions**

### **2.2 Incrustation Degree of a Spherical Particle**



### **3.1 Future Work: Application on Metasurfaces**



# The Finite Element Method

---





# Bibliography

---

- [1] R. S. Moirangthem, M. T. Yaseen, P.-K. Wei, J.-Y. Cheng, and Y.-C. Chang. Enhanced localized plasmonic detections using partially-embedded gold nanoparticles and ellipsometric measurements. *Biomedical Optics Express*, **3**(5):899, May 2012. ISSN: 2156-7085, 2156-7085. DOI: [10.1364/BOE.3.000899](https://doi.org/10.1364/BOE.3.000899). URL: <https://www.osapublishing.org/boe/abstract.cfm?uri=boe-3-5-899> (visited on 08/03/2021) (cited on page 3).
- [2] C. F. Bohren and D. R. Huffman. *Absorption and Scattering of Light by Small Particles*. English. Wiley Science Paperbak Series. John Wiley & Sons, 1st edition, 1983. ISBN: 0-471-029340-7 (cited on pages 3, 5, 10).
- [3] L. Tsang, J. A. Kong, and K.-H. Ding. *Scattering of Electromagnetic Waves: Theories and Applications*. en. John Wiley & Sons, Inc., New York, USA, July 2000. ISBN: 978-0-471-22428-0 978-0-471-38799-2. DOI: [10.1002/0471224286](https://doi.org/10.1002/0471224286). URL: <http://doi.wiley.com/10.1002/0471224286> (visited on 05/17/2020) (cited on pages 3–7, 10).
- [4] J. D. Jackson. *Classical electrodynamics*. Wiley, New York, 3rd ed edition, 1999. ISBN: 978-0-471-30932-1 (cited on page 5).
- [5] M. Pellarin, C. Bonnet, J. Lermé, F. Perrier, J. Laverdant, M.-A. Lebeault, S. Hermelin, M. Hillenkamp, M. Broyer, and E. Cottancin. Forward and Backward Extinction Measurements on a Single Supported Nanoparticle: Implications of the Generalized Optical Theorem. *The Journal of Physical Chemistry C*, **123**(24):15217–15229, June 2019. ISSN: 1932-7447, 1932-7455. DOI: [10.1021/acs.jpcc.9b03245](https://doi.org/10.1021/acs.jpcc.9b03245). URL: <https://pubs.acs.org/doi/10.1021/acs.jpcc.9b03245> (visited on 08/03/2021) (cited on page 6).
- [6] R. G. Newton. Optical theorem and beyond. en. *American Journal of Physics*, **44**(7):639–642, July 1976. ISSN: 0002-9505, 1943-2909. DOI: [10.1119/1.10324](https://doi.org/10.1119/1.10324). URL: <http://aapt.scitation.org/doi/10.1119/1.10324> (visited on 12/04/2020) (cited on page 6).
- [7] A. Zangwill. *Modern Electrodynamics*. Cambridge University Press, Cambridge, 2013. ISBN: 978-0-521-89697-9 (cited on pages 10, 11).
- [8] J. A. Stratton. *Electromagnetic theory*. McGraw-Hill, New York, 2012. ISBN: 978-1-4437-3054-9 (cited on page 10).

# List of Figures

1.1	Scattering plane unit vector systems . . . . .	4
1.2	. . . . .	8
1.3	. . . . .	8
1.4	Example of Figure title . . . . .	9
1.5	Example of Figure title . . . . .	9

# Index

---

- Extinction
  - Cross Section, [6](#)
- Faraday-Lenz
  - Law, [5](#)
- Finite Element Method, [3](#)
- Helmholtz
  - Equation, Scalar, [10](#)
  - Equation, Vectorial, [10](#)
- Irradiance, [5](#)
- Joule
  - Heating Law, [5](#)
- Law
  - Faraday-Lenz, [5](#)
- Maxwell
  - Equations, [5](#)
- Ohm
  - Law, [5](#)
- Optical
  - Theorem, [6](#)
- Plane
  - Scattering, [3](#)
- Plane Wave
  - Irradiance, [5](#)
- Plasmon
  - Localized Surface Plasmon Resonance (LSPR), [3](#)
- Poynting vector, [5](#)
- Scattering
  - Amplitude Matrix, [3](#)
  - Rayleigh, [10](#)
  - Thompson, [10](#)
- Spherical
  - Vectorial Harmonics, [10](#)
- Wave
  - Plane
    - Monochromatic, [3](#)