

PAPER • OPEN ACCESS

Reducing the numerical effort of finite-temperature density matrix renormalization group calculations

To cite this article: C Karrasch *et al* 2013 *New J. Phys.* **15** 083031

View the [article online](#) for updates and enhancements.

Related content

- [Hubbard-to-Heisenberg crossover \(and efficient computation\) of Drude weights at low temperatures](#)
C Karrasch
- [Quantum quenches in the anisotropic spin- \$\frac{1}{2}\$ Heisenberg chain: different approaches to many-body dynamics far from equilibrium](#)
Peter Barmettler, Matthias Punk, Vladimir Gritsev *et al.*
- [Precise evaluation of thermal response functions by optimized density matrix renormalization group schemes](#)
Thomas Barthel

Recent citations

- [Diffusion from convection](#)
Marko Medenjak *et al*
- [Numerical evaluation of two-time correlation functions in open quantum systems with matrix product state methods: a comparison](#)
Stefan Wolff *et al*
- [Finite-temperature density-matrix renormalization group method for electron-phonon systems: Thermodynamics and Holstein-polaron spectral functions](#)
David Jansen *et al*

Reducing the numerical effort of finite-temperature density matrix renormalization group calculations

C Karrasch^{1,2,3}, J H Bardarson^{1,2} and J E Moore^{1,2}

¹ Department of Physics, University of California, Berkeley, CA 95720, USA

² Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

E-mail: karrasch@berkeley.edu

New Journal of Physics **15** (2013) 083031 (21pp)

Received 26 March 2013

Published 13 August 2013

Online at <http://www.njp.org/>

doi:10.1088/1367-2630/15/8/083031

Abstract. Finite-temperature transport properties of one-dimensional systems can be studied using the time dependent density matrix renormalization group via the introduction of auxiliary degrees of freedom which purify the thermal statistical operator. We demonstrate how the numerical effort of such calculations is reduced when the physical time evolution is augmented by an additional time evolution within the auxiliary Hilbert space. Specifically, we explore a variety of integrable and non-integrable, gapless and gapped models at temperatures ranging from $T = \infty$ down to $T/\text{bandwidth} = 0.05$ and study both (i) linear response where (heat and charge) transport coefficients are determined by the current-current correlation function and (ii) non-equilibrium driven by arbitrary large temperature gradients. The modified density matrix renormalization algorithm removes an ‘artificial’ build-up of entanglement between the auxiliary and physical degrees of freedom. Thus, longer time scales can be reached.

³ Author to whom any correspondence should be addressed.



Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Contents

1. Introduction	2
2. Models and methods	5
2.1. Models	5
2.2. Transport properties	5
2.3. Density matrix renormalization group	7
3. Results	8
3.1. Prelude: time evolution of $ \Psi_T\rangle$	8
3.2. Reduction of the growth of entanglement: non-equilibrium	9
3.3. Reduction of the growth of entanglement: equilibrium	11
3.4. Spin correlation function	12
3.5. Optimization of U_Q	13
4. Summary	14
Acknowledgments	15
Appendix. Technical details	15
References	19

1. Introduction

A physical system is usually characterized by its response to perturbations. In transport setups, one studies charge or energy currents driven by voltage or temperature gradients. From the theoretical point of view this is complicated—computing the quantum mechanical time evolution of a system in non-equilibrium is one of the most active areas of research in condensed matter physics. When the external perturbations are small, one can simplify the problem by resorting to the Kubo formalism. For example, the charge conductivity $\sigma(\omega)$ describes the *linear response* current J induced by a *small* electric field (we will be more specific below):

$$\sigma(\omega) \sim \int e^{i\omega t} \langle J(t)J \rangle dt, \quad (1)$$

where the dynamical correlation function $\langle J(t)J \rangle$ is calculated in *thermal equilibrium*:

$$\langle A(t)B \rangle = \text{Tr}(\rho_T e^{iHt} A e^{-iHt} B), \quad \rho_T = \frac{e^{-H/T}}{\text{Tr} e^{-H/T}} \quad (2)$$

with H being the Hamiltonian of the system and T denoting the temperature. Unfortunately, computing transport coefficients such as $\sigma(\omega)$ is generally still difficult: even if one knows the exact thermal density matrix ρ_T (or the exact ground state), extracting correlation functions, which couple all excitations, remains a formidable task.

The key question posed and addressed in this work is: how can linear-response and non-equilibrium transport properties of one-dimensional (1D) systems at finite temperature be calculated *efficiently* using the density matrix renormalization group (DMRG) [1, 2]? DMRG was originally devised [3, 4] as tool to accurately determine *ground states* of 1D Hamiltonians. The reason for its success became understandable when it was formulated using matrix product states (MPS) [5–9]: the area law [10] stipulates that the ground states of 1D systems governed by local Hamiltonians are only entangled locally; this implies that they can be expressed efficiently

using a MPS with a small bond dimension χ (which encodes the amount of entanglement). The MPS describing a given system can be determined variationally—this is the very core of a ground state DMRG calculation [3, 4]. One way to compute *correlation functions* $\langle A(t)B \rangle$ at $T = 0$ is to directly simulate the time evolution (for other approaches see [1, 11–16])

$$e^{-iHt} B |\text{ground state}\rangle \quad (3)$$

using a time-dependent DMRG framework [17–25]. The corresponding algorithm can again be formulated elegantly using matrix product states. The physical growth of entanglement implies that the bond dimension needed to approximate $\langle A(t)B \rangle$ to a certain accuracy grows with time. This limits the accessible time scales.

Standard DMRG methods allow computing the time evolution of a *pure* state and are thus not directly applicable at $T > 0$. Various approaches for simulating finite-temperature dynamics using DMRG have been explored within the literature [26–40]. These include probabilistic sampling over an appropriately chosen set of pure states [30], schemes which time-evolve operators instead of states [31, 32], transfer-matrix DMRG [33–36] and exact representations through purification [37–39]. Purification expresses the thermal statistical operator ρ_T as a partial trace over a pure state $|\Psi_T\rangle$ living in an enlarged Hilbert space where auxiliary degrees of freedom Q encode the thermal bath [41]:

$$\rho_T = \text{Tr}_Q |\Psi_T\rangle \langle \Psi_T|. \quad (4)$$

One of the main advantages of this approach is that all the standard methods for time evolving quantum states within DMRG are directly applicable. In this work, we employ the time evolving block decimation [17, 22].

One of the first applications of finite-temperature ‘purification DMRG’ to dynamical problems was the calculation of spin–spin correlation functions of integrable spin-1/2 Heisenberg chains [38]. While DMRG yields data which are ‘numerically exact’ (this is verified by comparing with analytic results available for ‘non-interacting’ models [33, 34, 38]), the time scales accessible at finite temperatures are considerably smaller than at $T = 0$. This observation, which might be one reason why studies of finite- T dynamics [31–36, 38–40, 42–44] are rarer than their $T = 0$ counterparts, can be understood as follows. Assume we want to compute a ground state correlation function, i.e. evaluate equation (3). Under the time evolution, the entanglement grows *locally* around the region on which B acted. In contrast, at $T > 0$ where one needs to calculate (see below for details)

$$e^{-iHt} B |\Psi_T\rangle, \quad (5)$$

the entanglement generally increases *homogeneously* throughout the whole system. This holds true even for $B = 1$, i.e. when the state $|\Psi_T\rangle$ is exposed to a supposedly trivial time evolution [40]. The reason for this is that purification is not unique and various representations of the same ρ_T differ in their degree of entanglement between the auxiliary and physical degrees of freedom. Even when one starts out with a purification that minimizes this entanglement, it rapidly grows under the DMRG time evolution. This observation naturally leads to the question: Can that very same ‘freedom’ be used to our advantage to undo the non-physical growth of entanglement? In other words, is there a unitary transformation $U_Q(t)$ acting on the auxiliary Hilbert space Q which removes the artificial entanglement? In [40] we proposed to time-evolve Q backwards in time with the physical Hamiltonian acting on the auxiliary degrees of freedom:

$$U_Q(t) = e^{+i\tilde{H}t}. \quad (6)$$

This renders the time evolution of $|\Psi_T\rangle$ trivial, and the evaluation of equation (5) is therefore eventually only plagued by an entanglement building up around the region where B acts in complete analogy to the ground state calculation (the physical reason being quasi-locality [31, 32]). This generically leads to a slower increase of the bond dimension χ , and thus longer time scales can be reached.

In [40], the potential of the modified DMRG algorithm was demonstrated for the spin–spin correlation function $\langle S_n^z(t) S_m^z \rangle$ of the XXZ chain at zero anisotropy $\Delta = 0$, which maps to free fermions and thus allows for an exact (benchmark) solution. A more thorough comparison between the numerical effort of the standard [38] and modified [40] algorithms in calculating $\langle S_n^z(t) S_m^z \rangle$ for arbitrary Δ can be found in [32], where further optimizations using operator-space DMRG were explored. One of the ideas of [32] is to use time translation invariance to rewrite equation (2) as

$$\langle A(t) B \rangle = \langle A(t/2) B(-t/2) \rangle, \quad (7)$$

which allows accessing times *twice as large* without additional effort. Barthel *et al* and Barthel [31, 32] also show that for certain scenarios it can be more efficient to redistribute the evaluation of $\exp(-iHt)$ and $\exp(-\beta H)$ over two DMRG simulations.

From the point of view of physical applications, the modified DMRG schemes were used to investigate current correlation functions and the Drude weight of the integrable XXZ chain at intermediate to high temperatures [40, 44], scaling properties of the dc conductivity in presence of non-integrable perturbations [42], non-equilibrium induced by temperature gradients [43] and spectral functions of hardcore bosons [31]. For reasons of completeness, we mention other approaches to linear response transport properties of the XXZ model (and related ones). These include exact Bethe ansatz calculations [47–50], integrability arguments [51–56], field theories [35, 36, 57–59], quantum Monte Carlo [60–62], exact diagonalization [63–67], transfer matrix DMRG [35, 36] and dynamical DMRG [14]. For a non-exhaustive list of prior works on non-equilibrium thermal transport see [23, 68–78].

The first and foremost goal of this paper is to present extensive quantitative data on how the build-up of entanglement is reduced if the auxiliary Hilbert space Q is time-evolved with the physical Hamiltonian but reversed time. We focus on transport properties in linear response [40, 42, 44] and in thermal non-equilibrium [43]. Specifically, we study (i) homogeneous gapless and gapped spin-1/2 XXZ chains, also in presence of various perturbations which break integrability, (ii) the quantum Ising model, and (iii) impurity setups of a quantum dot connected to non-interacting leads. For temperatures from $T = \infty$ down to $T/\text{bandwidth} = 0.05$, the modified DMRG algorithm leads to a slower increase of the MPS dimension χ . Only at low T does the standard approach $U_Q = 1$ become more efficient (see [32] for details). For most (but not all; we will be more specific below) problems studied in this work, $T/\text{bandwidth} = 0.05$ is a low enough temperature to correspond to the $T = 0$ limit, which one can establish, e.g. by comparing with field theory results [77]. We show the typical behavior of χ on the relevant physical time scales for each problem at hand (e.g. on the time scale at which non-equilibrium currents generically reach their steady state). We reiterate how equation (7) can be implemented within the purification approach and illustrate (following [31, 32]) how it allows accessing times twice as large.

As the second purpose of this paper, we present technical details of the implementation of the algorithm. For example, we show how to time-evolve next-nearest neighbors (closely following [2]), which is necessary because physical degrees of freedom are separated by an

auxiliary site due to the purification. We discuss the numerical accuracy of our data, compare with exact results, investigate to what extent the choice of $U_Q(t) = \exp(i\tilde{H}t)$ is optimal [31, 32] and provide further evidence for the reliability of linear prediction extrapolation schemes [38, 42, 45, 46] for the spin–spin correlation function of the isotropic Heisenberg chain.

2. Models and methods

2.1. Models

As a prototypical model, we consider a chain of interacting spin-1/2 degrees of freedom $S_n^{x,y,z}$ governed by local Hamiltonians

$$h_n = J_n(S_n^x S_{n+1}^x + S_n^y S_{n+1}^y + \Delta_n S_n^z S_{n+1}^z) + b_n(S_n^z - S_{n+1}^z), \quad (8)$$

or equivalently spinless fermions through a Jordan–Wigner transformation. By choosing the couplings J_n , Δ_n and b_n appropriately:

$$J_n = \begin{cases} 1, & n \text{ odd,} \\ \lambda, & n \text{ even,} \end{cases} \quad \Delta_n = \Delta, \quad b_n = \frac{(-1)^n b}{2}, \quad (9)$$

we can study systems which are gapless or gapped, and investigate the role of integrability. For $\lambda = 1$ and $b = 0$, equation (8) can be diagonalized via Bethe ansatz [79]; the model is non-integrable otherwise. The energy spectrum is gapless for $|\Delta| \leq 1$ and gapped for $\Delta > 1$. A gap opens for $\lambda < \lambda_c$ or $b > b_c$, where $\lambda_c < 1$ and $b_c > 0$ if $-1 < \Delta < -1/\sqrt{2}$ [42, 80, 81]. In addition, we study the quantum Ising model

$$h_n = -4S_n^z S_{n+1}^z - g(S_n^x + S_{n+1}^x). \quad (10)$$

2.2. Transport properties

2.2.1. Linear response. We first consider a homogeneous system of size L governed by $H = \sum_{n=1}^{L-1} h_n$ within linear response. The optical charge (C) and energy (E) conductivities can be computed via the Kubo formula

$$\sigma(\omega) = \frac{1}{\omega L} \int_0^\infty e^{i\omega t} \langle [J(t), J] \rangle dt = \frac{1 - e^{-\omega/T}}{\omega L} \int_0^\infty e^{i\omega t} \langle J(t)J \rangle dt, \quad (11)$$

where the corresponding current operators $J = \sum_n j_n$ are defined through a continuity equation⁴:

$$\begin{aligned} \partial_t h_n &= j_{E,n} - j_{E,n+1} \Rightarrow J_E = i \sum_{n=2}^{L-1} [h_{n-1}, h_n], \\ \partial_t S_n^z &= j_{C,n} - j_{C,n+1} \Rightarrow J_C = i \sum_{n=2}^{L-1} [h_{n-1}, S_n^z]. \end{aligned} \quad (12)$$

One usually decomposes the real part of $\sigma(\omega)$ as

$$\text{Re}\sigma(\omega) = 2\pi D\delta(\omega) + \sigma_{\text{reg}}(\omega), \quad (13)$$

⁴ In the charge case we focus on $\lambda = 1$.




Figure 1. The non-equilibrium setups studied in this work. (A) two interacting chains of length $L/2$ which are initially in thermal equilibrium at temperatures T_L and T_R are coupled at time $t = 0$. (B) Two non-interacting chains are coupled at time $t = 0$ via an interacting resonant level model (IRLM).

where the so-called Drude weight D is the prefactor of the singular contribution. D can be determined from the long-time asymptote of the current–current correlation function

$$D = \lim_{t \rightarrow \infty} \lim_{L \rightarrow \infty} \frac{\text{Re} \langle J(t) J \rangle}{2LT}. \quad (14)$$

As already stated above, a time-dependent equilibrium correlation function is defined as

$$\langle A(t) B \rangle = \text{Tr} (\rho_T e^{iHt} A e^{-iHt} B), \quad \rho_T = \frac{e^{-H/T}}{Z_T} \quad (15)$$

with $Z_T = \text{Tr} \exp(-H/T)$ denoting the partition function.

2.2.2. Non-equilibrium. In addition to linear response, we study two thermal non-equilibrium setups. The first (labeled ‘non-equilibrium A’ and depicted in figure 1(A)) is introduced via the following protocol: we initially consider two separate chains

$$H_0 = H_L + H_R = \sum_{n=1}^{L/2-1} h_n + \sum_{n=L/2+1}^{L-1} h_n, \quad (16)$$

each being in thermal (grand-canonical) equilibrium at temperatures T_L and T_R . The corresponding density matrix factorizes

$$\rho_0 = \rho_L \otimes \rho_R, \quad \rho_i = \frac{\exp(-H_i/T_i)}{\text{Tr} \exp(-H_i/T_i)}, \quad i = L, R. \quad (17)$$

At time $t = 0$, the chains are coupled through $h_{L/2}$, and the time evolution of any observable A is computed using $H = H_0 + h_{L/2}$:

$$\langle A(t) \rangle = \text{Tr} \rho(t) A, \quad \rho(t) = e^{iHt} \rho_0 e^{-iHt}. \quad (18)$$

In the second setup (‘non-equilibrium B’; see figure 1(B)) we investigate two non-interacting chains $\Delta = b = 0$, $\lambda = 1$ of length $L/2$,

$$H_0 = H_L + H_R = \sum_{n=1}^{L/2-1} h_n + \sum_{n=L/2+2}^L h_n, \quad (19)$$

at different temperatures T_L and T_R . At $t = 0$, they are coupled via an interacting resonant level model:

$$h_{\text{IRLM}} = t' (S_{L/2}^x S_{L/2+1}^x + S_{L/2}^y S_{L/2+1}^y + U S_{L/2}^z S_{L/2+1}^z) + t' (S_{L/2+1}^x S_{L/2+2}^x + S_{L/2+1}^y S_{L/2+2}^y + U S_{L/2+1}^z S_{L/2+2}^z). \quad (20)$$

The site $n = L/2 + 1$ is initially in an equal superposition of up and down states (i.e. formally at infinite temperature).

2.3. Density matrix renormalization group

In this section, we give a brief overview of the DMRG method [1–4]. More details can be found in the [appendix](#). In order to evaluate equations (15) or (18) by a standard DMRG algorithm (which time evolves wave functions) one first needs to purify the thermal density matrix ρ_T by introducing an auxiliary Hilbert space Q such that $\rho_T = \text{Tr}_Q |\Psi_T\rangle\langle\Psi_T|$. This is analytically possible only at $T = \infty$ where ρ_T factorizes. However, $|\Psi_T\rangle$ can be obtained from $|\Psi_\infty\rangle$ by applying an imaginary time evolution, $|\Psi_T\rangle = e^{-H/(2T)} |\Psi_\infty\rangle$ [26, 37, 38]. The correlation function of equation (15) is *exactly* recast as

$$\langle A(t)B \rangle = \frac{\langle \Psi_\infty | e^{-\frac{H}{2T}} e^{iHt} A e^{-iHt} B e^{-\frac{H}{2T}} | \Psi_\infty \rangle}{\langle \Psi_\infty | e^{-\frac{H}{T}} | \Psi_\infty \rangle}, \quad (21)$$

and this object is directly accessible in standard time-dependent DMRG frameworks [17–22]. It is convenient to first express the initial state $|\Psi_\infty\rangle$ in terms of a matrix product state [5–8],

$$|\Psi_\infty\rangle = \sum_{\sigma_n, \sigma_{n_Q}} A^{\sigma_1} A^{\sigma_{1Q}} \dots A^{\sigma_L} A^{\sigma_{LQ}} |\sigma_1 \sigma_{1Q} \dots \sigma_L \sigma_{LQ}\rangle, \quad (22)$$

where

$$A_{a_i a_{i+1}}^{\sigma_i} = \Lambda_{a_i}^i \Gamma_{a_i a_{i+1}}^{\sigma_i}. \quad (23)$$

Here and in the following σ_i is a short hand for either a physical or auxiliary degrees of freedom: $\sigma_i \in \{\sigma_n, \sigma_{n_Q}\}$. The initial matrices associated with $|\Psi_\infty\rangle$ read

$$\begin{aligned} \Gamma^{\sigma_n = \uparrow} &= (1 \ 0), & \Gamma^{\sigma_n = \downarrow} &= (0 \ -1), \\ \Gamma^{\sigma_{n_Q} = \uparrow} &= (0 \ 1/\sqrt{2})^T, & \Gamma^{\sigma_{n_Q} = \downarrow} &= (1/\sqrt{2} \ 0)^T \end{aligned} \quad (24)$$

as well as $\Lambda^i = 1$. After factorizing the evolution operators $\exp(-\lambda H)$ using a second or fourth order Trotter decomposition, they can be successively applied to equation (23). At each time step $\Delta\lambda$, two singular value decompositions are carried out to update three consecutive matrices. The matrix dimension χ is dynamically increased such that at each time step the sum of all squared discarded singular values is kept below a threshold value ϵ .

An exact modification to the finite-temperature DMRG algorithm (which allows accessing longer time scales) was recently introduced in [40]. One has the analytic freedom to apply any time-dependent unitary transformation

$$U_Q(t) = \sum_{\substack{\sigma_{1Q} \dots \sigma_{LQ} \\ \sigma'_{1Q} \dots \sigma'_{LQ}}} C_{\sigma'_{1Q} \dots \sigma'_{LQ}}^{\sigma_{1Q} \dots \sigma_{LQ}}(t) |\sigma_{1Q} \dots \sigma_{LQ}\rangle \langle \sigma'_{1Q} \dots \sigma'_{LQ}| \quad (25)$$

to the in principle inert auxiliary sites σ_{n_Q} —physical quantities are determined by the trace over Q and are thus not affected by $U_Q(t)$:

$$[U_Q(t), \sigma_n] = 0. \quad (26)$$

For example, equation (21) can be rewritten as

$$\langle \Psi_T | e^{iHt} A e^{-iHt} e^{iHt'} B e^{-iHt'} | \Psi_T \rangle = \langle \Psi_T | U^\dagger(t) A U(t) U^\dagger(t') B U(t') | \Psi_T \rangle, \quad (27)$$

where

$$U(t) = e^{-iHt} U_Q(t). \quad (28)$$




Figure 2. MPS dimension during the time evolution of the state $|\Psi_T\rangle$ which purifies the thermal density matrix of a XXZ chain.

Put differently, purification is not unique. It turned out [31, 32, 40, 42, 43] that choosing

$$U_Q(t) = e^{+i\tilde{H}t}, \quad \tilde{H} = H(\sigma_n \rightarrow \sigma_{n_Q}, \text{ magnetic fields reversed}), \quad (29)$$

i.e. time-evolving the auxiliary sites with the physical Hamiltonian but reversed time, leads to a slower increase of χ and thus longer time scales can be reached. An intuitive way of understanding this will be given below. Only at low temperatures does using $U_Q = 1$ become more efficient [31, 32]. In practice it is most convenient to implement

$$e^{-iHt} e^{i\tilde{H}t} = e^{-iH\Delta t} e^{i\tilde{H}\Delta t} e^{-iH\Delta t} e^{i\tilde{H}\Delta t} \dots \quad (30)$$

3. Results

In this section we present our main results. We first quantify how the entanglement growth is reduced by time-evolving the auxiliaries with the physical Hamiltonian but reversed time. In an extension of earlier studies (see [31, 32, 40] and the discussion in section 1), we study gapless and gapped XXZ chains in presence of non-integrable perturbations and focus on transport problems both in linear response and out of equilibrium. We investigate to what extent $U_{\text{aux}}(t) = \exp(it\tilde{H})$ is optimal within our approach. In passing, we provide another example of the stability of linear prediction (see also [2, 38, 42, 45, 46]) and recast (one of) the ideas of [31, 32] in the language of purification.

3.1. Prelude: time evolution of $|\Psi_T\rangle$

It is instructive to first consider the trivial case of $A = B = 1$ in equation (21), i.e.

$$1 = \frac{\langle \Psi_T | e^{iHt} e^{-iHt} | \Psi_T \rangle}{\langle \Psi_T | \Psi_T \rangle} = \frac{\langle \Psi_T | e^{iHt} U_Q^\dagger(t) e^{-iHt} U_Q(t) | \Psi_T \rangle}{\langle \Psi_T | \Psi_T \rangle}. \quad (31)$$

Figure 2 shows the evolution of the bond dimension χ during the calculation of $e^{-iHt} U_Q(t) |\Psi_T\rangle$. For the standard approach $U_Q(t) = 1$, χ increases with time: the state $e^{-iHt} |\Psi_T\rangle$ which purifies

the density matrix becomes successively more entangled. Choosing $U_Q(t) = e^{i\tilde{H}t}$ removes this artifact. This immediately indicates why longer time scales can be reached in DMRG evaluations of, e.g. current correlation functions $e^{-iHt}U_Q(t)j_n|\Psi_T\rangle$. The entanglement only builds up locally around site n (the physical reason being quasi-locality [31, 32]), and likewise for our non-equilibrium setup it grows locally around the interface region over which the initial temperature gradient falls off. The artificial global build-up of entanglement is removed if the auxiliaries are evolved in time. Only at low T choosing $U_Q = 1$ becomes more efficient [31, 32] since the time evolution of the ground state is trivial (the latter is indicated in figure 2: χ grows more slowly at low $T = 0.2$ than it does at high T).

An intuitive understanding of why the particular choice of $U_Q(t) = e^{i\tilde{H}t}$ removes the ‘artificial’ entanglement growth was recently provided in [31, 32]: in an operator-space formulation, the modified DMRG algorithm corresponds to a Heisenberg time evolution of the matrix product operators representing A and B in equation (27). If A (and likewise for B) is local, most terms in the first Trotter step $e^{iH\Delta t} A e^{-iH\Delta t}$ cancel out. As time evolves,

$$\dots e^{iH\Delta t} e^{iH\Delta t} A e^{-iH\Delta t} e^{-iH\Delta t} \dots, \quad (32)$$

entanglement can only build-up gradually around the region on which A acted due to the so-called light-cone effect in non-relativistic systems: Lieb–Robinson bounds [84] generically stipulate that correlation functions $\langle A(t)B \rangle$ of local operators A and B fall off exponentially for $x > vt$, with x denoting the spatial distance between the regions on which A and B act, and v being some velocity with which excitations propagate.

3.2. Reduction of the growth of entanglement: non-equilibrium

In this section we illustrate quantitatively the effects of time-evolving the auxiliaries for the non-equilibrium setups sketched in figure 1. We start by studying an interacting XXZ chain ($\Delta \neq 0$) with two additional perturbations (dimerization $\lambda < 1$ and a staggered field $b > 0$) that both render the system non-integrable (see equation (8) for a precise definition). At time $t = 0$, two ‘semi-infinite’ chains (typical lengths being $L/2 \sim 100$ –200) prepared in thermal equilibrium at temperatures $T_{L,R}$ are coupled by $h_{L/2}$ to an overall ‘translationally-invariant’ chain. The temperature gradient drives an energy current $\langle j_{E,n}(t) \rangle$ whose typical behavior is shown in figure 3(a). For $b = 0$ the current at the interface $n = L/2$ saturates to a finite value on a scale $t \sim 1$ –10 irrespective of whether the system is gapless or gapped (indicating that either the non-integrable dimerized chain supports dissipationless transport or that its current decays on a hidden large scale; see [43] for further details). The steady-state current of the XXZ chain is of the simple ‘black-body’ form [43, 76, 77, 85]

$$\lim_{t \rightarrow \infty} \langle j_{E,n}(t) \rangle = f(T_L) - f(T_R), \quad (33)$$

implying that it is determined solely by the linear conductance $G(T) \sim \partial_T f(T)$ for arbitrary large $T_L - T_R$:

$$\lim_{t \rightarrow \infty} \langle j_{E,n}(t) \rangle \sim \int_{T_L}^{T_R} G(T) dT. \quad (34)$$

This agrees with a field theory result [77] at low temperatures; asymptotic low- T behavior sets in around $T \lesssim 0.2$.

The time evolution of the corresponding bond dimension χ is shown in figure 3(b) for the integrable XXZ chain both for parameters where it is gapless ($\Delta = 0.5$) and gapped ($\Delta = 2$).




Figure 3. DMRG time evolution of an XXZ chain of length $L = 200$ featuring an initial sharp temperature gradient $T_L \neq T_R$. The system is gapless for z -anisotropies $|\Delta| \leq 1$ and gapped otherwise; it becomes non-integrable in presence of a finite dimerization $\lambda < 1$. Note that for $\Delta = 0.5$ and $\lambda = 1$ asymptotic low-temperature behavior described by a field theory [77] sets in for $T \lesssim 0.2$. (a) Energy current at the interface. Choosing a discarded weight control parameter around $\epsilon \sim 10^{-6}$ at a Trotter stepsize of $\Delta t = 0.2$ is sufficient to accurately obtain its steady-state value. Inset: the *actual* discarded weights during a whole Trotter step and during one substep for $\Delta\chi = 20$. The difference to ϵ is explained in the main text. (b) Evolution of the dimension of the corresponding matrix product state. If the auxiliary degrees of freedom which purify the thermal density matrix are time-evolved with the physical Hamiltonian but reversed time (which is an exact modification to the standard algorithm), the build-up of entanglement is reduced. The computational cost of a singular value decomposition (which dominates the DMRG algorithm) scales as χ^3 .

Figure 4(a) shows the same in presence of non-integrable perturbations as well as for the quantum Ising model of equation (10) at criticality $g = 1$. In all cases and for all temperatures from $T_{L,R} = \infty$ down to $T_{L,R} = 0.1$ (which for this problem corresponds to zero temperature), time evolution of the auxiliaries leads to a slower increase of χ ; note that the computational cost of a singular value decomposition (which dominates the whole DMRG algorithm) scales as χ^3 .

The same reduction of χ holds for the non-equilibrium impurity problem depicted in figure 1(b) where an interacting resonant level model defined in equation (20) is coupled to two non-interacting leads ($\Delta = b = 0, \lambda = 1$) at different temperatures. An exemplary evolution of the MPS dimension is shown in figure 4(b); a discussion of the (transport) physics at small interactions or zero temperature can be found in [86]. Note that due to the appearance of a Kondo-like scale, $T/\text{bandwidth} = 0.05$ does not necessarily correspond to zero temperature for this problem [86].

The discarded weight $\epsilon = 10^{-6}$ in figure 3(b) is chosen small enough to ‘accurately’ determine the current and in particular its steady-state value. This is illustrated in figure 3(a) where ϵ is successively lowered from $\epsilon = 10^{-4}$ to 10^{-7} . Moreover, the non-interacting case $\Delta = b = 0, \lambda = 1$ allows for an exact evaluation of the steady-state current [76], which can be used to benchmark the accuracy of the DMRG data (a comparison can be found in [43]). The inset to figure 3(a) shows a generic example for the *actual* discarded weight (for the precise




Figure 4. (a) The same as in figure 3(b) but in presence of two perturbations (dimerization $\lambda < 1$ and a staggered magnetic field $b > 0$) rendering the model non-integrable. Inset: quantum Ising model at criticality $g = 1$. (b) Dimension of the MPS during the time evolution of two non-interacting XXZ chains of lengths $L/2 = 100$ which feature different temperatures T_L and T_R and are coupled at time $t = 0$ via an interacting resonant level model.

definition of ϵ see appendix A.2.4). It is always bound by 2ϵ ; the total discarded weight during a complete Trotter step $e^{-iH\Delta t}e^{i\tilde{H}\Delta t}$ is typically bound by 10ϵ . It is also instructive to carry out a calculation at a fixed MPS dimension χ rather than a fixed discarded weight. The result is shown in figure 3(a).

3.3. Reduction of the growth of entanglement: equilibrium

We now turn to current correlation functions in thermal equilibrium. Following equations (12) and (A.8), we need to evaluate

$$e^{-iHt}U_Q(t)j_n|\Psi_T\rangle. \quad (35)$$

The ac conductivity $\sigma(\omega)$ is determined by the Fourier transform of $\langle J(t)J \rangle$ via equation (11). The integrable gapless XXZ chain, on which we focus in this section, supports dissipationless transport at finite temperature, i.e. its Drude weight D in equation (13) is finite. D can be obtained from the long-time limit of the current–current correlation function via equation (14). The relevant time scale in this problem is thus the scale on which $\langle J(t)J \rangle$ saturate to their asymptote.

DMRG data for the charge–current correlation function at $\Delta = 0.71$ and for various discarded weights is shown in figure 5(a). At intermediate to high temperatures $T \gtrsim 0.5$, we can access time scales on which $\langle J(t)J \rangle$ saturates (we stop our simulation once the MPS dimension has reached values around $\chi \sim 1500$; see the numbers given in figure 5(a)). More details can be found in [40, 44]. A fairly large $\epsilon \sim 10^{-5}$ is sufficient to obtain D quantitatively. This is supported by the comparison with the exact solution for $\Delta = 0$ shown in the inset to figure 6(a). Note that $\langle J_C(t)J_C \rangle$ is constant for $\Delta = 0$ since J_C is conserved; however, every single term $\langle j_{C,n}(t)j_{C,m} \rangle$ that contributes to the sum is time-dependent. It is again instructive to carry out a calculation using a constant χ rather than a fixed discarded weight; the result is shown in the upper panel of figure 5(a).




Figure 5. (a) DMRG calculation of the global charge current correlation function $\langle J_C(t) J_C \rangle$ of the integrable XXZ chain (length $L = 100$) in thermal equilibrium. The calculation is stopped once χ reaches values around $\chi \sim 1500$; the numbers in the plot denote χ at time $t = 12$ for the different discarded weights (the auxiliaries are time-evolved). (b) Evolution of the corresponding MPS dimension for different anisotropies $\Delta = 0.71$ (gapless) and 3 (gapped).

For all temperatures from $T = \infty$ down to 0.125 time evolution of the auxiliaries leads to a slower increase of the dimension of the matrix product state. This is depicted in figure 5(b). One might expect low- T behavior of $D(T)$ to set in for $T = 0.125$ (see [35, 36, 44]), but the current correlation function decays so slowly that we cannot access its asymptotics without extrapolation. Exemplary data for the gapped phase is shown in the lower inset to figure 5(b).

3.4. Spin correlation function

We briefly present data for the spin–spin correlation function

$$S^{zz}(n, t) = \langle S_{n+L/2}^z(t) S_{L/2}^z \rangle, \quad S^{zz}(k, t) = \sum_n e^{ikn} S^{zz}(n, t), \quad (36)$$

which provides a standard testing ground for DMRG approaches to time-dependent correlation functions at finite temperature [33, 34, 38, 40]. The XX chain $\Delta = 0$ again allows for an exact solution by mapping the model to free fermions. Figure 6(a) shows a comparison of DMRG data with the exact result for $T = 1$ and 0.1 (which for this situation corresponds to low temperature). A large discarded weight $\epsilon \sim 10^{-5}$ is sufficient to reproduce the analytic result up to times $t \sim 30$. If the auxiliaries are time-evolved, the MPS dimension increases only moderately to $\chi \sim 100$ –200 in contrast to the standard approach (see [40] and also [33, 34] for transfer-matrix DMRG data). For finite Δ , the choice of $U_Q(t) = e^{i\tilde{H}t}$ still outperforms $U_Q = 1$ (except at low temperatures), but the difference becomes successively less pronounced [32]. For the isotropic chain $\Delta = 1$ and all $T \gtrsim 0.1$, the accessible time scales are about a factor 1.5–2 larger if the auxiliaries are evolved in time. In [38] (see also [2, 31, 42, 45, 46]), linear prediction was first established as an accurate tool to extrapolate correlation functions by considering exactly-solvable models and subsequently used to compute the Fourier transform of $S^{zz}(k, t)$ of the isotropic chain. For reasons of completeness, we revisit the isotropic chain and compare the result of linear prediction with DMRG data for larger times. Figure 6(b) illustrates that the agreement is good.




Figure 6. (a) Equilibrium spin and global charge–current correlation functions $S^{zz}(k, t) = \sum_n e^{ikn} \langle S_{n+L/2}^z(t) S_{L/2}^z \rangle$ and $\langle J_C(t) J_C \rangle$ of the XXZ chain at $\Delta = 0$. DMRG results are compared with the exact solution obtained by mapping the model to free fermions. Since J_C is conserved, $\langle J_C(t) J_C \rangle$ is constant (we dropped some DMRG data points to improve visibility). However, every single term $\langle j_{C,n}(t) j_{C,m} \rangle$ that contributes to the sum is time-dependent. (b) Demonstration of the stability of linear prediction (see also [38] and [2, 31, 42, 45, 46]) for the isotropic chain $\Delta = 1$. Dashed lines were obtained by fitting the DMRG data for times $t_{\text{fit}} \in [2.5, 5]$ and subsequent extrapolation to $t > 5$ using linear prediction. The extrapolated data almost coincides with the DMRG data for $t > 5$ (solid lines).

3.5. Optimization of U_Q

Finally, we shortly investigate to what extend our choice of $e^{i\tilde{H}t}$ for the time evolution of the auxiliaries is optimal (more details on the optimization of U_Q can be found in [31, 32]). To this end, we consider

$$U_Q^\eta(t) = e^{+i\tilde{H}t + i\eta N}, \quad (37)$$

with N begin an arbitrary Hermitian matrix. We compute the entanglement entropy

$$S_{\text{ent}}(t, \eta) = -2 \sum_{a_n} (\Lambda_{a_n}^{L/2})^2 \log_2 \Lambda_{a_n}^{L/2} - 2 \sum_{a_{n_Q}} (\Lambda_{a_{n_Q}}^{L/2_Q})^2 \log_2 \Lambda_{a_{n_Q}}^{L/2_Q} \quad (38)$$

between the left and right halves of the time-evolved state $e^{-iHt} U_Q^\eta(t) |\Psi_T\rangle$ which purifies the density matrix. We have verified that for arbitrary N coupling nearest neighbors, $S_{\text{ent}}(t, \eta)$ is at least quadratic in η . Figure 7 illustrates this for the two choices

$$N = 1 + 2 \sum_{n_Q=1}^{L-1} (S_{n_Q}^x S_{n_Q+1}^x + S_{n_Q}^y S_{n_Q+1}^y) \quad (39)$$

as well as

$$N = \sum_{n_Q=1}^{L-1} |\sigma_{n_Q} = \downarrow, \sigma_{n_Q+1} = \uparrow\rangle \langle \sigma_{n_Q} = \downarrow, \sigma_{n_Q+1} = \uparrow|. \quad (40)$$




Figure 7. Entanglement entropy of the state $|\Psi_T\rangle$ after time-evolving it up to a time $t = 4$. The physical degrees of freedom are time-evolved via $\exp(-iHt)$; the auxiliaries are time-evolved using $U_Q^\eta(t) = \exp(+i\tilde{H}t + it\eta N)$ where N is given by equation (39) (solid lines) and equation (40) (dashed lines).

As mentioned above, Barthel *et al* and Barthel [31, 32] present thorough details on how to further optimize finite-temperature dynamical DMRG. We here reiterate one of the main ideas using the language of purification. In thermal equilibrium, one can recast any correlation function exploiting time translation invariance:

$$\langle A(t)B \rangle = \langle A(t/2)B(-t/2) \rangle. \quad (41)$$

Thus, one only needs to carry out time evolutions up to times $|t/2|$ in order to compute $\langle A(t)B \rangle$. If $A = A^\dagger = B$ (e.g. for current–current correlation functions), it is sufficient to perform a single calculation

$$e^{iHt/2} U_Q^\dagger(t/2) A e^{-iHt/2} U_Q(t/2) |\Psi_T\rangle. \quad (42)$$

This implies that one can access a time scale twice as large without much additional effort. This is illustrated in figure 8(b) for the spin-autocorrelation function of the isotropic Heisenberg chain.

4. Summary

We presented extensive quantitative data for how the growth of entanglement in finite-temperature dynamical DMRG calculations can be reduced by time-evolving the auxiliary degrees of freedom which purify the thermal statistical operator. The time evolution of the auxiliaries is an exact modification to the standard algorithm. Our particular focus was on energy and charge transport properties both in linear response (described by current–current correlation functions) and in non-equilibrium driven by temperature gradients. We studied a variety of gapless and gapped integrable and non-integrable spin-1/2 chains (i.e. interacting spinless fermions), the quantum Ising model at criticality and impurity (quantum dot) setups. For all temperatures from $T = \infty$ down to $T/\text{bandwidth} = 0.05$, which for most problems investigated in this work corresponds to low temperatures, the build-up of entanglement is reduced. This speeds up numerics and allows to eventually access longer time scales.




Figure 8. (a) Equilibrium spin correlation functions $S^{zz}(n, t) = \langle S_{n+L/2}^z(t) S_{L/2}^z \rangle$ of the isotropic XXZ chain for various discarded weights. The calculations were stopped once the MPS dimension reached values around $\chi \sim 2000$. (b) By rewriting $S^{zz}(n = 0, t) = \langle S_{L/2}^z(t/2) S_{L/2}^z(-t/2) \rangle$ and time-evolving $S_{L/2}^z(\pm t/2)$ independently the time scale t can be accessed by a MPS of a smaller dimension χ (this idea was introduced in [31, 32]).

Acknowledgments

We are grateful to Frank Verstraete for very useful suggestions and thank Thomas Barthel, Fabian Heidrich-Meisner and Steve White for discussions. Support by the Deutsche Forschungsgemeinschaft via KA3360-1/1 (CK), by the DARPA TI program of UCLA (JHB) as well as by the Nanostructured Thermoelectrics program of LBNL (CK and JEM) is acknowledged.

Appendix. Technical details

A.1. Purification

One way to evaluate equations (15) and (18) within the DMRG is to purify [41] the thermal density matrix by introducing an auxiliary Hilbert space Q :

$$\rho_T = \frac{e^{-H/T}}{Z_T} = \text{Tr}_Q |\Psi_T\rangle \langle \Psi_T|. \quad (\text{A.1})$$

At infinite temperature, where ρ_T is proportional to the unit operator, one can analytically write down the purification:

$$\rho_\infty = \frac{1}{Z_\infty} = \frac{1}{2^L} = \text{Tr}_Q |\Psi_\infty\rangle \langle \Psi_\infty| = \text{Tr}_Q \prod_{n=1}^L |\Psi_\infty^n\rangle \langle \Psi_\infty^n|. \quad (\text{A.2})$$

In order to exploit conservation laws within the DMRG algorithm, it is convenient to choose

$$|\Psi_\infty^n\rangle = \frac{1}{\sqrt{2}} (|\sigma_n = \uparrow, \sigma_{n_Q} = \downarrow\rangle - |\sigma_n = \downarrow, \sigma_{n_Q} = \uparrow\rangle), \quad (\text{A.3})$$

where $\sigma_n = \uparrow, \downarrow$ denotes the eigenbasis of S_n^z , and Q is spanned by

$$Q = \text{span} \{ |\sigma_{1_Q} \dots \sigma_{L_Q} \rangle \}. \quad (\text{A.4})$$

One readily verifies that indeed

$$\frac{1}{2} = \frac{1}{2} \sum_{\sigma_n} |\sigma_n\rangle \langle \sigma_n| = \sum_{\sigma_n} \sum_{\sigma_{n_Q}} \langle \sigma_{n_Q} | \Psi_\infty^n \rangle \langle \Psi_\infty^n | \sigma_{n_Q} \rangle. \quad (\text{A.5})$$

The finite-temperature purified state $|\Psi_T\rangle$ is obtained from $|\Psi_\infty\rangle$ by applying an imaginary time evolution [2],

$$\begin{aligned} \rho_T &= \frac{e^{-H/T}}{Z_T} = \frac{Z_\infty}{Z_T} \text{Tr}_Q e^{-H/2T} |\Psi_\infty\rangle \langle \Psi_\infty| e^{-H/2T}, \\ &= \frac{1}{\langle \Psi_T | \Psi_T \rangle} \text{Tr}_Q |\Psi_T\rangle \langle \Psi_T|, \quad |\Psi_T\rangle = e^{-H/2T} |\Psi_\infty\rangle, \end{aligned} \quad (\text{A.6})$$

where we formally replaced

$$H \rightarrow H \otimes 1_Q. \quad (\text{A.7})$$

The correlation function of equation (15) is *exactly* recast as

$$\langle A(t)B \rangle = \frac{\langle \Psi_\infty | e^{-\frac{H}{2T}} e^{iHt} A e^{-iHt} B e^{-\frac{H}{2T}} | \Psi_\infty \rangle}{\langle \Psi_\infty | e^{-\frac{H}{T}} | \Psi_\infty \rangle}, \quad (\text{A.8})$$

and for two time arguments one similarly finds

$$\langle A(t)B(t') \rangle = \frac{\langle \Psi_T | e^{iHt} A e^{-iHt} e^{iHt'} B e^{-iHt'} | \Psi_T \rangle}{\langle \Psi_T | \Psi_T \rangle}. \quad (\text{A.9})$$

In order to evaluate equation (7) for $A = A^\dagger = B$, it is sufficient to compute

$$e^{iHt/2} A e^{-iHt/2} | \Psi_T \rangle. \quad (\text{A.10})$$

The expectation value of any observable A with respect to the time-dependent (non-equilibrium) density matrix of equation (18) is given by

$$\langle A(t) \rangle = \text{Tr} \rho(t) A = \frac{\langle \Psi_\infty | e^{-\frac{H_L}{2T_L} - \frac{H_R}{2T_R}} e^{iHt} A e^{-iHt} e^{-\frac{H_L}{2T_L} - \frac{H_R}{2T_R}} | \Psi_\infty \rangle}{\langle \Psi_\infty | e^{-\frac{H_L}{2T_L} - \frac{H_R}{2T_R}} | \Psi_\infty \rangle}. \quad (\text{A.11})$$

A.2. DMRG algorithm

A.2.1. Initial state. It is convenient to first express the initial state $|\Psi_\infty\rangle$ in terms of a matrix product state [5–8],

$$|\Psi_\infty\rangle = \sum_{\sigma_n, \sigma_{n_Q}} A^{\sigma_1} A^{\sigma_{1_Q}} \dots A^{\sigma_L} A^{\sigma_{L_Q}} |\sigma_1 \sigma_{1_Q} \dots \sigma_L \sigma_{L_Q}\rangle, \quad (\text{A.12})$$

where

$$A_{a_i a_{i+1}}^{\sigma_i} = \Lambda_{a_i}^i \Gamma_{a_i a_{i+1}}^{\sigma_i}. \quad (\text{A.13})$$

Here and in the following σ_i is a short hand for either a physical or auxiliary degrees of freedom: $\sigma_i \in \{\sigma_n, \sigma_{n_Q}\}$. The initial matrices corresponding to equation (A.3) read

$$\begin{aligned} \Gamma^{\sigma_n = \uparrow} &= (1 \ 0), & \Gamma^{\sigma_n = \downarrow} &= (0 \ -1), \\ \Gamma^{\sigma_{n_Q} = \uparrow} &= (0 \ 1/\sqrt{2})^T, & \Gamma^{\sigma_{n_Q} = \downarrow} &= (1/\sqrt{2} \ 0)^T, \end{aligned} \quad (\text{A.14})$$

as well as $\Lambda^i = 1$. Normalization generally stipulates

$$\sum_{\sigma_i a_i} (\Lambda_{a_i}^i \Gamma_{a_i a_{i+1}}^{\sigma_i})^\dagger (\Lambda_{a_i'}^i \Gamma_{a_i' a_{i+1}}^{\sigma_i}) = \delta_{a_{i+1} a_{i+1}'} \quad (\text{A.15})$$

as well as

$$\sum_{\sigma_i a_{i+1}} (\Gamma_{a_i a_{i+1}}^{\sigma_i} \Lambda_{a_{i+1}}^{i+1}) (\Gamma_{a_i' a_{i+1}}^{\sigma_i} \Lambda_{a_{i+1}}^{i+1})^\dagger = \delta_{a_i a_i'}. \quad (\text{A.16})$$

A.2.2. Trotter decomposition. In order to successively apply the imaginary and real time evolutions operators

$$e^{-\lambda H} = e^{-\Delta\lambda H} e^{-\Delta\lambda H} \dots, \quad \lambda = \Delta\lambda + \Delta\lambda + \dots \quad (\text{A.17})$$

to the initial state $|\Psi_\infty\rangle$, we factorize them either via a second or a fourth order Trotter decomposition [2]. The second order scheme reads

$$e^{-\Delta\lambda H} \approx e^{-\Delta\lambda/2 H_1} e^{-\Delta\lambda H_2} e^{-\Delta\lambda/2 H_1}, \quad (\text{A.18})$$

where the $H = H_1 + H_2$, and $H_{1,2}$ contain only terms which mutually commute:

$$H_1 = \sum_{n \text{ odd}} h_n, \quad H_2 = \sum_{n \text{ even}} h_n. \quad (\text{A.19})$$

In order to reduce the error, one can employ a fourth order decomposition [83],

$$e^{-\Delta\lambda H} \approx U(\Delta\lambda_1)^2 U(\Delta\lambda_2) U(\Delta\lambda_1)^2, \quad (\text{A.20})$$

where

$$U(\Delta\lambda_i) = e^{-\Delta\lambda_i H_1/2} e^{-\Delta\lambda_i H_2} e^{-\Delta\lambda_i H_1/2} \quad (\text{A.21})$$

and

$$\Delta\lambda_1 = \frac{1}{4 - 4^{1/3}} \Delta\lambda, \quad \Delta\lambda_2 = \Delta\lambda - 4\Delta\lambda_1. \quad (\text{A.22})$$

We typically employ a second order Trotter decomposition with a time step of $\Delta\beta = 0.005$ during the imaginary time evolution from $\beta = 0$ down to the physical temperature $\beta = 1/T$ and a fourth order decomposition with a time step of $\Delta t = 0.2$ during the real time evolution. This is sufficient for all problems studied in this paper (which we verify by comparing against data obtained for smaller $\Delta\beta = 0.0025$ and $\Delta t = 0.1$). The Trotter decomposition is typically not a significant source of error within time-dependent DMRG calculations [68].

A.2.3. DMRG step. The physical Hamiltonians of equations (8) and (10) couple matrices with indices σ_n and σ_{n+1} which in the matrix product state of equation (A.12) are next-nearest neighbors—they are separated by a matrix associated with an auxiliary degree of freedom σ_{n_Q} . Through equation (A.7), the local Hamiltonians of equations (8) and (10) are formally replaced by

$$h_n \rightarrow h_n \otimes \sum_{\sigma_{n_Q}} |\sigma_{n_Q}\rangle \langle \sigma_{n_Q}|. \quad (\text{A.23})$$

Thus, after each application of $\exp(-\Delta\lambda h_n)$, two singular value decompositions are carried out to update three consecutive matrices; to keep the notation transparent, let us denote those

matrices as:

$$\begin{aligned}\Gamma^{\sigma_0} &\longrightarrow \tilde{\Gamma}^{\sigma_0}, \\ \Gamma^{\sigma_1} &\longrightarrow \tilde{\Gamma}^{\sigma_1}, \quad \Lambda^1 \longrightarrow \tilde{\Lambda}^1, \\ \Gamma^{\sigma_2} &\longrightarrow \tilde{\Gamma}^{\sigma_2}, \quad \Lambda^2 \longrightarrow \tilde{\Lambda}^2,\end{aligned}\tag{A.24}$$

and furthermore abbreviate $h = h_n$. This ‘DMRG update’ can be achieved by a simple and straightforward generalization of the protocol for nearest neighbors outlined in [2]. Firstly, one forms the three-site tensor

$$\Psi_{a_0 a_3}^{\sigma_0 \sigma_1 \sigma_2} = \sum_{a_1 a_2} \Lambda_{a_0}^0 \Gamma_{a_0 a_1}^{\sigma_0} \Lambda_{a_1}^1 \Gamma_{a_1 a_2}^{\sigma_1} \Lambda_{a_2}^2 \Gamma_{a_2 a_3}^{\sigma_2} \Lambda_{a_3}^3,\tag{A.25}$$

which is then acted on by $\exp(-\Delta\lambda h)$:

$$\Phi_{a_0 a_3}^{\sigma_0 \sigma_1 \sigma_2} = \sum_{\sigma'_1 \sigma'_2 \sigma'_3} \Psi_{a_0 a_3}^{\sigma'_0 \sigma'_1 \sigma'_2} e^{-\Delta\lambda h} |_{\sigma_0 \sigma_1 \sigma_2}^{\sigma'_0 \sigma'_1 \sigma'_2}.\tag{A.26}$$

Now, two singular value decompositions (SVDs) are applied to appropriately reshaped tensors in order to obtain the new matrices $\tilde{\lambda}^i$ and $\tilde{\Gamma}^{\sigma_i}$:

$$\begin{aligned}\Phi_{a_0 a_3}^{\sigma_0 \sigma_1 \sigma_2} &= \Phi_{(a_0 \sigma_0), (a_3 \sigma_1 \sigma_2)}, \\ &= \sum_{a_1} U_{(a_0 \sigma_0), a_1} \tilde{\Lambda}_{a_1}^1 (V^\dagger)_{a_1, (\sigma_1 \sigma_2 a_3)}, \\ &= \sum_{a_1} \Lambda_{a_0}^0 \underbrace{\Lambda_{a_0}^{0,-1} U_{a_0 a_1}^{\sigma_0}}_{\tilde{\Gamma}_{a_0 a_1}^{\sigma_0}} \underbrace{\tilde{\Lambda}_{a_1}^1 (V^\dagger)_{a_1, (\sigma_1 \sigma_2 a_3)}}_{\tilde{\Phi}_{a_1, (\sigma_1 \sigma_2 a_3)}},\end{aligned}\tag{A.27}$$

and likewise

$$\begin{aligned}\tilde{\Phi}_{a_1, (\sigma_1 \sigma_2 a_3)} &= \tilde{\Phi}_{(a_1 \sigma_1), (\sigma_2 a_3)}, \\ &= \sum_{a_2} U_{(a_1 \sigma_1), a_2} \tilde{\Lambda}_{a_2}^2 (\tilde{V}^\dagger)_{a_2, (\sigma_2 a_3)}, \\ &= \sum_{a_2} \tilde{\Lambda}_{a_1}^1 \underbrace{\tilde{\Lambda}_{a_1}^{1,-1} U_{a_1 a_2}^{\sigma_1}}_{\tilde{\Gamma}_{a_1 a_2}^{\sigma_1}} \underbrace{\tilde{\Lambda}_{a_2}^2 (\tilde{V}^\dagger)_{a_2 a_3}^{\sigma_2} \Lambda_{a_3}^{3,-1}}_{\tilde{\Gamma}_{a_2 a_3}^{\sigma_2}} \Lambda_{a_3}^3.\end{aligned}\tag{A.28}$$

In case of a real time evolution, the normalization conditions of equations (A.15) and (A.16) are preserved automatically (up to errors associated with a potential truncation of the matrices; see below). For a non-unitary $\exp(-\Delta\beta h)$, however, the matrix product state needs to be brought back to its canonical form in order to simplify the evaluation of expectation values, and, more importantly, because otherwise the truncation below is not optimal. Normalization can be achieved approximately by successively acting with unit operators $\Delta\lambda = 0$ [22] or exactly using the procedure outlined in [82].

A.2.4. The discarded weight. Up to this point, the presented DMRG algorithm to evaluate equations (A.8)–(A.11) is exact (except for the Trotter error which can be made sufficiently small for all problems studied in this work). However, the dimension of the matrices in equation (A.12) increases during each of the singular value decompositions in equations (A.27) and (A.28), generically by a factor of two (corresponding to the number of local degrees of freedom). For brevity, let us neglect that for open boundary conditions the matrix dimension is

position-dependent and simply denote it by χ (a more thorough discussion can be found in [2]). The computational effort of the DMRG algorithm is dominated by the cost of the singular value decompositions, which scales as χ^3 . Increasing $\chi \rightarrow 2\chi$ during each sub-step of a Trotter step is numerically unfeasible (and fortunately generically unnecessary). Thus, we truncate the matrices to a dimension $\chi' < 2\chi$ by dropping the indices a_i which correspond to the smallest singular values $\tilde{\Lambda}_{a_i}$. For a normalized state (i.e. if equations (A.15) and (A.16) hold), this is the best approximation with respect to the Hilbert space norm. The error (i.e. the norm-distance of the states before and after truncation) is given by the sum of all squared singular values which are discarded during all sub-steps of one Trotter step. This so-called discarded weight is the key numerical control parameter; the DMRG algorithm becomes exact as it is sent to zero.

One can strictly enforce a fixed discarded weight by analyzing the singular values during all SVDs carried out within one Trotter substep and then retrospectively choosing χ' appropriately (or by simply repeating the substep if $\chi' > \chi$). More pragmatically, one can fix $\chi' = \chi$ and increase χ by $\Delta\chi$ after one substep if the total discarded weight exceeds a pre-defined threshold ϵ . While this simple approach certainly becomes inapplicable if one aims at strictly maintaining a discarded weight that is very small, it works for all problems studied in this paper, with $\Delta\chi \sim 10\text{--}20$ being a reasonable choice. We observe that the actual discarded weight is smaller than 2ϵ for all data presented in this paper (it is imperative to always monitor the real discarded weight which determines the actual error; this is illustrated in the inset to figure 3(a)). In the following we neglect this difference and simply refer to our control parameter ϵ as ‘discarded weight’. We successively lower ϵ from 10^{-8} – 10^{-13} during the imaginary time evolution and from 10^{-3} – 10^{-7} during the real time evolution, which turns out to be sufficient to obtain ‘accurate’ results for all physical quantities of interest. ‘Accurate’ means ‘sufficiently converged with respect to lowering ϵ ’, or ‘in reasonable agreement with exact data’ in case that the problem allows for an analytic solution; all this is illustrated in figures 3(a), 5(a), 6(a) and 8(a). We stop our simulations once the matrix dimension has reached values around $\chi \sim 1500\text{--}2500$.

References

- [1] Schollwöck U 2005 *Rev. Mod. Phys.* **77** 259
- [2] Schollwöck U 2011 *Ann. Phys.* **326** 96
- [3] White S R 1992 *Phys. Rev. Lett.* **69** 2863
- [4] White S R 1993 *Phys. Rev. B* **48** 10345
- [5] Fannes M, Nachtergaele B and Werner R F 1991 *J. Phys. A: Math. Gen.* **24** L185
- [6] Östlund S and Rommer S 1995 *Phys. Rev. Lett.* **75** 3537
- [7] Verstraete F, Cirac J I and Murg V 2008 *Adv. Phys.* **57** 143
- [8] Verstraete F and Cirac J I 2006 *Phys. Rev. B* **73** 094423
- [9] Schuch N, Wolf M M, Verstraete F and Cirac J I 2008 *Phys. Rev. Lett.* **100** 030504
- [10] Eisert J, Cramer M and Plenio M B 2010 *Rev. Mod. Phys.* **82** 277
- [11] Hallberg K 1995 *Phys. Rev. B* **52** 9827
- [12] Ramasesha S, Pati S K, Krishnamurthy H, Shuai Z and Bredas J 1997 *Synth. Met.* **85** 1019
- [13] Kühner E D and White S R 1999 *Phys. Rev. B* **60** 335
- [14] Jeckelmann E 2002 *Phys. Rev. B* **66** 045114
- [15] Holzner A, Weichselbaum A, McCulloch I P, Schollwöck U and von Delft J 2011 *Phys. Rev. B* **83** 195115
- [16] Dargel P E, Honecker A, Peters R, Noack R M and Pruschke T 2011 *Phys. Rev. B* **83** 161104
- [17] Vidal G 2004 *Phys. Rev. Lett.* **93** 040502

- [18] White S R and Feiguin A E 2004 *Phys. Rev. Lett.* **93** 076401
- [19] Daley A, Kollath C, Schollwöck U and Vidal G 2004 *J. Stat. Mech.: Theory Exp.* **P04005**
- [20] Feiguin A E and White S R 2005 *Phys. Rev. B* **72** 020404
- [21] Schmitteckert P 2004 *Phys. Rev. B* **70** 121302
- [22] Vidal G 2007 *Phys. Rev. Lett.* **98** 070201
- [23] Prosen T and Žnidarič M 2009 *J. Stat. Mech.* **P02035**
- [24] Bañuls M C, Hastings M B, Verstraete F and Cirac J I 2009 *Phys. Rev. Lett.* **102** 240603
- [25] Muth D, Unanyan R G and Fleischhauer M 2011 *Phys. Rev. Lett.* **106** 077202
- [26] Verstraete F, García-Ripoll J J and Cirac J I 2004 *Phys. Rev. Lett.* **93** 207204
- [27] Zwolak M and Vidal G 2004 *Phys. Rev. Lett.* **93** 207205
- [28] Wang X Q and Xiang T 1997 *Phys. Rev. B* **56** 5061
- [29] Shibata N 1997 *J. Phys. Soc. Japan* **66** 2221
- [30] White S R 2009 *Phys. Rev. Lett.* **102** 190601
- [31] Barthel T, Schollwöck U and Sachdev S 2012 arXiv:1212.3570
- [32] Barthel T 2013 arXiv:1301.2246
- [33] Sirker J and Klümper A 2005 *Phys. Rev. B* **71** 241101
- [34] Sirker J 2006 *Phys. Rev. B* **73** 224424
- [35] Sirker J, Pereira R G and Affleck I 2009 *Phys. Rev. Lett.* **103** 216602
- [36] Sirker J, Pereira R G and Affleck I 2011 *Phys. Rev. B* **83** 035115
- [37] Feiguin A E and White S 2005 *Phys. Rev. B* **72** 220401
- [38] Barthel T, Schollwöck U and White S R 2009 *Phys. Rev. B* **79** 245101
- [39] Feiguin A E and Fiete G A 2010 *Phys. Rev. B* **81** 075108
- [40] Karrasch C, Bardarson J H and Moore J E 2012 *Phys. Rev. Lett.* **108** 227206
- [41] Uhlmann A 1976 *Rep. Math. Phys.* **9** 273
- Uhlmann A 1986 *Rep. Math. Phys.* **24** 229
- [42] Huang Y, Karrasch C and Moore J E 2012 arXiv:1212.0012
- [43] Karrasch C, Ilan R and Moore J E 2012 arXiv:1211.2236
- [44] Karrasch C, Hauschild J, Langer S and Heidrich-Meisner F 2013 *Phys. Rev. B* **87** 245128
- [45] White S and Affleck I 2008 *Phys. Rev. B* **77** 134437
- [46] Kjäll J, Pollmann F and Moore J E 2011 *Phys. Rev. B* **83** 020407
- [47] Shastry B S and Sutherland B 1990 *Phys. Rev. Lett.* **65** 243
- [48] Zotos X 1999 *Phys. Rev. Lett.* **82** 1764
- [49] Benz J, Fukui T, Klümper A and Scheeren C 2005 *J. Phys. Soc. Japan Suppl.* **74** 181
- [50] Klümper A and Sakai K 2002 *J. Phys.: A: Math* **35** 2173
- [51] Castella H, Zotos X and Prelovšek P 1995 *Phys. Rev. Lett.* **74** 972
- [52] Zotos X, Naef F and Prelovšek P 1997 *Phys. Rev. B* **55** 11029
- [53] Prosen T 2011 *Phys. Rev. Lett.* **106** 217206
- [54] Znidarič M 2011 *Phys. Rev. Lett.* **106** 220601
- [55] Prosen T and Znidarič M 2012 *Phys. Rev. B* **86** 125118
- [56] Znidarič M 2013 *Phys. Rev. Lett.* **110** 070602
- [57] Rosch A and Andrei N 2000 *Phys. Rev. Lett.* **85** 1092
- [58] Saito K 2003 *Phys. Rev. B* **67** 064410
- [59] Orignac E, Chitra R and Citro R 2003 *Phys. Rev. B* **67** 134426
- [60] Alvarez J V and Gros C 2002 *Phys. Rev. Lett.* **88** 077203
- [61] Alvarez J V and Gros C 2002 *Phys. Rev. Lett.* **89** 156603
- Heidrich-Meisner F, Honecker A, Cabra D C and Brenig W 2004 *Phys. Rev. Lett.* **92** 069703
- Gros C and Alvarez J V 2004 *Phys. Rev. Lett.* **92** 069704
- [62] Heidarian D and Sorella S 2007 *Phys. Rev. B* **75** 241104
- [63] Narozhny B N, Millis A J and Andrei N 1998 *Phys. Rev. B* **58** R2921

- [64] Heidrich-Meisner F, Honecker A, Cabra D C and Brenig W 2002 *Phys. Rev. B* **66** 140406
- [65] Heidrich-Meisner F, Honecker A, Cabra D C and Brenig W 2003 *Phys. Rev. B* **68** 134436
- [66] Jung P, Helmes R W and Rosch A 2006 *Phys. Rev. Lett.* **96** 067202
- [67] Jung P and Rosch A 2007 *Phys. Rev. B* **76** 245108
- [68] Gobert D, Kollath C, Schollwöck U and Schütz G 2005 *Phys. Rev. E* **71** 036102
- [69] Michel M, Hess O, Wichterich H and Gemmer J 2008 *Phys. Rev. B* **77** 104303
- [70] Langer S, Heidrich-Meisner F, Gemmer J, McCulloch I P and Schollwöck U 2009 *Phys. Rev. B* **79** 214409
- [71] Langer S, Heyl M, McCulloch I P and Heidrich-Meisner F 2011 *Phys. Rev. B* **84** 205115
- [72] Prosen T 2011 *Phys. Rev. Lett.* **107** 137201
- [73] Steinigeweg R, Ogiewa M and Gemmer J 2009 *Europhys. Lett.* **87** 10002
- [74] Ajisaka S, Barra F, Mejia-Monasterio C and Prosen T 2012 arXiv:1204.1321
- [75] Arrachea L, Lozano G S and Aligia A A 2009 *Phys. Rev. B* **80** 014425
- [76] Bruneau L, Jaksic V and Pillet C 2012 arXiv:1201.3190
- [77] Bernard D and Doyon B 2012 *J. Phys. A: Math. Gen.* **45** 362001
- [78] Bernard D and Doyon B 2013 arXiv:1302.3125
- [79] Haldane F D M 1980 *Phys. Lett. A* **81** 153
- [80] Alcaraz F C and Malvezzi A L 1998 *J. Phys. A: Math. Gen.* **28** 1521
- [81] Takayoshi S and Sato M 2010 *Phys. Rev. B* **82** 214420
- [82] Shi Y-Y, Duan L-M and Vidal G 2006 *Phys. Rev. A* **74** 022320
- [83] Suzuki M 1976 *Prog. Theor. Phys.* **56** 1454
- [84] Lieb E H and Robinson D W 1972 *Commun. Math. Phys.* **28** 251
- [85] Cardy J 2010 *J. Stat. Mech.* **P10004**
- [86] Mehta P and Andrei N 2006 *Phys. Rev. Lett.* **96** 216802
 - Mehta P, Chao S-P and Andrei N 2007 arXiv:cond-mat/0703426 (erratum)
 - Doyon B 2007 *Phys. Rev. Lett.* **99** 076806
 - Borda L, Vladár K and Zawadowski A 2007 *Phys. Rev. B* **75** 125107
 - Boulat E, Saleur H and Schmitteckert P 2008 *Phys. Rev. Lett.* **101** 140601
 - Karrasch C, Pletyukhov M, Borda L and Meden V 2010 *Phys. Rev. B* **81** 125122
 - Karrasch C, Andergassen S, Pletyukhov M, Schuricht D, Borda L, Meden V and Schoeller H 2010 *Europhys. Lett.* **90** 30003
 - Andergassen S, Pletyukhov M, Schuricht D, Schoeller H and Borda L 2011 *Phys. Rev. B* **83** 205103
 - Andergassen S, Pletyukhov M, Schuricht D, Schoeller H and Borda L 2011 *Phys. Rev. B* **84** 039905 (erratum)
 - Kennes D M, Jakobs S G, Karrasch C and Meden V 2012 *Phys. Rev. B* **85** 085113
 - Kennes D M and Meden V 2012 arXiv:1210.1340