

Mammalian musculoskeletal regeneration is associated with reduced inflammatory cytokines and an influx of T cells.

Thomas R. Gawriluk^{1,*}, Jennifer Simkin², Corin K. Hacker¹, John M. Kimani³, Stephen G. Kiama³, Vanessa O. Ezenwa^{4,5}, and Ashley W. Seifert^{1,3,*}

¹*Department of Biology, University of Kentucky, Lexington, KY, 40506 USA.*

²*Department of Orthopedics, Louisiana State University Health Sciences Center, New Orleans, LA, 70112 USA.*

³*Department of Veterinary Anatomy and Physiology, University of Nairobi, Nairobi, Kenya.*

⁴*Odom School of Ecology, University of Georgia, Athens, GA, 30602 USA.*

⁵*Department of Infectious Disease, College of Veterinary Medicine, University of Georgia, Athens, GA, 30602 USA.*

Correspondence:

Thomas R. Gawriluk: tgawriluk@uky.edu

Ashley W. Seifert: awseifert@uky.edu

ABSTRACT

Whether the immune response to injury contributes to tissue regeneration is not well understood. We quantified systemic and local cytokines during ear pinna repair to provide the first comprehensive comparison of the immune response to injury between mammalian regeneration (*A. cahirinus* and *A. percivali*) and fibrotic repair (*M. musculus*). Importantly, by comparing laboratory-reared and wild-caught animals we identified responses specifically associated with healing outcome. Fibrotic repair showed a greater local release of IL-6, CCL2 and CXCL1. Conversely, regeneration showed decreased circulating IL-5, IL-6, IL-17, CCL3 and CXCL1 and increased local IL-12 and IL-17. The differential IL-6 response was substantiated by increased pSTAT3 during the inflammatory phase of fibrotic repair and with blastema formation and tissue morphogenesis in *Acomys*. COX-2 inhibition was not sufficient to induce regeneration. Interestingly, a unique influx of lymphocytes was coupled with regeneration and RNA-expression analysis suggested they were regulatory T cells. Together, the data support regeneration-specific inflammation and T cell responses in *Acomys*.

Keywords: spiny mouse, *Acomys*, immune response, regeneration, blastema, inflammation, T cell, wild animal

INTRODUCTION

Epimorphic regeneration in response to tissue damage occurs as a chronological and overlapping series of processes that includes hemostasis, inflammation, re-epithelialization, activation of local progenitor cells, tissue morphogenesis, and replacement of the injured tissue. The outcome of these processes is scar-free healing. In contrast, most human injuries heal by fibrotic repair, which is characterized by limited cellular proliferation and intense collagen deposition that results in a scar repairing the damaged tissue [1]. As with any trauma or infection that disrupts tissue architecture, regeneration and fibrotic repair are concomitant with a multiphasic immune response that promotes hemostasis, creates inflammation, protects against microbial infection, initiates re-epithelialization, and stimulates a local fibrotic response [2]. During most instances of regeneration (e.g., limb, fin, digit tip, etc.) there is an apparent resolution of acute inflammation that coincides with the accumulation of resident cells, which subsequently re-enter the cell cycle and self-organize to undergo morphogenesis [3, 4]. This transition from an inflammatory environment to morphogenesis is synonymous with regenerative blastema formation [5]. As such, the injured tissue must precisely coordinate dynamic interactions between cells and factors (i.e., cytokines, chemokines, growth factors etc.) within the injury microenvironment to resolve the inflammatory response and promote blastema formation.

Despite a rich literature describing the effects of immune cells and their products in non-regenerating wounds [6, 7], our knowledge of the immune response during epimorphic regeneration remains poor [8-10]. Recent studies in fish, frogs, salamanders, and spiny mice support that immune cells and their products are required for blastema formation and successful regeneration. For instance, complete or timed depletion of macrophages or regulatory T cells (T_{REG}) prevents re-epithelialization and subsequent blastema formation [11-18] and blocking

reactive oxygen species (ROS) production elicits a similar outcome [19-21]. Perhaps not unexpectedly, similar experiments in non-regenerating systems cause incomplete wound closure and angiogenesis, suggesting that the same immune signals initiate fibrotic repair and regeneration [22-27]. Moreover, comparing the immune response to injury between fetal and adult mammals [28-30], pre- and post-metamorphic amphibians [31-33], closely-related regenerating and non-regenerating vertebrates [34], and regeneration-competent and scarring tissues in the same animal [35-37] all support that reduced inflammation and a muted immune response promotes epimorphic regeneration over fibrotic repair. However, an important series of studies support the idea that some immune cells are passive participants during tissue regeneration. For example, removal of the spleen [38], or induction of leukopenia [39] during newt limb regeneration demonstrate that a severely reduced leukocyte response does not prevent blastema formation or regeneration.

These contrasting viewpoints raise several unanswered questions. (1) Are there specific factors produced by immune cells that polarize local cell phenotypes and specifically promote regeneration or fibrotic repair? (2) Does the inflammatory response impede blastema formation and subsequent regeneration in adult mammals? (3) Are the initial stages of fibrotic repair and regeneration driven by different immune responses, such that altering the immune response could stimulate regeneration in lieu of fibrotic repair? In order to address these questions and to test how immune cell factors specifically affect resident cells in injured tissue, there is a need to establish the timing and extent to which these factors are deployed during regeneration and fibrotic repair.

In this study, we provide insight of the immune mechanisms that coincide with mammalian regeneration by following-up on the relatively recent discovery that multiple species

of spiny mice (e.g., *Acomys cahirinus*, *A. percivali*, *A. kempfi*) regenerate skin and musculoskeletal tissue [15, 40-42]. Previous work in spiny mice supports that epimorphic regeneration is associated with resident cell activation, cell cycle progression, cell proliferation, and an inflammatory response distinguished by a prolonged signature of NADPH-oxidase-derived ROS [15, 40, 42]. Additionally, macrophages are required for blastema formation in spiny mice [15], and there is additional evidence that macrophage subtypes differentially drive regeneration or fibrotic repair in rodents [15-17]. Therefore, we sought to characterize and compare the cytokine response to injury during fibrotic repair and regeneration to specifically test if the immediate immune response to injury is different between these two healing outcomes.

We compared the injury response using a 4 mm ear punch assay among three species (*A. cahirinus*, *A. percivali* and *M. musculus*) and two source populations (wild-caught *A. percivali* and *M. musculus*, and laboratory-reared *A. cahirinus* and *M. musculus*) using a panel of sixteen cytokines. Our results showed that injury across all groups induced a common set of pro-inflammatory cytokines (IL-1 β , IL-6, TNF α) and leukocyte chemotactic factors (GM-CSF/CSF2, and MIP-1 α /CCL3) with a similar timed resolution, supporting that some signals of acute inflammation are a shared feature of regeneration and fibrotic repair. Supporting a difference between regeneration and fibrotic repair, we observed differential responses for several pro-inflammatory cytokines during the acute inflammatory phase. Surprisingly, we found a faster, stronger and prolonged adaptive immune response characterized by T cell influx during regeneration.

RESULTS

Cross-species validation of cytokine detection in rodent serum and tissue

To begin characterizing the mammalian immune response during epimorphic regeneration, we analyzed sixteen cytokines (Interleukin 1-alpha (IL-1 α), IL-1 β , IL-2, IL-4, IL-5, IL-6, IL-10, IL-12p70, IL-17, chemokine (C-C motif) ligand 2 (CCL2) (a.k.a. monocyte chemoattractant protein 1 or MCP-1), CCL3 (a.k.a. macrophage inflammatory protein 1 α or MIP-1 α), CCL5 (a.k.a. regulated on activation, normal T cell expressed and secreted or RANTES), colony-stimulating factor 2 (CSF2) (a.k.a. granulocyte-macrophage colony-stimulatory factor or GM-CSF), tumor necrosis factor-alpha (TNF α), interferon-gamma (IFN γ) and chemokine (C-X-C motif) ligand 1 (CXCL1) (a.k.a. KC)) using a custom-designed sandwich ELISA array. We used this assay to compare five groups: three at the University of Kentucky, (1) laboratory-reared, outbred *Mus musculus* (Mm-UKY), (2) wild-caught *M. musculus* (Mm-Wild), (3) laboratory-reared *Acomys cahirinus* (Ac), and two at the University of Nairobi, (4) outbred *M. musculus* reared by a local breeder (Mm-Kenya) and (5) wild-caught *A. percivali* (Ap). Our experimental design allowed us to compare cytokine responses between regenerating and non-regenerating species (Ac and Ap compared to Mm-UKY, Mm-Kenya and Mm-Wild), and between immune-challenged and laboratory-reared animals (Mm-Kenya, Mm-Wild and Ap compared to Mm-UKY and Ac).

Parallelism analysis showed comparable slopes between *Mus* and *Acomys* serum and tissue samples with the mouse standard curve for a majority of cytokines (Table 1 and Supplementary Figure 1). Given non-parallel slopes, we did not validate using this assay to compare IL-10 or CCL5 between species (Figure S1). Several cytokines not present in *Acomys* serum were quantified in tissue lysate (Ac: IL-1 β , IL-4, IL-6, IL-17, CSF2, CCL2; Ap: IL-17,

CSF2, CXCL1) supporting that the serum concentration was below the limit of detection and that the antibody binding epitopes were conserved between species. Therefore, if the cytokine was detected in one tissue source or one *Acomys* species, we concluded it could be detected in the other source or species. This provided us with a way to determine if the cytokine was present or absent. A comparison of full-length predicted amino acid sequences between *A. cahirinus* and *M. musculus* indicated conservation—minimum of 56.8% (IFN γ) to a maximum of 95.8% (TNF α) (Table 2 and Supplemental Figure 2A-Q). Given that CXCL1 was the only cytokine not detected in the Ac samples, this supports that it was likely not being present versus a failure to detect it. Together, these results supported that the ELISA could be used to directly compare changes in most of the cytokines between species.

Fibrotic repair is associated with elevated amounts of circulating IL-5, IL-6 and CCL3.

Using our cytokine assay, we first compared circulating serum cytokine concentrations from uninjured animals among groups (species and source population) to establish a systemic baseline for each group (Figure 1A). A total of 13 cytokines were compared as CSF2 was not present in the serum of any species. While many baseline concentrations were similar between groups, immune-challenged animals (i.e., wild) exhibited higher IL-4, IL-6, CCL2, and TNF α compared to laboratory-reared animals (Figure 1A). Interestingly, the Mm-Kenya animals were a transitional group between Mm-UKY and Mm-Wild for TNF α and IL-4 (Figure 1A). Heightened concentrations of IL-6, TNF α and IL-4 support previous pathogen exposure and a possibility of current infection [43, 44]. Thus, Mm-Kenya, Mm-Wild and Ap had a relatively activated immune system, while Mm-UKY and Ac possessed a more naïve immune system [45]. There were no consistent differences between regenerators and non-regenerators (Figure 1A).

Next, we quantified the systemic injury response for each cytokine compared to its baseline, beginning 24 hr (D1) after injury and over the next twenty days (Figure 1B). In most cases (except IL-2, IL-6, IL-17 and CXCL1), there was no effect of day (Supplemental Table 2), indicating that the immediate systemic response persisted for 20 days. Animals with a more naïve immune system showed increased IL-2 and TNF α , and decreased IL-1 α compared to a relatively activated immune system (Figure 1B: solid lines compared to intermittent lines and Supplemental Data File). Regeneration showed decreased IL-5, IL-6, IL-17, CCL3 and CXCL1 compared to fibrotic repair (Figure 1B: red lines compared to black lines and Supplemental Data File). This latter result supported that animals healing by fibrotic repair and regeneration could be separated by their systemic response to injury.

A regenerative microenvironment is marked by induction of T-cell-associated cytokines and a dampened pro-inflammatory cytokine response.

Resident cells and infiltrating immune cells secrete cytokines that likely polarize the injury microenvironment to support regeneration or fibrotic repair [12, 13, 15, 16, 34, 46, 47]. Thus, to quantify local cytokine concentrations we assayed tissue lysate collected throughout the healing response. IL-1 α could not be compared because baseline concentrations were above the upper limit of quantification in more than 80% of samples, indicating that IL-1 α in the ear pinna was at least two orders of magnitude greater than the other cytokines measured. Although ear pinna tissue is structurally similar across species [40], local cytokines in *Acomys* were consistently detected at lower concentrations compared to *M. musculus*. Moreover, because the concentration of cytokine may not be as important as the dynamics of the cytokine, we compared the change in cytokine compared to baseline over 20 days (Figure 2). Injury elicited significant

changes over time compared to baseline for all cytokines measured in tissue lysate supporting a dynamic response (Supplemental Table 3). Furthermore, while most cytokines shared similar trajectories over time, there was an effect of Group and the Group*Day interaction for all cytokines supporting significant differences in the magnitude of change among the groups (Figure 2 and Supplemental Table 3). Supporting an inflammatory response in all groups, several pro-inflammatory cytokines (IL-6, TNF α) and myeloid chemotactic factors (CCL3, CSF2 and CXCL1) showed an increase compared to baseline between D1 and D3 that then decreased to baseline or below between D5 and D20 (Figure 2). There was also an overall decrease compared to baseline for IL-5 and a small but significant change from baseline for IL-2 and IL-4 across all groups (Figure 2). Animals with naïve immune responses had a stronger increase in CCL3 and a smaller increase for CCL2 and CXCL1 compared to activated immune responses (Figure 2: solid compared to intermittent lines). We also identified several cytokines that showed differential changes between regeneration and fibrotic repair that we describe below (Figure 2: red compared to black lines).

During the acute inflammatory phase (D1 and D2), CCL2 and CXCL1 were increased 9 and 12-fold during fibrotic repair compared to regeneration, respectively (Figure 3). IL-6 showed a similar result at D2 where Mm-UKY, Mm-Kenya and Mm-Wild were increased 10-fold compared to Ac and Ap (Figure 3). Additionally, IL-17 was increased in Ap, but decreased in Mm-UKY and Mm-Kenya and IL-12 was increased in Ap compared to all *Mus* (Figure 3). Interestingly, the TNF α response—a biomarker of inflammation—could not reliably separate fibrotic repair and regeneration (Figure 3).

Regardless of healing outcome, re-epithelialization occurs by D10 [40, 42] coincident with resolution of many pro-inflammatory cytokine responses (Figure 1B and 2: yellow bars).

While there were some differences among groups for the timing of resolution, IL-1 β , TNF α , and CCL2 were similar to or below baseline at D10 for each species (Figure 3). IL-6 also followed this pattern; however, there was a differential response where Ac remained elevated through D20 while all other species decreased below baseline (Figure 3).

At D20, during tissue morphogenesis, the only cytokines that showed a differential response were IL-12 and IL-17 that were increased during regeneration compared to fibrotic repair (Figure 3). The anti-inflammatory cytokine IL-4 did not differ over time with respect to regenerative ability, suggesting that the differences in pro-inflammatory cytokine release is likely not an IL-4 mediated response. Our results suggest that subtle differences in how cytokines are deployed in the injury microenvironment can distinguish regeneration or fibrotic repair. These data suggest that strong, acute increases in the pro-inflammatory cytokines IL-6, CCL2 and CXCL1 are associated with fibrosis, while the release of IL-12 and IL-17 during tissue morphogenesis is associated with regeneration.

Regeneration is associated with an early burst of T cell influx to the injury site.

The release of IL-12 and IL-17 into the regenerative microenvironment suggested enhanced T cell activation during regeneration [48, 49]. Therefore, we quantified T cell influx into uninjured and healing tissue from our laboratory populations of *Mus* (Mm-UKY) and *Acomys* (Ac) using flow cytometry with an antibody to the extracellular portion of the T cell marker CD3. We observed significant differences in CD3⁺ cells in injured tissue between species over time (two-way ANOVA, n=57; species: Df=1, F=49.49 $P<0.001$; day: Df=6, F=89.07, $P<0.001$; species*day: Df=6, F=21.49 $P<0.001$) (Figure 4A). In uninjured tissue, Mm-UKY had 10-times more CD3⁺ cells compared to Ac (Tukey-Kramer HSD post-hoc test,

Df=6, $t=6.21$, $P<0.001$) (Figure 4A). While the total number of CD3⁺ cells that infiltrated the wound was higher in Mm-UKY compared to Ac, there was a greater fold change relative to D0 for CD3⁺ cells during regeneration compared to fibrotic repair (Figure 4B). Ac exhibited a monophasic response to injury starting on D1 with a 78-fold influx of T cells that peaked on D3 and remained above baseline at D15. Mm-UKY showed a biphasic response with peak influx of 10-fold at D7 that returned to baseline at D15 (Figure 4B). Importantly, at D15, when IL-12 was increased (Figure 2), the influx of CD3⁺ cells remained high in Ac compared to Mm-UKY (Figure 4B).

We next used immunohistochemistry with an antibody specific to the intracellular portion of the CD3 receptor to assess the spatial distribution of T cells during acute inflammation and morphogenesis (Figure 4C-E). First, these data confirm the greater influx of CD3⁺ cells at D5 and D15 in Ac compared to Mm-UKY (Figure 4C-E). Second, in Mm-UKY most CD3⁺ cells were associated with the epidermis and were rarely observed distal to the amputation plane (Figure 4C). On the other hand, CD3⁺ cells in Ac were present in the epidermis and dermis, and regularly observed in healing tissue distal to the amputation plane (Figure 4E). At D15, CD3⁺ cells were found in the epidermis and dermis of both species (Figure 4D, F). Interestingly, CD3⁺ cells associated with the epidermis in Mm-UKY (Figure 4G) exhibited a spindle-shape morphology compared to a rounded shape in Ac (Figure 4H). There also appeared to be more CD3⁺ cells in the dermis of Ac compared to Mm-UKY (Figure 4D, F), and the CD3⁺ cells tended to localize near regenerating hair follicles in Ac (Figure 4I). Attempts to characterize individual T cell phenotypes during regeneration using flow cytometry and IHC using 19 commercially available antibodies, supported significant differences in antibody-epitope binding between species that prevented further T cell phenotyping by receptor subtype in *Acomys*

(Supplemental Table 5). Therefore, we interrogated a comparative injury RNAseq dataset for differential expression of T cell associated transcripts between *Mus* and *Acomys* [40]. While expression for genes associated with non-lymphocyte immune cell populations were generally similar between species, several transcripts associated with T cells and natural killer cells were increased in *Acomys* and decreased in *Mus* in response to injury (Figure 4J). Increased expression of *Cd8*, *Ctla4*, *Il2ra*, *Foxp3*, and *Tnfrsf4* specifically suggested an activated cytotoxic and regulatory T cell response during regeneration but not fibrotic repair (Figure 4J). During fibrotic repair, *Cd4* was differentially increased at D5 and D10 suggesting the presence of CD4 helper T cells not present during regeneration (Figure 4J). Together, these data demonstrate that regeneration was associated with a proportionally greater influx of CD3+ cells that accumulate quickly at the injury site and that specific subtypes of activated T cells were preferentially associated with regeneration.

STAT3 is activated independently from IL-6 during blastema formation

We also sought to test our observation that strong induction of the pro-inflammatory cytokine IL-6 was associated with the acute inflammatory phase of fibrotic repair. To do this, we assayed for IL-6 signaling using STAT3 phosphorylation (Figure 5A-F). STAT3 is phosphorylated in response to the ligand IL-6 binding its membrane receptor, which activates signal transduction in target cells [50]. Corroborating our ELISA quantification for IL-6 in the tissue microenvironment, we found that pSTAT3 increased 8-fold in response to injury in Mm-UKY during the acute inflammatory phase (Figure 5A, B). Similarly, during fibrosis when IL-6 concentrations resolved in Mm-UKY, pSTAT3 began to decline toward baseline (Figure 5A, B). In *Acomys*, pSTAT3 was significantly elevated at D1, although to a lesser extent than compared

to Mm-UKY (Figure 5A, B). Moreover, during blastema formation (D10-15) when our ELISA data showed increased IL-6 compared to baseline in Ac (Figure 3), analysis of pSTAT3 showed further induction of pSTAT3 in Ac (Figure 5A, B).

To determine the cellular localization of STAT3 phosphorylation, we assayed for pSTAT3 using immunohistochemistry during the acute inflammatory phase (D2) and new tissue formation (D15) (Figure 5C-F). Supporting the immunoblot data, both species showed extensive nuclear staining for pSTAT3 at D2 in the epidermis and mesenchymal compartments (Figure 5C', D'). Positive staining in the epidermis > 200 μ M proximal to the amputation plane suggested STAT3 activation was a pervasive response to injury within the ear pinna in both species (Figure 5C, D). Supporting the 2-fold difference in pSTAT3 we observed between Mm-UKY and Ac (Figure 5A, B), we found that nearly every epidermal cell in Mm-UKY appeared positive for pSTAT3 whereas less than half of the epidermal cells were positive in Ac (Figure 5C', D'). The internal tissue compartments (e.g., dermis, cartilage, muscle and adipose) at D2 were similar between species with approximately half of the total cells positive for pSTAT3. At D15, only a few pSTAT3 positive cells were present in Mm-UKY and they were isolated to the epidermis distal to the amputation plane (Figure 5E, E'). In contrast, pSTAT3 positive cells were widespread throughout the blastema in Ac (Figure 5F, F'). Together, these data support stronger IL-6 mediated STAT3 activation in Mm-UKY compared to Ac during the acute inflammatory phase and increased STAT3 activation during blastema formation.

Greater increases in IL-6 and CXCL1 during the acute inflammatory phase of fibrotic repair in *M. musculus* suggested that these molecules might antagonize a potential regenerative response. Previous studies have shown that a balance in these molecules regulate wound healing as IL-6 and CXCL1 are potent pro-inflammatory molecules and hyper-elevated concentrations

after injury are attributed to aberrant healing and chronic inflammation [51-53]. Additionally, genetic ablation of *IL-6*, the *IL-6 receptor*, or the *CXCL1 receptor (CXCR2)*, causes severely delayed re-epithelialization, scab formation and abhorrent wound healing in cutaneous and incisional wounds [54-57]. *IL-6* signaling activates several downstream mediators of inflammation including cyclooxygenase-2 (*COX-2*), and its enzymatic products can amplify the inflammatory response [58]. To test if *COX-2* activity promotes fibrosis in the ear pinna, we used our ear punch assay in Mm-UKY treated healing tissue with Celecoxib, a specific and potent *COX-2* inhibitor [59] (Figure 6A). Comparing ear-hole closure between celecoxib- and vehicle-treated animals there was no support for a difference in the rate of closure (D5 through D30) (repeated measures two-way ANOVA, $n=31$; treatment: $Df=1$, $F=2.42$, $P=0.137$; day: $Df=5$, $F=179.62$, $P<0.001$; treatment*day: $Df=5$, $F=0.36$, $P=0.875$) (Figure 6B). Similarly, comparison of ear-hole area at D64 showed no support for a difference between treatment and control ears (unpaired t-test; $t=0.671$, $P=0.512$) (Figure 6C). One out of five celecoxib-treated animals had not completed re-epithelialization by D10 (Figure 6D-F). Lastly, while the intensity of Picrosirius stain appeared to be lower in celecoxib treated animals compared to controls, there was no difference in the area of collagen deposition at D64 (unpaired t-test, $t=0.104$, $P=0.918$) (Figure 6G-I). Thus, these data support that systemic inhibition of *COX-2* activity is not sufficient to reduce fibrosis or induce a regenerative response after ear pinna injury in *M. musculus*.

DISCUSSION

In this study, we performed a comprehensive cytokine characterization of the immune response to injury where identical injuries in closely-related species undergo two different healing responses: regeneration or fibrotic repair. Importantly, our experimental design leveraged a comparison of animals with an activated and naïve immune system in order to identify species-specific cytokine changes that were associated with regeneration and not due to an environment-immunity interaction. Our analyses showed that regardless of healing outcome, injury induced a common set of pro-inflammatory factors (IL-6, and TNF α) and chemokines (CCL3, CSF2 and CXCL1) during the acute inflammatory phase of fibrosis and regeneration. Additionally, we observed similar responses between healing outcomes for IL-2, IL-4 and IL-5 in the tissue microenvironment. While this supports that fibrotic repair and regeneration share inflammation during the healing process, we also found significantly greater responses for IL-6, CCL2 and CXCL1 during fibrotic repair compared to regeneration. In contrast, regeneration was uniquely associated with local increases in IL-12 and IL-17. Supporting our cytokine analysis, we found that regeneration was associated with a strong influx of T cells during acute inflammation compared to fibrotic repair and that regeneration-competent T cells were closely associated with the dermis during blastema formation. This latter point suggests that T cells may influence local inflammation and contribute to the local regulation of tissue morphogenesis in spiny mice.

Recent studies comparing immune profiles between laboratory-reared and pet-store or wild-caught *M. musculus* demonstrate that non-laboratory strains have more CD44⁺ effector T cells, memory T cells and circulating neutrophils [60, 61]. Although neither group directly measured serum cytokines from the different populations, the elevated baseline concentrations of

IL-4, IL-6, CCL2 and TNF α that we measured in circulation from immune-challenged animals support larger active populations of effector and memory T cells. These data support that our immune-challenged group have been exposed to more pathogens than the laboratory-reared mice which is undoubtedly the case. In addition to the increased baseline concentrations of these cytokines, we also found significant differences in the response to injury for IL-1 α , IL-2 and TNF α between animals with an activated or naïve immune system. Studying wild-caught populations enabled us to identify responses that accurately reflected phenotypic differences between species, rather than differences that could be explained by immune status. Of particular importance was our inclusion of wild-caught *A. percivali* that indicate increases in TNF α , CCL2 and CXCL1 are not inhibitory to regeneration. Additionally, we observed high variation in cytokine concentrations across our dataset, indicating that the immune response to injury could be confounded by individual variation. In other words, researchers should not expect to identify clear transition phases based on time after injury with small sample sizes. Ultimately these data support that injury elicits a local cytokine response that is dependent of baseline immune status with respect to release of cytokines that in turn effects the timing of events but does not change healing outcome.

Acute inflammation is a necessary component of the innate immune response designed to fight invading microbes by recruiting leukocytes from circulation and activating local myeloid and lymphoid cells. Our analyses demonstrate that injury induces an acute inflammatory response regardless of healing outcome that resolves within ~10 days; a timeframe in line with human and rodent wound healing studies [7, 62]. In particular, we found the local release of CCL3, CSF2 and CXCL1 in all groups, which are known to be potent chemokines for monocytes, macrophages and neutrophils. Moreover, the local release of IL-6 and TNF α

supports the presence of activated macrophages and neutrophils as a common injury response. Supporting our previous work that neutrophils infiltrate injured spiny mouse tissue slower compared to laboratory mice [15], regeneration was associated with delayed release and a reduced maximal fold change in IL-6 and CXCL1 compared to fibrotic repair. Additionally, we did not detect a CCL2 response during regeneration, and IL-6, CXCL1 and CCL2 are known to positively regulate the speed of re-epithelialization [54-56], which we find to be delayed at most five days in *A. cahirinus* compared to *M. musculus* [40]. Thus, while acute inflammation is a component of regeneration and fibrotic repair, the cellular differences are likely attributed to reduced pro-inflammatory cytokines released into the microenvironment.

Corroborating our observation that the IL-6 response was weaker during regeneration compared to fibrotic repair, we found diminished activation of STAT3 during acute inflammation (D1-10) in spiny mouse epidermis compared to mouse. Interestingly, we observed an increase in pSTAT3 during blastema formation, whereas pSTAT3 levels declined during fibrotic repair. Furthermore, during tissue morphogenesis at D15 many blastemal cells were STAT3 positive. Given that IL-6 concentrations did not appreciably increase during blastema formation or tissue morphogenesis the increase in STAT3 activity is likely independent of IL-6. STAT3 is activated through multiple pathways (e.g. leukemia inhibitory factor, epidermal growth factor, palette derived growth factor, IL-10, IL-17, etc.). Although IL-17 increased in *A. percivali* after D10, it did not increase in *A. cahirinus* suggesting it is not responsible for the late phase of STAT3 phosphorylation. Given that STAT3 signaling is multifaceted, one potential biological link is that STAT3 activity is necessary for satellite-cell activation and axon regeneration in mammals [63-65]. Interestingly, the expression of *Sal4*—a factor necessary for blastema maintenance in *Xenopus* and *Ambystoma*—is regulated by pSTAT3 [66-68]. While

Sal4 does not have a mammalian homolog, this data supports that activation of STAT3 in regenerating tissue is an evolutionary conserved mechanism and interrogating unique STAT3 targets in spiny mice may uncover mechanisms that regulate blastema formation in mammals.

Inhibition of downstream signaling induced by IL-6 / CXCL1, such as arachidonic acid metabolism by COX-2, has been shown to reduce fibrosis post epidermal injury (e.g. incisional, cutaneous and chronic pressure wounds) [69-71]. Celecoxib treatment to inhibit COX-2 in the present study may have slowed re-epithelialization. Additionally, while the total area of fibrosis was not different between celecoxib- and vehicle-treated animals there appeared to be a small reduction in the total amount of collagen produced in celecoxib-treated animals from reduced intensity of picrosirius staining. However, similar to previous reports, reduction in COX-2 activity did not induce regeneration, supporting that inflammation is not the ultimate inhibitory barrier.

In addition to the magnitude increase in IL-6 and CXCL1, our analyses found that increased local CCL2 was specific to fibrotic repair. CCL2 was first identified as a monocyte-specific chemoattractant to sites of injury and infection, and activates macrophages [72, 73]. CCL2 also attracts neutrophils and supports neutrophil-dependent tissue damage [72]. As such, the amount of CCL2 that is released into an injury microenvironment regulates the healing response and studies support there is a positive relationship between CCL2 and the amount of fibrosis during fibrotic repair [74-76]. However, a careful balance must be maintained as CCL2 knockout mice do not heal wounds [77]. Thus, it is possible that the reduced IL-6, CCL2 and CXCL1 responses are responsible for reduced fibrosis in spiny mice. Although these key factors appear to interact in the hierarchy of the progression of fibrotic repair, the paracrine mechanism

of how they would activate dermal fibroblasts remains unknown. It is likely another cell-type, such as a macrophage or T cell, is mediating the signal.

In addition to this study, two studies have quantified cytokines during regeneration—one in axolotl limbs [12] and the other in spiny mouse dorsal skin wounds [34]. Godwin et al. (2013) used a mouse cytokine array to analyze regenerating salamander limbs and found that all but two cytokines detected reached peak amounts within 48hrs of injury and that every cytokine returned to baseline by D15 after a blastema had formed. Brant et al. (2016) used the same cytokine array to assess cytokines during the first 14 days of spiny mouse dorsal skin regeneration and observed a similar phenomenon and all detected cytokines resolve to baseline by D14. Despite study-specific differences in the ability to detect antigens and a lack of parallelism validation, these studies support that release of CCL3 and TNF α in tandem with a differential inflammatory response occurs prior to tissue regeneration. However, our comparative analyses also suggested that the magnitude of the increase in IL-6 and CXCL1 might serve as early indicators of a fibrotic repair trajectory. For example, the IL-6 response to injury, although present, was small and CXCL1 did not respond during both axolotl limb and spiny mouse skin regeneration.

Finally, our cellular analysis uncovered a surprisingly rapid adaptive immune response measured as an early influx of T cells in regenerating compared to non-regenerating species. Importantly, our findings support that the arrival of T cells in spiny mice is concurrent with the arrival and proliferation of monocytes [15], which suggests there is a regenerative-competent T cell response that is different from a fibrotic T cell response. Contrary to hypotheses suggesting that a strong adaptive immune response reduces regenerative ability [78, 79], our findings suggest T cells may positively regulate regeneration in spiny mice. Similar to macrophages, T cells can differentiate into a number of functional subpopulations which differentially affect cells

in the inflammatory microenvironment [80]. Our analysis of the transcriptional response to injury between *M. musculus* and *A. cahirinus* suggests that fibrotic repair is associated with an accumulation of inactivated T_H cells, while during regeneration there is an accumulation of activated cytotoxic T and regulatory T (T_{REG}) cells. Studies have shown that loss of cytotoxic CD8⁺ T cells inhibits skeletal muscle regeneration, accelerates bone fracture healing, and increases fibrosis in incisional wounds [81-87]. Additionally, recent work showed that T_{REG} populations infiltrate injured muscle quickly after injury and are necessary to regulate the ratio of MHC-class II positive and negative macrophages present in the injured tissue. When the T_{REG} population is ablated subsequent regeneration is impaired [17]. Moreover, spiny mice have a greater NADPH oxidase induced ROS response [15], which can be partially controlled by T_{REG} cells [88-90]. Thus, these studies support an anti-fibrotic role for T cells and suggest the action of cytotoxic T and T_{REG} cells may have a positive role during spiny mouse epimorphic regeneration.

Together, the data presented here support that tissue regeneration in *Acomys* occurs in cooperation with an adaptive immune response and that lymphocyte phenotype might play a key role in facilitating a regenerative or fibrotic response. Ongoing studies in our laboratory are aimed at characterizing the macrophage and T cell populations that are associated with the injured tissue during regeneration and fibrotic repair. These future datasets (and the present one) will create a framework to begin testing how the immune response functions during complex tissue regeneration in a mammalian model. We believe that modulating the immune response at the injury microenvironment will be an essential piece to inducing epimorphic regeneration in tissues that naturally heal by fibrotic repair.

AUTHOR CONTRIBUTIONS

Conceptualization, T.R.G., V.O.E. S.G.K. and A.W.S.; Methodology, T.R.G., J.S., and A.W.S.; Validation, T.R.G. and J.S.; Investigation, T.R.G, J.S., J.M.K., C.K.H., V.O.E. and A.W.S.; Resources, S.G.K., V.O.E. and A.W.S.; Writing – Original Draft, T.R.G.; Writing – Review & Editing, T.R.G., J.S., V.O.E. and A.W.S.; Visualization, T.R.G. and A.W.S.; Supervision, T.R.G., S.G.K., V.O.E. and A.W.S., Project Administration, T.R.G., V.O.E. and A.W.S.; Funding Acquisition, S.G.K., V.O.E. and A.W.S.

ACKNOWLEDGEMENTS

We thank Adam Cook, Chanung Wang, Bailey Gensheimer, Malik Guidry, John Ewoi and Stanley Marete for help with live-trapping, animal care and data collection. We also thank Peter Jessel and the Tomlinson family for kindly allowing us continued access to their property in Kenya. We acknowledge David Higginbotham, Jim Monegue and rest of the swine unit for their help and enthusiasm with trapping the wild *Mus musculus*. Funding for this work was provided by the National Science Foundation (NSF) and the Office for International Science and Engineering (OISE) to A.W.S. (IOS-1353713) and V.O.E (IOS-1353857). The University of Kentucky provided additional funding to A.W.S.

DECLARATION OF INTERESTS

The authors declare no competing interests.

492 FIGURE LEGENDS

493 **Figure 1 Ability to regenerate and immunity-status are associated with specific systemic**
 494 **immune profiles and immune responses to injury.** (A) Comparison of cytokine
 495 concentrations in serum from uninjured animals showed higher concentrations of IL-4, IL-6,
 496 CCL2 and TNF α in wild-caught animals compared to laboratory-reared animals indicated that
 497 wild animals have a “primed” immune system. No difference was found between non-
 498 regenerators (Mm-UKY, Mm-Kenya and Mm-Wild) and regenerators (Ac and Ap). Data
 499 represent box and whiskers with median, interquartile range and individual data points. N/A
 500 denotes concentrations could not be quantified in any animal of the group. The dashed line in
 501 each graph represents the lower limit of detection for the specific cytokine. N/S denotes $P>0.05$
 502 for One-way ANOVA (See Supplemental Table 1) and different letters above the data denotes
 503 $P\leq 0.05$ for Tukey-Kramer pairwise comparisons (See Supplemental File). (B) The change in
 504 systemic cytokines compared to D0 over 20 days after ear hole punch. A naïve immune system
 505 response (solid lines) was consistent with increased IL-2 and TNF α , and decreased IL-1 α
 506 compared to a primed response (intermittent lines) (See Supplemental Table 2). Fibrotic repair
 507 (black) was associated with increases in IL-5, IL-6, IL-17, CCL3 and CXCL1 compared to
 508 regeneration (red). Data represent mean and S.E.M. for at least $n=5$ animals per species per
 509 timepoint. The dashed line at $Y=1$ represents no change compared to D0 and the yellow boxes
 510 represent the inflammation resolution window.

511 **Figure 2 The tissue microenvironment is dynamic, and the ability to regenerate is**
 512 **associated with a specific immune response to injury.** The comparison of the change in
 513 cytokines from tissue lysate compared to D0 over 20 days after ear hole punch. All cytokines
 514 varied over time and between groups (See Supplemental Table 3). There was a stronger increase
 515 in CCL3 a muted increase for IL-12 and CXCL1 in immune-naïve (solid lines) compared to
 516 immune-primed animals (intermittent lines). Non-regenerators (black) had stronger increases for
 517 IL-6, CCL2 and CXCL1 compared to regenerators (red). Additionally, IL-17 decreased in non-
 518 regenerators and increased in regenerators. Data represent mean and S.E.M. for at least $n=5$
 519 animals per species per timepoint. The dashed line at $Y=1$ represents no change compared to D0
 520 and the yellow boxes represent the inflammation resolution window.

521 **Figure 3 Regeneration is associated with specific temporal cytokine responses in the tissue**
 522 **microenvironment.** The comparison of the change in cytokines from tissue lysate compared to
 523 baseline. Non-regenerators (Mm-UKY, Mm-Kenya, and Mm-Wild: Black) show a stronger
 524 response compared to regenerators (Ac and Ap: Red) for several inflammatory cytokines during
 525 the immediate response, acute inflammatory phase. These differences mostly resolved at day 10
 526 and 20. In contrast, regenerators showed a stronger IL-12 and IL-17 response during tissue
 527 morphogenesis. Data represent box and whiskers with median, interquartile range and individual
 528 data points. N/A denotes concentrations could not be quantified. The dashed line at $Y=1$
 529 represents no change compared to D0. Each graph showed $P<0.05$ for effect of group using a
 530 One-way ANOVA on log-transformed data (See Supplemental Table 4) and different letters
 531 above each group denotes $P<0.05$ for Tukey-Kramer pairwise comparisons (See Supplemental
 532 File).

Figure 4 The regeneration microenvironment is primed by greater T cell influx and activated T_{REG} signature. (A) Comparison of total CD3⁺ cells quantified by flow cytometry from disassociated ear pinna and (B) the ratio relative to uninjured tissue for *M. musculus* (black) and *A. cahirinus* (red). Data represent mean and S.E.M. and n=4 or 5. An * denotes $P < 0.05$ for pairwise comparison within the day between species for Tukey-Kramer HSD posthoc test. (C-I) Representative immunohistochemistry for CD3 (red) counterstained with DAPI (gray) at the proximal wound margin (amputation plane can be determined from the end of the cartilage—indicated by the dotted line) from D5 and D15 after injury of *M. musculus* (C, D) and *A. cahirinus* (E, F). More T cells (yellow arrowhead) were present throughout the wound bed and were mainly found in the dermis of *A. cahirinus* compared to *M. musculus*. The T cells associated with epidermis (boundaries indicated by the dotted line) tended to be spindle-like in *M. musculus* (G), while rounded in *A. cahirinus* (H). The dermal T cells in *A. cahirinus* also tended to be in close proximity to regenerating epidermal appendages (I). N=4 and bar equals 200 μ m (C-F) or 20 μ m (G-I). (J) Heatmap of differential gene expression compared to uninjured tissue suggests that the regeneration microenvironment contains a substantial NK, CD8⁺ and T_{REG} cell response while fibrotic repair has a CD4⁺ cell response. Data comes from a previously published analysis [40].

Figure 5 Time-dependent STAT3 activation in the blastema is associated with regeneration. (A-B) Comparison of representative immunoblots for pSTAT3 and ACTB for indicated time and species from injured tissue homogenate from ear-hole punch assay (A). Fibrotic repair is associated with strong early STAT3 activation while regeneration is associated with a weak early and strong postponed activation (B). Data represent mean fold change of pSTAT3 intensity normalized to ACTB intensity and relative to D0 and S.E.M. n=4 individuals per time point and species (B). (C-F) Representative images for immunohistochemistry for pSTAT3 (Brown) counterstained with hematoxylin (Blue) for *M. musculus* (C, C', E and E') and *A. cahirinus* (D, D', F and F') at the indicated time points. On D2 after injury, nearly every epidermal cell in *M. musculus* was positive compared to about half in *A. cahirinus* (C, D). On D15 after injury, only a small population of epidermal cells in *M. musculus* were positive compared to about half in *A. cahirinus* (E, F). Additionally, about half of the cells in the forming blastema remain positive at D15. Data represent n=3, and bar equals 200 μ m.

Figure 6 COX-2 inhibition does not liberate regeneration in *M. musculus*. (A) Celecoxib treatment during the first 10 days of injury reduces secondary inflammation caused by IL-6 and CXCL1. (B-C) Celecoxib treatment did not affect the rate of ear-hole closure from D5 to D30 (B), or ear-hole area at D64 (C). (D-F) Mason's trichrome stain on D10 ear tissue from control (D) and treatment (E) showed 20% of the distal injury plane from treated ears were not re-epithelialized, while 100% of control treated ears and the proximal injury plane were re-epithelialized (F). (G-I) Picrosirius stain on D64 ear tissue from control (G) and treatment (H) showed no difference in the total collagen fibrotic area distal to the amputation plane (I). Data represent mean $\pm$ S.E.M. and the lines represent cubic regression for n=10 per treatment (B), OR individual data (red dots) and median and interquartile range for n=10 per treatment (C, I).

TABLES

Table 1: Comparison of cytokine slopes from parallelism test of cytokine assay. This was used to determine which species and source comparisons could be made for each cytokine in the Comparison column.

Antigen	Standard	Sample	<i>Mus musculus (Mm)</i>	<i>Acomys cahirinus (Ac)</i>	<i>Acomys percivali (Ap)</i>	Comparison
IL-1 α	-0.89 $\pm$ 0.03	Serum: Tissue:	-1.05 $\pm$ 0.03* <i>Too few points</i> [^]	-0.82 $\pm$ 0.14* -1.03 $\pm$ 0.04*	-1.37 $\pm$ 0.10* -0.97 $\pm$ 0.18*	Mm, Ac, Ap Mm, Ac, Ap
IL-1 β	-0.93 $\pm$ 0.10	Serum: Tissue:	<i>Too few points</i> ^{&} -0.94 $\pm$ 0.02*	<i>Not detected</i> -0.86 $\pm$ 0.03*	-1.63 $\pm$ 0.21* -0.65 $\pm$ 0.07*	Mm, Ap Mm, Ac, Ap
IL-2	-1.34 $\pm$ 0.05	Serum: Tissue:	<i>Too few points</i> ^{&} -1.21 $\pm$ 0.10*	<i>Not detected</i> -0.59 $\pm$ 0.22*	-0.87 $\pm$ 0.00* -0.57 $\pm$ 0.20*	Mm, Ap Mm, Ac, Ap
IL-4	-0.89 $\pm$ 0.06	Serum: Tissue:	-0.88 $\pm$ 0.18* -0.69 $\pm$ 0.09*	<i>Too few points</i> ^{&} -0.68 $\pm$ 0.39*	-2.16 $\pm$ 0.32* -0.75 $\pm$ 0.12*	Mm, Ap Mm, Ac, Ap
IL-5	-1.13 $\pm$ 0.01	Serum: Tissue:	-1.63 $\pm$ 0.20* -0.73 $\pm$ 0.05*	-1.53 $\pm$ 0.96* <i>Too few points</i> ^{&}	-2.52 $\pm$ 0.28* -0.87 $\pm$ 0.03*	Mm, Ac, Ap Mm, Ap
IL-6	-1.07 $\pm$ 0.02	Serum: Tissue:	-0.74 $\pm$ 0.12* -0.80 $\pm$ 0.14*	<i>Too few points</i> ^{&} -0.47 $\pm$ 0.06*	-1.81 $\pm$ 0.11* -0.63 $\pm$ 0.11*	Mm, Ap Mm, Ac, Ap
IL-10	-1.56 $\pm$ 0.03	Serum: Tissue:	<i>Not detected</i> <i>Not detected</i>	<i>Not detected</i> 0.52 $\pm$ 0.25	<i>Not detected</i> 0.24 $\pm$ 0.52	None None
IL-12	-1.10 $\pm$ 0.02	Serum: Tissue:	-0.68 $\pm$ 0.16* -0.73 $\pm$ 0.17*	-1.55 $\pm$ 0.49* -0.68 $\pm$ 0.16*	-1.26 $\pm$ 0.18* -0.63 $\pm$ 0.15*	Mm, Ac, Ap Mm, Ac, Ap
IL-17	-1.03 $\pm$ 0.02	Serum: Tissue:	-0.72 $\pm$ 0.25* -1.28 $\pm$ 0.04*	-0.16 $\pm$ 0.23 -0.85 $\pm$ 0.34*	-0.49 $\pm$ 0.16 -0.81 $\pm$ 0.03*	Mm Mm, Ac, Ap
CSF2	-1.45 $\pm$ 0.05	Serum: Tissue:	<i>Not detected</i> -0.71 $\pm$ 0.11*	<i>Not detected</i> -0.68 $\pm$ 0.25*	<i>Too few points</i> ^{&} -0.70 $\pm$ 0.10*	None Mm, Ac, Ap
CCL2	-1.17 $\pm$ 0.00	Serum: Tissue:	-1.23 $\pm$ 0.14* -0.97 $\pm$ 0.02*	<i>Too few points</i> ^{&} -1.23 $\pm$ 0.14*	-1.56 $\pm$ 0.13* -0.63 $\pm$ 0.10*	Mm, Ap Mm, Ac, Ap
CCL3	-0.98 $\pm$ 0.06	Serum: Tissue:	-2.05 $\pm$ 0.03* -0.93 $\pm$ 0.01*	-1.25 $\pm$ 0.09* -0.95 $\pm$ 0.01*	-1.01 $\pm$ 0.09* -0.94 $\pm$ 0.06*	Mm, Ac, Ap Mm, Ac, Ap
CCL5	-0.90 $\pm$ 0.07	Serum: Tissue:	-0.78 $\pm$ 0.03* -0.90 $\pm$ 0.16*	<i>Not detected</i> 0.34 $\pm$ 0.13	0.33 $\pm$ 0.32 0.07 $\pm$ 0.27	Mm Mm
TNF α	-1.10 $\pm$ 0.06	Serum: Tissue:	-1.36 $\pm$ 0.38* -1.14 $\pm$ 0.10*	-2.15 $\pm$ 0.01* -0.93 $\pm$ 0.01*	-1.09 $\pm$ 0.17* -1.21 $\pm$ 0.03*	Mm, Ac, Ap Mm, Ac, Ap
IFN γ	-0.79 $\pm$ 0.10	Serum: Tissue:	-0.60 $\pm$ 0.16* -0.81 $\pm$ 0.31*	-1.48 $\pm$ 0.72* -0.72 $\pm$ 0.05*	-1.21 $\pm$ 0.22* -0.84 $\pm$ 0.05*	Mm, Ac, Ap Mm, Ac, Ap
CXCL1	-1.20 $\pm$ 0.02	Serum: Tissue:	-1.15 $\pm$ 0.27* -1.11 $\pm$ 0.05*	<i>Not detected</i> <i>Too few points</i> ^{&}	<i>Not detected</i> -1.15 $\pm$ 0.29*	Mm Mm, Ap

Slopes in *italics* are not different from zero ($P > 0.05$ for regression test)

* denotes slope is similar to the standard and can be reliably quantified

& denotes values are below the assay's lower limit of detection

[^] denotes values are above the assay's upper limit of quantification

586 **Table 2:** Comparison of *A. cahirinus* predicted peptide sequences used in cytokine analysis

Gene ID	Peptide ID	% similar to <i>M. musculus</i>	% similar to <i>R. norvegicus</i>	% similar to <i>H. sapiens</i>
<i>Il1a</i>	IL-1 α	82.90	86.90	66.90
<i>Il1b</i>	IL-1 β	87.71	87.71	72.35
<i>Il2</i>	IL-2	70.41	84.61	71.59
<i>Il4</i>	IL-4	66.45	77.21	56.32
<i>Il5</i>	IL-5	90.15	84.84	68.93
<i>Il6</i>	IL-6	75.23	78.97	48.13
<i>Il10</i>	IL-10	85.05	87.93	74.13
<i>Il12a</i>	IL-12 p35	70.42	80.54	55.25
<i>Il12b</i>	IL-12 p40	73.58	74.10	63.33
<i>Il17a</i>	IL-17	84.88	83.13	75.58
<i>Cxcl1</i>	KC	80.37	81.30	69.15
<i>Ccl2</i>	MCP-1	73.50	70.86	43.70
<i>Ccl3</i>	MIP-1 α	93.54	94.62	81.72
<i>Ccl5</i>	RANTES	88.04	84.78	85.86
<i>Csf2</i>	GM-CSF	65.24	73.75	61.70
<i>IFNγ</i>	IFN γ	56.77	57.05	41.56
<i>Tnf</i>	TNF α	95.81	96.65	84.93

587

MATERIALS AND METHODS

Animals

Mus musculus (Mm-UKY) and *Acomys cahirinus* (Ac) were maintained at our animal facility at the University of Kentucky, Lexington, KY, USA. Mm-UKY were sexually mature (10- to 12-week old), female, outbred Swiss Webster (ND4, Envigo, Indianapolis, IN). They were housed at a density of 2-4 individuals in static IVC cages with pine shavings and given autoclaved water and 18% protein mouse chow (Tekland Global 2918, Envigo). Ac were sexually mature (12- to 28-weeks old), males and females and were housed at a density of 10-15 individuals in large metal wire cages (24 inch x 12 inch x 14 inch, height x width x depth, Quality Cage Company, Portland, OR) with pelleted pine bedding (Southern States Cooperative, Inc., Richmond, VA) and given autoclaved water and a 3:1 mixture by volume of 14% protein mouse chow (Teklad Global 2014, Envigo) and black-oil sunflower seeds (Pennington Seed Inc., Madison, GA) [91]. Additionally, the air within facility was filtered, and the animals were exposed to natural light through large windows (approximately 12h:12h L:D light cycle during the experiment). All Mm-UKY and Ac samples were collected between 9/20/2015 and 10/28/2015.

Wild *M. musculus* (Mm-Wild) were live trapped at the C. Oran Little Research Center in Versailles, KY (38°4'N, 84°44'W) and maintained in an alternate animal facility at the University of Kentucky. Mm-Wild were housed at a density of 10-12 individuals in large metal wire cages with pelleted pine bedding and given autoclaved water and 18% protein mouse chow. The animals acclimated to captivity for at least twenty-one days before any experiments were started. The air within the facility was filtered and the animals were exposed to a 12:12h L:D

cycle by fluorescent lights. All Mm-Wild samples were collected between 4/1/2017 and 6/21/2017 and 12/12/2017 and 3/6/2018.

The Kenyan *M. musculus* (Mm-Kenya) were sexually mature (10- to 12-week old), female, outbred Swiss Webster mice obtained from a local breeder in Nairobi, Kenya and maintained in an animal facility at the University of Nairobi, Kenya. Sexually mature *Acomys percivali* (Ap) were live-captured at Mpala Research Centre in Laikipia, Kenya (0°17'N, 37°52'E), and transported to the University of Nairobi for study. Each species was separated by sex and housed at a density of 10-15 animals in large metal wire cages, given tap water, fed mouse pencils (Argroside Inc., Nairobi, Kenya) 1x per day and exposed to natural light through windows (equivalent 12h:12h L:D cycle). The animals acclimated to captivity for at least twenty-one days before any experiments were started. Additionally, the facility was open to the natural environment (i.e., the mice were exposed to Nairobi air) and cool nighttime temperatures were supplemented with ceramic heaters. Mm-Kenya samples were collected between 6/04/2015 and 7/04/2015 and Ap samples were collected between 5/04/2015 and 7/04/2015, and between 5/02/2016 and 6/04/2016.

All animal trapping and procedures were approved by the University of Kentucky Institutional Animal Care and Use Committee (IACUC) under protocol 2013-1119, Kenyan Wildlife Service (KWS), and the University of Nairobi Faculty of Veterinary Medicine Animal Care and Use Committee (FVM ACUC). Research in Kenya was approved by the Kenyan National Council for Science and Technology (NACOSTI). All wild species trapped were species of least concern.

Sample collection and preparation

We used a 4 mm biopsy punch to create a hole through the ear pinna, as previously described [40]. Healing ear tissue was collected on D0, 1, 2, 3, 5, 10, 15 and 20. To minimize circadian effects, animals were injured between 10:00 and 12:00, and samples were collected between 11:00 and 15:00. Animals were deeply anesthetized with 5% (v/v) isoflurane and a maximal amount of blood was collected by cardiac puncture using a 25-gauge needle. An 8 mm biopsy punch was used to harvest healing ear tissue.

To isolate serum, blood was collected into a serum separator tube (#454243, Greiner bio-one, Kremsmünster, Austria) and allowed to clot for at least 45 minutes, followed by centrifuging at 3,000 x g for 10 minutes. Serum was aliquoted and stored at -80°C or on dry ice until analysis. The tissue was used for two downstream assays, histology and cytokine quantification. For histology, one of the 8 mm biopsies was placed into 10% (v/v) neutral buffered formalin (American MasterTech, McKinney, TX) overnight, dehydrated, embedded in paraffin and cut to 5 µm thickness on a rotary microtome. For cytokine quantification, a ring of tissue closest to the injury approximately 1 mm wide was snap frozen in liquid nitrogen or a slurry of dry ice and ethanol, and then stored at -80°C or on dry ice. Next, the tissue was homogenized in RIPA buffer supplemented with protease and phosphatase inhibitors (#24948, Santa Cruz Biotechnology, Inc. Dallas, TX; #78427, Thermo Scientific) using ceramic beads (Matrix D, MP Biomedicals, LLC, Solon, OH) and a bead mill for 5 minutes (Next Advance, Inc., Troy, NY), centrifuged at 10,000 x g for 15 minutes to pellet insoluble protein, and the soluble protein was separated into a new tube. The total protein was quantified by bicinchoninic acid assay (#23225, Thermo Scientific) with a standard curve created from the same stock of bovine serum albumin, and then the protein lysate was stored at -80°C or on dry ice until analysis.

Cytokine assay

To assess the immune response to injury in multiple species, we evaluated methods that: 1) used minimal sample, 2) measured local (tissue lysate) and systemic (serum) samples, 3) measured several cytokines at once, 4) differentiated the magnitude and type of immune response during an ear punch assay, and 5) exhibited cross-reactivity among the study species. We used a custom-designed, multiplexed, sandwich ELISA array (Quansys Biosciences, Logan UT). This platform meets the above requirements and the experiments can be performed in multiple locations (i.e., Kentucky and Kenya) because the imager and reagents can be easily transported. Importantly, the imager does not require specialized calibration after being moved, and the reagents do not need remain frozen. The custom assay was designed to measure 16 antigens including IL-1 α , IL-1 β , IL-2, IL-4, IL-5, IL-6, IL-10, IL-12p70, IL-17, CCL2, CCL3, CCL5, CSF2, IFN γ , TNF α and CXCL1.

Initial testing identified that antigens from serum could be quantified by diluting serum in the supplied mouse specific diluent 1:1 and from tissue lysate using 5, 40, and 80 μ g total protein in RIPA buffer for *M. musculus*, *A. cahirinus*, and *A. percivali*, respectively. The samples were run in duplicate using a protocol modified from the manufacturer's instructions, as follows: All serum samples were diluted 1:1 (serum : diluent) and all tissue samples were diluted 1:1 (RIPA + lysate : mouse sample diluent) to a volume of 50 μ L per well. The diluted samples were then loaded onto a new assay plate with an appropriate standard curve (1:3 to 1:59049) and four blanks. Samples were incubated at 4°C for 8 hours on a plate shaker set to 500 rpm to capture antigen in each well. After washing the plate 4 times with wash buffer, the primary antibody cocktail was loaded and the plate was incubated at 4°C for 8 hours on a plate shaker set to 500

rpm to allow binding of the biotinylated detection antibodies to the captured antigens. After washing the plate 4 times, streptavidin-HRP conjugated secondary antibody was loaded and the plate was incubated at room temperature for 30 minutes on a plate shaker set to 500 rpm. The plate was washed 8 times, chemiluminescent reagent was added, and the plate was immediately imaged with a chemiluminescent plate imager set to the manufacturer recommended image capture settings (Q-view imager, Quansys Biosciences).

We verified that cytokine concentrations derived from the Quansys multiplex array were comparable between *Mus* and *Acomys* by testing for parallelism of the mouse standards with *Acomys* serum and tissue lysate. We also evaluated the peptide-level similarity between *Mus* and *Acomys* for each gene represented on the array. Parallelism was examined using standard protocols [92, 93]. Briefly, samples from species and source were randomly pooled to provide a representative cytokine concentration and were run in triplicate at serial dilutions (1:2, 1:6, 1:18, 1:54, and 1:162). To determine parallelism, linear regressions were calculated for samples that had at least 3 dilutions above the lower limit of detection and compared the slopes to the standard curve. For peptide comparisons, the *A. cahirinus* genomic and/or transcribed sequences corresponding to the 16 cytokines of interest were identified by using TBLASTX with inputted *M. musculus* peptide sequence into previously published spiny mouse transcriptomes [40, 94] and an unpublished draft genome. The mRNA sequence was then translated and aligned to peptide sequences for *M. musculus*, *Rattus norvegicus* and *Homo sapiens* using MAFFT [95, 96]. Total similarity and identity was calculated using the Sequence Identity and Similarity (SIAS) tool (<http://imed.med.ucm.es/Tools/sias.html>).

Individual cytokine concentrations were obtained using image analysis software (Q-view v3.09, Quansys Biosciences). First, the standard curve pixel intensity values were observed and

pixel intensity values greater than 60,000 were masked to remove saturated data points. Sample concentrations were calculated from standard curves created by a five parameter logistic regression (5PL) with $\sqrt{y}$ weighting. The average value from each duplicate was then used for subsequent analyses. If the average value was above the lower limit of detection and the pixel intensity co-efficient of variation between duplicates was greater than 15%, the sample was re-assayed on another plate and a new average calculated. Initially, we re-assayed tissues samples below the limit of detection with a greater amount of total protein, but in most cases, additional protein did not equate to quantifiable antigen, suggesting that there was a minimal amount of antigen in those samples. Thus, to maximize use of the plates, we opted to quantify a greater total number of samples and assayed each sample at one dilution. Antigens below the lower limit of detection were recorded as “not present”, and to calculate ratios they were assigned the largest value of the lower limit of detection for that antigen across all plates assayed [97].

Cox-2 inhibition

Mm-UKY were subjected to a routine ear punch assay and randomly split into two groups: A) 100 mg/kg celecoxib, a potent and specific COX-2 inhibitor or B) vehicle. A Celecoxib capsule was opened and mixed into 0.5% (w/v) methyl cellulose to the appropriate concentration and a 200 μ L dose was administered (100 mg active drug / kg body weight) by oral gavage using a 20x30 mm gavage needle tipped with a sugar solution each morning beginning 1 day before injury through 20 days after injury. Ear holes were measured and ear hole area was calculated for every 5 days post injury, as previously described [40]. On D10 and 64 entire ears were harvested from a different set of animals and used for histology and stained with Mason’s Trichrome or Picrosirius red, as previously described [40]. Re-epithelialization

was confirmed by the presence of a connected and complete epidermis distal to the amputation plane by examining two tissue sections from the proximal and distal wound sites for each animal at D10. Fibrosis was determined by quantifying the area of collagen deposition in the dermis distal from the amputation plane from two sections from the proximal and distal wound sites using circular polarized light microscopy and the thresholding function in Image J after removing the epidermis, epidermal appendages and tissue artifacts.

Flow Cytometry

To quantify the number of CD3⁺ cells present in healing ear tissue, tissue was harvested from a separate group of Mm-UKY and Ac females at D0, 1, 3, 7 and 15 using an 8 mm biopsy punch. Harvested tissue from both ears was combined and a single-cell suspension was created using combination of enzymatic and mechanical digestion, as previously described [15]. Total cells were counted by hemacytometer and incubated with PE-conjugated-anti-CD3 (Clone 17A2, BioLegend, San Diego, CA) at a concentration of 1 μg / 10^6 cells for 1 hour at room temperature, washed and suspended in cell staining buffer (Cat#420201, BioLegend). Flow cytometry was carried out at the University of Kentucky Flow Cytometry Core using the iCyt Synergy sorter system (Sony Biotechnology Inc., San Jose, CA). Laser calibration and compensation was performed for each experiment using unstained and single fluorescent control samples. Analysis was done using FlowJo (Version 10, FlowJo, LLC, Ashland, OR) to identify CD3-positive lymphocytes by PE fluorescence and forward- and side-scatter. The same gating strategies between species were used (n = 4 or 5 animals per timepoint).

Immunohistochemistry

To identify the locations of STAT3 responsive cells and CD3⁺ cells, tissue sections were de-paraffinized, rehydrated, and prepared for examination by light- or fluorescent-microscopy, respectively. For light-microscopy, resident peroxidase was quenched by H₂O₂, antigens were exposed by heat-mediated retrieval with sodium citrate buffer, pH=6.0, blocked with serum, incubated with primary antibody (anti-pSTAT3, Cell Signaling Technology Cat#9145, 1:200) overnight at 4°C, incubated with a horseradish peroxidase conjugated secondary (Santa Cruz Biotechnology, Cat#sc-2030, 1:1000) for 1 hour at room temp, treated with 3,3'-Diaminobenzidine (SK-4100, Vector Laboratories, Burlingame, CA) until a visible brown precipitate was observed, counter-stained with hematoxylin, dehydrated and cover-slipped. For fluorescent-microscopy, antigens were exposed by heat mediated retrieval with sodium citrate buffer, pH=6.0, resident avidin and biotin was blocked, sections blocked with serum, incubated with primary antibody (anti-CD3, DAKO, Cat#A0452, 1:500) overnight at 4°C, incubated with a biotin conjugated secondary antibody (Vector Laboratories, Cat#PK-6101, 1:400) for 1 hour at room temp, incubated with streptavidin conjugated AlexaFluor-594 (Molecular Probes, Cat#S11227, 1:5,000), counter-stained with 4',6-Diamidino-2-Phenylindole, Dihydrochloride (Molecular Probes, Cat#D1306, 1:10,000) and cover-slipped. Images were acquired using a compound epi-fluorescence microscope (IX-51, Olympus Corporation, Tokyo, Japan) equipped with a CCD camera (DP-74, Olympus Corporation) and software (cellSens v1.12, Olympus Corporation).

Immunoblot

To quantify the STAT3 response to injury, 30 or 40 µg of total protein from tissue lysate was denatured and separated by SDS polyacrylamide gel electrophoresis and transferred to a

PVDF membrane (IB401002, Life Technologies). Membranes were cut along the 55 kDa ladder marker and blocked with either 5% BSA or 5% dry skim milk in TBST for 1 hour at room temperature, incubated with primary antibody (pSTAT3, Cell Signaling, Cat#9145, 1:2000; ACTB, Cell Signaling, Cat#4967, 1:5000), washed, incubated with secondary antibody (Santa Cruz Biotechnologies, Cat#sc-2030, 1:10,000), and visualized by chemiluminescence (Cat#RPN2235, GE Healthcare) using a digital CCD camera (UVP LLC, Upland, CA). Total pixel intensity was quantified using regions of interest and normalized to background and uninjured tissue using ImageJ2 [98].

Statistical Analysis

To compare the baseline serum cytokine concentrations we used a one-way ANOVA and Tukey-Kramer HSD posthoc tests to test for individual group differences. To compare the dynamics of cytokine concentration over time in serum and tissue, a ratio of the injured concentration mean to the uninjured concentration mean was calculated for each cytokine by group (Mm-UKY, Mm-Kenya, Mm-Wild, Ac and Ap) and time point (D1-D20). To normalize cytokine fold-change distributions, data were log transformed. A two-way ANOVA was then used to test for effects of time and group on tissue and serum separately. Pairwise comparisons were tested using the Tukey-Kramer HSD method. In the event that several undetected values existed at an individual timepoint and log transformed data still did not meet normality, we used non-parametric Wilcoxon rank sum tests with Steel Dwass post-hoc tests for pairwise comparisons. Datasets for which non-parametric analyses were performed are indicated in figure legends.

To compare the immunoblot data, pixel intensity was calculated for the bands of interest using an identical sized region of interest with ImageJ [98]. The pixel intensity of pSTAT3 was normalized to ACTB and a two-way ANOVA with time and species was used to compare values and pairwise comparisons were made using the Tukey-Kramer HSD method. To compare the flow cytometry results, we used a two-way ANOVA with time and species on log-transformed data and pairwise comparisons were made using the Tukey-Kramer HSD method. To compare the ear-hole closure rate between control and celecoxib-treated animals we used a repeated-measures ANOVA and cubic regression, as previously published [40]. To compare the ear-hole area and area of tissue positive for Picrosirius, we used a student's t-test. All statistical tests were done using JMP Pro 14 (SAS Institute Inc., Cary, NC) or Prism 5.0 (GraphPad Software, Inc., San Diego, CA). A *P*-value < 0.05 was used to determine significance for each test. All graphs were created in Prism 5.0 and placed into figures using Illustrator CS5 (Adobe Systems, Inc. San Jose, CA).

SUPPLEMENTAL INFORMATION

Supplemental Table 1: One-way ANOVA analyses of log-transformed uninjured serum data. Groups listed in comparison column represent groups for which data was quantified and could be compared. *P*-values indicate where at least one group is significantly different from another group. Tukey-Kramer HSD post-hoc tests were used for pairwise comparisons are summarized in Figure 1A (See Supplemental File).

Antigen	Comparison	N	F	<i>P</i> -value
IL-1 α	All groups	5	2.69	0.051
IL-1 β	All groups	5	11.15	<0.001*
IL-2	All groups	5	10.27	<0.001*
IL-4	All groups	5	6.00	0.001*
IL-5	All groups	5	2.63	0.055
IL-6	All groups	5	6.31	0.001*
IL-12	All groups	5	5.21	0.003*
IL-17	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	4	1.15	0.349
CSF2	None – all undetectable	n/a	n/a	n/a
CCL2	All groups	5	16.03	<0.001*
CCL3	All groups	5	2.59	0.058
IFN γ	All groups	5	3.25	0.026*
TNF α	All groups	5	23.80	<0.001*
CXCL1	All groups	5	4.07	0.010*

Supplemental Table 2: Two-way ANOVA analyses of log-transformed serum time series data. Groups listed in comparison column represent groups for which data was quantified throughout the time series. Tukey-Kramer HSD post-hoc tests were used for pairwise comparisons and the results are summarized in Figure 1B (See Supplemental File).

Antigen	Comparison	N	Group		Day		Group*Day	
			F	P-value	F	P-value	F	P-value
IL-1 α	All groups	255	32.51	<0.001*	0.93	0.471	1.62	0.039*
IL-1 β	Mm-UKY, Mm-Wild, Ap	157	3.02	0.052	1.06	0.391	1.08	0.386
IL-2	All groups	255	27.82	<0.001*	2.46	0.025*	1.96	0.006*
IL-4	All groups	255	34.54	<0.001*	1.47	0.188	1.55	0.053
IL-5	All groups	255	63.22	<0.001*	0.94	0.469	0.82	0.706
IL-6	All groups	255	64.61	<0.001*	2.29	0.037*	2.63	<0.001*
IL-12	All groups	255	33.25	<0.001*	2.21	0.660	0.91	0.583
IL-17	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	205	7.64	<0.001*	2.78	0.013*	2.60	<0.001*
CSF2	None – all undetectable	0	n/a	n/a	n/a	n/a	n/a	n/a
CCL2	All groups	255	5.72	<0.001*	1.46	0.193	1.16	0.278
CCL3	All groups	255	38.75	<0.001*	0.91	0.486	1.47	0.079
IFN γ	All groups	255	37.81	<0.001*	1.22	0.296	1.77	0.018*
TNF α	All groups	255	31.88	<0.001*	0.85	0.535	0.91	0.583
CXCL1	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	205	69.43	<0.001*	4.33	<0.001*	1.47	0.104

Supplemental Table 3: Two-way ANOVA analyses of log-transformed tissue lysate time series data. Groups listed in comparison column represent groups for which data was quantified throughout the time series. *P*-values indicate where at least one group is significantly different from another group. Tukey-Kramer HSD post-hoc tests were used for pairwise comparisons and the results are summarized in Figure 2 (See Supplemental File).

Antigen	Comparison	N	Group		Day		Group*Day	
			F	P-value	F	P-value	F	P-value
IL-1 α	None – all above limit of detection	0	n/a	n/a	n/a	n/a	n/a	n/a
IL-1 β	All groups	261	77.88	<0.001*	39.47	<0.001*	5.33	<0.001*
IL-2	All groups	261	190.93	<0.001*	6.68	<0.001*	13.59	<0.001*
IL-4	All groups	261	29.09	<0.001*	11.24	<0.001*	6.76	<0.001*
IL-5	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	211	22.73	<0.001*	10.03	<0.001*	10.70	<0.001*
IL-6	All groups	261	76.89	<0.001*	104.11	<0.001*	15.06	<0.001*
IL-12	All groups	261	123.20	<0.001*	10.42	<0.001*	18.28	<0.001*
IL-17	Mm-UKY, Mm-Kenya, Ap	177	598.63	<0.001*	4.42	<0.001*	17.55	<0.001*
CSF2	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	211	104.44	<0.001*	21.36	<0.001*	8.51	<0.001*
CCL2	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	211	72.96	<0.001*	38.71	<0.001*	5.23	<0.001*
CCL3	All groups	261	8.30	<0.001*	160.11	<0.001*	6.24	<0.001*
IFN γ	Mm-Kenya, Mm-Wild, Ap	161	40.47	<0.001*	2.42	0.030*	3.00	<0.001*
TNF α	All groups	261	149.87	<0.001*	81.32	<0.001*	11.84	<0.001*
CXCL1	Mm-UKY, Mm-Kenya, Mm-Wild, Ap	211	137.76	<0.001*	201.75	<0.001*	26.64	<0.001*

Supplemental Table 4: One-way ANOVA analysis of tissue cytokine data to test for an effect of group at D1, 2, 10 and 20. *P*-values indicate where at least one group is significantly different from another group. Tukey-Kramer HSD post-hoc tests were used for pairwise comparisons and are summarized in Figure 3 (See Supplemental File).

Antigen	Comparison	N	Day	F-Ratio	P-value
IL-1 β	All groups		1	5.39	0.002*
			2	32.52	<0.001*
			10	20.70	<0.001*
			20	19.05	<0.001*
IL-4	All groups		1	9.71	<0.001*
			2	4.8	0.004*
			10	5.23	0.003*
			20	18.89	<0.001*
IL-6	All groups		1	14.56	<0.001*
			2	47.24	<0.001*
			10	23.30	<0.001*
			20	21.50	<0.001*
IL-12	All groups		1	57.33	<0.001*
			2	14.22	<0.001*
			10	7.66	<0.001*
			20	26.33	<0.001*
IL-17	Mm-UKY, Mm-Kenya and Ap		1	45.87	<0.001*
			2	93.60	<0.001*
			10	441.78	<0.001*
			20	116.93	<0.001*
CCL2	Mm-UKY, Mm-Kenya, Mm-Wild, Ap		1	21.55	<0.001*
			2	33.43	<0.001*
			10	7.04	0.002*
			20	11.68	<0.001*
TNF α	All groups		1	39.61	<0.001*
			2	54.97	<0.001*
			10	27.95	<0.001*
			20	67.67	<0.001*
CXCL1	All groups		1	60.99	<0.001*
			2	210.02	<0.001*
			10	15.23	<0.001*
			20	9.54	<0.001*

Supplemental Table 5: List of T cell phenotyping antibodies tried using flow cytometry in *Acomys cahirinus*.

Antigen	Clone	Supplier	Catalog#	Status
CD45	OX-1	BioLegend	202207	No – no separation
	30-F11	BioLegend	103106	No – no separation
	I3/2.3	Molecular Probes	A15395	No – no separation
CD3	17A2	BioLegend	100236	Yes – good population
TCRab	H57-597	BioLegend	109207	No – poor separation
	R73	BioLegend	201107	No – no separation
TCRgd	GL3	BioLegend	118107	No – poor separation
	VC7-13D5	BioLegend	107507	No – poor separation
CD4	GK1.5	BioLegend	100433	No – no separation
	RM4-4	BioLegend	116011	No – poor separation
CD8	53-6.7	BioLegend	100726	No – no separation
CD25	P61	BioLegend	102015	No – poor separation
	3C7	BioLegend	101915	No – no separation
CD69	H1.2F3	BioLegend	104511	No – poor separation
CD196	29-2L17	BioLegend	129813	No – poor separation
	140706	BD Pharmingen	561753	Yes – good population
CD206	C068C2	BioLegend	141705	Yes – good population
CD49b	HMa2	BioLegend	103515	Yes – good population
	DX5	BioLegend	108909	No – poor separation

Supplemental Figure 1 Comparison of parallelism for cytokine assays demonstrates ability to quantify cytokines across three species. Serial dilution of serum (triangles, solid lines) and tissue lysate (squares, dashed lines) shows similar negative slopes for *M. musculus* (green), *A. percivali* (red) and *A. cahirinus* (blue) compared to the standard (black, circles, solid line). Data represent the mean for duplicates and lines are linear regressions calculated from all data that was above the lower limit of detection.

Supplemental Figure 2 Comparisons of amino acid sequence alignments for antigens in study. Predicted amino acid sequence was translated from mRNA (*A. cahirinus*) or obtained from NCBI (*H. sapiens*, *M. musculus*, *R. rattus*) and aligned together using MAFFT. Figures were created using BOXSHADE where solid black shading is equivalent, gray shading is similar and no shading is dissimilar.

Supplemental Files

Supplemental File 1 contains summary tables of post-hoc multiple comparison tests used in Figure 1B, Figure 2 and Figure 3.

857 REFERENCES

- 858 1. Seifert, A.W., and Maden, M. (2014). New insights into vertebrate skin regeneration. In
859 International review of cell and molecular biology, Volume 310. (Elsevier), pp. 129-169.
- 860 2. Martin, P. (1997). Wound healing--aiming for perfect skin regeneration. *Science* 276, 75-81.
- 861 3. Carlson, B.M. (2007). Principles of regenerative biology, (Amsterdam ; Burlington, Mass.:
862 Elsevier/Academic Press).
- 863 4. Harty, M., Neff, A.W., King, M.W., and Mescher, A.L. (2003). Regeneration or scarring: an
864 immunologic perspective. *Dev Dyn* 226, 268-279.
- 865 5. Seifert, A.W., and Muneoka, K. (2018). The blastema and epimorphic regeneration in mammals.
866 *Dev Biol* 433, 190-199.
- 867 6. Eming, S.A., Wynn, T.A., and Martin, P. (2017). Inflammation and metabolism in tissue repair
868 and regeneration. *Science* 356, 1026-1030.
- 869 7. Stoecklein, V.M., Osuka, A., and Lederer, J.A. (2012). Trauma equals danger--damage control by
870 the immune system. *J Leukoc Biol* 92, 539-551.
- 871 8. Mescher, A.L. (2017). Macrophages and fibroblasts during inflammation and tissue repair in
872 models of organ regeneration. *Regeneration (Oxf)* 4, 39-53.
- 873 9. Simkin, J., and Seifert, A.W. (2018). Concise Review: Translating Regenerative Biology into
874 Clinically Relevant Therapies: Are We on the Right Path? *Stem Cells Transl Med* 7, 220-231.
- 875 10. Godwin, J.W., and Brookes, J.P. (2006). Regeneration, tissue injury and the immune response. *J*
876 *Anat* 209, 423-432.
- 877 11. Aurora, A.B., Porrello, E.R., Tan, W., Mahmoud, A.I., Hill, J.A., Bassel-Duby, R., Sadek, H.A.,
878 and Olson, E.N. (2014). Macrophages are required for neonatal heart regeneration. *J Clin Invest*
879 124, 1382-1392.
- 880 12. Godwin, J.W., Pinto, A.R., and Rosenthal, N.A. (2013). Macrophages are required for adult
881 salamander limb regeneration. *Proc Natl Acad Sci U S A* 110, 9415-9420.
- 882 13. Godwin, J.W., Debuque, R., Salimova, E., and Rosenthal, N.A. (2017). Heart regeneration in the
883 salamander relies on macrophage-mediated control of fibroblast activation and the extracellular
884 landscape. *NPJ Regen Med* 2.
- 885 14. Petrie, T.A., Strand, N.S., Yang, C.T., Rabinowitz, J.S., and Moon, R.T. (2014). Macrophages
886 modulate adult zebrafish tail fin regeneration. *Development* 141, 2581-2591.
- 887 15. Simkin, J., Gawriluk, T.R., Gensel, J.C., and Seifert, A.W. (2017). Macrophages are necessary for
888 epimorphic regeneration in African spiny mice. *Elife* 6.
- 889 16. Simkin, J., Sammarco, M.C., Marrero, L., Dawson, L.A., Yan, M., Tucker, C., Cammack, A., and
890 Muneoka, K. (2017). Macrophages are required to coordinate mouse digit tip regeneration.
891 *Development*.
- 892 17. Panduro, M., Benoist, C., and Mathis, D. (2018). Treg cells limit IFN-gamma production to
893 control macrophage accrual and phenotype during skeletal muscle regeneration. *Proc Natl Acad*
894 *Sci U S A*.
- 895 18. Hui, S.P., Sheng, D.Z., Sugimoto, K., Gonzalez-Rajal, A., Nakagawa, S., Hesselson, D., and
896 Kikuchi, K. (2017). Zebrafish Regulatory T Cells Mediate Organ-Specific Regenerative
897 Programs. *Dev Cell* 43, 659-672 e655.
- 898 19. Love, N.R., Chen, Y., Ishibashi, S., Kritsiligkou, P., Lea, R., Koh, Y., Gallop, J.L., Dorey, K.,
899 and Amaya, E. (2013). Amputation-induced reactive oxygen species are required for successful
900 *Xenopus* tadpole tail regeneration. *Nat Cell Biol* 15, 222-228.
- 901 20. Gauron, C., Rampon, C., Bouzaffour, M., Ipendey, E., Teillon, J., Volovitch, M., and Vriza, S.
902 (2013). Sustained production of ROS triggers compensatory proliferation and is required for
903 regeneration to proceed. *Sci Rep* 3, 2084.
- 904 21. Ferreira, F., Luxardi, G., Reid, B., and Zhao, M. (2016). Early bioelectric activities mediate
905 redox-modulated regeneration. *Development* 143, 4582-4594.

22. He, L., and Marneros, A.G. (2013). Macrophages are essential for the early wound healing response and the formation of a fibrovascular scar. *Am J Pathol* 182, 2407-2417.
23. Nosbaum, A., Prevel, N., Truong, H.A., Mehta, P., Ettinger, M., Scharschmidt, T.C., Ali, N.H., Pauli, M.L., Abbas, A.K., and Rosenblum, M.D. (2016). Cutting Edge: Regulatory T Cells Facilitate Cutaneous Wound Healing. *J Immunol* 196, 2010-2014.
24. Mirza, R., DiPietro, L.A., and Koh, T.J. (2009). Selective and specific macrophage ablation is detrimental to wound healing in mice. *Am J Pathol* 175, 2454-2462.
25. Goren, I., Allmann, N., Yogev, N., Schurmann, C., Linke, A., Holdener, M., Waisman, A., Pfeilschifter, J., and Frank, S. (2009). A transgenic mouse model of inducible macrophage depletion: effects of diphtheria toxin-driven lysozyme M-specific cell lineage ablation on wound inflammatory, angiogenic, and contractive processes. *Am J Pathol* 175, 132-147.
26. Lucas, T., Waisman, A., Ranjan, R., Roes, J., Krieg, T., Muller, W., Roers, A., and Eming, S.A. (2010). Differential roles of macrophages in diverse phases of skin repair. *J Immunol* 184, 3964-3977.
27. Huo, Y., Qiu, W.Y., Pan, Q., Yao, Y.F., Xing, K., and Lou, M.F. (2009). Reactive oxygen species (ROS) are essential mediators in epidermal growth factor (EGF)-stimulated corneal epithelial cell proliferation, adhesion, migration, and wound healing. *Exp Eye Res* 89, 876-886.
28. Hopkinson-Woolley, J., Hughes, D., Gordon, S., and Martin, P. (1994). Macrophage recruitment during limb development and wound healing in the embryonic and foetal mouse. *J Cell Sci* 107 (Pt 5), 1159-1167.
29. Whitby, D.J., and Ferguson, M.W. (1991). Immunohistochemical localization of growth factors in fetal wound healing. *Dev Biol* 147, 207-215.
30. Martin, P., Dickson, M.C., Millan, F.A., and Akhurst, R.J. (1993). Rapid induction and clearance of TGF beta 1 is an early response to wounding in the mouse embryo. *Dev Genet* 14, 225-238.
31. Mescher, A.L., Neff, A.W., and King, M.W. (2013). Changes in the inflammatory response to injury and its resolution during the loss of regenerative capacity in developing *Xenopus* limbs. *PLoS One* 8, e80477.
32. Robert, J., and Ohta, Y. (2009). Comparative and developmental study of the immune system in *Xenopus*. *Dev Dyn* 238, 1249-1270.
33. Bertolotti, E., Malagoli, D., and Franchini, A. (2013). Skin wound healing in different aged *Xenopus laevis*. *J Morphol* 274, 956-964.
34. Brant, J.O., Yoon, J.H., Polvadore, T., Barbazuk, W.B., and Maden, M. (2016). Cellular events during scar-free skin regeneration in the spiny mouse, *Acomys*. *Wound Repair Regen* 24, 75-88.
35. Mak, K., Manji, A., Gallant-Behm, C., Wiebe, C., Hart, D.A., Larjava, H., and Hakkinen, L. (2009). Scarless healing of oral mucosa is characterized by faster resolution of inflammation and control of myofibroblast action compared to skin wounds in the red Duroc pig model. *J Dermatol Sci* 56, 168-180.
36. Szpaderska, A.M., Zuckerman, J.D., and DiPietro, L.A. (2003). Differential injury responses in oral mucosal and cutaneous wounds. *J Dent Res* 82, 621-626.
37. Chen, L., Arbieva, Z.H., Guo, S., Marucha, P.T., Mustoe, T.A., and DiPietro, L.A. (2010). Positional differences in the wound transcriptome of skin and oral mucosa. *BMC Genomics* 11, 471.
38. Fini, M.E., and Sicard, R.E. (1980). Limb regeneration of the adult newt (*Notophthalmus viridescens*) in the absence of the spleen. *Wilehm Roux Arch Dev Biol* 189, 77-79.
39. Schotte, O.E., and Sicard, R.E. (1982). Cyclophosphamide-induced leukopenia and suppression of limb regeneration in the adult newt, *notophthalmus viridescens*. *J Exp Zool* 222, 199-202.
40. Gawriluk, T.R., Simkin, J., Thompson, K.L., Biswas, S.K., Clare-Salzler, Z., Kimani, J.M., Kiama, S.G., Smith, J.J., Ezenwa, V.O., and Seifert, A.W. (2016). Comparative analysis of ear-hole closure identifies epimorphic regeneration as a discrete trait in mammals. *Nat Commun* 7, 11164.

- 956 41. Seifert, A.W., Kiama, S.G., Seifert, M.G., Goheen, J.R., Palmer, T.M., and Maden, M. (2012).
957 Skin shedding and tissue regeneration in African spiny mice (*Acomys*). *Nature* *489*, 561-565.
- 958 42. Matias Santos, D., Rita, A.M., Casanellas, I., Brito Ova, A., Araujo, I.M., Power, D., and
959 Tiscornia, G. (2016). Ear wound regeneration in the African spiny mouse *Acomys cahirinus*.
960 *Regeneration (Oxf)* *3*, 52-61.
- 961 43. Leshner, A., Li, B., Whitt, P., Newton, N., Devalapalli, A.P., Shieh, K., Solow, J.S., and Parker,
962 W. (2006). Increased IL-4 production and attenuated proliferative and pro-inflammatory
963 responses of splenocytes from wild-caught rats (*Rattus norvegicus*). *Immunol Cell Biol* *84*, 374-
964 382.
- 965 44. Titus, R.G., Sherry, B., and Cerami, A. (1991). The involvement of TNF, IL-1 and IL-6 in the
966 immune response to protozoan parasites. *Immunol Today* *12*, A13-16.
- 967 45. Viney, M., and Riley, E.M. (2017). The Immunology of Wild Rodents: Current Status and Future
968 Prospects. *Front Immunol* *8*, 1481.
- 969 46. Julier, Z., Park, A.J., Briquez, P.S., and Martino, M.M. (2017). Promoting tissue regeneration by
970 modulating the immune system. *Acta Biomater* *53*, 13-28.
- 971 47. Panduro, M., Benoist, C., and Mathis, D. (2018). Treg cells limit IFN-gamma production to
972 control macrophage accrual and phenotype during skeletal muscle regeneration. *Proc Natl Acad*
973 *Sci U S A* *115*, E2585-E2593.
- 974 48. Trinchieri, G., Pflanz, S., and Kastelein, R.A. (2003). The IL-12 family of heterodimeric
975 cytokines: new players in the regulation of T cell responses. *Immunity* *19*, 641-644.
- 976 49. Weaver, C.T., Hatton, R.D., Mangan, P.R., and Harrington, L.E. (2007). IL-17 family cytokines
977 and the expanding diversity of effector T cell lineages. *Annu Rev Immunol* *25*, 821-852.
- 978 50. Heinrich, P.C., Behrmann, I., Muller-Newen, G., Schaper, F., and Graeve, L. (1998). Interleukin-
979 6-type cytokine signalling through the gp130/Jak/STAT pathway. *Biochem J* *334 (Pt 2)*, 297-
980 314.
- 981 51. Hunter, C.A., and Jones, S.A. (2017). Corrigendum: IL-6 as a keystone cytokine in health and
982 disease. *Nat Immunol* *18*, 1271.
- 983 52. Tanaka, T., Narazaki, M., and Kishimoto, T. (2014). IL-6 in inflammation, immunity, and
984 disease. *Cold Spring Harb Perspect Biol* *6*, a016295.
- 985 53. Kroeze, K.L., Boink, M.A., Sampat-Sardjoepersad, S.C., Waaijman, T., Scheper, R.J., and Gibbs,
986 S. (2012). Autocrine regulation of re-epithelialization after wounding by chemokine receptors
987 CCR1, CCR10, CXCR1, CXCR2, and CXCR3. *J Invest Dermatol* *132*, 216-225.
- 988 54. Lin, Z.Q., Kondo, T., Ishida, Y., Takayasu, T., and Mukaida, N. (2003). Essential involvement of
989 IL-6 in the skin wound-healing process as evidenced by delayed wound healing in IL-6-deficient
990 mice. *J Leukoc Biol* *73*, 713-721.
- 991 55. Gallucci, R.M., Simeonova, P.P., Matheson, J.M., Kommineni, C., Guriel, J.L., Sugawara, T.,
992 and Luster, M.I. (2000). Impaired cutaneous wound healing in interleukin-6-deficient and
993 immunosuppressed mice. *FASEB J* *14*, 2525-2531.
- 994 56. Devalaraja, R.M., Nanney, L.B., Du, J., Qian, Q., Yu, Y., Devalaraja, M.N., and Richmond, A.
995 (2000). Delayed wound healing in CXCR2 knockout mice. *J Invest Dermatol* *115*, 234-244.
- 996 57. McFarland-Mancini, M.M., Funk, H.M., Paluch, A.M., Zhou, M., Giridhar, P.V., Mercer, C.A.,
997 Kozma, S.C., and Drew, A.F. (2010). Differences in wound healing in mice with deficiency of
998 IL-6 versus IL-6 receptor. *J Immunol* *184*, 7219-7228.
- 999 58. Wu, K.K. (1996). Cyclooxygenase 2 induction: molecular mechanism and pathophysiologic
1000 roles. *J Lab Clin Med* *128*, 242-245.
- 1001 59. Penning, T.D., Talley, J.J., Bertenshaw, S.R., Carter, J.S., Collins, P.W., Docter, S., Graneto,
1002 M.J., Lee, L.F., Malecha, J.W., Miyashiro, J.M., et al. (1997). Synthesis and biological evaluation
1003 of the 1,5-diarylpyrazole class of cyclooxygenase-2 inhibitors: identification of 4-[5-(4-
1004 methylphenyl)-3-(trifluoromethyl)-1H-pyrazol-1-yl]benzene nesulfonamide (SC-58635, celecoxib).
1005 *J Med Chem* *40*, 1347-1365.

60. Beura, L.K., Hamilton, S.E., Bi, K., Schenkel, J.M., Odumade, O.A., Casey, K.A., Thompson, E.A., Fraser, K.A., Rosato, P.C., Filali-Mouhim, A., et al. (2016). Normalizing the environment recapitulates adult human immune traits in laboratory mice. *Nature* 532, 512-516.
61. Abolins, S., King, E.C., Lazarou, L., Weldon, L., Hughes, L., Drescher, P., Raynes, J.G., Hafalla, J.C.R., Viney, M.E., and Riley, E.M. (2017). The comparative immunology of wild and laboratory mice, *Mus musculus domesticus*. *Nat Commun* 8, 14811.
62. Martin, P., and Leibovich, S.J. (2005). Inflammatory cells during wound repair: the good, the bad and the ugly. *Trends Cell Biol* 15, 599-607.
63. Pellegrino, M.J., and Habecker, B.A. (2013). STAT3 integrates cytokine and neurotrophin signals to promote sympathetic axon regeneration. *Mol Cell Neurosci* 56, 272-282.
64. Qin, S., Zou, Y., and Zhang, C.L. (2013). Cross-talk between KLF4 and STAT3 regulates axon regeneration. *Nat Commun* 4, 2633.
65. Zhu, H., Xiao, F., Wang, G., Wei, X., Jiang, L., Chen, Y., Zhu, L., Wang, H., Diao, Y., Wang, H., et al. (2016). STAT3 Regulates Self-Renewal of Adult Muscle Satellite Cells during Injury-Induced Muscle Regeneration. *Cell Rep* 16, 2102-2115.
66. Bard, J.D., Gelebart, P., Amin, H.M., Young, L.C., Ma, Y., and Lai, R. (2009). Signal transducer and activator of transcription 3 is a transcriptional factor regulating the gene expression of SALL4. *FASEB J* 23, 1405-1414.
67. Grow, M., Neff, A.W., Mescher, A.L., and King, M.W. (2006). Global analysis of gene expression in *Xenopus* hindlimbs during stage-dependent complete and incomplete regeneration. *Dev Dyn* 235, 2667-2685.
68. Neff, A.W., King, M.W., and Mescher, A.L. (2011). Dedifferentiation and the role of sall4 in reprogramming and patterning during amphibian limb regeneration. *Dev Dyn* 240, 979-989.
69. Fairweather, M., Heit, Y.I., Buie, J., Rosenberg, L.M., Briggs, A., Orgill, D.P., and Bertagnolli, M.M. (2015). Celecoxib inhibits early cutaneous wound healing. *J Surg Res* 194, 717-724.
70. Romana-Souza, B., Santos, J.S., Bandeira, L.G., and Monte-Alto-Costa, A. (2016). Selective inhibition of COX-2 improves cutaneous wound healing of pressure ulcers in mice through reduction of iNOS expression. *Life Sci* 153, 82-92.
71. Wilgus, T.A., Vodovotz, Y., Vittadini, E., Clubbs, E.A., and Oberyshyn, T.M. (2003). Reduction of scar formation in full-thickness wounds with topical celecoxib treatment. *Wound Repair Regen* 11, 25-34.
72. Hildebrand, F., Hubbard, W.J., Choudhry, M.A., Frink, M., Pape, H.C., Kunkel, S.L., and Chaudry, I.H. (2006). Kupffer cells and their mediators: the culprits in producing distant organ damage after trauma-hemorrhage. *Am J Pathol* 169, 784-794.
73. Deshmane, S.L., Kremlev, S., Amini, S., and Sawaya, B.E. (2009). Monocyte chemoattractant protein-1 (MCP-1): an overview. *J Interferon Cytokine Res* 29, 313-326.
74. Low, Q.E., Drugea, I.A., Duffner, L.A., Quinn, D.G., Cook, D.N., Rollins, B.J., Kovacs, E.J., and DiPietro, L.A. (2001). Wound healing in MIP-1alpha(-/-) and MCP-1(-/-) mice. *Am J Pathol* 159, 457-463.
75. Ziraldo, C., Vodovotz, Y., Namas, R.A., Almahmoud, K., Tapias, V., Mi, Q., Barclay, D., Jefferson, B.S., Chen, G., Billiar, T.R., et al. (2013). Central role for MCP-1/CCL2 in injury-induced inflammation revealed by in vitro, in silico, and clinical studies. *PLoS One* 8, e79804.
76. Sierra-Filardi, E., Nieto, C., Dominguez-Soto, A., Barroso, R., Sanchez-Mateos, P., Puig-Kroger, A., Lopez-Bravo, M., Joven, J., Ardavin, C., Rodriguez-Fernandez, J.L., et al. (2014). CCL2 shapes macrophage polarization by GM-CSF and M-CSF: identification of CCL2/CCR2-dependent gene expression profile. *J Immunol* 192, 3858-3867.
77. Lu, B., Rutledge, B.J., Gu, L., Fiorillo, J., Lukacs, N.W., Kunkel, S.L., North, R., Gerard, C., and Rollins, B.J. (1998). Abnormalities in monocyte recruitment and cytokine expression in monocyte chemoattractant protein 1-deficient mice. *J Exp Med* 187, 601-608.
78. Godwin, J.W., and Rosenthal, N. (2014). Scar-free wound healing and regeneration in amphibians: immunological influences on regenerative success. *Differentiation* 87, 66-75.

- 1057 79. Mescher, A.L., Neff, A.W., and King, M.W. (2017). Inflammation and immunity in organ
1058 regeneration. *Dev Comp Immunol* 66, 98-110.
- 1059 80. Hirahara, K., and Nakayama, T. (2016). CD4+ T-cell subsets in inflammatory diseases: beyond
1060 the Th1/Th2 paradigm. *Int Immunol* 28, 163-171.
- 1061 81. Barbul, A., Breslin, R.J., Woodyard, J.P., Wasserkrug, H.L., and Efron, G. (1989). The effect of
1062 in vivo T helper and T suppressor lymphocyte depletion on wound healing. *Ann Surg* 209, 479-
1063 483.
- 1064 82. Barbul, A., Shawe, T., Rotter, S.M., Efron, J.E., Wasserkrug, H.L., and Badawy, S.B. (1989).
1065 Wound healing in nude mice: a study on the regulatory role of lymphocytes in fibroplasia.
1066 *Surgery* 105, 764-769.
- 1067 83. Barbul, A., Sisto, D., Rettura, G., Levenson, S.M., Seifter, E., and Efron, G. (1982). Thymic
1068 inhibition of wound healing: abrogation by adult thymectomy. *J Surg Res* 32, 338-342.
- 1069 84. Fishel, R., Barbul, A., Wasserkrug, H.L., Penberthy, L.T., Rettura, G., and Efron, G. (1983).
1070 Cyclosporine A impairs wound healing in rats. *J Surg Res* 34, 572-575.
- 1071 85. Peterson, J.M., Barbul, A., Breslin, R.J., Wasserkrug, H.L., and Efron, G. (1987). Significance of
1072 T-lymphocytes in wound healing. *Surgery* 102, 300-305.
- 1073 86. Zhang, J., Xiao, Z., Qu, C., Cui, W., Wang, X., and Du, J. (2014). CD8 T cells are involved in
1074 skeletal muscle regeneration through facilitating MCP-1 secretion and Gr1(high) macrophage
1075 infiltration. *J Immunol* 193, 5149-5160.
- 1076 87. Reinke, S., Geissler, S., Taylor, W.R., Schmidt-Bleek, K., Juelke, K., Schwachmeyer, V., Dahne,
1077 M., Hartwig, T., Akyuz, L., Meisel, C., et al. (2013). Terminally differentiated CD8(+) T cells
1078 negatively affect bone regeneration in humans. *Sci Transl Med* 5, 177ra136.
- 1079 88. Kwon, J., Devadas, S., and Williams, M.S. (2003). T cell receptor-stimulated generation of
1080 hydrogen peroxide inhibits MEK-ERK activation and I κ B serine phosphorylation. *Free Radic Biol*
1081 *Med* 35, 406-417.
- 1082 89. Kim, H.R., Lee, A., Choi, E.J., Hong, M.P., Kie, J.H., Lim, W., Lee, H.K., Moon, B.I., and Seoh,
1083 J.Y. (2014). Reactive oxygen species prevent imiquimod-induced psoriatic dermatitis through
1084 enhancing regulatory T cell function. *PLoS One* 9, e91146.
- 1085 90. Kraaij, M.D., Savage, N.D., van der Kooij, S.W., Koekkoek, K., Wang, J., van den Berg, J.M.,
1086 Ottenhoff, T.H., Kuijpers, T.W., Holmdahl, R., van Kooten, C., et al. (2010). Induction of
1087 regulatory T cells by macrophages is dependent on production of reactive oxygen species. *Proc*
1088 *Natl Acad Sci U S A* 107, 17686-17691.
- 1089 91. Haughton, C.L., Gawriluk, T.R., and Seifert, A.W. (2016). The Biology and Husbandry of the
1090 African Spiny Mouse (*Acomys cahirinus*) and the Research Uses of a Laboratory Colony. *J Am*
1091 *Assoc Lab Anim Sci* 55, 9-17.
- 1092 92. Plikaytis, B.D., Holder, P.F., Pais, L.B., Maslanka, S.E., Gheesling, L.L., and Carlone, G.M.
1093 (1994). Determination of parallelism and nonparallelism in bioassay dilution curves. *J Clin*
1094 *Microbiol* 32, 2441-2447.
- 1095 93. Tu, J., and Bennett, P. (2017). Parallelism experiments to evaluate matrix effects, selectivity and
1096 sensitivity in ligand-binding assay method development: pros and cons. *Bioanalysis* 9, 1107-
1097 1122.
- 1098 94. Mamrot, J., Legaie, R., Ellery, S.J., Wilson, T., Seemann, T., Powell, D.R., Gardner, D.K.,
1099 Walker, D.W., Temple-Smith, P., Papenfuss, A.T., et al. (2017). De novo transcriptome assembly
1100 for the spiny mouse (*Acomys cahirinus*). *Sci Rep* 7, 8996.
- 1101 95. Katoh, K., Misawa, K., Kuma, K., and Miyata, T. (2002). MAFFT: a novel method for rapid
1102 multiple sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 30, 3059-3066.
- 1103 96. Katoh, K., Rozewicki, J., and Yamada, K.D. (2017). MAFFT online service: multiple sequence
1104 alignment, interactive sequence choice and visualization. *Brief Bioinform.*
- 1105 97. Whitcomb, B.W., and Schisterman, E.F. (2008). Assays with lower detection limits: implications
1106 for epidemiological investigations. *Paediatr Perinat Epidemiol* 22, 597-602.

1107 98. Rueden, C.T., Schindelin, J., Hiner, M.C., DeZonia, B.E., Walter, A.E., Arena, E.T., and Eliceiri,
1108 K.W. (2017). ImageJ2: ImageJ for the next generation of scientific image data. BMC
1109 Bioinformatics 18, 529.
1110




Figure 1 Ability to regenerate and immunity-status are associated with a specific systemic immune profile and immune response to injury. (a) Comparison of cytokine concentrations in serum from uninjured animals showed higher concentrations of IL-4, IL-6, CCL2 and TNF α in wild-caught animals compared to laboratory-reared animals indicated that wild animals have a “primed” immune system. No difference was found between non-regenerators (Mm-UKY, Mm-Kenya and Mm-Wild) and regenerators (Ac and Ap). Data represent box and whiskers with median, interquartile range and individual data points. N/A denotes concentrations could not be quantified in any animal of the group. The dashed line in each graph represents the lower limit of detection for the specific cytokine. N/S denotes $P > 0.05$ for One-way ANOVA (See Supplemental Table 1) and different letters above the data denotes $P \leq 0.05$ for Tukey-Kramer pairwise comparisons (See Supplemental File). (b) The change in systemic cytokines compared to D0 over 20 days after ear hole punch. A naïve immune system response (solid lines) was consistent with increased IL-2 and TNF α , and decreased IL-1 α compared to a primed response (intermittent lines) (See Supplemental Table 2). Non-regenerators (black) showed increases in IL-5, IL-6, IL-17, CCL3 and CXCL1 compared to regenerators (red). Data represent mean and S.E.M. for at least $n=5$ animals per timepoint. The dashed line at $Y=1$ represents no change compared to D0 and the yellow boxes represent the inflammation resolution window.




Figure 2 The tissue microenvironment is dynamic, and the ability to regenerate is associated with a specific immune response to injury. The comparison of the change in cytokines from tissue lysate compared to D0 over 20 days after ear hole punch. All cytokines varied over time and between groups (See Supplemental Table 3). There was a stronger increase in CCL3 a muted increase for IL-12 and CXCL1 in immune-naïve (solid lines) compared to immune-primed animals (intermittent lines). Non-regenerators (black) had stronger increases for IL-6, CCL2 and CXCL1 compared to regenerators (red). Additionally, IL-17 decreased in non-regenerators and increased in regenerators. Data represent mean and S.E.M. for at least n=5 animals per species per timepoint. The dashed line at Y=1 represents no change compared to D0 and the yellow boxes represent the inflammation resolution window.




Figure 3 Regeneration is associated with specific temporal cytokine responses in the tissue microenvironment. The comparison of the change in cytokines from tissue lysate compared to baseline. Non-regenerators (Mm-UKY, Mm-Kenya, and Mm-Wild: Black) show a stronger response compared to regenerators (Ac and Ap: Red) for several inflammatory cytokines during the immediate response, acute inflammatory phase. These differences mostly resolved at day 10 and 20. In contrast, regenerators showed a stronger IL-12 and IL-17 response during tissue morphogenesis. Data represent box and whiskers with median, interquartile range and individual data points. N/A denotes concentrations could not be quantified. The dashed line at Y=1 represents no change compared to D0. Each graph showed P<0.05 for effect of group using a One-way ANOVA on log-transformed data (See Supplemental Table 4) and different letters above each group denotes P<0.05 for Tukey-Kramer pairwise comparisons (See Supplemental File).




Figure 4 The regeneration microenvironment is primed by greater T cell influx and TREG signature.

(A) Comparison of total CD3⁺ cells quantified by flow cytometry from dissociated ear pinna and (B) the ratio relative to uninjured tissue for *M. musculus* (black) and *A. cahirinus* (red). Data represent mean and S.E.M. and n=4 or 5. An * denotes P<0.05 for pairwise comparison within the day between species for Holm-Sidak posthoc test.

(C-I) Representative immunohistochemistry for CD3 (red) counterstained with DAPI (gray) at the proximal wound margin (amputation plane can be determined from the end of the cartilage—indicated by the dotted line) from D5 and D15 after injury of *M. musculus* (C, D) and *A. cahirinus* (E, F). More T cells (yellow arrowhead) were present throughout the wound bed and were mainly found in the dermis of *A. cahirinus* compared to *M. musculus*. The T cells associated with epidermis (boundaries indicated by the dotted line) tended to be spindle-like in *M. musculus* (G), while rounded in *A. cahirinus* (H). The dermal T cells in *A. cahirinus* also tended to be in close proximity to regenerating epidermal appendages (I). N=4 and bar equals 200 μ m (C-F) or 20 μ m (G-I). (J) Heatmap of differential gene expression compared to uninjured tissue suggests that the regeneration microenvironment contains a substantial NK, CD8⁺ and TREG cell response while fibrotic repair has a CD4⁺ cell response. Data comes from a previously published analysis [34].




Figure 5 Postponed STAT3 activation in the blastema is associated with regeneration. (A-B) Comparison of representative immunoblots for pSTAT3 and ACTB for indicated time and species from injured tissue homogenate from ear-hole punch assay (A). Fibrotic repair is associated with strong early STAT3 activation while regeneration is associated with a weak early and strong postponed activation (B). Data represent mean fold change of pSTAT3 intensity normalized to ACTB intensity and relative to D0 and S.E.M. n=4 individuals per time point and species (B). (C-F) Representative images for immunohistochemistry for pSTAT3 (Brown) counterstained with hematoxylin (Blue) for *M. musculus* (C, C', E and E') and *A. cahirinus* (D, D', F and F') at the indicated time points. On D2 after injury, nearly every epidermal cell in *M. musculus* was positive compared to about half in *A. cahirinus* (C, D). On D15 after injury, only a small population of epidermal cells in *M. musculus* were positive compared to about half in *A. cahirinus* (E, F). Additionally, about half of the cells in the forming blastema remain positive at D15. Data represent n=3, and bar equals 200 μ m.





Figure 6 COX-2 Inhibition does not liberate regeneration in *M. musculus*.

(A) Celecoxib treatment during the first 10 days of injury reduces secondary inflammation caused by IL-6 and CXCL1. (B-C) Celecoxib treatment did not affect the rate of ear-hole closure from D5 to D30 (B), or ear-hole area at D64 (C). (D-F) Mason's trichrome stain on D10 ear tissue from control (D) and treatment (E) showed 20% of the distal injury plane from treated ears were not re-epithelialized, while 100% of control treated ears and the proximal injury plane were re-epithelialized (F). (G-I) Picro-sirius stain on D64 ear tissue from control (G) and treatment (H) showed no difference in the total collagen fibrotic area distal to the amputation plane (I). N.S. indicates not significant for comparison between groups. Data represent mean $\pm$ S.E.M. and the lines represent cubic regression for $n=10$ per treatment (B), OR individual data (red dots) and median and interquartile range for $n=10$ per treatment (C, I).



Supplemental Figure 1 Comparison of parallelism for cytokine assays demonstrates ability to quantify cytokines across three species. Serial dilution of serum (triangles, solid lines) and tissue lysate (squares, dashed lines) shows similar negative slopes for *M. musculus* (green), *A. percivali* (red) and *A. cahirinus* (blue) compared to the standard (black, circles, solid line). Data represent the mean for duplicates and lines are linear regressions calculated from all data that was above the lower limit of detection.

Supplemental Figure 1a

IL-1α

MAFFT Alignment

Homo	1	MAKVPMFEDLKNCYSENEEDSSSIDHLSLNQKSFYHVSYGELHEGCMQDSVSLSISETS
Mus	1	MAKVPDLFEDLKNCYSENEEYSSAIDHLSLNQKSFYDASYGSLHETCTDQFVSLRTSETS
Rattus	1	MAKVPDLFEDLKNCYSENEEYSSAIDHLSLNQKSFYDASYGSLHENCTDKFVSLRTSETS
Acomys	1	MAKVPDLFEDLKNCYSENEEYSSAIDHLSLNQKSFYDASYGSGPEHCTAKLVSRASETS
Homo	61	KTSKLTfKESMVVV---ATNGKVLKKRRLSLSOSITDDDLAIAANDSEEEIHKPRSAFSS
Mus	61	KMSNFTFKESRVTVSATSSNGKILKKRRLSFSETFEDDLQSTHD-LEETIQPRSAFYT
Rattus	61	KMSNFTFKESRVTVSATSNKGKILKKRRLSFNQPFTEDDLAIAHD-LEETIQPRSAFHS
Acomys	61	KKSNTFKESLVTVSATANEKGKVLKKRRLSFNQAFDEDDLEAIAHN-LEETIQPRSAFYS
Homo	118	FLSNVKYNFMRIIKYEFILNDALNQSIIRANDQ-YLTAAALHNLDEAVKFDMGAYKSSKD
Mus	120	YQSDLRKLMKLVQKFVMNDSLNTIYQDVDKHYLSTTWLNDLQQEVKFDMYAYSSGGD
Rattus	120	FQNNLRKLIIRIVKQEFIMNDSLQNIIYVDMDRTHLKAASLNDLQLEVKFDMYAYSSGGD
Acomys	120	FQHNVRKLIIRIIRKQEFILNDPLNQNIYLDPNVHLKAASLTDLQHEVKFDMYAYSS-GD
Homo	177	DAKITVILRISKTLQLYVTAQDEDQPVLLKEMPEIPKTIITGSETNLIFFWETHGKKNYFTS
Mus	180	DSKYPVTLKISDSQLFVSAQGEDQPVLLKEIPETPKLITGSETDLIFFWKSINSKNYFTS
Rattus	180	DSKYPVTLKISNTQLFVSAQGEDKPVLLKEIPETPKLITGSETDLIFFWEKINSKNYFTS
Acomys	179	DSKYPVTLKISNTQLFVSAQGEDQPVLLKEMPEIPKLIITGSETNLIFFWKSINSKNYFTS
Homo	237	VAHPNLFIAATKQDYWVCLAGGPPSTTDFQILENQA
Mus	240	AAYPELFIAATKEQSRVHLARGLPSTDFQI----S
Rattus	240	AAYPELFIATKEQSQVHLARGLPSTDFQI----S
Acomys	239	AAYPELFIAATKEQSRVHLARGLPSTDFQI----A

Supplemental Figure 1b

IL-1 β

MAFFT Alignment

Homo	1	MAEVP	ELASE	MMAY	YSGN	EDDL	FF	EADG	PKQ	MKCS	FQDL	DLCP	LDGG	IQLR	ISDH	YSKG
Mus	1	MATVP	ELNCE	MPF	DS-	DEND	LF	FEVD	GPQ	KMG	CFQT	FDLG	CPDE	SIQL	QISQ	QHINKS
Rattus	1	MATVP	ELNCE	IAAF	DS-	EEND	LF	FEAD	RPQ	KIK	DCFQ	ALDL	GCPD	ESIQL	QISQ	QHLDKS
Acomys	1	MATVP	ELNSE	VTAF	HS-	DKND	LF	FEVD	RPQ	KMK	SCLQ	TLDL	GS	PDESI	QLQIS	QHFNKS
Homo	61	FRQA	ASV	VVAM	DKL	RKML	VP	CP	QTF	QEND	LST	FF	PF	FIFEE	EPIFF	DTWDNEA--YVHDAP
Mus	60	FRQAV	SLIV	AVEK	LWQL	PVSE	FP	WTF	QDE	DM	ST	FFS	FIFEE	EPI	LCDS	WDDDDNLLVCDVP
Rattus	60	FRKAV	SLIV	AVEK	LWQL	PMSC	PWS	FQ	DE	DP	ST	FFS	FIFEE	EPI	LCDS	WDDDDNLLVCDVP
Acomys	60	FRQV	VSLF	VAVE	KLWN	IPVA	CP	WTF	QDE	DL	GT	FFS	FIFEE	EH	ILSD	SWDDEQ--LVCDVA
Homo	119	VRSL	NCT	LRDS	QOKS	SLV	MSG	PYEL	KALH	LQGD	ME	QQV	VFS	MSFV	QGEES	NDKIPVALGL
Mus	120	IRQL	HYR	LRDE	QOKS	SLV	SDP	PYEL	KALH	LNQ	NIN	QQV	IFS	MSFV	QGEPS	NDKIPVALGL
Rattus	119	IRQL	HCRL	RDE	QOKS	CLV	SDP	CEL	KALH	LNQ	NIS	QQV	VFS	MSFV	QGETS	NDKIPVALGL
Acomys	118	IRQL	HCRL	RDE	QOKS	CLV	SDP	CEL	KALH	LN	GEN	NIN	QQV	VFS	MSFV	HGETSINKIPVALGL
Homo	179	KE	KNLY	LSCV	IKD	DK	PTL	QLES	VDP	KNY	P	KKK	MEK	RFV	NKIE	NNKLEFESAQFPNWI
Mus	180	KG	KNLY	LSCV	MKD	GT	PTL	QLES	VDP	KQY	P	KKK	MEK	RFV	NKIE	VKSKVEFESAQFPNWI
Rattus	179	KG	KNLY	LSCV	MKD	GT	PTL	QLES	VDP	KQY	P	KKK	MEK	RFV	NKIE	VKSKVEFESAQFPNWI
Acomys	178	KG	KNLY	LSCV	MKD	GK	PTL	QLES	VDP	KQY	P	KKK	MEK	RFV	NKIE	TEKSKVEFESAQFPNWI
Homo	239	STSQA	ENMP	VFL	GG	TKG	GQDI	T	DFT	M	Q	FVSS				
Mus	240	STSQA	EHKP	VFL	-	G	NNSG	QDI	T	DFT	M	ESVSS				
Rattus	239	STSQA	EH	RPVFL	-	G	NSNG	R	DIV	DFT	M	EPVSS				
Acomys	238	STSQA	EHKP	VFL	-	G	NNSG	QDI	V	DFT	M	ESVSS				

Supplemental Figure 1c

IL-2

MAFFT Alignment

Homo	1	MYRMQLLSCTIALSLALVTNSAPTSSST-----KKTQLQLEHLLLDLQMIIN
Mus	1	MYSMQLASCVTTLTLVLLVNSAPTSSSTSSSTAQAQQQQQQQQQQHLEQLLMDLQELLS
Rattus	1	MYSMQLASCVALTLVLLVNSAPTSSPA-----KETQQHLEQLLLDLQVLLR
Acomys	1	MYSMQLASCVALTLVLLVNSAPTSSST-----EETGQHLEQLLLDLQVLER

Homo	47	GINNYKNPKLTRMLTFKFYMPKKATELKHLQCLEEELKPLEEVINLAQSKNFHLR-PRDL
Mus	61	RMENYRNKLPRMLTFKFYLPKQATELKDLQCLEDELGPIRVLDLTQSKSFQLEDAENF
Rattus	47	GIDNYKNLKLPMMLTFKFYLPKQATELKHLQCLENELGALQSVLDLTQSKSFHLEDAGNF
Acomys	47	GINNYKNPKLPMMLRFKFYMPKATELKHLQCLEEELGALQSVLDLNQSKSFYLEDTGNF

Homo	106	ISNINIVIVLELKGSETTFMCEYADETATIVEFLNRWITFCQSIISTLT-
Mus	121	ISNIRVTVVKLKGSNTFECQFDDESATVVDFLRRWIAFCQSIISTSPQ
Rattus	107	ISNIRVTVVKLKGSENKFECQFDDEPATVVEFLRRWIAICQSIISTMTQ
Acomys	107	INNIRVTVVKLKGSNTFKCKFDDEPVTVVELLSRWIAFCQSAISTMIQ

Supplemental Figure 1d

IL-4

MAFFT Alignment

Homo	1	MGLTSQLLPPLFFLLACAGNFVHGKCDITLQEIICKTLNSITEQKTLCTELTVTDIFAAS
Mus	1	MGLNPQLVVILLFFLECTRSHIHGCD-KNHLREIIGILNEVTGEGTPCTEMDVPNVLTAT
Rattus	1	MGLSPHLAVTLFCFLICTGNGIHGCN-DSPLREIINTLNQVTEKGPCTEMFVPDVLAT
Acomys	1	MGLRPQLATVLLCFLACTGDYIHGHN-DTALKEIHTLNQVTEKGPCTEMVVPDVLAT
Homo	61	KNTTEKETFCRAATVLRQFYSHHEKDTRCLGATAQQFHRHKQLIRFLKRLDRNLWGLAGL
Mus	60	KNTTESELVCRASKVLRIFYLKHGK-TPCL-----KKNSSVLMELQRLFRFRCLDSS
Rattus	60	RNTTENELICRASRVLRKFYFPRDV-PPCL-----KNKSGVLGELRKLCRGVSGLNSL
Acomys	60	KNSTEKELLCRASRVLRKFYFPHEV-TLCL-----KNNPKVLKDLKKLSRGISSLYPL
Homo	121	NSCPVKEANQSTLENFLERLKIIMREKYSKCS---S
Mus	112	ISCTMNEKSTSLKDFLESLSIMQMDYS-----
Rattus	112	RSCTVNESTLTTLKDFLESLSIRGKYLQSCTSMS
Acomys	112	ESCTVNESSYTTLKDFLERLRIRVQKKYCQC-----

Supplemental Figure 1e

IL-5

MAFFT Alignment

Homo	1	M-RMLLHLSLLALGAAYVYAIPTTEIPTSAIVKETLALLSTHRTLLIANETLRIPVPVHKN
Mus	1	MRRMLLHLSVLTLSCL--VWATAMEIPMSTVVKETLTQLSAHRALLTSNETMRLPVPTHKN
Rattus	1	M-RMLLCLNVLTLSCL--VWAIAMEIPMSTVVKETLIQLSTHRALLTSNETMRLPVPTHKN
Acomys	1	M-RVLLHLSFELTLPCL--VWAVAMEIPMSAVVKETLIQLSAHRALLTSNETVRLPVPTHKN

Homo	60	HQLCTEEIFQGTIGTLESQTVQGGTVERLFKNLSLIKKYIDGQKKKCGEERRRVNQFLDYL
Mus	59	HQLCIGEIFQGLDILKNQTVRGGTVEMLFQNLSLIKKYIDRQKEKCGEERRRTRQFLDYL
Rattus	58	HQLCIGEIFQGLDILKNQTVRGGTVEIPFQNLSLIKKYIDGQKEKCGEERRTRHFLDYL
Acomys	58	HQLCIAEIFQGLDILKNQTARGGTVETLFQNLSLIKKYIDRQKEKCGEERRRTRQFLDYL

Homo	120	QEFLGVMNTEWITE-
Mus	119	QEFLGVMSTEWAMEG
Rattus	118	QEFLGVMSTEWAMEV
Acomys	118	QEFLGVISTEWTMDG

Supplemental Figure 1f

IL-6

MAFFT Alignment

Homo	1	MNSEFSTSAFGPVAEFS	LGILLVLPAAFP-APVPPGEDSKDVAAPHRQPLTSSERIDKQIRY
Mus	1	MKFLSARDFHPVAE-	LGLMLVTTTAFPTSQVRRGDF-TEDTTPNRPVYTTS-QVGGLITH
Rattus	1	MKFLSARDFQPVAE-	LGLMLITATAFPTSQVRRGDF-TEDTTHNRPVYTTS-QVGGLITY
Acomys	1	MKFLSARDLHPVVF-	LGLMLVTAAAFPTSQVRRGDLATADTTPNRPVYTTSQQVGGLVTN
Homo	60	ILDGISAIRKETCNKSNMCESSKE	ALAENNLNLPKMAEKDGCFOSGFNEETCLVKIITGL
Mus	58	VLWEIVEMRKELCNGNSDCMNND	DALAENNLKLPEIQRNDGCVQTGYNQEICLLKISSGL
Rattus	58	VLREILEMRKELCNGNSDCMNSD	DALSENNLKLPEIQRNDGCFQTGYNQEICLLKICSGL
Acomys	60	VLKEVLEMRKELCNGSSDCMNND	DALSENRLNFPVIQINDGCFETKHDWEICLLKITSGL
Homo	120	LEFEVYLEYLQNRFE-SSEEQARAVQ	MSTKVLIIQFLOKKAKNLDAITTPDPTTNASLLTK
Mus	118	LEYHSYLEYMKNNLKDNKKDKARV	LQRTETLIHIFNQEVKDLHKIVLPTPTSNALLTDK
Rattus	118	LEERFYLEFVKNNLQDNKKDKARV	IQSNTETLVHIFKQEIKDSYKIVLPTPTSNALLMEK
Acomys	120	LEYETIYLEYVKNNIQDSKKEKARV	IQISTQALINILRQEVKEPGKIISPGETSTALQMET
Homo	179	LQAQNOWLQDMTTHLILRSFKEFL	QSSIRALROM
Mus	178	LESQKEWLRTKTIQFILKSLEEFL	KVTLRSTRQT
Rattus	178	LESQKEWLRTKTIQLILKALEEFL	KVTMRSTRQT
Acomys	180	LKPQNEWLRTKITQLILKALEEFL	KDTMRSTRKS

Supplemental Figure 1g

IL-10

MAFFT Alignment

Homo	1	MHSSALLCCLVLLTGVRASPGQGTQSENSCTHFPGNLPNMLRLRLDAFSRVKTFQMKDQ
Mus	1	MPGSALLCCLLLLTGVRISRGQYSREDNNCTHFVPGQSHMLLELRTAFSQVKTFQTKDQ
Rattus	1	MPGSALLCCLLLLAGVKTSKGHSIRGDNNTCTHFVPSQTHMLRELRAAFSQVKTFQKKDQ
Acomys	1	MPGSALLCCLLLLAGVGTSRG-----EYNTCTHFAVSQTHMLRELRAAFSQVKTFQKRDQ
Homo	61	LDNILLKESLIEDFKGYLGCAALSEMIQFYLEEVMPQAENQDPDIKAHVNSLGENLKTLR
Mus	61	LDNILLTDSLMDQDFKGYLGCAALSEMIQFYLVEVMPQAEKHGPEIKEHLNSLGEKLTLR
Rattus	61	LDNILLTDSLMDQDFKGYLGCAALSEMIKFYLVVMPQAENHGPEIKEHLNSLGEKLTW
Acomys	56	LDNILLTDSLMDQDFKGYLGCAALSEMIKFYLVVMPQAENHGPEIKEHLNSLGEKLTLR
Homo	121	LRRLRRCHRFLPCENKSKAVEQVKNAFNKLQEKGYKAMSEFDIFINYTEAYMTMKIRN
Mus	121	MRLRRCHRFLPCENKSKAVEQVKSDFNKLQDQGVYKAMNEFDIFINCIEAYMTMKKS
Rattus	121	LRRLRRCHRFLPCENKSKAVEQVKNDFNKLQDQGVYKAMNEFDIFINCIEAYMTMKKN
Acomys	116	MRLRRCHRFLPCENKSKAVEQVKSDFNKLQEKGVYKAMSEFDIFINCIEAYMTMKKN

Supplemental Figure 1i

IL-12p40

MAFFT Alignment

```

Homo      1  MCHQQLVISWFSIVFLASPLVAIWELKKDQVYVVELDWYPDAPGEMVVLTCDTPEEDGITW
Mus       1  MCPQKLTISWFAIVLLVSPLMANWELEKDQVYVVEVDWTPDAPGETVNLTCDTPEEDDITW
Rattus    1  MCHQKLTISWFAIVLLVSPLMANWELEKDQVYVVEVDWRPDAPGETVTLTCDSPPEEDDITW
Acomys     1  MCHQKLTISWFAIVLLASPLMAIWELKDQVYVVEVDWSPGAPGESVALTCDTPEEDDITW

Homo      61  TLDQSSEVIGSGKTLTIQVKEFGDAGQYTCHKGGEVLSHSLLLLHKKEDGIWSTDILKDQ
Mus       61  TSDQRHGVIGSGKTLTITVKEFLDAGQYTCHKGGETLSHSHLLLHKKENGIWSTEILKN-
Rattus    61  TSDQRHGVIGSGKTLTITVREFLDAGQYTCHFGGETLSHSHLLLHKKENGIWSTEILKN-
Acomys     61  TSDQRIDFKEGSGKTVTFEVKEFIHAGQYTCHKGDETLSHSRLLLLHKKENGIWSTDILKD-

Homo      121  KEPKNTFLRCEAKNYSGRFTCWLLTTISTDLTFESVKSSRGSSDPQGVTCGAATLSAERV
Mus       120  --FKNKTFLKCEAPNYSGRFTCSWLVRNMDLKENIKSSSSSPDSRAVTCGMASLSAEKV
Rattus    120  --FKNKTFLKCEAPNYSGRFTCSWLVRNMDLKENIKSSSSSPDSRAVTCGRASLSAEKV
Acomys     120  --SKNKTFLRCEAPNYSGRFTCSWLAER-TDLKFSIKSGSSSPDSRAVTCGAASLSSEKV

Homo      181  RGDNKEYE-YSVEQCQEDSACPAAEESLPIEVMDAVHKLKYENYTSFFFIRDIKPDPPK
Mus       178  TLDQRDYEKYSVSCQEDVTCPTAEETLPIELALEARQQNKYENYSTSFFFIRDIKPDPPK
Rattus    178  TLDQRDYEKYSVSCQEDVTCPTAEETLPIELVLEAQQNKYENYSTSFFFIRDIKPDPPK
Acomys     177  RVDEQDFEQYSVSCQEDVACPTAEETLPIELVLDAEQNKYENYSTSFFFIRDIKPDPPK

Homo      240  NLQKPLKNSRQVEVSWEYPTIWSTPHSYFSLTFVQVQVQKSKREK-----
Mus       238  NLQKPLKNSRQVEVSWEYPTISWSTPHSYFSLKFFVRIQRKKEKMK-----ETEEGC
Rattus    238  NLQKPLKNSRQVEVSWEYPTISWSTPHSYFSLKFFVRIQRKKEKTK-----ETEEEC
Acomys     237  NVQVQKPLKND-EVEVSWEYPTISWSTPHSYFSLKFFVQIQCNKEKTKKKRKKTEVEIERKC

Homo      286  --KDRVFTDKTSATVICKRKNASTSVRAQDRYYSSSWSEWASVPCS-----
Mus       289  NQKGAFLEKTSSTEVQC-KGCNVCVQAQDRYYNSSCSKWACVPCRVR-----
Rattus    289  NQKGAFLEKTSSTAEVQC-KGANVCVQAQDRYYNSSCSKWTCVPCRGRS-----
Acomys     296  KQKEPFLVDKTSATVECKGAKVCVQAQDRYYNSSFSKWEVCPCKIQSQDATLEGKEVED

Homo      -----
Mus       -----
Rattus    -----
Acomys     355  NRKEKFHLMMEETPKAIFCLAFFQFLGS

```

Supplemental Figure 1j

IL-17

MAFFT Alignment

Homo	1	MIPGKTSLVSL--LLLLLSLEATVKAGITIPRNPGCPNSEDKNEPRTVMVNLNTHNR-NTN
Mus	1	MSPGRASSVSLMLLLLLSLAATVKAATIPQSSACPNTAKDFLQNVKVNLIKVFNSLGAK
Rattus	1	MSPRRIPSMCLMLLLLLNLEATVKAATIPQSSVCPNAEANNFLQNVKVNLIKVLNSLSSK
Acomys	1	MSLGKTSVSLLLLLLSLEAAVNAGTLVPQGVCPNTDTKSFLQDVKNLKIILNSFSPR
Homo	58	TNPKRSSDYYNRSTSPWNLHRNEDPERYPYSVIWEAKCRHLGCTNADGNVDYHMNSVPIQQ
Mus	61	VSSRRPSDYLNLRSTSPWTLHRNEDPDYPSVIWEAQCRHQRCVNAEGKLDHMHNSVLIQQ
Rattus	61	ASSRRPSDYLNLRSTSPWTLNRNEDPDYPSVIWEAQCRHQRCVNAEGKLDHMHNSVLIQQ
Acomys	61	VNSRRPSDYLNLRSTSPWTLHRNEDPDYPPVIWEARCRHQRCVNAEGKLDHMHNSVLIQQ
Homo	118	EILVLRREPPHCPNSFRLEKILVSVGCTCVTFIVHHVA
Mus	121	EILVLKREPESCPFTFRVEKMLVGVGCTCVASIVRQAA
Rattus	121	EILVLKREPEKCPFTFRVEKMLVGVGCTCVSSIVRHAS
Acomys	121	EILVLRREAEEKCPLSFRLEKMLVGVGCTCVSSIVRHVA

Supplemental Figure 1k

CCL2

MAFFT Alignment

Homo	1	MKVS	AVLL	CLLL	MTAA	FN	PQGL	AQPD	ALNV	PS	TCC	FTF	SS	KK	IS	LQ	RL	KS	YV	-IT	IS	RCP
Mus	1	MQVP	VMLL	GLL	ETV	AGW	SIHV	LAQP	DAVN	APLT	CCY	SFT	SK	MI	PM	SR	LE	SY	KR	IT	SS	RCP
Rattus	1	MQV	SVTL	LGL	ETV	AACS	SIHV	LSQP	DAVN	APLT	CCY	SFT	GK	MI	PM	SR	LE	NY	KR	IT	SS	RCP
Acomys	1	MQV	SAKLL	CLLL	TAA	ASSP	MLAQ	PD	SVTS	PERT	CCY	SFT	SK	RI	PL	QR	LE	SY	KR	IT	SS	KCP

Homo	60	QKAV	IFRT	KL	GKE	ICAD	PK	KEW	VQNY	MKHL	GRK	-----					AH	TL	KT	-----				
Mus	61	KEAV	VFVT	KL	KRE	V	CAD	PK	KEW	VQNY	IKNL	DRN	QMR	SEPT	TI	EK	TAS	AL	RS	SAP	LN	-VKL		
Rattus	61	KEAV	VFVT	KL	KRE	ICAD	PN	KEW	VQNY	IRKL	DQNR	SEPT	TVF	YKI	AST	LR	TS	SAP	LN	-VNL				
Acomys	61	KEA	IFVT	KL	KRE	ICAD	PTMD	WVQ	SYIQ	KLD	QNR	SEAT	AV	EK	TAS	SP	GS	SAS	LN	AANS				

Homo		-----																								
Mus	120	TR	KSE	AN	AS	T	-	T	F	S	T	T	S	S	T	S	V	G	V	T	S	V	T	V	N	
Rattus	120	TH	KSE	AN	AS	T	-	T	F	S	T	T	S	S	T	S	V	E	V	T	S	M	T	E	N	
Acomys	121	TH	KPS	AN	AS	T	A	T	E	P	T	A	T	S	S	T	S	V	G	V	T	S	V	T	V	N

Supplemental Figure 11

CCL3

MAFFT Alignment

Homo	1	M	Q	V	S	T	A	A	L	A	V	L	L	C	T	M	A	L	C	N	Q	V	L	S	A	P	L	A	A	D	T	P	T	A	C	C	F	S	Y	T	S	R	Q	I	P	Q	N	F	I	A	D	Y	F	E	T	S	S	Q	C	S	K
Mus	1	M	K	V	S	T	T	A	L	A	V	L	L	C	T	M	T	L	C	N	Q	V	F	S	A	P	Y	G	A	D	T	P	T	A	C	C	F	S	Y	-	S	R	K	I	P	R	Q	F	I	V	D	Y	F	E	T	S	S	L	C	S	Q
Rattus	1	M	K	V	S	T	A	A	L	A	V	L	L	C	T	M	A	L	W	N	E	V	F	S	A	P	Y	G	A	D	T	P	T	A	C	C	F	S	Y	-	G	R	Q	I	P	R	K	F	I	A	D	Y	F	E	T	S	S	L	C	S	Q
Acomys	1	M	K	V	P	T	A	A	L	A	V	L	L	C	T	M	A	L	C	N	Q	V	F	S	A	P	Y	G	A	D	T	P	T	A	C	C	F	S	Y	-	S	R	Q	I	P	R	K	F	I	V	D	Y	F	E	T	S	S	L	C	S	Q

Homo	61	P	S	V	I	F	L	T	K	R	G	R	Q	V	C	A	D	P	S	E	E	W	V	Q	K	Y	V	S	D	L	E	L	S	A
Mus	60	P	G	V	I	F	L	T	K	R	N	R	Q	I	C	A	D	S	K	E	T	W	V	Q	E	Y	I	T	D	L	E	L	N	A
Rattus	60	P	G	V	I	F	L	T	K	R	N	R	Q	I	C	A	D	P	K	E	T	W	V	Q	E	Y	I	T	E	L	E	L	N	A
Acomys	60	P	G	V	I	F	L	T	K	R	N	R	Q	I	C	A	D	P	K	E	T	W	V	Q	E	Y	I	T	D	L	E	L	N	A

Supplemental Figure 1m

CCL5

MAFFT Alignment

Homo	1	MKVS-AAALAVILTATALCAPASASPYSSDTPCCFAYIARPLPRAHIKEYFYTSGKCSN
Mus	1	MKIS-AAALTIIILTAAALCTPAPASPYGSDTPCCFAYLSLALPRAHVKEYFYTSSKCSN
Rattus	1	MKISTAASLTIVILVAAALCTPAPASPYGSDTPCCFAYLSLALPRAHVKEYFYTSSKCSN
Acomys	1	MKIS-AAALAVIVTAAATCAPASASPYGSDTAPCCFAYHSRVLPRDHVKEYFYTSSKCSN

Homo	60	PAVVFVTRKNRQVCANPEKKWVREYINSLEMS
Mus	60	LAVVFVTRRNQVCANPEKKWVQYINYLEMS
Rattus	61	LAVVFVTRRNQVCANPEKKWVQYINYLEMS
Acomys	60	LAVIFVTRRNQVCANPEKKWVRKYINYLEMK

Supplemental Figure 1n

CSF2

MAFFT Alignment

Homo	1	MWLQSLLLLGTVACSI	SAPARSPSPSTQ	PEHVNAIQEARLLNLSRDTAA---EMNETV
Mus	1	MWLQNLLFLGIVVYSL	SAPTRSPPTVTRPWKHV	EAIKEALNLLD---DMPV---TLNEEV
Rattus	1	MWLQNLLFLGIVVYSF	SAPTRSPNPVTRPWKHVD	AIKEALSLN---DMRALENEKNEVD
Acomys	1	MWLQSLFLSIVVCS	SAPTRSPVPVTRPWKHVD	AIKEALSLN---EMPV---TVTENV
Homo	58	EVISEMFDIQEPTCL	QTRLELYKQGLRGS	ITKLKGPLTMMASHYKQHC
Mus	55	EVVSNEFSFKKLT	CVQTRLKTFEQLGRGN	FTKLKGALNMTASYQTY
Rattus	58	DIISNEFSIQRP	TCVQTRLKLYKQGLRGN	LTKLNGALTMIAHYQTNC
Acomys	55	GVVSNDFSIQN	PTCVQTRLDLKQGLRGN	FTRLGDLTVIASHYKKNC
Homo	118	ITTESFKENLKDFL	LVI	PFDCWEPVQE
Mus	115	TTYADFDISLKT	FLTDIPFECKKPGQK	
Rattus	118	TTEDFIKNLKGFL	FDIPFDCWKPVQK	
Acomys	115	IIVYEFIEENLKGFL	MV	IPFDCWKSQK

Supplemental Figure 1o

IFN γ

MAFFT Alignment

Homo	1	MKYTSYILAFQLCIVLGS	LG	GCYCQDPYVKEAENLKKYFNAGHSDVADNGTLEFLGILKNWK			
Mus	1	MNATHCILALQLELMAVS-	GCYCHGTVIESLESLNNYFNSSGID-VEEKSLEFLDIWRNWQ				
Rattus	1	MSATRRVLVLQLCLMAIS-	GCYCQGTIIIESLESLKNYFNSSMDAMEGKSLLLDIWRNWQ				
Acomys	1	MNAKHCILALQFCLMAIY-	GCYCQGTVIEETTNLKEYF-ASSISVSNGEDLLHILRNWQ				
Homo	61	EE	SDRKIMQSQIISFYFKLFKNEKDDQSIQKS	VE	TIKEDMNVKFFNSNKKKRD	DFEKL	TN
Mus	59	KD	GDMKILQSQIISFYLRLEVLKDNQAISNNISVIESHLIT	TF	FSNSKAKKDAFMSIAK		
Rattus	60	KD	GNTKILESQIISFYLRLEVLKDNQAISNNISVIESHLIT	NF	FSNSKAKKDAFMSIAK		
Acomys	59	QD	GDTKTIDIQIISFYFKLFEALKGHKAIQRS	SD	TIRADLIANFFNNSEEKYN	GF	MRIAK
Homo	121	YS	VTDLNVQRKAIHELIQVMAELSPA	AKTGKRKRSQMLF	-----	RGRRA	SQ
Mus	119	FE	VNNPQVQRQAENELIRVVHQLLP	ESSLRKRKRSRC	-----		
Rattus	120	FE	VNNPQIQHKANLIRVIHQLSP	ESSLRKRKRSRC	-----		
Acomys	119	IE	VNDPQNQRKAINELVTVM	SHLSPKSKQRKRKRSRC	CF	GAGDRLNKNNPAS	TI

Supplemental Figure 1p

TNF α

MAFFT Alignment

Homo	1	MSTESMIRDVELAEEALPKKTGGPQGSRRCLFSLFSFLIVAGATTFLCLLHFGVIGPQR
Mus	1	MSTESMIRDVELAEEALPQKMGGFQNSRRCLCLSLFSFLLVAGATTFLCLLNFGVIGPQR
Rattus	1	MSTESMIRDVELAEEALPKKMGGLQNSRRCLCLSLFSFLLVAGATTFLCLLNFGVIGPNK
Acomys	1	MSTESMIRDVELAEEALPQKAGSPQNSRRCLCLSLFSFLLVAGATLFLCLLNFGVIGPQR
Homo	61	EEFPRLDLSLISPLAQ--AVRSSSRTPSDKPVAVHVVANPQAEGLQWLNRANALLANGV
Mus	61	DEKFPNGLPLISSMAQTTLRSSSQNSDDKPVAVHVVANHQVEEQLEWLSQRANALLANGM
Rattus	61	EEKFPNGLPLISSMAQTTLRSSSQNSDDKPVAVHVVANHQAEEQLEWLSQRANALLANGM
Acomys	61	EEKFPNGLPIIGSMAQTTLRSSSQNSDDKPVAVHVVANHQVEEQLEWLSRRANALLANGM
Homo	118	ELRDNQLVVPSEGLYLIIYSQVLFKGQGCPSSTHVLLTHTISRIVSYQTKVNLLSAIKSPC
Mus	121	DLKDNQLVVPADGLYLIVYSQVLFKGQGCPSYVLLTHTVSRFATSYQEKVNLLSAVKSPC
Rattus	121	DLKDNQLVVPADGLYLIIYSQVLFKGQGCPSYVLLTHTVSRFATSYQEKVSLLSAIAIKSPC
Acomys	121	DLKDNQLVVPSSDGLYLIVYSQVLFKGQGCPSYVLLTHTVSRFATSYEDKVNLLSAIKSPC
Homo	178	QRETPEGAEAKPWYEPIYLGGVFQLEKGDRLSAEINRPDYLDFAESGQVYFGTIAL
Mus	180	PKDTPEGAEALKPWYEPIYLGGVFQLEKGDQLSAEVNLPKYLDFAESGQVYFGVIAL
Rattus	180	PKDTPEGAEALKPWYEPMYLGGVFQLEKGDLLSAEVNLPKYLDITTESGQVYFGVIAL
Acomys	180	PKDTPEGAEALKPWYEPIYLGGVFQLEKGDRLSAEVNLPKYLDFAESGQVYFGVIAL

Supplemental Figure 1q

CXCL1

MAFFT Alignment

Homo	1	MARATLSAAPS	NPRL	LRVALL	LLLLVA	ASRR	AAGAP	LATE	ELRCQCLQT	LQ	GIHLKNIQSV	
Mus	1	M-----	IPATR	SLLCAA	---	LLL	LATSRL	ATGAPI	ANELRCQCLQT	M	AGIHLKNIQSL	
Rattus	1	M-----	VSATR	SLLCAA	---	LPV	LATS	RQATGAPV	ANELRCQCLQT	V	AGIHLKNIQSL	
Acomys	1	M-----	APAT	SPLLR	GT	LLLL	LLLAT	SPQATGAPV	ASELRCQCLQT	V	AGIHLKNIQSL	
Homo	61	KVKSP	GP	HC	AQTE	VIAT	LN	KGQK	AC	NPAS	PMVKKI	IEKMLKNGKSN
Mus	51	KVLPS	GP	HCTQ	TE	VIAT	LN	GREAC	LDPEAP	VQKIVQ	KMLK-GVPK	
Rattus	51	KVMPP	GP	HCTQ	TE	VIAT	LN	GREAC	LDPEAP	PMVQKIVQ	KMLK-GVPK	
Acomys	54	KVTTP	GP	HC	IQTE	VIAT	LN	GREAC	LDPEAP	MRKVVQ	KMLN-GISK	