A F&B manager wants to determine whether there is any significant difference in the diameter of the cutlet between two units. A randomly selected sample of cutlets was collected from both units and measured? Analyze the data and draw inferences at 5% significance level. Please state the assumptions and tests that you carried out to check validity of the assumptions.

Minitab File: Cutlets.mtw

A hospital wants to determine whether there is any difference in the average Turn Around Time (TAT) of reports of the laboratories on their preferred list. They collected a random sample and recorded TAT for reports of 4 laboratories. TAT is defined as sample collected to report dispatch.

Analyze the data and determine whether there is any difference in average TAT among the different laboratories at 5% significance level.

Minitab File: LabTAT.mtw

Sales of products in four different regions is tabulated for males and females. Find if male-female buyer rations are similar across regions.

	East	West	North	South
Males	50	142	131	70
Females	550	351	480	350

Ha

• All proportions are equal

1. Check p-value
2. If p-Value < alpha, we reject
Null Hypothesis

Buyer Ratio.mtw

TeleCall uses 4 centers around the globe to process customer order forms. They audit a certain % of the customer order forms. Any error in order form renders it defective and has to be reworked before processing. The manager wants to check whether the defective % varies by centre. Please analyze the data at 5% significance level and help the manager draw appropriate inferences

Minitab File: CustomerOrderForm.mtw