

Extensiones del modelo lineal: regresión local

Contents

1	Algoritmo	1
2	Estimacion del modelo	1
3	Prediccion	2
3.1	Predicción puntual	2
3.2	Intervalo de confianza	2
3.3	Intervalo de predicción	3

1 Algoritmo

- Para x_i , $i = 1, \dots, n$:
 - Se eligen un total de $k = s * n$ puntos alrededor de x_i .
 - Se ajusta un modelo de regresión lineal en x_i utilizando los k puntos.
 - Los modelos más utilizados son la recta y el polinomio de segundo orden.
 - El valor predicho en cada punto es $x_i \Rightarrow \hat{f}(x_i) = X\hat{\beta}$.
- El parámetro s controla la suavidad de la curva ($s \in [0, 1]$).
- Se pueden estimar otras funciones polinómicas distintas a la recta.

2 Estimacion del modelo

```
library(ISLR)
datos = Wage
datos = datos[datos$wage<250,]
```

```
m1 = loess(wage ~ age, data = datos, span = 0.2, degree = 2)
m2 = loess(wage ~ age, data = datos, span = 0.5, degree = 2)
summary(m1)
```

```
## Call:
## loess(formula = wage ~ age, data = datos, span = 0.2, degree = 2)
##
## Number of Observations: 2921
## Equivalent Number of Parameters: 16.27
## Residual Standard Error: 30.18
## Trace of smoother matrix: 17.99 (exact)
##
## Control settings:
##   span      : 0.2
##   degree    : 2
##   family    : gaussian
##   surface   : interpolate      cell = 0.2
##   normalize : TRUE
```


```
## parametric: FALSE
## drop.square: FALSE
```

3 Prediccion

3.1 Predicción puntual

```
age_grid = seq(from = min(datos$age), to = max(datos$age), by = 1)
# con loess hay que utilizar se = T, ya que interval = "" no funciona
yp1 = predict(m1, newdata = data.frame(age = age_grid), se = T)
yp2 = predict(m2, newdata = data.frame(age = age_grid), se = T)

plot(datos$age, datos$wage, cex = 0.5, col = "darkgrey", ylab = "wage (x 1000 $)", xlab = "age")
#
lines(age_grid, yp1$fit, col = "blue", lwd = 2)
lines(age_grid, yp2$fit, col = "red", lwd = 2)
#
legend(60, 200, legend = c("s = 0.2", "s = 0.5"), col = c("blue", "red"), lty = 1, lwd = 2)
```



3.2 Intervalo de confianza

Recordemos que:

$$x_p^T \hat{\beta} - t_{\alpha/2} se(x_p^T \beta) \leq (x_p^T \beta) \leq x_p^T \hat{\beta} + t_{\alpha/2} se(x_p^T \beta)$$

dónde

$$se(x_p^T \beta) = \hat{s}_R \sqrt{x_p^T (X^T X)^{-1} x_p}$$


Por tanto:

```

alfa = 0.05
yp11 = yp1$fit + qnorm(alfa/2)*yp1$se.fit # utilizamos la normal en lugar de la t-student
yp12 = yp1$fit + qnorm(1-alfa/2)*yp1$se.fit

plot(datos$age,datos$wage, cex = 0.5, col = "darkgrey", ylab = "wage (x 1000 $)", xlab = "age")
#
lines(age_grid, yp1$fit, col = "blue", lwd = 2)
lines(age_grid, yp11, col = "blue", lty = 3)
lines(age_grid, yp12, col = "blue", lty = 3)

```



3.3 Intervalo de predicción

$$x_p^T \hat{\beta} - t_{\alpha/2} \hat{s}_R \sqrt{1 + x_p^T (X^T X)^{-1} x_p} \leq y_p \leq x_p^T \hat{\beta} + t_{\alpha/2} \hat{s}_R \sqrt{1 + x_p^T (X^T X)^{-1} x_p}$$

```

sR = m1$s
alfa = 0.05
yp13 = yp1$fit + qnorm(alfa/2)*sqrt(sR^2 + yp1$se.fit^2)
yp14 = yp1$fit + qnorm(1-alfa/2)*sqrt(sR^2 + yp1$se.fit^2)

plot(datos$age,datos$wage, cex = 0.5, col = "darkgrey", ylab = "wage (x 1000 $)", xlab = "age")
#
lines(age_grid, yp1$fit, col = "blue", lwd = 2)
lines(age_grid, yp11, col = "blue", lty = 3)
lines(age_grid, yp12, col = "blue", lty = 3)
#
lines(age_grid, yp13, col = "red", lty = 3)
lines(age_grid, yp14, col = "red", lty = 3)

```

