

# Homework 1

● Graded

13 Hours, 59 Minutes Late

## Student

Javier Herrera

 [View or edit group](#)

## Total Points

24.5 / 30 pts

## Question 1

### Problem 1

 7.5 / 10 pts

✓ + 1.5 pts (a) First expression is correct

✓ + 1.5 pts (a) Second expression is correct

✓ + 1 pt (b) Correctly applied formula for the sum of first  $k$  integers:

$$\sum_{i=1}^k i = \frac{k(k+1)}{2}$$

$$\sum_{i=2}^{3n+1} i = \frac{1}{2}(3n+1)(3n+2) - 1$$

✓ + 2 pts (b) Correctly applied formula for the sum of first  $k$  squares

$$\sum_{i=1}^k i^2 = \frac{k(k+1)(2k+1)}{6}$$

$$\sum_{i=1}^{2n} 9i^2 = \frac{9}{6}(2n)(2n+1)(4n+1)$$

+ 1 pt (b) Partial: formula for the sum of first  $k$  squares, small mistake

✓ + 3 pts (b) Correct derivation of  $Z(n)$

$$Z(n) = 24n^3 + \frac{69}{2}n^2 + \frac{31}{2}n$$

+ 2 pts (b) Partial: Mostly correct derivation of  $Z(n)$ , small mistakes

+ 1 pt (b) Partial: Incomplete derivation of  $Z(n)$  with correct ideas, did not simplify, contains small mistakes

✓ + 1 pt (c) Correct answer, brief justification

+ 0 pts Incorrect/Not submitted

– 0.01 pts [Click here to replace this description.](#)

 – 2.5 pts Late submission, 10% off.

## Question 2

### Problem 2

10 / 10 pts

✓ + 0.5 pts (a) Correct Proof

✓ + 1.5 pts (b) Correct proof

+ 1 pt (b) Partial: correct approach, incomplete

✓ + 1 pt (c) Base Case

✓ + 1 pt (c) Correctly stated assumption and what to prove

✓ + 1 pt (c) Correctly used assumption

✓ + 3 pts (c) Correctly finished

+ 2 pts (c) Partial: the remaining part of the proof has a small error

+ 1 pt (c) Partial: remaining proof, correct idea

✓ + 0.5 pts (d) Correct definition (maybe implicitly)

✓ + 1 pt (d) Stated the values of  $c$  and  $n_0$  and drew the conclusion

✓ + 0.5 pts (d) Showing connection between parts d and c

+ 0 pts Incorrect/Not submitted

### Question 3

#### Problem 3

6 / 10 pts

✓ + 0.5 pts a) Correct value

✓ + 1 pt a) Correct justification for big-O

+ 0.5 pts a) Partial: justification for big-O, incomplete

✓ + 0.5 pts (a) Correct justification for big-Omega

✓ + 0.5 pts b) Correct value

+ 1 pt b) Correct justification for big-O

✓ + 0.5 pts b) Partial: justification for big-O, incomplete

+ 0.5 pts b) Correct justification for big-Omega

✓ + 0.5 pts c) Correct value

+ 1 pt c) Correct justification for big-O

✓ + 0.5 pts c) Partial: justification for big-O, incomplete

+ 0.5 pts c) Correct justification for big-Omega

✓ + 0.5 pts d) Correct value

+ 1 pt d) Correct justification for big-O

✓ + 0.5 pts d) Partial: justification for big-O, incomplete

+ 0.5 pts d) Correct justification for big-Omega

✓ + 0.5 pts e) Correct value

+ 1 pt e) Correct justification for big-O

✓ + 0.5 pts e) Partial: justification for big-O, incomplete

+ 0.5 pts e) Correct justification for big-Omega

+ 0 pts Incorrect/Not submitted

Question 4

Academic integrity

1 / 0 pts

✓ + 0.5 pts Matched questions

+ 0 pts Didn't match questions.

✓ + 0.5 pts AI submitted

– 1 pt AI is missing

+ 0 pts Missing group members in submission or both group members made separate submissions.

+ 0 pts Not Latex submission

Question assigned to the following page: [1](#)

## CS111 Fall'25 ASSIGNMENT 1

**Problem 1:** Let  $B(n)$  be the number of letters “B” printed by Algorithm PRINTBS below. Give the asymptotic estimate for function  $B(n)$ . Your solution *must* consist of the following steps:

- (a) First express  $B(n)$  using the summation notation  $\sum$ .
- (b) Next, derive a closed-form expression<sup>1</sup> for  $B(n)$ .
- (c) Finally, give the asymptotic value of  $B(n)$  using the  $\Theta$ -notation.

Show your work and include a justification for each step.

**Algorithm PRINTBS** ( $n$  : integer)  
  **for**  $i \leftarrow 2$  **to**  $3n + 1$  **do**  
    **for**  $j \leftarrow i + 1$  **to**  $2i$  **do** print(“B”)  
  **for**  $i \leftarrow 1$  **to**  $2n$  **do**  
    **for**  $j \leftarrow 1$  **to**  $(3i + 1)^2$  **do** print(“B”)

*Note:* If you need any summation formulas for this problem, you are allowed to look them up. You do not need to prove them, you can just state in the assignment when you use them.

**Solution 1:**

(a) **For**  $B_1(n)$ : The inner loop runs  $2i - (i + 1) + 1 = i$  times.

$$B_1(n) = \sum_{i=2}^{3n+1} i$$

**For**  $B_2(n)$ : The inner loop runs  $(3i + 1)^2 - 1 + 1 = (3i + 1)^2$  times.

$$B_2(n) = \sum_{i=1}^{2n} (3i + 1)^2$$

**Total**  $B(n)$ :

$$B(n) = \sum_{i=2}^{3n+1} i + \sum_{i=1}^{2n} (3i + 1)^2$$

(b) **For**  $B_1(n)$ : Using  $\sum_{k=1}^m k = \frac{m(m+1)}{2}$ :

$$\begin{aligned} B_1(n) &= \left( \sum_{i=1}^{3n+1} i \right) - 1 \\ &= \frac{(3n+1)(3n+2)}{2} - 1 \\ &= \frac{9n^2 + 9n + 2}{2} - \frac{2}{2} \\ &= \frac{9n^2 + 9n}{2} = \frac{9}{2}n^2 + \frac{9}{2}n \end{aligned}$$

---

<sup>1</sup>A closed-form expression is a formula that can be evaluated in some fixed number of arithmetic operations, independent of  $n$ . For example,  $3n^5 + n - 1$  and  $n2^n + 5n^2$  are closed-form expressions, but  $\sum_{i=1}^n i^2$  is not, as it involves  $n - 1$  additions.

Question assigned to the following page: [2](#)

**For  $B_2(n)$ :** Using  $\sum_{k=1}^m k^2 = \frac{m(m+1)(2m+1)}{6}$ ,  $\sum_{k=1}^m k = \frac{m(m+1)}{2}$ , and  $\sum_{k=1}^m 1 = m$ , with  $m = 2n$ .

$$\begin{aligned}
 B_2(n) &= \sum_{i=1}^{2n} (9i^2 + 6i + 1) \\
 &= 9 \sum_{i=1}^{2n} i^2 + 6 \sum_{i=1}^{2n} i + \sum_{i=1}^{2n} 1 \\
 &= 9 \left( \frac{2n(2n+1)(4n+1)}{6} \right) + 6 \left( \frac{2n(2n+1)}{2} \right) + 2n \\
 &= 3n(8n^2 + 6n + 1) + 6n(2n + 1) + 2n \\
 &= (24n^3 + 18n^2 + 3n) + (12n^2 + 6n) + 2n \\
 &= 24n^3 + 30n^2 + 11n
 \end{aligned}$$

**Total  $B(n)$ :**

$$\begin{aligned}
 B(n) &= B_1(n) + B_2(n) \\
 &= \left( \frac{9}{2}n^2 + \frac{9}{2}n \right) + (24n^3 + 30n^2 + 11n) \\
 &= 24n^3 + (30 + 4.5)n^2 + (11 + 4.5)n \\
 &= 24n^3 + 34.5n^2 + 15.5n \\
 &= 24n^3 + \frac{69}{2}n^2 + \frac{31}{2}n
 \end{aligned}$$

(c) The asymptotic growth of a polynomial is determined by its highest-degree term,  $24n^3$ . Therefore:

$$B(n) = \Theta(n^3)$$

**Problem 2:** Let  $f(n) = n4^n + 10n^2$ . Prove, directly from the definition, that  $f(n) = O(5^n)$ . Your solution *must* consist of the following steps.

- (a) Solve the inequality  $\frac{5}{4}x \geq x + 1$  to show that it holds for all reals  $x \geq 4$ .
- (b) Solve the inequality  $5x^2 \geq (x + 1)^2$  to show that it holds for all reals  $x \geq 4$ .
- (c) Use mathematical induction to prove that  $2 \cdot 5^n \geq n4^n + 10n^2$  for all integers  $n \geq 4$ . Here you can use the inequalities from parts (a) and (b).
- (d) Using the inequality from part (c), prove that  $f(n) = O(5^n)$ . You need to give a rigorous proof derived directly from the definition of the  $O$ -notation, without using any theorems from class. (First, give a complete statement of the definition. Next, show how  $f(n) = O(5^n)$  follows from this definition.)

**Solution 2:**

(a)

$$\begin{aligned}
 \frac{5}{4}x &\geq x + 1 \\
 \frac{1}{4}x &\geq 1 \\
 x &\geq 4
 \end{aligned}$$



Question assigned to the following page: [2](#)

The inequality holds if and only if  $x \geq 4$ . Thus, it holds for all reals  $x \geq 4$ .

(b)

$$\begin{aligned} 5x^2 &\geq x^2 + 2x + 1 \\ 4x^2 - 2x - 1 &\geq 0 \end{aligned}$$

The quadratic  $4x^2 - 2x - 1 = 0$  has roots at  $x = \frac{1 \pm \sqrt{5}}{4}$ . The positive root is  $x_2 \approx 0.809$ . Since  $4x^2 - 2x - 1$  is an upward-opening parabola, the inequality holds for all  $x \geq x_2$ . Since  $4 > 0.809$ , it holds for all reals  $x \geq 4$ .

(c) Let  $P(n)$  be the statement  $2 \cdot 5^n \geq n4^n + 10n^2$ .

**Base Case** ( $n = 4$ ):

$$\begin{aligned} 2 \cdot 5^4 &= 1250 \\ 4 \cdot 4^4 + 10 \cdot 4^2 &= 1024 + 160 = 1184 \end{aligned}$$

Since  $1250 \geq 1184$ , the base case holds.

**Inductive Step:** Assume  $P(k)$  is true for  $k \geq 4$ :  $2 \cdot 5^k \geq k4^k + 10k^2$ . We need to prove  $P(k+1)$ :  $2 \cdot 5^{k+1} \geq (k+1)4^{k+1} + 10(k+1)^2$ .

$$\begin{aligned} 2 \cdot 5^{k+1} &= 5 \cdot (2 \cdot 5^k) \\ &\geq 5(k4^k + 10k^2) \quad (\text{by IH}) \\ &= 5k4^k + 50k^2 \quad (*) \end{aligned}$$

The target RHS is  $(k+1)4^{k+1} + 10(k+1)^2 = 4(k+1)4^k + 10(k^2 + 2k + 1) = 4k4^k + 4 \cdot 4^k + 10k^2 + 20k + 10$ . We must show that  $5k4^k + 50k^2$  is greater than or equal to the RHS:

$$\begin{aligned} (5k4^k + 50k^2) - (4k4^k + 4 \cdot 4^k + 10k^2 + 20k + 10) \\ &= (k4^k - 4 \cdot 4^k) + (40k^2 - 20k - 10) \\ &= (k-4)4^k + (40k^2 - 20k - 10) \end{aligned}$$

Since  $k \geq 4$ : 1.  $(k-4)4^k \geq 0$ . 2.  $40k^2 - 20k - 10 > 0$  (since the polynomial is positive for  $k \geq 4$ , as shown in part b). Thus, the difference is  $\geq 0$ , which means  $5k4^k + 50k^2 \geq (k+1)4^{k+1} + 10(k+1)^2$ . Combining this with (\*),  $P(k+1)$  is true.

(d) We say that  $\mathbf{f}(\mathbf{n}) = \mathbf{O}(\mathbf{g}(\mathbf{n}))$  if there exist positive constants  $C$  and  $n_0$  such that:

$$|f(n)| \leq C|g(n)| \quad \text{for all } n \geq n_0$$

**Proof:** We have  $f(n) = n4^n + 10n^2$  and  $g(n) = 5^n$ . From part (c), we know that for all integers  $n \geq 4$ :

$$n4^n + 10n^2 \leq 2 \cdot 5^n$$

Since  $n \geq 4$ ,  $f(n)$  and  $5^n$  are positive, satisfying:

$$|n4^n + 10n^2| \leq 2|5^n| \quad \text{for all } n \geq 4$$

We choose the positive constants:

- $C = 2$
- $n_0 = 4$

Question assigned to the following page: [3](#)

The condition  $|f(n)| \leq C|g(n)|$  is satisfied for all  $n \geq n_0$ . Therefore,  $f(n) = O(5^n)$ .

---

**Problem 3:** Give asymptotic estimates, using the  $\Theta$ -notation, for the following functions:

- (a)  $3n^6 + 5n^3 - 3n^2 + 1$
- (b)  $4n^3 \log^3 n + 4n^2 \log^4 n + 2n^4$
- (c)  $3n^4 \log^5 n + 2n^3 \sqrt{n} \cdot \log^2 n + 2\sqrt{n^9}$
- (d)  $n^3 \sqrt{n^5} + n \cdot (1.2)^n + 4n^5 \log^3 n$
- (e)  $n^7 + n^3 \cdot 4^n + n^2 \cdot \left(\frac{9}{2}\right)^n$

For part (a), give a proof derived directly from the definition of the  $O$ -notation. For (b) - (e), justify your answers using the properties of asymptotic notations covered in class, including relations between the basic reference functions:  $n^b$ ,  $\log n$ , and  $c^n$ .

**Solution 3:**

(a)  $3n^6 + 5n^3 - 3n^2 + 1$

$$\Theta(n^6)$$

**Proof derived directly from the definition of the  $O$ -notation (for  $O(n^6)$ ).** We show  $f(n) = 3n^6 + 5n^3 - 3n^2 + 1 = O(n^6)$ . We need  $C, n_0 > 0$  such that  $|f(n)| \leq C|n^6|$  for  $n \geq n_0$ . For  $n \geq 1$ :

$$|3n^6 + 5n^3 - 3n^2 + 1| \leq 3n^6 + 5n^3 + 1$$

Since  $n^6 \geq n^3$  and  $n^6 \geq 1$  for  $n \geq 1$ :

$$3n^6 + 5n^3 + 1 \leq 3n^6 + 5n^6 + 1n^6 = 9n^6$$

By choosing  $\mathbf{C} = 9$  and  $\mathbf{n}_0 = 1$ , we satisfy the definition:  $f(n) = O(n^6)$ . (To fully show  $\Theta(n^6)$ , we also need to show  $\Omega(n^6)$ . For  $n \geq 1$ ,  $f(n) \geq 3n^6 + 5n^3 - 3n^2$ . Since  $n^3 \geq n^2$ ,  $f(n) \geq 3n^6$ . Choosing  $C' = 3, n'_0 = 1$  gives  $\Omega(n^6)$ .)

(b)  $4n^3 \log^3 n + 4n^2 \log^4 n + 2n^4$  The terms are polynomial/poly-logarithmic. The dominant term is determined by the highest polynomial degree.

- Term 1:  $\Theta(n^3 \log^3 n)$
- Term 2:  $\Theta(n^2 \log^4 n)$
- Term 3:  $\Theta(n^4)$

Since  $\mathbf{n}^4$  grows faster than any term of the form  $n^c \log^b n$  where  $c < 4$ , the third term is dominant.

$$\Theta(n^4)$$

(c)  $3n^4 \log^5 n + 2n^3 \sqrt{n} \cdot \log^2 n + 2\sqrt{n^9}$  First, rewrite the terms with clear polynomial exponents:

- Term 1:  $\Theta(n^4 \log^5 n)$
- Term 2:  $2n^{3.5} \log^2 n = \Theta(n^{3.5} \log^2 n)$
- Term 3:  $2n^{4.5} = \Theta(n^{4.5})$

Question assigned to the following page: [4](#)

The highest polynomial degree is  $n^{4.5}$ . This dominates  $n^4 \log^5 n$  because  $n^{4.5} = n^4 \cdot n^{0.5}$ , and the factor  $n^{0.5}$  grows faster than  $\log^5 n$ .

$$\Theta(n^{4.5}) \text{ or } \Theta(\sqrt{n^9})$$

(d)  $n^3 \sqrt{n^5} + n \cdot (1.2)^n + 4n^5 \log^3 n$  The terms are  $n^{5.5}$ ,  $n \cdot (1.2)^n$ , and  $4n^5 \log^3 n$ . \*\*Exponential growth dominates polynomial growth.\*\* Since  $1.2 > 1$ , the term  $n \cdot (1.2)^n$  is dominant.

$$\Theta(n(1.2)^n)$$

(e)  $n^7 + n^3 \cdot 4^n + n^2 \cdot \left(\frac{9}{2}\right)^n$  The function is dominated by the term with the \*\*largest exponential base\*\*. The bases are 4 and  $\frac{9}{2} = 4.5$ . Since  $4.5 > 4$ , the last term is dominant.

$$\Theta(n^2 \cdot \left(\frac{9}{2}\right)^n) \text{ or } \Theta(n^2 \cdot (4.5)^n)$$

**Academic integrity declaration.** The homework papers must include at the end an academic integrity declaration. In this statement each group member must summarize briefly, in their own words, their contributions to the homework. Each group member must also state that he/she verified and has full understanding of all solutions in the submission. In addition, you need to state all external help and resources you used.

**Submission.** To submit the homework, you need to upload the pdf file to Gradescope. If you submit for a group, you need to put all member names on the assignment and submit it as a group assignment.

**Reminders.** Remember that only L<sup>A</sup>T<sub>E</sub>X papers are accepted.

**Assignment 1 Declaration:** In this assignment, I used a small bunch of resources to help me write and answer these questions including:

- Asymptotic notation notes and slides from Canvas
- Quiz 1 samples
- Google Gemini AI to help me with how to format my LaTeX document neatly