Preoperative Atelectasis

Part 2: Descriptive characteristics and map

Javier Mancilla Galindo

2024-04-08

Table of contents

Setup	1
State of residence of participants	3
Distribution of numerical variables	5
Characteristics of participants Table 1	5
Package references	8

Setup

Packages used

```
if (!require("pacman", quietly = TRUE)) {
   install.packages("pacman")
}

pacman::p_load(
   tidyverse, # Used for basic data handling and visualization.
   table1, #Used to create table of descriptive characteristics of sample.
   RColorBrewer, #Color palettes for data visualization.
   gridExtra, #Used to arrange multiple ggplots in a grid.
   grid, #Used to arrange multiple ggplots in a grid.
```

```
rnaturalearth, #Used to extract geographical data to create maps.
rnaturalearthhires, #Used together with the prior package to create map.
sf, #Used together with the prior package to create map.
plotly, #Used together with prior two packages to create map.
reticulate, #Required together with plotly
flextable, #Used to export tables.
officer, #Used to export tables.
report, #Used to cite packages used in this session.
webshot2 #Required to render this document as pdf.
```

Session and package dependencies


```
R version 4.3.3 (2024-02-29 ucrt)
Platform: x86_64-w64-mingw32/x64 (64-bit)
Running under: Windows 11 x64 (build 22631)
Matrix products: default
locale:
[1] LC_COLLATE=Spanish_Mexico.utf8 LC_CTYPE=Spanish_Mexico.utf8
[3] LC_MONETARY=Spanish_Mexico.utf8 LC_NUMERIC=C
[5] LC_TIME=Spanish_Mexico.utf8
time zone: Europe/Berlin
tzcode source: internal
attached base packages:
[1] grid
                    graphics grDevices datasets utils
              stats
                                                                methods
[8] base
other attached packages:
 [1] remotes_2.5.0
                                   webshot2_0.1.1
 [3] report_0.5.8
                                  officer 0.6.5
 [5] flextable_0.9.5
                                 reticulate_1.35.0
 [7] plotly 4.10.4
                                   sf 1.0-15
 [9] rnaturalearthhires_1.0.0.9000 rnaturalearth_1.0.1
[11] gridExtra 2.3
                                   RColorBrewer 1.1-3
[13] table1_1.4.3
                                   lubridate_1.9.3
[15] forcats_1.0.0
                                   stringr_1.5.1
```

[17] dplyr_1.1.4	purrr_1.0.2
[19] readr_2.1.5	$tidyr_1.3.1$
[21] tibble_3.2.1	ggplot2_3.5.0
[23] tidyverse_2.0.0	pacman_0.5.1

State of residence of participants

Map generated with the accompanying script $Map_USA_Canada.R$

This map was built by partly using code adapted from contribution by cpsievert.



Distribution of numerical variables

Distributions were examined with the accompanying sourced function distribution_numerical_variables.R

Near normal distribution:

- Age: light tails
- height: heavy right tail, 4 outliers right
- hb: heavy tails, bilateral outliers
- hct: heavy tails, bilateral outliers
- leu: near normal, bilateral outliers
- neu absolute: heavy right tail, two right outliers
- linf_absolute: heavy right tail, bilateral outliers (more right)
- mon_absolute: heavy right tail, bilateral outliers (more right)
- platelets: two right outliers
- urea: four right outliers
- creatinine: three right outliers

Distribution not normal:

- Weight: right-skewed, outliers are verified observations of extreme weight.
- BMI: right-skewed, outliers are verified observations of extreme BMI.
- spo2_VPO: Left-skewed
- neu percent: left-skewed
- linf percent: right-skewed
- glucose: right-skewed
- mon_percent: observations around only 5 data points. Will not use this variable, only absolute monocytes will be used.
- altitude: distribution not clear as values are quite apart an concentrate around single states with differing mean altitudes. Will attempt to model a smooth term or categorical term in subsequent analyses.

Outcome variable:

- atelectasis_percent: Zero-inflated. Would be difficult to manage as categorical ordinal due to low number of patients in some categories. Will re-assess alongside subsequent analyses to decide.

Characteristics of participants

Table 1 generated with the accompanying sourced script table 1 arguments.R

Characteristics of participants are shown for the total sample and by obesity class category as defined by the World Health Organization:

- Class 1, BMI (30-35) kg/m2
- Class 2, BMI (35-40) kg/m2
- Class 3, BMI >40 kg/m2

Characteristics of participants according to BMI class are shown in ${\bf Table}~{\bf 1}.$

Table 1

		Class 1	Class 2	Class 3
	Total	Obesity	Obesity	Obesity
	(N=236)	(N=63)	(N=53)	(N=120)
Sex			, ,	
Woman	$214\ (90.7\%)$	60 (95.2%)	48~(90.6%)	106~(88.3%)
Man	22 (9.3%)	3(4.8%)	5 (9.4%)	$14\ (11.7\%)$
Age (years)				
Mean (SD)	40.3 (9.87)	42.1 (10.2)	40.8 (9.25)	39.1 (9.85)
Weight (kilograms (kg))				
Median [Q1, Q3]	111 [97.4, 130]	89.2 [84.5,	107 [102, 112]	128 [114, 142]
111001011 [0[1, 0[0]	111 [0,11, 100]	95.9]	10. [102, 112]	120 [111, 112]
Height (meters (m))		1		
Mean (SD)	1.67 (0.0822)	1.67 (0.0674)	1.69 (0.0856)	1.67 (0.0876)
Body mass index				
(kg/m^2)				
Median [Q1, Q3]	40.3 [34.6,	33.0 [31.5,	38.3 [36.6,	45.8 [42.4,
	46.0]	33.8]	39.1]	51.2]
Surgical procedure				
LBGS	$31\ (13.1\%)$	5(7.9%)	9~(17.0%)	17~(14.2%)
OAGB	5(2.1%)	1 (1.6%)	1 (1.9%)	3~(2.5%)
RYGB	6~(2.5%)	1 (1.6%)	1 (1.9%)	4 (3.3%)
SG	189 (80.1%)	53 (84.1%)	41~(77.4%)	95~(79.2%)
ARISCAT risk group				
Intermediate Risk	61~(25.8%)	18 (28.6%)	12~(22.6%)	$31\ (25.8\%)$
Low Risk	175~(74.2%)	45~(71.4%)	$41 \ (77.4\%)$	89 (74.2%)
CO-RADS				

	Total	Class 1 Obesity	Class 2 Obesity	Class 3 Obesity
CO-RADS 1	230 (97.5%)	62 (98.4%)	51 (96.2%)	117 (97.5%)
CO-RADS 2	6(2.5%)	1 (1.6%)	2(3.8%)	3~(2.5%)
Oxygen saturation				
(SpO2) (%)				
Median [Q1, Q3]	96.0 [93.0,	97.0 [95.0,	96.0 [94.0,	94.0 [92.0,
	97.0]	97.5]	97.0]	97.0]
Mean altitude (meters)				
Median [Q1, Q3]	519 [519, 806]	519 [382, 806]	519 [519, 885]	519 [519, 806]
Acute Myocardial				
Infarction				
No	$210 \ (89.0\%)$	62 (98.4%)	52 (98.1%)	96~(80.0%)
Yes	26 (11.0%)	1 (1.6%)	1 (1.9%)	24 (20.0%)
Hypertension				
No	$177 \ (75.0\%)$	53 (84.1%)	$40 \ (75.5\%)$	$84 \ (70.0\%)$
Yes	59 (25.0%)	$10 \ (15.9\%)$	13~(24.5%)	36 (30.0%)
Diabetes				
No	$211 \ (89.4\%)$	58 (92.1%)	48 (90.6%)	105~(87.5%)
Yes	25~(10.6%)	5(7.9%)	5 (9.4%)	15~(12.5%)
Asthma				
No	216 (91.5%)	56~(88.9%)	46~(86.8%)	$114 \ (95.0\%)$
Yes	20~(8.5%)	7 (11.1%)	7 (13.2%)	6 (5.0%)
COPD				
No	$228 \ (96.6\%)$	62 (98.4%)	53 (100%)	113~(94.2%)
Yes	8 (3.4%)	1 (1.6%)	0 (0%)	7(5.8%)
Obstructive sleep apnea				
No	203~(86.0%)	60 (95.2%)	50 (94.3%)	93~(77.5%)
Yes	$33\ (14.0\%)$	3(4.8%)	3(5.7%)	$27\ (22.5\%)$
Oxygen use				
No	206~(87.3%)	60~(95.2%)	50 (94.3%)	96~(80.0%)
Yes	$30\ (12.7\%)$	3(4.8%)	3(5.7%)	$24\ (20.0\%)$
CPAP				
No	203~(86.0%)	60 (95.2%)	50 (94.3%)	93~(77.5%)
Yes	$33\ (14.0\%)$	3(4.8%)	3(5.7%)	$27\ (22.5\%)$
Hypothyroidism				
No	213 (90.3%)	56 (88.9%)	50 (94.3%)	107 (89.2%)
Yes	23 (9.7%)	7 (11.1%)	3 (5.7%)	13 (10.8%)
Dyslipidemia	` '	, ,	` '	, ,
No	218 (92.4%)	59 (93.7%)	48 (90.6%)	111 (92.5%)
	, ,	` ,	,	, ,

	Total	Class 1 Obesity	Class 2 Obesity	Class 3 Obesity
Yes Antidepressants use	18 (7.6%)	4 (6.3%)	5 (9.4%)	9 (7.5%)
No Yes	142 (60.2%) 94 (39.8%)	37 (58.7%) 26 (41.3%)	33 (62.3%) 20 (37.7%)	72 (60.0%) 48 (40.0%)

NOTE: The **ASA** physical status variable has not been included in analyses since the updated version of ASA consulted in October 2023 classifies obesity (30<BMI<40) as ASA 2 and obesity (BMI 40) as ASA 3. The distribution of frequencies of ASA~obesity class in this dataset does not match such definition. This occurred since an outdated version of ASA that did not include obesity was likely used by clinicians when writing the preoperative assessment medical note:

		${\tt Class}$	1 Obesity	Class 2	Obesity	Class 3	Obesity
ASA	1		31		18		3
ASA	2		30		34		84
ASA	3		0		0		32

Package references

- Auguie B (2017). gridExtra: Miscellaneous Functions for "Grid" Graphics. R package version 2.3, https://CRAN.R-project.org/package=gridExtra.
- Chang W (2023). webshot2: Take Screenshots of Web Pages. R package version 0.1.1, https://CRAN.R-project.org/package=webshot2.
- Gohel D, Moog S (2024). officer: Manipulation of Microsoft Word and PowerPoint Documents. R package version 0.6.5, https://CRAN.R-project.org/package=officer.
- Gohel D, Skintzos P (2024). *flextable: Functions for Tabular Reporting*. R package version 0.9.5, https://CRAN.R-project.org/package=flextable.
- Grolemund G, Wickham H (2011). "Dates and Times Made Easy with lubridate." *Journal of Statistical Software*, 40(3), 1-25. https://www.jstatsoft.org/v40/i03/.
- Makowski D, Lüdecke D, Patil I, Thériault R, Ben-Shachar M, Wiernik B (2023). "Automated Results Reporting as a Practical Tool to Improve Reproducibility and Methodological Best Practices Adoption." CRAN. https://easystats.github.io/report/.
- Massicotte P, South A (2023). rnaturalearth: World Map Data from Natural Earth. R package version 1.0.1, https://CRAN.R-project.org/package=rnaturalearth.
- Müller K, Wickham H (2023). *tibble: Simple Data Frames*. R package version 3.2.1, https://CRAN.R-project.org/package=tibble.
- Neuwirth E (2022). *RColorBrewer: ColorBrewer Palettes*. R package version 1.1-3, https://CRAN.R-project.org/package=RColorBrewer.

- Pebesma E, Bivand R (2023). Spatial Data Science: With applications in R. Chapman and Hall/CRC. doi:10.1201/9780429459016 https://doi.org/10.1201/9780429459016, https://r-spatial.org/book/. Pebesma E (2018). "Simple Features for R: Standardized Support for Spatial Vector Data." The R Journal, 10(1), 439-446. doi:10.32614/RJ-2018-009 https://doi.org/10.32614/RJ-2018-009.
- R Core Team (2024). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/.
- Rich B (2023). table1: Tables of Descriptive Statistics in HTML. R package version 1.4.3, https://CRAN.R-project.org/package=table1.
- Rinker TW, Kurkiewicz D (2018). pacman: Package Management for R. version 0.5.0, http://github.com/trinker/pacman.
- Sievert C (2020). Interactive Web-Based Data Visualization with R, plotly, and shiny. Chapman and Hall/CRC. ISBN 9781138331457, https://plotly-r.com.
- South A, Michael S, Massicotte P (????). rnaturalearthhires: High Resolution World Vector Map Data from Natural Earth used in rnaturalearth. R package version 1.0.0.9000, https://github.com/ropensci/rnaturalearthhires, https://docs.ropensci.org/rnaturalearthhires.
- Ushey K, Allaire J, Tang Y (2024). reticulate: Interface to 'Python'. R package version 1.35.0, https://CRAN.R-project.org/package=reticulate.
- Wickham H (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York. ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.
- Wickham H (2023). forcats: Tools for Working with Categorical Variables (Factors). R package version 1.0.0, https://CRAN.R-project.org/package=forcats.
- Wickham H (2023). stringr: Simple, Consistent Wrappers for Common String Operations. R package version 1.5.1, https://CRAN.R-project.org/package=stringr.
- Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Grolemund G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019). "Welcome to the tidyverse." *Journal of Open Source Software*, 4(43), 1686. doi:10.21105/joss.01686 https://doi.org/10.21105/joss.01686.
- Wickham H, François R, Henry L, Müller K, Vaughan D (2023). dplyr: A Grammar of Data Manipulation. R package version 1.1.4, https://CRAN.R-project.org/package=dplyr.
- Wickham H, Henry L (2023). purr: Functional Programming Tools. R package version 1.0.2, https://CRAN.R-project.org/package=purr.
- Wickham H, Hester J, Bryan J (2024). readr: Read Rectangular Text Data. R package version 2.1.5, https://CRAN.R-project.org/package=readr.
- Wickham H, Vaughan D, Girlich M (2024). *tidyr: Tidy Messy Data*. R package version 1.3.1, https://CRAN.R-project.org/package=tidyr.