

Analysis of the current trigger and selection efficiency for the process $H \rightarrow J/\psi \gamma \rightarrow \mu\mu\gamma t$ CMS and comparison with new trigger proposals

Javier Mariño Villadamigo

Research Activities

Academic year: 2021/2022

Università degli studi di Padova

October 25, 2022

1222-2022
800
ANNI



UNIVERSITÀ
DEGLI STUDI
DI PADOVA

Outline

- 1 Introduction
- 2 Objectives
- 3 Trigger and selection efficiencies
- 4 Results
- 5 Conclusions

Outline

- 1 Introduction
- 2 Objectives
- 3 Trigger and selection efficiencies
- 4 Results
- 5 Conclusions

Introduction

A word on physics motivations

The Higgs boson is considered to be discovered in 2012, when a new particle with the properties predicted by Standard Model (SM) was observed by CMS and ATLAS collaborations. Yukawa couplings to first and second-generation quarks are still to be measured.

- Rare **exclusive decays** of the Higgs boson to mesons in association with a photon can be used to explore these couplings.
- $H \rightarrow J/\psi\gamma$ can be used to explore the Higgs boson **coupling to the charm quark** \Rightarrow **test of SM predictions**.
- $Z \rightarrow J/\psi\gamma$ can be used as an **experimental benchmark**.

Introduction

- Both decays receive contributions from **direct** and **indirect** processes





Figure: Lowest order Feynman diagrams for the Z (or H) $\rightarrow J/\psi\gamma$ decay. The left-most diagram shows the direct and the remaining diagrams the indirect processes[†].

- The cumulative yield of the decay is **proportional to the branching ratio (BR)**.

[†]Sirunyan et al., “Search for rare decays of Z and Higgs bosons to J/ψ and a photon in proton-proton collisions at $\sqrt{s} = 13$ TeV”.

Introduction

- Experimental efforts have only established an **upper bound** 200 times the SM predicted value.
- The main challenge is the **discrimination** between background and signal \Rightarrow higher luminosity and **more efficient trigger** needed.



[‡]Sirunyan et al., “Search for rare decays of Z and Higgs bosons to J/ψ and a photon in proton-proton collisions at $\sqrt{s} = 13$ TeV”.

Outline

- 1 Introduction
- 2 Objectives**
- 3 Trigger and selection efficiencies
- 4 Results
- 5 Conclusions

Objectives

Trigger

In order to filter the amount of data that is recorded from each collision at the LHC, triggers are used. Current High Level Trigger requires the presence of a muon and a photon exceeding 17 and 30 GeV in the final state.

Goal

Design a new trigger that can outperform *HLT_Mu17_Photon30*.

Outline

- 1 Introduction
- 2 Objectives
- 3 Trigger and selection efficiencies
- 4 Results
- 5 Conclusions

Trigger and selection efficiencies

The **trigger efficiency**:

$$\varepsilon_{\text{trigger}} = \frac{\# \text{ events passing HLT \& selection}}{\# \text{ events passing selection}}$$

The **selection efficiency**:

$$\varepsilon_{\text{selection}} = \frac{\# \text{ events passing selection}}{\# \text{ total events}}$$

Muons	Photons
nMuon ≥ 2	nPhoton ≥ 1
Opposite charges: μ^- , μ^+	-
$ \eta < 2.4$	$ \eta < 2.4$
Retain pair with min. $\Delta R = \sqrt{(\Delta\phi)^2 + (\Delta\eta)^2}$	-
$p_T^{\mu_1} > 10 \text{ GeV}$, $p_T^{\mu_2} > 5 \text{ GeV}$	$p_T^\gamma > 15 \text{ GeV}$
mediumId	mvaID_WP90
-	pixel_Seed = 0

Figure: Kinematic, charge and quality selection criteria imposed to the muons and photons in the final state.

Example calculation of ε for *HLT_Mu17_Photon30*

	<i>HLT_Mu17_Photon30</i>			Efficiencies (%)	
	# total events	# passing sel.	# passing HLT & sel.	Trigger	Selection
QCD	21 335 910	34	26	76(7)	0.00016(3)
Z	459 000	98 630	81 829	83.0(1)	21.49(6)
H	448 000	160 587	142 102	88.49(8)	35.85(7)

Figure: Example calculation of the trigger and selection efficiencies for the trigger *HLT_Mu17_Photon30* and selection cuts: $p_T^\mu > 18$ GeV and $p_T^\gamma > 32$ GeV.

Outline

- 1 Introduction
- 2 Objectives
- 3 Trigger and selection efficiencies
- 4 Results**
- 5 Conclusions

Muon1: $p_T > 15$ GeV, muon2: $p_T > 10$ GeV, photon: $p_T > 15$ GeV

	Selection efficiencies (%)	
	$p_T^{\mu 1} > 18$ GeV, $p_T^\gamma > 32$ GeV	$p_T^{\mu 1} > 15$ GeV, $p_T^{\mu 2} > 10$ GeV, $p_T^\gamma > 15$ GeV
QCD	$(1.6 \pm 0.3) \cdot 10^{-4}$	$(4.4 \pm 0.5) \cdot 10^{-4}$
Z	21.49 ± 0.06	20.55 ± 0.06
H	35.85 ± 0.07	28.65 ± 0.07

Figure: Selection efficiencies obtained for the selection cuts: $p_T^{\mu 1} > 15$ GeV, $p_T^{\mu 2} > 10$ GeV, $p_T^\gamma > 15$ GeV (right column) compared to the ones obtained for $p_T^{\mu 1} > 18$ GeV, $p_T^\gamma > 32$ GeV (left column).

- Efficiency of QCD **increases** while Z and Higgs efficiencies **decrease** \Rightarrow scheme is **discarded**.

Muon1: $p_T > 18$ GeV, photon: $p_T > 24$ GeV, $\Delta R(\mu - \mu) < 0.4$




Figure: Normalized histogram for angular separation ΔR between the two muons reconstructed in the final state.

Muon1: $p_T > 18$ GeV, photon: $p_T > 24$ GeV, $\Delta R(\mu - \mu) < 0.4$

	Selection efficiencies (%)	
	$p_T^{\mu 1} > 18$ GeV, $p_T^\gamma > 32$ GeV	$p_T^{\mu 1} > 18$ GeV, $p_T^\gamma > 24$ GeV, $\Delta R(\mu - \mu) < 0.4$
QCD	$(1.6 \pm 0.3) \cdot 10^{-4}$	$(1.4 \pm 0.3) \cdot 10^{-4}$
Z	21.49 ± 0.06	24.73 ± 0.06
H	35.85 ± 0.07	37.65 ± 0.07

Figure: Selection efficiencies obtained for the selection cuts: $p_T^{\mu 1} > 18$ GeV, $p_T^\gamma > 24$ GeV, $\Delta R(\mu - \mu) < 0.4$ (right column) compared to the ones obtained for $p_T^{\mu 1} > 18$ GeV, $p_T^\gamma > 32$ GeV (left column).

- QCD selection efficiency: -10%
- Z selection efficiency: $+15\%$
- H selection efficiency: $+5\%$

Muon1: $p_T > 15$ GeV, photon: $p_T > 20$ GeV, $2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$




Figure: Normalized histogram for invariant mass of the muon pair system reconstructed in the final state.

Muon1: $p_T > 15$ GeV, photon: $p_T > 20$ GeV, $2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$

	Selection efficiencies (%)	
	$p_T^{\mu 1} > 18 \text{ GeV}, p_T^\gamma > 32 \text{ GeV}$	$p_T^{\mu 1} > 15 \text{ GeV}, p_T^\gamma > 20 \text{ GeV}, 2 < m_{\mu\mu} < 4 \text{ GeV}$
QCD	$(1.6 \pm 0.3) \cdot 10^{-4}$	$(1.3 \pm 0.2) \cdot 10^{-4}$
Z	21.49 ± 0.06	27.42 ± 0.07
H	35.85 ± 0.07	38.75 ± 0.07

Figure: Selection efficiencies obtained for the selection cuts: $p_T^{\mu 1} > 15 \text{ GeV}, p_T^\gamma > 20 \text{ GeV}, 2 < m_{\mu\mu} < 4 \text{ GeV}$ (right column) compared to the ones obtained for $p_T^{\mu 1} > 18 \text{ GeV}, p_T^\gamma > 32 \text{ GeV}$ (left column).

- QCD selection efficiency: **-25%**
- Z selection efficiency: **+28%**
- H selection efficiency: **+8%**

Muon1: $p_T > 10$ GeV, muon2: $p_T > 5$ GeV, photon: $p_T > 15$ GeV,
 $2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$

	Selection efficiencies (%)	
	$p_T^{\mu 1} > 18 \text{ GeV}, p_T^{\gamma} > 32 \text{ GeV}$	$p_T^{\mu 1} > 10 \text{ GeV}, p_T^{\mu 2} > 5 \text{ GeV},$ $p_T^{\gamma} > 15 \text{ GeV}, 2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$
QCD	$(1.6 \pm 0.3) \cdot 10^{-4}$	$(4.5 \pm 0.5) \cdot 10^{-4}$
Z	21.49 ± 0.06	29.83 ± 0.07
H	35.85 ± 0.07	39.81 ± 0.07

Figure: Selection efficiencies obtained for the selection cuts: $p_T^{\mu 1} > 10$ GeV, $p_T^{\mu 2} > 5$ GeV, $p_T^{\gamma} > 15$ GeV, $2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$ (right column) compared to the ones obtained for $p_T^{\mu 1} > 18$ GeV, $p_T^{\gamma} > 32$ GeV (left column).

- QCD selection efficiency: **+181%**
- Z selection efficiency: **+38%**
- H selection efficiency: **+11%**


Muon1: $p_T > 10$ GeV, muon2: $p_T > 5$ GeV, photon: $p_T > 22.7$ GeV,
 $2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$

	Selection efficiencies (%)	
	$p_T^{\mu 1} > 18 \text{ GeV}, p_T^{\gamma} > 32 \text{ GeV}$	$p_T^{\mu 1} > 10 \text{ GeV}, p_T^{\mu 2} > 5 \text{ GeV},$ $p_T^{\gamma} > 22.7 \text{ GeV}, 2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$
QCD	$(1.6 \pm 0.3) \cdot 10^{-4}$	$(1.6 \pm 0.3) \cdot 10^{-4}$
Z	21.49 ± 0.06	28.22 ± 0.07
H	35.85 ± 0.07	38.90 ± 0.07

Figure: Selection efficiencies obtained for the selection cuts: $p_T^{\mu 1} > 10$ GeV, $p_T^{\mu 2} > 5$ GeV, $p_T^{\gamma} > 22.7$ GeV, $2 \text{ GeV} < m_{\mu\mu} < 4 \text{ GeV}$ (right column) compared to the ones obtained for $p_T^{\mu 1} > 18$ GeV, $p_T^{\gamma} > 32$ GeV (left column).

- QCD selection efficiency: **+3%**
- Z selection efficiency: **+31%**
- H selection efficiency: **+9%**

Summary of results



Outline

- 1 Introduction
- 2 Objectives
- 3 Trigger and selection efficiencies
- 4 Results
- 5 Conclusions**

Conclusions

- **Comparison** between the standard selection associated to *HLT_Mu17_Photon30* and other possibilities has been made.
- Increases on efficiency of the Higgs boson signal are **very limited**.
- Selections including criteria beyond purely kinematical cuts are more efficient.
- Rising slightly the photon momentum threshold decreases QCD efficiency.
- 3-4 selection criteria are more efficient than the current one.
- The trigger efficiency corresponding to these cuts should be obtained in dedicated simulations.

References



Sirunyan, Albert M et al. “Search for rare decays of Z and Higgs bosons to J/ψ and a photon in proton-proton collisions at $\sqrt{s} = 13$ TeV”. In: *The European Physical Journal C* 79.2 (2019), pp. 1–27.



Thanks for your attention!