Cosmic Watch: a live, explorable observatory that applies computer vision and machine learning to public astronomy data streams

Jay Gupta[a,*]

[a] *SVKM's NMIMS Mukesh Patel School of Technology Management & Engineering, Mumbai, India*

Abstract

Public astronomy data now arrive within minutes to days of being taken, but they are spread across many archives, formats and interfaces. We present Cosmic Watch, an open-source Python and WebGL system that continuously collects newly released data from thirteen public feeds (Rubin and ZTF alerts, JWST and Hubble images, solar, space-weather, gravitational-wave, Mars rover and near-Earth-object data), analyses it with computer vision and neural networks, cross-matches the results against archival catalogues, and presents them in one interactive 3D environment. On calibrated JWST imaging the source detector controls purity with sign-flipped images and measures completeness by injecting synthetic galaxies: on a NIRCam F460M mosaic it recovered 12,029 sources with spurious rates of 0.24% and 0.42% and a 5σ point-source depth of AB 27.3. A galaxy morphology classifier fine-tuned from Zoobot reaches 88.7% ten-class accuracy on Galaxy10 DECaLS, and the catalogue matcher recovers 175 of 177 reference sources across 23 fields. We also audit the 744 anomaly candidates raised in the first twelve days of operation, show that most were stars, long-lived active nuclei, asteroids or relayed events, and describe vetting rules that retract 248 of 313 transient and outlier candidates. No discoveries are claimed. The software runs on a laptop, needs no API keys, and is released under the MIT licence.

Keywords: astronomical software, machine learning, galaxy morphology, time-domain astronomy, source detection, cross-matching

*ORCID: 0009-0002-7127-6146

Email address: jay4link@gmail.com (Jay Gupta)

1. Introduction

Most observational astronomy data are now public soon after they are taken. The Vera C. Rubin Observatory (Ivezić et al., 2019) and the Zwicky Transient Facility (Bellm et al., 2019) publish transient alerts in near real time through brokers such as ALeRCE (Förster et al., 2021); new JWST and Hubble observations appear in MAST; the Solar Dynamics Observatory (Pesnell et al., 2012; Lemen et al., 2012) and SOHO/LASCO (Brueckner et al., 1995) stream solar images; and space-weather, gravitational-wave and planetary-mission data are released through their own services. These data are spread across many archives, formats and interfaces. A student, educator or amateur astronomer who wants to follow the sky (what JWST released today, whether the Sun is flaring, which transients were flagged last night) must visit many services and has no easy way to run modern analysis on what they find. Brokers and archive tools excel at their own data type but are not built to put many data types side by side or to explain a result to a non-specialist.

Cosmic Watch addresses this with one reproducible pipeline that runs on a laptop with a consumer GPU and needs no API keys. It collects new data, analyses it with computer vision and trained neural networks, checks the results against archival catalogues, and presents everything in an interactive 3D environment (Fig. 1). A design principle throughout is a clear line between measurement and model output: for example, super-resolved pixels from SwinIR (Liang et al., 2021) or Real-ESRGAN (Wang et al., 2021) are always labelled as invented detail and never reach a detector or classifier. The system is aimed at teaching and outreach, at student and citizen-science projects that need a readable end-to-end example, and at quick-look triage of new releases and alerts.

This paper describes the design (Section 2), the validation of its main components (Section 3), an audit of the anomaly candidates it raised in its first days of operation together with the vetting rules that audit motivated (Section 4), and its limitations (Section 5).

2. System design

2.1. Data feeds and architecture

A FastAPI server and a scheduled pipeline write to a single SQLite database. Each source has a fetcher and each data type a detector. The thirteen live feeds are: Rubin and ZTF alerts via ALeRCE, new JWST




Figure 1: The running application, from a live screen recording. (a) Observatory overview: each feed and detector is a node. (b) Inspect mode for a JWST detection: CNN probabilities beside the real pixels. (c) SDO/AIA 171 Å with detected active regions and live NOAA space weather. (d) A Perseverance image classified by the Mars CNN, with rock candidates outlined.

and Hubble images from MAST, SDO solar images, SOHO/LASCO coronagraph data, NOAA space-weather products, gravitational-wave events from GraceDB, NASA GCN circulars, Mars rover images, and NASA/JPL near-Earth-object data. Every derived claim (a classification, a catalogue match, an anomaly flag) is stored together with the evidence that supports it, so that any result shown to a user can be traced back to its inputs.

2.2. Source detection on calibrated images

Deep-field sources are extracted with SEP (Barbary, 2016), the library form of SExtractor (Bertin and Arnouts, 1996). On calibrated JWST and HST mosaics detection runs in several passes: a cold pass for extended sources, a hot pass for compact sources, and binned low-surface-brightness passes. Each pass is also run on the sign-flipped image, in which any detection must be noise; detection thresholds are raised until the spurious rate measured in this way falls below 3%, and a pass that cannot meet that limit is dropped. Completeness is measured by injecting synthetic galaxies of known magnitude and size and recovering them with the same pipeline.

2.3. Classification models

The galaxy classifier is a ConvNeXt-nano (Liu et al., 2022) from Zoobot (Walmsley et al., 2023), pre-trained on Galaxy Zoo DECaLS (Walmsley et al., 2022) and fine-tuned on the ten-class Galaxy10 DECaLS data set distributed with astroNN (Leung and Bovy, 2019). A ResNet-18 (He et al., 2016) trained on the same data serves as a baseline. Rover images are classified by a ConvNeXt-Tiny trained on the MSL Curiosity image set of Lu and Wagstaff (2020). Solar images are searched for sunspots, active regions and coronal holes with classical image processing.

2.4. Active learning

Users can open any detection in a tiled deep-zoom view of the real pixels and label it. These labels, together with weak labels and an eleventh "not a galaxy" class, are used to train challenger models. A challenger replaces the current champion only if (i) its Galaxy10 accuracy falls by at most 1.5 percentage points, (ii) it rejects held-out stars and gas with at least 60% recall, and (iii) it rejects at most 10% of held-out galaxies.

2.5. Cross-matching

Detections are cross-matched against Gaia DR3 (Gaia Collaboration et al., 2023), Pan-STARRS1 (Chambers et al., 2016), SDSS, the DESI Legacy Imaging Surveys (Dey et al., 2019) and SIMBAD (Wenger et al., 2000). Matches are scored with a Bayesian likelihood ratio that compares the probability of the observed separation for a true counterpart with that for a chance alignment given the local source density, and Gaia proper motions are applied before matching.

2.6. Anomaly ranking and interface

An Isolation Forest (Liu et al., 2008) scores deep-field outliers, broker alerts and cross-survey verdicts provide transient candidates, and a Thompson-sampling bandit (Thompson, 1933) ranks findings for the user. The front end uses Three.js and OpenSeadragon, with optional in-browser hand-gesture control through MediaPipe (Lugaresi et al., 2019).

3. Validation

All numbers in this section come from model cards and result files in the repository, and a script in the repository (`paper/make_figures.py`) rebuilds every figure from them.




Figure 2: Galaxy CNN confusion matrix on the Galaxy10 DECaLS validation split, normalised by row.

3.1. Galaxy morphology classifier

On the 2,656-galaxy Galaxy10 validation split the classifier reaches 88.7% ten-class accuracy, 97.2% top-2 accuracy and 92.7% coarse-class accuracy, against 82.8%, 93.3% and 89.5% for the ResNet-18 baseline. The hardest classes are *Disturbed* (57% recall) and *Unbarred loose spiral* (78%; Fig. 2). Accuracy on JWST and HST imaging is expected to be lower because of the shift in resolution, wavelength and depth from the training images; Fig. 3 shows a public NIRCam image in which many small, faint sources receive merger or disturbed labels. Of three active-learning rounds, two challengers were correctly rejected by the promotion rules and one was promoted, with 88.1% Galaxy10 accuracy, 82% recall on held-out stars and gas, and 4.9% of real galaxies rejected.

3.2. Detection on calibrated data

On a JWST NIRCam F460M mosaic the detector found 12,029 sources, with spurious rates measured on the sign-flipped image of 0.24% and 0.42%




Figure 3: A public JWST NIRCam F150W image (ESO 350-IG038) processed by the pipeline. Right: the boxed region with each CNN-classified detection coloured by coarse class; the many merger/disturbed calls on small, faint sources show the domain shift from the training images.

for the cold and hot passes. The low-surface-brightness passes failed the purity test and were dropped automatically. The 5σ point-source depth is AB 27.3, and the 50% completeness limit falls from AB 25.6 for the smallest injected galaxies to AB 21.7 for the largest (Fig. 4).

3.3. Cross-matching

Using Gaia DR3's Pan-STARRS1 best-neighbour table as an answer key, the matcher recovered 175 of 177 sources in 23 fields spanning Galactic latitudes from $-74°$ to $+76°$, a precision of 98.9% and a recall of 100%.

3.4. Mars terrain classifier

The Mars classifier reaches 80.0% accuracy (89.0% top-3, 61.1% balanced) on the test split of Lu and Wagstaff (2020), which uses sols not seen in training.

4. Operation and audit of anomaly candidates

In its first twelve days of operation (30 October to 10 November 2026) the pipeline ingested 824 images, 1,870 Rubin and ZTF alerts and 46,455




Figure 4: Injection-recovery completeness on a calibrated JWST NIRCam F460M mosaic, by size class (half-light radius r_{50}).

detections, built 3,857 cross-survey objects from 29,191 archive matches, and raised 744 candidate anomalies.

These candidates are not discoveries, and an audit of them shows why (Table 1). Only 394 concern individual astronomical objects; the rest are solar-activity changes or relays of events announced by other facilities, and only 56 (7.5%) had been reviewed by a person. Of the 211 deep-field outliers, 96 are point sources and 26 more are rejected by the CNN as "not a galaxy", so the Isolation Forest mostly selected the brightest and largest sources. Of 29 broker tidal-disruption candidates, 23 were first detected two or more years earlier, which a flare lasting months cannot explain. Half of the "hostless" transients (8 of 16) are single-detection alerts that the broker classes as asteroids, and 29 of 57 supernova candidates carry an AGN, QSO or blazar class.

We therefore added vetting rules, implemented in one module with unit tests and shared by the live pipeline and the re-measurement: deep-field "normal" is defined over extended sources only; tidal-disruption candidates must be young and not identified as stars or AGN by the archives; a "new, hostless" verdict needs repeat detections, a successful check against known asteroids using JPL ephemerides, and a recent first detection; and supernova candidates must not carry an AGN-type broker class. Re-applied to the same data (Table 1), the rules retracted 248 of 313 transient and outlier candidates.

Table 1: Anomaly candidates raised from 30 October to 10 November 2026, before and after the vetting rules.

Candidate type	Raised	After vetting	Main reason for retraction
Deep-field outliers	211	33 (+51 newly selected)	star (96), CNN "not a galaxy" (28)
Tidal-disruption candidates	29	4	first detected years earlier (23)
Supernova candidates	57	28	broker class AGN/QSO/blazar (29)
"New, hostless" transients	16	0	broker asteroid (8), not new (6)
CNN mergers, nuclear transients	81	81	not yet vetted
Solar, relayed and other events	350	350	not object discovery
Total	**744**	**496 + 51**	56 had been reviewed by a person

In a visual inspection of 24 randomly chosen outliers before and after vetting, the fraction that were real galaxies rose from 9 of 24 to 14 of 24; the remaining contaminants are nebulae, satellite trails and detector artefacts. None of the surviving candidates has yet been confirmed as scientifically unusual. The next steps are adjudication against spectroscopic classifications from the Transient Name Server and a measured precision on a larger, independently vetted sample.

5. Limitations

Classifications are probabilistic, and on JWST and HST data they are applied outside the training domain. Detections made on preview (non-calibrated) images are indicative only. Rubin full frames are not public in real time, so Rubin coverage is limited to alerts. The anomaly candidates have not been validated against spectroscopic classifications. Cosmic Watch complements, and does not replace, the archives and brokers it draws on.

6. Conclusions

Cosmic Watch brings thirteen public astronomy data streams into one pipeline and one interactive environment, with measured purity and completeness on calibrated JWST data, a validated galaxy classifier and cata-

logue matcher, and an evidence trail for every claim. Its first days of operation also show that an unvetted anomaly stream is dominated by stars, long-lived active nuclei, asteroids and relayed events, and that simple, tested vetting rules remove most of them. The code, model cards and figure scripts are public so that every number in this paper can be reproduced.

Code availability

The source code is available under the MIT licence at `https://github.com/jay4linkh/cosmic-watch-ai`.

AI usage disclosure

Generative AI was used in this work. Anthropic's Claude (Claude Opus models, accessed through the Claude Code assistant in 2026) was used to help write, refactor and test parts of the software and its documentation, and to help draft and edit this manuscript. The author designed the system and chose the methods, ran every experiment, checked every number reported here against the repository's result files, and reviewed, modified and validated all AI-assisted code and text. The author takes full responsibility for the content of this paper.

Acknowledgements

This work uses public data and services from NASA, ESA, STScI/MAST, the Vera C. Rubin Observatory, ZTF and ALeRCE, LIGO, Virgo and KAGRA via GraceDB, NASA GCN, NASA/SDO and the AIA and HMI teams, SOHO/LASCO, NOAA SWPC, NASA/JPL-Caltech, SDSS, Pan-STARRS, ESA Gaia, CDS/SIMBAD, NED, the DESI Legacy Imaging Surveys and the NOIRLab Astro Data Lab, and builds on Zoobot and the Galaxy Zoo volunteers. This project is dedicated to the author's late mother.

References

Barbary, K., 2016. SEP: Source extractor as a library. Journal of Open Source Software 1, 58. doi:10.21105/joss.00058.

Bellm, E.C., et al., 2019. The Zwicky Transient Facility: System overview, performance, and first results. Publications of the Astronomical Society of the Pacific 131, 018002. doi:10.1088/1538-3873/aaecbe.

Bertin, E., Arnouts, S., 1996. SExtractor: Software for source extraction. Astronomy and Astrophysics Supplement Series 117, 393–404. doi:10.1051/aas:1996164.

Brueckner, G.E., et al., 1995. The Large Angle Spectroscopic Coronagraph (LASCO). Solar Physics 162, 357–402. doi:10.1007/BF00733434.

Chambers, K.C., et al., 2016. The Pan-STARRS1 surveys. arXiv:1612.05560.

Dey, A., et al., 2019. Overview of the DESI Legacy Imaging Surveys. The Astronomical Journal 157, 168. doi:10.3847/1538-3881/ab089d.

Förster, F., et al., 2021. The automatic learning for the rapid classification of events (ALeRCE) alert broker. The Astronomical Journal 161, 242. doi:10.3847/1538-3881/abe9bc.

Gaia Collaboration, Vallenari, A., et al., 2023. Gaia Data Release 3: Summary of the content and survey properties. Astronomy and Astrophysics 674, A1. doi:10.1051/0004-6361/202243940.

He, K., Zhang, X., Ren, S., Sun, J., 2016. Deep residual learning for image recognition, in: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition (CVPR), pp. 770–778. doi:10.1109/CVPR.2016.90.

Ivezić, Ž., et al., 2019. LSST: From science drivers to reference design and anticipated data products. The Astrophysical Journal 873, 111. doi:10.3847/1538-4357/ab042c.

Lemen, J.R., et al., 2012. The Atmospheric Imaging Assembly (AIA) on the Solar Dynamics Observatory (SDO). Solar Physics 275, 17–40. doi:10.1007/s11207-011-9776-8.

Leung, H.W., Bovy, J., 2019. Deep learning of multi-element abundances from high-resolution spectroscopic data. Monthly Notices of the Royal Astronomical Society 483, 3255–3277. doi:10.1093/mnras/sty3217. astroNN; the Galaxy10 DECaLS data set is distributed with astroNN.

Liang, J., Cao, J., Sun, G., Zhang, K., Van Gool, L., Timofte, R., 2021. SwinIR: Image restoration using Swin transformer, in: Proceedings of the IEEE/CVF International Conference on Computer Vision Workshops (ICCVW), pp. 1833–1844. doi:10.1109/ICCVW54120.2021.00210.

Liu, F.T., Ting, K.M., Zhou, Z.H., 2008. Isolation forest, in: 2008 Eighth IEEE International Conference on Data Mining, pp. 413–422. doi:10.1109/ICDM.2008.17.

Liu, Z., Mao, H., Wu, C.Y., Feichtenhofer, C., Darrell, T., Xie, S., 2022. A ConvNet for the 2020s, in: Proceedings of the IEEE/CVF Conference on Computer Vision and Pattern Recognition (CVPR), pp. 11976–11986. doi:10.1109/CVPR52688.2022.01167.

Lu, S., Wagstaff, K.L., 2020. MSL Curiosity Rover images with science and engineering classes. doi:10.5281/zenodo.4033453.

Lugaresi, C., et al., 2019. MediaPipe: A framework for building perception pipelines. arXiv:1906.08172.

Pesnell, W.D., Thompson, B.J., Chamberlin, P.C., 2012. The Solar Dynamics Observatory (SDO). Solar Physics 275, 3–15. doi:10.1007/s11207-011-9841-3.

Thompson, W.R., 1933. On the likelihood that one unknown probability exceeds another in view of the evidence of two samples. Biometrika 25, 285–294. doi:10.1093/biomet/25.3-4.285.

Walmsley, M., et al., 2022. Galaxy Zoo DECaLS: Detailed visual morphology measurements from volunteers and deep learning for 314 000 galaxies. Monthly Notices of the Royal Astronomical Society 509, 3966–3988. doi:10.1093/mnras/stab2093.

Walmsley, M., et al., 2023. Zoobot: Adaptable deep learning models for galaxy morphology. Journal of Open Source Software 8, 5312. doi:10.21105/joss.05312.

Wang, X., Xie, L., Dong, C., Shan, Y., 2021. Real-ESRGAN: Training real-world blind super-resolution with pure synthetic data, in: Proceedings of the IEEE/CVF International Conference on Computer Vision Workshops (ICCVW), pp. 1905–1914. doi:10.1109/ICCVW54120.2021.00217.

Wenger, M., et al., 2000. The SIMBAD astronomical database: The CDS reference database for astronomical objects. Astronomy and Astrophysics Supplement Series 143, 9–22. doi:10.1051/aas:2000332.