Gaia DR3 4277855016732107520: A Strong Dormant Black Hole Candidate from Astrometric Orbital Solution with an RGB Primary

Joel Ayala-Baez[1]⋆
[1] *Independent Researcher*

Accepted XXX. Received XXX; in original form XXX

ABSTRACT

We present a multi-wavelength analysis of Gaia DR3 4277855016732107520, an astrometric binary in the DR3 non-single-star (NSS) catalogue with a full three-dimensional orbital solution. Under the Gaia NSS pipeline assumptions, the companion mass is $M_2 = 12.32^{+1.73}_{-1.47}\,\mathrm{M}_\odot$ (median and 68 per cent credible interval from 5×10^5 Monte Carlo draws). The system comprises an RGB primary ($M_1 = 1.34 \pm 0.40\,\mathrm{M}_\odot$) orbiting the unseen companion with $P = 424.4\,\mathrm{d}$, $e = 0.34$, and $a = 2.64\,\mathrm{AU}$ at 657 pc. The 75.4σ NSS detection and 3753σ astrometric excess noise establish robust orbital photo-centre motion. The entire posterior lies above $5\,\mathrm{M}_\odot$; no luminous companion is detected, and ROSAT and XMM-Newton non-detections are consistent with a dormant system. Of seven non-BH scenarios evaluated, four are firmly excluded, two (artefact and chance alignment) are strongly disfavoured pending radial-velocity (RV) confirmation, and one (stripped He star) remains disfavoured pending UV spectroscopy. The predicted $K_1 \approx 65\,\mathrm{km\,s^{-1}}$ makes spectroscopic confirmation feasible with 2–4 m class facilities. Independent multi-epoch RV monitoring is required before this candidate can be considered a bona fide black hole. All analysis code and data are publicly available.

Key words: binaries: visual – stars: black holes – astrometry – stars: individual: Gaia DR3 4277855016732107520

1 INTRODUCTION

Theoretical models predict 10^8–10^9 stellar-mass black holes (BHs) in the Milky Way (Agol & Kamionkowski 2002; Shapiro & Teukolsky 1983), yet fewer than ~ 70 are dynamically confirmed, nearly all in X-ray binaries undergoing active accretion (Remillard & McClintock 2006). The vast majority are expected to reside in non-interacting (dormant) systems, producing no electromagnetic signature beyond gravitational influence on a luminous companion.

Gaia Data Release 3 (DR3; Gaia Collaboration et al. 2023a) has transformed the search for dormant BHs. The non-single-star (NSS) catalogue (Gaia Collaboration et al. 2023b) provides orbital solutions for $\sim 180\,000$ binaries, including astrometric orbits that — under the modelling assumptions of the pipeline — yield direct companion mass estimates rather than the minimum masses available from spectroscopic (SB1) solutions alone. This critical advantage eliminates the inclination degeneracy that limits radial-velocity surveys.

The landmark discoveries of Gaia BH1 ($M_2 \approx 9.6\,\mathrm{M}_\odot$; El-Badry et al. 2023a), Gaia BH2 ($M_2 \approx 8.9\,\mathrm{M}_\odot$; El-Badry et al. 2023b), and Gaia BH3 ($M_2 \approx 33\,\mathrm{M}_\odot$; Gaia Collaboration et al. 2024) have established the efficacy of astrometric BH detection. Systematic searches have since expanded the candidate list (Jayasinghe et al. 2021; Thompson et al. 2019; Shenar et al. 2022; Mahy et al. 2022). In a companion paper (Ayala-Baez 2026, in prep.; GRAVITAS OmniScan v13 catalogue) we identify 614 compact-object candidates from the Gaia DR3 NSS catalogue, 163 of which are GOLD-tier BH candidates.

In this paper we focus on one of these — Gaia DR3 4277855016732107520 — a GOLD-tier candidate ranked among the highest-priority targets for ground-based follow-up. The system is a wide binary ($P = 424\,\mathrm{d}$) comprising an RGB primary ($M_1 = 1.34\,\mathrm{M}_\odot$) and a dark companion whose mass from the full astrometric orbit is $M_2 \simeq 12.3\,\mathrm{M}_\odot$ (under Gaia NSS modelling assumptions). Located at 657 pc in Ophiuchus ($l = 32.1°$, $b = 9.3°$), the system benefits from moderate extinction, excellent parallax precision (10 per cent), and accessibility from both hemispheres.

The paper is structured as follows. Section 2 describes the observational data. Section 3 presents the analysis methodology. Section 4 reports results. Section 5 systematically tests non-BH explanations. Section 6 discusses implications and follow-up strategy, and Section 7 summarises our findings.

⋆ E-mail: jayalabaez@gmail.com

© 2026 The Authors

2 DATA

2.1 Gaia DR3 astrometry and NSS orbital solution

Gaia DR3 4277855016732107520 appears in both the main astrometric catalogue and the NSS orbital supplement with a full three-dimensional orbital solution (model type: `Orbital`). Key astrometric parameters include parallax $\varpi = 1.5228 \pm 0.1549$ mas (distance $d = 656.7$ pc), proper motion $(\mu_\alpha^*, \mu_\delta)$ as measured by Gaia, and photometry $G = 11.25$, $G_{\rm BP} = 11.66$, $G_{\rm RP} = 10.66$ ($G_{\rm BP} - G_{\rm RP} = 0.993$). The renormalised unit-weight error RUWE = 9.31 is extremely elevated, as expected for an unresolved binary with large photo-centre motion. The astrometric excess noise significance is 3753σ, among the highest in the NSS catalogue. While exceptionally high excess-noise values can in principle flag poorly resolved sources or pipeline failures, the combination of a coherent Keplerian orbital solution at 75σ significance, internally consistent period and eccentricity, and a well-behaved goodness-of-fit (18.3) together indicate that the excess noise reflects genuine photo-centre motion of a luminous primary orbiting a massive companion rather than an instrumental artefact.

The NSS orbital solution yields: period $P = 424.403 \pm 1.159$ d, eccentricity $e = 0.3427 \pm 0.0194$, and solution significance 75.4σ (far exceeding the threshold of 5 required for catalogue inclusion). Crucially, because this is an `Orbital` solution (full astrometric orbit), the companion mass $M_2 = 12.31\,{\rm M}_\odot$ reported in the Gaia `binary_masses` table is derived under specific modelling assumptions (see Section 3.1), rather than being a minimum value subject to the $\sin i$ degeneracy.

The Gaia DR3 GSP-Phot pipeline provides stellar parameters: $T_{\rm eff} = 5922$ K and $\log g = 2.93$, placing the primary firmly on the red giant branch. The systemic radial velocity is $v_{\rm sys} = -29.49 \pm 0.57\,{\rm km\,s^{-1}}$.

2.2 Multi-wavelength photometry

We compile multi-band photometry from 2MASS ($J = 9.94$, $H = 9.54$, $K_s = 9.41$; Cutri et al. 2003) and AllWISE ($W1 = 9.34$, $W2 = 9.36$; Cutri et al. 2014). GALEX (Bianchi et al. 2017) provides a near-UV detection (NUV = 17.52) but no far-UV counterpart, consistent with an RGB star without a hot companion.

2.3 X-ray archival data

We query the ROSAT All-Sky Survey (2RXS; Boller et al. 2016) and XMM-Newton serendipitous catalogue (4XMM-DR13; Webb et al. 2020) via VizieR and find no counterpart within 30″. Upper limits from typical ROSAT survey exposure times constrain the 0.1–2.4 keV X-ray luminosity to $L_X \lesssim 10^{30}\,{\rm erg\,s^{-1}}$ at the source distance — consistent with a non-accreting (dormant) compact companion.

2.4 SIMBAD classification

SIMBAD (Wenger et al. 2000) lists no spectral type or specific object classification for this source, underscoring the need for spectroscopic follow-up. The Gaia GSP-Phot parameters ($T_{\rm eff} = 5922$ K, $\log g = 2.93$) provide the primary characterisation.

3 METHODS

3.1 Gaia NSS mass derivation chain

The companion mass M_2 reported for this source originates from two distinct Gaia DR3 tables. The `gaiadr3.nss_two_body_orbit` table provides the astrometric orbital elements (period, eccentricity, angular semi-major axis, inclination, etc.) fitted to the along-scan residuals. The `gaiadr3.binary_masses` table then combines these elements with the parallax and an external primary-mass estimate to derive M_2 via Kepler's third law. Importantly, the primary mass entering the `binary_masses` derivation is itself model-dependent: it relies on the Gaia GSP-Phot atmospheric parameters ($T_{\rm eff}$, $\log g$, metallicity) and isochrone-based mass estimation. Thus, while the `Orbital` solution eliminates the $\sin i$ degeneracy inherent to spectroscopic (SB1) orbits — a critical advantage — the resulting M_2 is not a purely empirical "true mass" but depends on the modelling assumptions of the NSS pipeline, the parallax, and the adopted primary mass. We therefore quote the catalogue value $M_2 = 12.31\,{\rm M}_\odot$ as a *derived mass under Gaia modelling assumptions* and propagate the relevant uncertainties independently via Monte Carlo (Section 3.6).

Recent follow-up work by El-Badry et al. (2024) has shown that a subset of extreme-mass-ratio candidates in the Gaia NSS catalogue are spurious when tested against independent radial velocities, typically due to incorrectly resolved astrometric solutions. We note that the present source has *not* been flagged by any published vetting study, but caution that the same failure modes could in principle apply. Independent spectroscopic confirmation therefore remains essential (Section 6.3).

3.2 Orbital dynamics

Given the full astrometric orbit from the Gaia NSS pipeline (model `Orbital`), the derived companion mass M_2 follows once the orbital solution, parallax, and adopted primary-mass estimate are combined — rather than from a spectroscopic mass function. The total system mass is obtained via Kepler's third law:

$$M_{\rm tot} = M_1 + M_2 = \frac{4\pi^2 a^3}{GP^2}\,, \tag{1}$$

where a is the true semi-major axis derived from the angular semi-major axis and parallax. With M_1 estimated independently from the primary's position on the HR diagram, M_2 is then inferred.

3.3 Primary-mass estimation

The primary mass is estimated from the Gaia GSP-Phot parameters ($T_{\rm eff} = 5922$ K, $\log g = 2.93$). We note that GSP-Phot effective temperatures for RGB stars carry known systematic uncertainties of ∼100–200 K (Andrae et al. 2023), which propagate into the primary-mass estimate. Crucially, no independent spectroscopic classification currently exists for this source (SIMBAD lists no spectral type; Section 2.4), so the RGB identification rests entirely on the pipeline photometric parameters. We therefore treat the primary's evolutionary state as a *working assumption* rather than a spectroscopically confirmed fact. Under this assumption, the at-

mospheric parameters place the star on the RGB, consistent with an evolved star of $M_1 = 1.34 \pm 0.40\,\mathrm{M}_\odot$ (adopting a 30 per cent fractional uncertainty to account for age, metallicity, and systematic uncertainties).

3.4 Extinction analysis

The source lies at moderate Galactic latitude ($b = 9.3°$), where extinction is present but manageable. We derive the colour excess from the discrepancy between the observed $G_{\rm BP} - G_{\rm RP} = 0.993$ and the intrinsic colour for $T_{\rm eff} = 5922\,\mathrm{K}$ ($(G_{\rm BP} - G_{\rm RP})_0 \approx 0.76$):

$$E(G_{\rm BP} - G_{\rm RP}) = 0.993 - 0.76 = 0.233\,\mathrm{mag}\,, \tag{2}$$

yielding $E(B-V) = 0.175\,\mathrm{mag}$ and $A_V = 0.543\,\mathrm{mag}$ via the Cardelli et al. (1989) extinction law with $R_V = 3.1$. This modest extinction is consistent with the moderate latitude and ∼657 pc distance.

3.5 SED fitting

We construct the spectral energy distribution from 8 photometric bands (Gaia G, $G_{\rm BP}$, $G_{\rm RP}$; 2MASS J, H, K_s; WISE $W1$, $W2$) after applying band-dependent extinction corrections. The dereddened SED is compared to a single-temperature blackbody model at $T_{\rm eff} = 5922\,\mathrm{K}$. Residuals below 5 per cent in all bands confirm that the photometry is consistent with a single RGB star plus a dark (non-luminous) companion.

We note that a single-temperature blackbody is a first-order consistency check rather than a full spectral fit. Synthetic atmosphere models (e.g. PHOENIX, ATLAS9/MARCS; Husser et al. 2013; Castelli & Kurucz 2003) would provide a more rigorous assessment of the primary's SED shape and improve sensitivity to faint secondary contributions, particularly in the UV where a stripped helium star or hot subdwarf could contribute detectable flux even when overwhelmed in the optical. We strongly encourage such modelling in future work. However, for a companion whose expected luminosity exceeds the primary by a factor > 100 in every optical/IR band, the blackbody check suffices to demonstrate that no main-sequence companion of the required mass is present. It does not, by itself, exclude a hot, underluminous companion visible only in the UV.

3.6 Monte Carlo mass posterior

We draw $N = 500\,000$ Monte Carlo realisations of the companion mass, propagating uncertainties in:

(i) Parallax ϖ: Gaussian ($1.5228 \pm 0.1549\,\mathrm{mas}$).
(ii) Period P: Gaussian ($424.403 \pm 1.159\,\mathrm{d}$).
(iii) Primary mass M_1: log-normal prior centred on $1.34\,\mathrm{M}_\odot$ with $\sigma = 0.40\,\mathrm{M}_\odot$.
(iv) Model systematic: 10 per cent Gaussian.

Because M_2 is derived from the full astrometric orbit (not a minimum mass), no inclination marginalisation is needed — a significant advantage over SB1 analyses. However, the resulting mass depends on the assumptions and priors entering the Gaia NSS pipeline (Section 3.1). The fraction of draws with $M_2 > 5\,\mathrm{M}_\odot$ gives the posterior probability $P(\mathrm{BH})$.

3.7 Companion-light exclusion

For a hypothetical main-sequence companion of mass M_c, we compute the expected luminosity using empirical mass–luminosity relations ($L \propto M^{3.5}$) and compare the resulting flux contribution to the primary's luminosity ($L_\star = 22.0\,\mathrm{L}_\odot$). A companion is detectable if its flux exceeds 1 per cent of the primary in any band.

4 RESULTS

4.1 Orbital and dynamical solution

Table 1 summarises the observed and derived properties. The orbital solution is among the most significant in the DR3 NSS catalogue (75.4σ), and the extreme astrometric excess noise significance (3753σ) is consistent with genuine orbital photo-centre motion. We note that such extreme excess-noise values can in principle arise from poorly resolved blends or pipeline failures; however, the coherent Keplerian fit, the well-constrained period precision (0.27 per cent), and the goodness-of-fit (18.3) jointly argue against an instrumental origin. Independent RV confirmation is nevertheless essential before the companion mass can be regarded as secure.

The semi-major axis $a = 2.64\,\mathrm{AU}$ places the companion well outside the primary's Roche lobe ($R_L = 0.56\,\mathrm{AU}$; filling factor 3.7 per cent), confirming a fully detached, non-interacting system. The eccentricity $e = 0.34$ is tidally consistent: the circularisation timescale $\tau_{\rm circ} \sim 3 \times 10^{15}\,\mathrm{yr}$ exceeds the age of the Universe, so the orbit retains its natal eccentricity.

The predicted RV semi-amplitude is $K_1 \approx 65\,\mathrm{km\,s^{-1}}$, readily detectable with ground-based spectrographs (Section 6.3).

4.2 Stellar characterisation

The extinction analysis reveals modest reddening ($A_V = 0.54\,\mathrm{mag}$). The dereddened absolute magnitude $M_G = 1.68\,\mathrm{mag}$ and bolometric luminosity $L_\star = 22.0\,\mathrm{L}_\odot$ are consistent with an RGB star of $\sim 1.3\,\mathrm{M}_\odot$ with radius $R_\star = 4.5\,\mathrm{R}_\odot$. The dereddened SED is well-fit by a single-temperature model with residuals < 5 per cent in all bands. No secondary SED component is required.

4.3 Companion-light exclusion

A main-sequence star of $M_2 = 12.3\,\mathrm{M}_\odot$ would have $L \approx 6550\,\mathrm{L}_\odot$ and $T_{\rm eff} \approx 24\,200\,\mathrm{K}$, contributing $\sim 29\,750$ per cent of the primary flux — overwhelmingly detectable in every photometric band. We test luminous companions over the range 3–15 $\mathrm{M}_\odot$ and find that no MS star above $\sim 1.5\,\mathrm{M}_\odot$ can remain hidden in the photometry (Fig. 4).

The companion must therefore be optically faint at the level probed here.

4.4 Mass posterior and BH probability

The Monte Carlo posterior (Fig. 2) yields:

- Median companion mass: $\tilde{M}_2 = 12.32\,\mathrm{M}_\odot$
- 68 per cent CI: $[10.85, 14.05]\,\mathrm{M}_\odot$
- 90 per cent CI: $[9.99, 15.38]\,\mathrm{M}_\odot$

- Under adopted assumptions, the entire posterior lies above $5\,M_\odot$
- $P(M_2 > 10\,M_\odot) = 94.9$ per cent

A sensitivity analysis over primary mass (Table 3) shows that the posterior remains entirely above $5\,M_\odot$ for all tested M_1 values from 0.80 to $2.50\,M_\odot$. The BH interpretation is robust to the choice of primary mass prior, owing to the large companion mass inferred from the orbital solution.

5 ALTERNATIVE SCENARIOS

We systematically evaluate seven non-BH explanations for the dark companion. Table 2 summarises the results.

5.0.0.1 Main-sequence companion. A $12.3\,M_\odot$ MS star would have $L \sim 6550\,L_\odot$ and $T_{\rm eff} \sim 24\,200$ K, contributing ∼30 000 per cent of the primary flux. The SED shows no secondary component (Section 4.3). **Excluded.**

5.0.0.2 White dwarf. $M_2 = 12.3\,M_\odot$ exceeds the Chandrasekhar limit ($1.44\,M_\odot$) by a factor of 8.6. No white dwarf can have this mass. **Excluded.**

5.0.0.3 Neutron star. $M_2 = 12.3\,M_\odot$ exceeds the most generous NS mass ceiling ($\sim 2.3\,M_\odot$; Rezzolla et al. 2018) by a factor of 5.3. **Excluded.**

5.0.0.4 Hierarchical triple. Two $6.2\,M_\odot$ MS stars would have combined luminosity $\sim 1160\,L_\odot$ — easily detectable. The Mardling–Aarseth stability criterion (Mardling & Aarseth 2001) requires $P_{\rm inner} \lesssim 42$ d. No secondary spectral signature is detected. **Excluded.**

5.0.0.5 Stripped helium star. A $12.3\,M_\odot$ stripped He star would have $T_{\rm eff} > 50\,000$ K and $L \sim 10^4\,L_\odot$, producing strong UV and X-ray emission. At the source distance of 657 pc and with $A_V = 0.54$ mag ($A_{\rm NUV} \approx 2.2$ mag), such a companion would have an apparent NUV magnitude of ~ 9 — roughly 8.5 mag brighter than the observed GALEX NUV = 17.52, corresponding to a flux excess of $\sim 2700\times$. The faint observed NUV is therefore inconsistent with a luminous stripped He star, though a cooler or lower-luminosity subdwarf cannot be fully excluded. No X-ray source is detected in ROSAT/XMM. This scenario is **highly disfavoured** but not formally excluded without UV spectroscopy.

5.0.0.6 Astrometric artefact. The NSS significance (75.4σ), RUWE (9.31), astrometric excess noise significance (3753σ), and goodness-of-fit (18.3) collectively support the orbital solution as astrophysically robust. The period precision of 0.27 per cent rules out a spurious signal. Nevertheless, El-Badry et al. (2024) have shown that some high-significance NSS solutions fail independent RV checks; this scenario is therefore **strongly disfavoured** but formally pending RV confirmation.

5.0.0.7 Chance alignment. Gaia's orbital solution represents a coherent Keplerian signal over 424 d with 75σ significance. Chance alignments produce scattered astrometry, not phase-locked periodic motion. As above, we consider this **strongly disfavoured** pending independent RV verification.

Summary: Four of seven alternative scenarios are firmly excluded on dynamical, photometric, and astrometric grounds; two more (artefact and chance alignment) are strongly disfavoured pending independent RV data. The stripped helium star scenario is highly disfavoured but cannot be formally ruled out without UV spectroscopy (Section 6.3). A black hole of $\sim 12.3\,M_\odot$ remains the strongly preferred explanation.

6 DISCUSSION

6.1 Context among Gaia BH discoveries

Table 4 places this system alongside the three confirmed Gaia dormant BH systems. At $M_2 = 12.3\,M_\odot$, it would rank as the second most massive after Gaia BH3 ($33\,M_\odot$). Critically, it benefits from a full 3D astrometric orbit, eliminating the $\sin i$ ambiguity that affects purely spectroscopic candidates; under the Gaia NSS modelling assumptions, the inferred companion mass is not subject to the spectroscopic $\sin i$ lower-limit degeneracy.

The long period ($P = 424$ d) and moderate eccentricity ($e = 0.34$) place it in a similar orbital regime to Gaia BH1 ($P = 185.6$ d, $e = 0.45$) and Gaia BH2 ($P = 1277$ d, $e = 0.52$), consistent with a formation channel involving limited or no mass transfer.

6.2 Binary evolution and BH formation

The BH progenitor was a massive star with zero-age main-sequence mass $M_{\rm ZAMS} \gtrsim 25\,M_\odot$, based on the remnant mass (Fryer et al. 2012). The wide orbit ($a = 2.64$ AU) survived the supernova event, constraining the natal kick velocity to moderate values. The normal chemical composition of the RGB primary (consistent with no peculiar enrichment) suggests minimal mass transfer from the progenitor prior to collapse, favouring a direct-collapse or failed-supernova formation channel (Langer et al. 2020).

6.3 Follow-up strategy

The predicted RV semi-amplitude $K_1 \approx 65\,{\rm km\,s^{-1}}$ is remarkably large, making spectroscopic confirmation feasible with modest ground-based resources:

(i) **Radial-velocity monitoring**: At $G = 11.25$, the source is within reach of echelle spectrographs on 2–4 m class telescopes (HARPS, FEROS, CHIRON). We estimate that ∼6–8 RV epochs distributed over one orbital period (∼424 d) would confirm the period, semi-amplitude, and eccentricity to high precision. Even a single epoch would detect the $\sim 65\,{\rm km\,s^{-1}}$ amplitude.

(ii) **Spectroscopic classification**: High-resolution spectroscopy would yield independent $T_{\rm eff}$, $\log g$, metallicity, and chemical abundances.

(iii) **UV spectroscopy**: HST/COS or Swift/UVOT observations would definitively test the stripped He star scenario.

The target is located at RA = 18.28^h, Dec = $+3.5°$ (Ophiuchus), accessible from both hemispheres, and is an evening target in the northern summer.

We stress that radial-velocity follow-up is the single most important observational step for this system. Given the large predicted semi-amplitude ($K_1 \approx 65\,\mathrm{km\,s^{-1}}$) and the bright apparent magnitude ($G = 11.25$), even a single high-resolution spectrum could meaningfully test the Gaia astrometric solution; a complete RV orbit would either confirm or rule out the BH interpretation independently of the pipeline assumptions.

6.4 Limitations and required confirmation

We emphasise that several limitations prevent this system from being classified as a confirmed black hole at present:

- **Gaia NSS pipeline assumptions.** The companion mass depends on the NSS orbital model, the parallax, and the adopted primary mass (Section 3.1). El-Badry et al. (2024) have shown that a fraction of extreme-mass-ratio Gaia NSS solutions are spurious when tested against independent radial velocities. Although this source has not been flagged by any published vetting study, the same failure modes could in principle apply.
- **No independent multi-epoch RV orbit exists.** The orbital solution relies entirely on the Gaia astrometric pipeline. Until an independent RV orbit confirms the period, eccentricity, and semi-amplitude, the companion mass remains contingent on the Gaia solution.
- **SIMBAD provides no spectral classification.** The primary characterisation and its classification as an RGB star rest entirely on Gaia GSP-Phot parameters, which have known systematic uncertainties at the 100–200 K level for RGB stars. Until an independent spectrum confirms the evolutionary state and chemical abundances of the primary, this classification should be treated as an assumption.
- **The stripped He star scenario remains formally open**, requiring UV spectroscopy (e.g. HST/COS) for definitive exclusion.
- **Parallax uncertainty of** *sim***10 per cent** propagates into the semi-major axis and thence into M_2. Gaia DR4 will tighten this substantially.
- **Single-temperature SED check.** Our blackbody comparison establishes consistency with a single luminous source but does not substitute for a full atmospheric model fit.

7 CONCLUSIONS

We have presented a comprehensive multi-wavelength analysis of Gaia DR3 4277855016732107520, an astrometric binary with a full 3D orbital solution. Our principal findings are:

(i) The companion mass from our Monte Carlo posterior is $M_2 = 12.32^{+1.73}_{-1.47}\,\mathrm{M}_\odot$ (median; 90 per cent CI: $[10.0, 15.4]\,\mathrm{M}_\odot$), well above the BH threshold of $5\,\mathrm{M}_\odot$ under the adopted modelling assumptions.

(ii) Under the adopted priors, the Monte Carlo posterior gives $P(\mathrm{BH}) > 99.9$ per cent across all tested primary-mass configurations (0.8–2.5 $\mathrm{M}_\odot$).

(iii) No luminous companion is detected: a MS star of the required mass would exceed the photometric detection threshold by a factor of $\sim 30\,000$. The companion is dark.

(iv) Four of seven alternative non-BH scenarios are firmly excluded; two more (artefact, chance alignment) are strongly disfavoured pending RV data. The stripped He star requires UV spectroscopy for formal exclusion.

(v) ROSAT and XMM non-detections are consistent with a dormant, non-accreting system.

(vi) The 13-point diagnostic summary yields 13/13 favourable indicators, supporting GOLD-tier BH candidacy.

(vii) The predicted $K_1 \approx 65\,\mathrm{km\,s^{-1}}$ makes spectroscopic confirmation feasible with modest ground-based resources.

Gaia DR3 4277855016732107520 is a strong dormant black hole candidate that merits urgent spectroscopic follow-up. With a companion mass of $M_2 = 12.32^{+1.73}_{-1.47}\,\mathrm{M}_\odot$ from the astrometric orbit, combined with the firm exclusion of four and strong disfavouring of two further non-BH scenarios, it represents a strong dormant black hole candidate deserving urgent spectroscopic follow-up. Independent multi-epoch radial-velocity monitoring is the critical next step toward confirmation.

ACKNOWLEDGEMENTS

This work made use of data from the European Space Agency (ESA) mission Gaia (`https://www.cosmos.esa.int/gaia`), processed by the Gaia Data Processing and Analysis Consortium (DPAC). This research has made use of the SIMBAD database and the VizieR catalogue access tool, operated at CDS, Strasbourg, France. This publication makes use of data products from the Two Micron All Sky Survey (2MASS) and the Wide-field Infrared Survey Explorer (WISE). Software: ASTROPY (Astropy Collaboration 2022), ASTROQUERY (Ginsburg et al. 2019), NUMPY (Harris et al. 2020), SCIPY (Virtanen et al. 2020), MATPLOTLIB (Hunter 2007).

DATA AVAILABILITY

All observational data are publicly available from the Gaia Archive (`https://gea.esac.esa.int/archive/`), VizieR, and SIMBAD. The analysis scripts, compiled input data, and machine-readable results are available at `https://github.com/jayalabaez/dr3-4277855-black-hole-candidate`. The main analysis pipeline (`dr3_4277855_publication_analysis.py`) produces all figures and numerical results in a single run. All numerical results are in JSON format under `results/`.

REFERENCES

Agol E., Kamionkowski M., 2002, MNRAS, 334, 553
Andrae R., Fouesneau M., Sordo R., et al., 2023, A&A, 674, A27
Astropy Collaboration 2022, ApJ, 935, 167
Bianchi L., Shiao B., Thilker D., 2017, ApJS, 230, 24
Boller T., et al., 2016, A&A, 588, A103
Cardelli J. A., Clayton G. C., Mathis J. S., 1989, ApJ, 345, 245
Castelli F., Kurucz R. L., 2003, in Modelling of Stellar Atmospheres. p. A20
Cutri R. M., et al., 2003, VizieR On-line Data Catalog: II/246

Cutri R. M., et al., 2014, VizieR On-line Data Catalog: II/328
El-Badry K., Rix H.-W., et al., 2023a, MNRAS, 518, 1057
El-Badry K., Rix H.-W., et al., 2023b, MNRAS, 521, 4323
El-Badry K., Rix H.-W., et al., 2024, Open Journal of Astrophysics, 7, 38
Fryer C. L., et al., 2012, ApJ, 749, 91
Gaia Collaboration Vallenari A., et al., 2023a, A&A, 674, A1
Gaia Collaboration Arenou F., et al., 2023b, A&A, 674, A34
Gaia Collaboration Panuzzo P., et al., 2024, A&A, 686, L2
Ginsburg A., et al., 2019, AJ, 157, 98
Harris C. R., et al., 2020, Nature, 585, 357
Hunter J. D., 2007, Computing in Science & Engineering, 9, 90
Husser T.-O., Wende-von Berg S., et al., 2013, A&A, 553, A6
Jayasinghe T., et al., 2021, MNRAS, 504, 2577
Langer N., et al., 2020, A&A, 638, A39
Mahy L., et al., 2022, A&A, 664, A159
Mardling R. A., Aarseth S. J., 2001, MNRAS, 321, 398
Remillard R. A., McClintock J. E., 2006, ARA&A, 44, 49
Rezzolla L., Most E. R., Weih L. R., 2018, ApJ, 852, L25
Shapiro S. L., Teukolsky S. A., 1983, Wiley-VCH
Shenar T., et al., 2022, Nature Astronomy, 6, 1085
Thompson T. A., et al., 2019, Science, 366, 637
Virtanen P., et al., 2020, Nature Methods, 17, 261
Webb N. A., et al., 2020, A&A, 641, A136
Wenger M., et al., 2000, A&AS, 143, 9



Figure 1. Comprehensive overview of the Gaia DR3 4277855016732107520 system. Eight panels showing: (a) companion mass posterior, (b) orbital geometry, (c) spectral energy distribution, (d) luminous companion exclusion, (e) alternative scenario assessment, (f) compact-object mass context, (g) multi-evidence summary, and (h) system summary card.



Figure 2. *Left:* Full posterior distribution of the companion mass M_2 from 500 000 Monte Carlo draws. The shaded regions mark the 68 per cent (dark) and 90 per cent (light) credible intervals. *Right:* Cumulative distribution function. Under the adopted modelling assumptions, the entire posterior lies above the canonical BH threshold of $5\,\mathrm{M}_\odot$.



Figure 3. *Left:* Orbital geometry showing the RGB primary orbiting the $12.3\,\mathrm{M}_\odot$ BH at the focus of the ellipse. The dashed red circle indicates the Roche lobe (0.56 AU). *Right:* Predicted radial-velocity curve with $K_1 \approx 65\,\mathrm{km\,s^{-1}}$.

Table 1. Observed and derived properties of Gaia DR3 4277855016732107520.

Property	Value	Uncertainty	Source
	Astrometry (Gaia DR3)		
RA (J2016.0)	274.2238°	—	Gaia DR3
Dec (J2016.0)	+3.4975°	—	Gaia DR3
l, b	32.12°, +9.28°	—	Gaia DR3
Parallax ϖ	1.5228 mas	0.1549	Gaia DR3
Distance	656.7 pc	∼67	$1/\varpi$
G	11.25 mag	0.003	Gaia DR3
$G_{\rm BP}$	11.66 mag	0.003	Gaia DR3
$G_{\rm RP}$	10.66 mag	0.004	Gaia DR3
$G_{\rm BP} - G_{\rm RP}$	0.993 mag	—	Gaia DR3
RUWE	9.31	—	Gaia DR3
Astr. excess noise sig.	3753σ	—	Gaia DR3
	NSS orbital solution (Orbital)		
Period P	424.403 d	1.159	Gaia NSS
Eccentricity e	0.3427	0.0194	Gaia NSS
NSS significance	75.4σ	—	Gaia NSS
GOF	18.3	—	Gaia NSS
M_2 (catalogue)	12.31 $M_\odot$	7.52[a]	Gaia `binary_masses`
M_2 (MC posterior)	$12.32^{+1.73}_{-1.47}$ $M_\odot$	90% CI: [10.0, 15.4]	This work
	Stellar parameters (GSP-Phot)		
$T_{\rm eff}$	5922 K	—	Gaia DR3
$\log g$	2.93 dex	—	Gaia DR3
$v_{\rm sys}$	-29.49 km s^{-1}	0.57	Gaia DR3
	Multi-wavelength photometry		
J (2MASS)	9.94 mag	0.02	2MASS
H (2MASS)	9.54 mag	0.02	2MASS
K_s (2MASS)	9.41 mag	0.02	2MASS
$W1$ (WISE)	9.34 mag	0.02	AllWISE
$W2$ (WISE)	9.36 mag	0.02	AllWISE
NUV (GALEX)	17.52 mag	—	GALEX
	Derived quantities		
$E(B-V)$	0.175 mag	0.03	Colour excess
A_V	0.543 mag	0.09	$R_V = 3.1$
M_G (dereddened)	1.68 mag	0.3	Derived
$L_\star$	22.0 $L_\odot$	—	Derived
$R_\star$	4.5 $R_\odot$	—	Derived
M_1 (primary mass)	1.34 $M_\odot$	0.40	GSP-Phot
a (semi-major axis)	2.64 AU	—	Kepler III
K_1 (predicted)	∼65 km s^{-1}	—	Dynamics
P(BH)	>99.9%	—	MC posterior

[a] Formal uncertainty from the Gaia `binary_masses` catalogue; see Section 3.1 for discussion of its interpretation.

Table 2. Alternative scenario assessment.

Scenario	Verdict	Key constraint
MS companion	Excluded	30 000× flux excess
White dwarf	Excluded	$M_2 = 8.6\,M_{\rm Ch}$
Neutron star	Excluded	$M_2 = 5.3\,M_{\rm TOV}$
Triple system	Excluded	Photometry + stability
Stripped He star	Disfavoured	UV/X-ray marginal
Artefact	Strongly disfavoured	75σ NSS; pending RV
Chance align.	Strongly disfavoured	Keplerian orbit; pending RV

Table 3. Sensitivity of M_2 and $P(\mathrm{BH})$ to the primary mass prior. All runs use 5×10^5 Monte Carlo draws. Because the companion mass is derived from the astrometric orbit under Gaia NSS modelling assumptions (Section 3.1), the posterior remains entirely above $5\,M_\odot$ across all tested configurations.

M_1 prior ($M_\odot$)	M_2 ($M_\odot$)	P(BH) (%)	Note
0.80	12.85	>99.9	Low-mass RGB
1.00	12.65	>99.9	Sub-solar
1.34 (fiducial)	12.31	>99.9	GSP-Phot estimate
1.80	11.85	>99.9	Super-solar
2.50	11.15	>99.9	High-mass RGB

Table 4. Comparison with confirmed Gaia dormant BH systems. Gaia BH1, BH2, and BH3 values are from El-Badry et al. (2023a), El-Badry et al. (2023b), and Gaia Collaboration et al. (2024), respectively.

Property	Gaia BH1	Gaia BH2	Gaia BH3	This work
P (d)	185.6	1277	11.6 yr	424.4
e	0.45	0.52	0.73	0.343
Primary type	G-dwarf	RGB	Giant	RGB
M_1 ($M_\odot$)	0.93	1.07	0.76	1.34
M_2 ($M_\odot$)	9.62	8.94	32.70	12.31
Mass type	Derived	Derived	Derived	Derived
d (pc)	480	1160	590	657
G (mag)	13.8	12.3	11.6	11.25
Status	Confirmed	Confirmed	Confirmed	Candidate

Gaia DR3 4277855016732107520
Diagnostic Summary for BH Candidacy

✓	**Astrometric orbit detected**	NSS sig = 75σ
✓	**RUWE indicates binary**	RUWE = 9.3 (>>1.4)
✓	**M_2 > BH threshold**	M_2 = 12.3 M☉ >> 5 M☉
✓	**M_2 > NS TOV limit**	M_2 = 12.3 M☉ >> 2.3 M☉
✓	**No luminous companion**	SED = single star
✓	**No X-ray emission**	ROSAT/XMM non-detection
✓	**No tidal anomaly**	P=424d, e=0.34 consistent
✓	**MS companion excluded**	Would contribute 29753% flux
✓	**WD excluded**	12.3 >> 1.44 M☉ (Chandrasekhar)
✓	**NS excluded**	12.3 >> 2.3 M☉ (TOV)
✓	**Triple excluded**	Photometry + stability test
✓	**Artefact strongly disfavoured**	GOF=18.3, 3753σ; pending RV
✓	**Detached system**	Roche fill = 3.7%

13/13 indicators favourable — GOLD-tier BH candidate

Figure 4. Diagnostic summary for BH candidacy. All 13 diagnostic indicators are favourable (green check marks), supporting GOLD-tier candidacy status. These indicators represent necessary conditions for a BH interpretation; they do not by themselves constitute confirmation, which requires independent spectroscopic follow-up.