GW170814 : A three-detector observation of gravitational waves from a binary black hole coalescence

The LIGO Scientific Collaboration and The Virgo Collaboration

On August 14, 2017 at 10:30:43 UTC, the Advanced Virgo detector and the two Advanced LIGO detectors coherently observed a transient gravitational-wave signal produced by the coalescence of two stellar mass black holes, with a false-alarm-rate of $\lesssim 1$ in 27000 years. The signal was observed with a three-detector network matched-filter signal-to-noise ratio of 18. The inferred masses of the initial black holes are $30.5^{+5.7}_{-3.0}\,\mathrm{M}_{\odot}$ and $25.3^{+2.8}_{-4.2}\,\mathrm{M}_{\odot}$ (at the 90% credible level). The luminosity distance of the source is $540^{+130}_{-210}\,\mathrm{Mpc}$, corresponding to a redshift of $z=0.11^{+0.03}_{-0.04}$. A network of three detectors improves the sky localization of the source, reducing the area of the 90% credible region from 1160 deg 2 using only the two LIGO detectors to 60 deg 2 using all three detectors. For the first time, we can test the nature of gravitational wave polarizations from the antenna response of the LIGO-Virgo network, thus enabling a new class of phenomenological tests of gravity.

PACS numbers:

INTRODUCTION

The era of gravitational wave (GW) astronomy began with the detection of binary black hole (BBH) mergers, by the Advanced Laser Interferometer Gravitational Wave Observatory (LIGO) detectors [1], during the first of the Advanced Detector Observation Runs. Three detections, GW150914 [2], GW151226 [3], and GW170104 [4], and a lower significance candidate, LVT151012 [5], have been announced so far. The Advanced Virgo detector [6] joined the second observation run on August 1, 2017.

On August 14, 2017, GWs from the coalescence of two black holes at a luminosity distance of 540^{+130}_{-210} Mpc, with masses of $30.5^{+5.7}_{-3.0}\,\mathrm{M}_{\odot}$ and $25.3^{+2.8}_{-4.2}\,\mathrm{M}_{\odot}$, were observed in all three detectors. The signal was first observed at the LIGO Livingston detector at 10:30:43 UTC, and at the LIGO Hanford and Virgo detectors with a delay of $\sim\!8$ ms and $\sim\!14$ ms, respectively.

The signal-to-noise ratio (SNR) time series, the time-frequency representation of the strain data and the time series data of the three detectors together with the inferred GW waveform, are shown in Fig. 1. The different sensitivities and responses of the three detectors result in the GW producing different values of matched-filter SNR in each detector.

Three methods were used to assess the impact of the Virgo instrument on this detection: (a) Using the best fit waveform obtained from analysis of the LIGO detectors' data alone, we find that the probability, in 5000 s of data around the event, of a peak in SNR from Virgo data due to noise and as large as the one observed, within a time window determined by the maximum possible time of flight, is 0.3%. (b) A search for un-modelled GW transients, demonstrates that adding Advanced Virgo improves the false-alarm-rate by an order of magnitude over the two-detector network. (c) We compare the matched-filter marginal likelihood for a model with a coherent BBH signal in all three detectors to that for a model assuming pure Gaussian noise

in Virgo and a BBH signal only in the LIGO detectors: the three detector BBH signal model is preferred with a Bayes factor of more than 1600. We note that our parameter estimation assumes that the noise in the detectors is Gaussian in the 4 s window around the event. An Anderson-Darling test on the whitened residuals, after subtraction of the best-fit waveform, is consistent with this assumption.

Until Advanced Virgo became operational, typical GW position estimates were highly uncertain compared to the fields of view of most telescopes. The baseline formed by the two LIGO detectors allowed us to localize most mergers to roughly annular regions spanning hundreds to about a thousand square degrees at the 90% credible level [7–9]. Virgo adds additional independent baselines, which in cases such as GW170814 can reduce the positional uncertainty by an order of magnitude or more [8].

Tests of general relativity (GR) in the strong field regime have been performed with the signals from the BBH mergers detected by the LIGO interferometers [2–5, 10]. In GR, GWs are characterized by two tensor (spin-2) polarizations only, whereas generic metric theories may allow up to six polarizations [11, 12]. As the two LIGO instruments have similar orientations, little information about polarizations can be obtained using the LIGO detectors alone. With the addition of Advanced Virgo we can probe, for the first time, gravitational-wave polarizations geometrically by projecting the wave's amplitude onto the three detectors. As an illustration, we perform a test comparing tensor-only mode with scalar-only and vector-only modes. We find that purely tensor polarization is strongly favored over purely scalar or vector polarizations. With this, and additional tests, we find that GW170814 is consistent with GR.




FIG. 1: The GW event GW170814 observed by LIGO Hanford, LIGO Livingston and Virgo. Times are shown from August 14, 2017, 10:30:43 UTC. *Top row*: SNR time series produced in low latency and used by the low-latency localization pipeline on August 14, 2017. The time series were produced by time-shifting the best-match template from the online analysis and computing the integrated SNR at each point in time. The single-detector SNRs in Hanford, Livingston and Virgo are 7.3, 13.7 and 4.4, respectively. *Second row*: Time-frequency representation of the strain data around the time of GW170814. *Bottom row*: Time-domain detector data (in color), and 90% confidence intervals for waveforms reconstructed from a morphology-independent wavelet analysis [13] (light gray) and BBH models described in the Source Properties section (dark gray), whitened by each instrument's noise amplitude spectral density between 20 Hz and 1024 Hz. For this figure the data were also low-passed with a 380 Hz cutoff to eliminate out-of-band noise. The whitening emphasizes different frequency bands for each detector, which is why the reconstructed waveform amplitude evolution looks different in each column. The left ordinate axes are normalized such that the physical strain of the wave form is accurate at 130 Hz. The right ordinate axes are in units of whitened strain, divided by the square root of the effective bandwidth (360 Hz), resulting in units of noise standard deviations.

DETECTORS

LIGO operates two 4 km long detectors in the US, one in Livingston, LA and one in Hanford, WA [14], while Virgo consists of a single 3 km long detector near Pisa, Italy [15]. Together with GEO600 located near Hanover, Germany [16], several science runs of the initialera gravitational wave network were conducted through 2011. LIGO stopped observing in 2010 for the Advanced LIGO upgrade[1]. The Advanced LIGO detectors have been operational since 2015 [17]. They underwent a series of upgrades between the first and second observation runs [4], and began observing again in November 2016.

Virgo stopped observing in 2011 for the Advanced Virgo upgrade, during which many parts of the detector were replaced or improved [6]. Among the main changes are an increase of the finesse of the arm-cavities, the use of heav-

ier test mass mirrors that have lower absorption and better surface quality [18]. To reduce the impact of the coating thermal noise [19], the size of the beam in the central part of the detector was doubled, which required modifications of the vacuum system and the input/output optics [20, 21]. The recycling cavities are kept marginally stable as in the initial Virgo configuration. The optical benches supporting the main readout photodiodes have been suspended and put under vacuum to reduce impact of scattered light and acoustic noise. Cryogenic traps have been installed to improve the vacuum level. The vibration isolation and suspension system, already compliant with the Advanced Virgo requirement [22, 23], has been further improved to allow for a more robust control of the last-stage pendulum and the accommodation of baffles to mitigate the effect of scattered light. The test mass mirrors are currently suspended with metallic wires. Following one year of commissioning,

Advanced Virgo joined LIGO in August 2017 for the last month of the second observation run.

For Virgo, the noises that are currently limiting the sensitivity at low frequencies are thermal noise of the test mass suspension wires, control noise, the 50 Hz mains line and harmonics, and scattered light driven by seismic noise. At high frequencies, the largest contribution comes from shot noise of the main interferometer beam, with smaller contributions coming from scattered light, and shot noise of a secondary beam used to control the laser frequency. The noise sources that limit LIGO's sensitivity are described in [24] and [25]. For both LIGO and Virgo, commissioning will continue to reach their ultimate designed sensitivities [26].

Several noise sources that are linearly coupled to the GW data channel can be subtracted in post-processing, using auxiliary sensors (e.g. photodiodes monitoring beam motion) and coupling transfer functions calculated via optimal Wiener filters. This technique was used in the initial detector era [27-29]. For LIGO, we remove calibration lines, power mains and harmonics, the effect of some length and angular controls, and the effect of laser beam motion. This noise removal can improve the sensitivity of the LIGO detectors by approximately 20 % [30]. For Virgo, we remove the effect of some length controls, and the laser frequency stabilization control. The search pipelines described in the Searches section use the calibrated strain data which were produced in low-latency and which have not undergone post-processing noise subtraction. They also use data quality flags which were produced offline. The source properties however, described in the Source Properties section, are inferred using the post-processing noisesubtracted data. Figure 2 shows the sensitivity of the Advanced LIGO - Advanced Virgo network around the time of GW170814, after the post-processing removal of several noise sources.

Detection validation procedures at LIGO [2, 31], and similar checks performed at Virgo found no evidence that instrumental or environmental disturbances could account for GW170814. Tests quantifying the detectors' susceptibility to external environmental disturbances, such as electromagnetic fields [32], indicated that any disturbance strong enough to account for the signal would be clearly detected by the array of environmental sensors. None of the environmental sensors recorded any disturbances consistent with a signal that evolved in time and frequency like GW170814. A noise transient with a central frequency around 50 Hz occurs in the Virgo detector 50 ms after GW170814. This falls outside the window expected due to the light travel time between the detectors, and has therefore no effect on the interpretation of the GW signal.

LIGO is calibrated by inducing test-mass motion using photon pressure from modulated auxiliary lasers [33, 34], and Virgo is calibrated using electromagnetic actuators [35, 36]. Frequency-dependent calibration uncertainties are determined for both LIGO detectors for GW170814 using the method in [37], and used for estimation of the properties




FIG. 2: Amplitude spectral density of strain sensitivity of the Advanced LIGO – Advanced Virgo network, estimated using 4096 s of data around the time of GW170814. Here, several known linearly coupled noise sources have been removed from the data.

of this event; the maximum 1- σ uncertainty for the strain data in the frequency range 20 - 1024 Hz is 7% in amplitude and 4° in phase. The maximum 1- σ uncertainties for Virgo are 8% in amplitude and 3° in phase over the same frequency range. The estimation of properties of GW170814 use these maximum values for the Virgo uncertainty over the whole frequency range. Uncertainties in the time stamping of the data are 10 μ s for LIGO and 20 μ s for Virgo, which does not limit the sky localization.

SEARCHES

GW170814 was first identified with high confidence ~ 30 s after its arrival by two independent low-latency matched-filter pipelines [38–44] that filter the data against a collection of approximate gravitational-wave templates [45–53], triggering an alert that was shared with partners for electromagnetic follow-up [54].

The significance estimates for this event were found by the two matched-filter pipelines, and a fully coherent unmodelled search pipeline [55], analyzing 5.9 days of coincident strain data from the Advanced LIGO detectors spanning August 13, 2017 to August 21, 2017. The matched-filter pipelines do not currently use data from Virgo for significance estimates. Coherent searches, however, use the Virgo data to improve significance estimates.

The analysis was performed over the same source parameter space as the GW170104 matched-filter analysis [4] and with additional data quality information unavailable in low-latency [5, 31], although the noise-subtracted data described in the Detectors section was not used. Both pipelines identified GW170814 with a Hanford-Livingston

network SNR of 15, with ranking statistic values from the two pipelines corresponding to a false-alarm rate of 1 in 140 000 years in one search [38, 39] and 1 in 27 000 years in the other search [40–44, 56], clearly identifying GW170814 as a GW signal. The difference in significance is due to the different techniques used to rank candidate events and measure the noise background in these searches, however both report a highly significant event.

The significance of GW170814 was confirmed on the full network of three detectors by an independent coherent analysis that targets generic gravitational-wave transients with increasing frequency over time [55]. This more generic search reports a false alarm rate < 1 in 5 900 years. By comparison, when we limit this analysis to the two LIGO detectors only, the false alarm rate is approximately 1 in 300 years; the use of the data from Virgo improves significance by more than an order of magnitude. Moreover, this independent approach recovers waveforms and SNRs at the three detectors which are compatible with respect to the coherent analyses used to infer source properties (see Source Properties section).

LOCALIZATION

Some compact object mergers are thought to produce not just GWs but also broadband electromagnetic emission. LIGO and Virgo have been distributing low-latency alerts and localizations of GW events to a consortium now consisting of ground- and space-based facilities who are searching for gamma-ray, X-ray, optical, near-infrared, radio, and neutrino counterparts [57–59].

For the purpose of position reconstruction, the LIGO–Virgo GW detector network can be thought of as a phased array of antennas. Any single detector provides only minimal position information, its slowly varying antenna pattern favoring two broad regions perpendicular to the plane of the detectors' arms [60, 61]. However, with a network of detectors, sky position can be inferred by triangulation employing the time differences [62, 63], phase differences, and amplitude ratios on arrival at the sites [64].

An initial rapid localization was performed by coherent triangulation of the matched-filter estimates of the times, amplitudes, and phases on arrival [65]. The localization was then progressively refined by full coherent Bayesian parameter estimation [66], using more sophisticated waveform models and treatment of calibration systematics, as described in the next section.

The localization of GW170814 is shown in Fig. 3. For the rapid localization from Hanford and Livingston, the 90% credible area on the sky is $1160\,\mathrm{deg^2}$ and shrinks to $100\,\mathrm{deg^2}$ when including Virgo data. The full parameter estimation further constrains the position to a 90% credible area of $60\,\mathrm{deg^2}$ centered at the maximum a posteriori position of right ascension R.A.= $03^\mathrm{h}11^\mathrm{m}$ and declination Dec.= $-44^\circ57^\mathrm{m}$ (J2000). The shift between the rapid lo-

calization and the full parameter estimation is partly due to the noise removal and final detector calibration, described in a previous section, that was applied for the full parameter estimation but not the rapid localization.

Incorporating Virgo data also reduces the luminosity distance uncertainty from $570^{+300}_{-230}\,\mathrm{Mpc}$ (rapid localization) to $540^{+130}_{-210}\,\mathrm{Mpc}$ (full parameter estimation). As with the previous paragraph, the three-dimensional credible volume and number of possible host galaxies also decreases by an order of magnitude [67–69], from $71\times10^6\,\mathrm{Mpc}^3$, to $3.4\times10^6\,\mathrm{Mpc}^3$, to $2.1\times10^6\,\mathrm{Mpc}^3$.

Follow-up observations of GW170814 were conducted by 25 facilities in neutrinos [70–72], gamma-rays [73–81], X-rays [82–85], and in optical and near-infrared [86–98]. No counterpart has been reported so far.

SOURCE PROPERTIES

The parameters of the source are inferred through a coherent Bayesian analysis [66, 99] of offline noise-subtracted data for the LIGO and Virgo detectors using two independently developed waveform models.

Both of these models are calibrated to partially overlapping sets of numerical-relativity simulations of binary black hole coalescences, following from the initial breakthroughs reported in [100–102]. One model includes the full two-spin inspiral-dynamics in the absence of precession [53, 103], whereas the other model includes an effective treatment of the spin-precession dynamics through a rotation of an originally nonprecessing model [104–107]. Previous studies [108–111] have investigated the effect of systematic waveform modelling in sources we believe to be similar to GW170814, albeit for a different detector network configuration. Based on these analogous investigations, and due to the brevity and amplitude of GW170814, we expect systematic biases to be significantly smaller than the statistical error reported in this work.

In addition to a waveform model, the coherent Bayesian analysis also incorporates the detectors' noise power spectral densities at the time of the event [13, 112] and marginalizes over the calibration uncertainties described in the Detector section, as in [4, 5, 113]. The coherent Bayesian analysis recovers the maximum matched-filter SNR across the LIGO-Virgo network of 18.3 [66, 114], with individual detector matched-filter SNRs of 9.7, 14.8 and 4.8 in LIGO Hanford, LIGO Livingston and Virgo respectively.

Table I shows source parameters for GW170814, where we quote the median value and the symmetric 90% credible intervals. The final mass (or equivalently the energy radiated), final spin and peak luminosity are computed from averages of fits to numerical relativity simulations [115–119]. The reported uncertainties account for both statistical and systematic uncertainties from averaging over the two waveform models used. An independent calculation using direct




FIG. 3: Localization of GW170814. The rapid localization using data from the two LIGO sites is shown in yellow, with the inclusion of data from Virgo shown in green. The full Bayesian localization is shown in purple. The contours represent the 90% credible regions. The left panel is an orthographic projection and the inset in the center is a gnomonic projection; both are in equatorial coordinates. The inset on the right shows the posterior probability distribution for the luminosity distance, marginalized over the whole sky.

TABLE I: Source parameters for GW170814: median values with 90% credible intervals. We quote source-frame masses; to convert to the detector frame, multiply by (1+z) [120, 121]. The redshift assumes a flat cosmology with Hubble parameter $H_0=67.9~{\rm km\,s}^{-1}\,{\rm Mpc}^{-1}$ and matter density parameter $\Omega_{\rm m}=0.3065$ [122].


Primary black hole mass m_1	$30.5^{+5.7}_{-3.0}\mathrm{M}_{\odot}$
Secondary black hole mass m_2	$25.3^{+2.8}_{-4.2}\mathrm{M}_{\odot}$
Chirp mass \mathcal{M}	$24.1^{+1.4}_{-1.1}\mathrm{M}_{\odot}$
Total mass M	$55.9^{+3.4}_{-2.7}\mathrm{M}_{\odot}$
Final black hole mass M_{f}	$53.2^{+3.2}_{-2.5}\mathrm{M}_{\odot}$
Radiated energy E_{rad}	$2.7^{+0.4}_{-0.3}{ m M}_{\odot}{ m c}^2$
Peak luminosity $\ell_{\rm peak}$	$3.7^{+0.5}_{-0.5} \times 10^{56} \mathrm{erg s^{-1}}$
Effective inspiral spin parameter $\chi_{\rm eff}$	$0.06^{+0.12}_{-0.12}$
Final black hole spin a_{f}	$0.70^{+0.07}_{-0.05}$
Luminosity distance D_{L}	$540^{+130}_{-210} \mathrm{Mpc}$
Source redshift z	$0.11^{+0.03}_{-0.04}$

comparison to numerical relativity, similar to [108], gives consistent parameters.

The inferred posterior distributions for the two black hole masses m_1 and m_2 are shown in Fig. 4. GW170814 allows for measurements of comparable accuracy of the total binary mass $M=m_1+m_2$, which is primarily

governed by the merger and ringdown, and the chirp mass $\mathcal{M} = (m_1 m_2)^{3/5}/M^{1/5}$, determined by the binary inspiral [64, 125–131], similarly to both GW150914 [99] and GW170104 [4].

The orbital evolution is dominated by the black hole masses and the components of their spins $S_{1,2}$ perpendicular to the orbital plane, and other spin components affect the GW signal on a subdominant level. The dominant spineffects are represented through the effective inspiral spin parameter $\chi_{\text{eff}} = (m_1 a_1 \cos \theta_{LS_1} + m_2 a_2 \cos \theta_{LS_2})/M$ which is approximately conserved throughout the evolution of the binary orbit [132–135]. Here θ_{LS_i} is the angle between the black hole spin S_i and the Newtonian orbital angular momentum L for both the primary (i = 1) and secondary (i = 2) black holes, and $a_i = |c\mathbf{S}_i/Gm_i^2|$ is the dimensionless spin magnitude of the initial (i = 1, 2) and final (i = f) black holes. For $a_{1,2}$, this analysis assumed a uniform prior distribution between 0 and 0.99, with no restrictions on the spin orientations. As with GW150914 and GW170104, $\chi_{\rm eff}$ is consistent with having a arbitrarily small value [4, 5]. The spin-components orthogonal to L are interesting, as they lead to precession of the binary orbit [136, 137] and are here quantified by the effective precession spin parameter χ_p [106, 137]. As for previous events [4, 5, 110, 124], the $\chi_{\rm p}$ posterior distribution is dominated by assumptions about the prior, as shown in Fig. 4. Given these assumptions, as well as statistical and systematic uncertainties, we cannot draw further robust conclusions about the transverse components of the spin.






FIG. 4: Posterior probability density for the source-frame masses m_1 and m_2 (top) and the effective inspiral and precession spin parameters, $\chi_{\rm eff}$ and $\chi_{\rm P}$ (bottom) measured at a gravitational-wave frequency of 20 Hz, well before the merger. The dashed lines mark the 90% credible interval for the one-dimensional marginalized distributions. The two-dimensional plots show the contours of the 50% and 90% credible regions plotted over a color-coded posterior density function. For GW170814, both $\chi_{\rm eff}$ and $\chi_{\rm P}$ are influenced by their respective prior distributions, shown in green. While the GW observation provides additional constraints for the $\chi_{\rm eff}$ posterior, there is only a marginal information gain for $\chi_{\rm P}$. (Kullback–Leibler divergence between the prior and posterior distribution of 0.08 nat [4, 123, 124])

The event, GW170814, is consistent with the population of BBHs, physical parameters and merger rate reported in previous BBH papers [5, 138, 139].

TESTS OF GENERAL RELATIVITY

To determine the consistency of the signal with GR, we allowed the post-Newtonian and additional coefficients describing the waveform to deviate from their nominal values [140–142], as was done for previous detections [2–5, 10]. In addition to previously tested coefficients, these analyses were expanded to also explicitly consider phase contributions at effective -1PN order, *i.e.* with a frequency dependence of $f^{-7/3}$. Additionally, as in [2–4], we check that the inspiral and merger-ringdown regimes are mutually consistent, and check for possible deviations from GR in the propagation of GWs due to a massive graviton and/or Lorentz invariance violation. Preliminary results of all these tests show no evidence for disagreement with the predictions of GR; detailed investigations are still ongoing, and full results will be presented at a later date.

GRAVITATIONAL-WAVE POLARIZATIONS

One of the key predictions of GR is that metric perturbations possess two tensor degrees of freedom [143, 144]. These two are only a subset of the six independent modes allowed by generic metric theories of gravity, which may in principle predict any combination of tensor (spin-2), vector (spin-1), or scalar (spin-0) polarizations [11, 12].

So far, some evidence that GWs are described by the tensor (spin-2) metric perturbations of GR has been obtained from measurements of the rate of orbital decay of binary pulsars, in the context of specific beyond-GR theories (see e.g. [145, 146], or [147, 148] for reviews), and from the rapidly changing GW phase of BBH mergers observed by LIGO, in the framework of parameterized models [2, 4, 10]. The addition of Advanced Virgo provides us with another, more compelling, way of probing the nature of polarizations by studying GW geometry directly through the projection of the metric perturbation onto our detector network [149–151].

The coherent Bayesian analysis described in the Source Properties section is repeated after replacing the standard tensor antenna response functions with those appropriate for scalar or vector polarizations [152]. In our analysis, we are interested in the geometric projection of the GW onto the detector network therefore the details of the phase model itself are less relevant so long that it is a faithful representation of the fit to the data in Fig 1. Hence, we assume a GR phase model. We find Bayes factors of more than 200 and 1000 in favour of the purely tensor polarization against purely vector and purely scalar, respectively. We also find

that, as expected, the reconstructed sky location and distance change significantly depending on the polarization content of the source, with non-overlapping 90% credible regions for tensor, vector and scalar. Similar tests were inconclusive for previous events [10] because the two LIGO detectors are very nearly coaligned, and record the same combination of polarizations.

CONCLUSION

On August 1 2017, Advanced Virgo joined the two Advanced LIGO detectors in the second of the Advanced Detector Observation Runs. On August 14, 2017, a GW signal coming from the merger of two stellar mass black holes was observed with the Virgo and LIGO detectors. The three-detector detection of GW170814 allowed for a significant reduction in the search volume for the source. The black hole characteristics of GW170814 are similar to GW150914 and GW170104, and are found to be consistent with the astrophysical population and merger rate determined with previous detections. The addition of Virgo has allowed us to probe the polarization content of the signal for the first time; we find that the data strongly favor pure tensor polarization of gravitational waves, over pure scalar or pure vector polarizations.

ACKNOWLEDGMENTS

The authors gratefully acknowledge the support of the United States National Science Foundation (NSF) for the construction and operation of the LIGO Laboratory and Advanced LIGO as well as the Science and Technology Facilities Council (STFC) of the United Kingdom, the Max-Planck-Society (MPS), and the State of Niedersachsen/Germany for support of the construction of Advanced LIGO and construction and operation of the GEO600 detector. Additional support for Advanced LIGO was provided by the Australian Research Council. The authors gratefully acknowledge the Italian Istituto Nazionale di Fisica Nucleare (INFN), the French Centre National de la Recherche Scientifique (CNRS) and the Foundation for Fundamental Research on Matter supported by the Netherlands Organisation for Scientific Research, for the construction and operation of the Virgo detector and the creation and support of the EGO consortium. The authors also gratefully acknowledge research support from these agencies as well as by the Council of Scientific and Industrial Research of India, the Department of Science and Technology, India, the Science & Engineering Research Board (SERB), India, the Ministry of Human Resource Development, India, the Spanish Agencia Estatal de Investigación, the Vicepresidència i Conselleria d'Innovació, Recerca i Turisme and the Conselleria d'Educació i Universitat del Govern de les Illes Balears, the Conselleria d'Educació, Investigació, Cultura

i Esport de la Generalitat Valenciana, the National Science Centre of Poland, the Swiss National Science Foundation (SNSF), the Russian Foundation for Basic Research, the Russian Science Foundation, the European Commission, the European Regional Development Funds (ERDF), the Royal Society, the Scottish Funding Council, the Scottish Universities Physics Alliance, the Hungarian Scientific Research Fund (OTKA), the Lyon Institute of Origins (LIO), the National Research, Development and Innovation Office Hungary (NKFI), the National Research Foundation of Korea, Industry Canada and the Province of Ontario through the Ministry of Economic Development and Innovation, the Natural Science and Engineering Research Council Canada, the Canadian Institute for Advanced Research, the Brazilian Ministry of Science, Technology, Innovations, and Communications, the International Center for Theoretical Physics South American Institute for Fundamental Research (ICTP-SAIFR), the Research Grants Council of Hong Kong, the National Natural Science Foundation of China (NSFC), the Leverhulme Trust, the Research Corporation, the Ministry of Science and Technology (MOST), Taiwan and the Kavli Foundation. The authors gratefully acknowledge the support of the NSF, STFC, MPS, INFN, CNRS and the State of Niedersachsen/Germany for provision of computational resources.

- [1] J. Aasi et al. (LIGO Scientific Collaboration), Class. Quantum Grav. **32**, 074001 (2015), 1411.4547.
- [2] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. 116, 061102 (2016), 1602.03837.
- [3] B. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. 116, 241103 (2016), 1606.04855.
- [4] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. 118, 221101 (2017), 1706.01812.
- [5] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. X 6, 041015 (2016), 1606.04856.
- [6] F. Acernese et al. (Virgo Collaboration), Class. Quantum Grav. 32, 024001 (2015), 1408.3978.
- [7] M. M. Kasliwal and S. Nissanke, ApJ 789, L5 (2014), 1309.1554.
- [8] L. P. Singer, L. R. Price, B. Farr, A. L. Urban, C. Pankow, et al., Astrophys. J. 795, 105 (2014), 1404.5623.
- [9] C. P. L. Berry et al., Astrophys. J. 804, 114 (2015), 1411.6934.
- [10] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. 116, 221101 (2016), 1602.03841.
- [11] D. M. Eardley, D. L. Lee, A. P. Lightman, R. V. Wagoner, and C. M. Will, Phys. Rev. Lett. 30, 884 (1973).
- [12] D. M. Eardley, D. L. Lee, and A. P. Lightman, Phys. Rev. D8, 3308 (1973).
- [13] N. J. Cornish and T. B. Littenberg, Class. Quantum Grav.

- **32**, 135012 (2015), 1410.3835.
- [14] B. Abbott et al. (LIGO Scientific Collaboration), Rept. Prog. Phys. 72, 076901 (2009), 0711.3041.
- [15] T. Accadia et al. (Virgo Collaboration), JINST 7, P03012 (2012).
- [16] K. L. Dooley et al., Class. Quant. Grav. 33, 075009 (2016), 1510.00317.
- [17] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. 116, 131103 (2016), 1602.03838.
- [18] L. Pinard, C. Michel, B. Sassolas, L. Balzarini, J. Degallaix, V. Dolique, R. Flaminio, D. Forest, M. Granata, B. Lagrange, et al., Appl. Opt. 56, C11 (2017).
- [19] M. Granata, E. Saracco, N. Morgado, A. Cajgfinger, G. Cagnoli, J. Degallaix, V. Dolique, D. Forest, J. Franc, C. Michel, et al., Phys. Rev. D 93, 012007 (2016).
- [20] C. Buy, E. Genin, M. Barsuglia, R. Gouaty, and M. Tacca, Classical and Quantum Gravity 34, 095011 (2017).
- [21] M. Tacca, F. Sorrentino, C. Buy, M. Laporte, G. Pillant, E. Genin, P. L. Penna, and M. Barsuglia, Appl. Opt. 55, 1275 (2016).
- [22] R. Del Fabbro, A. Di Virgilio, A. Giazotto, H. Kautzky, V. Montelatici, and D. Passuello, Phys. Lett. A124, 253 (1987)
- [23] F. Acernese et al., Astropart. Phys. 33, 182 (2010).
- [24] D. V. Martynov et al. (LIGO Scientific Collaboration), Phys. Rev. 93, 112004 (2016), 1604.00439.
- [25] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Supplement: GW170104: Observation of a 50-Solar-Mass Binary Black Hole Coalescence at Redshift 0.2, http://link.aps.org/supplemental/10.1103/PhysRevLett.118.
- [26] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Living Rev. Relat. **19**, 1 (2016), 1304.0670.
- [27] J. C. Driggers, M. Evans, K. Pepper, and R. Adhikari, Rev. Sci. Instrum. 83, 024501 (2012), 1112.2224.
- [28] G. D. Meadors, K. Kawabe, and K. Riles, Class. Quant. Grav. 31, 105014 (2014), 1311.6835.
- [29] V. Tiwari et al., Class. Quant. Grav. 32, 165014 (2015), 1503.07476.
- [30] J. C. Driggers, Tech. Note, LIGO-P1700260 (2017), URL https://dcc.ligo.org/P1700260/public.
- [31] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Class. Quant. Grav. 33, 134001 (2016), 1602.03844.
- [32] A. Effler, R. M. S. Schofield, V. V. Frolov, G. González, K. Kawabe, J. R. Smith, J. Birch, and R. McCarthy, Class. Quant. Grav. 32, 035017 (2015), 1409.5160.
- [33] S. Karki et al., Rev. Sci. Instrum. 87, 114503 (2016), 1608.05055.
- [34] B. P. Abbott et al. (LIGO Scientific Collaboration), Phys. Rev. D 95, 062003 (2017), 1602.03845.
- [35] T. Accadia et al. (Virgo Collaboration), Class. Quant. Grav. **28**, 025005 (2011), [Erratum: Class. Quant. Grav.28,079501(2011)], 1009.5190.
- [36] T. Accadia et al. (Virgo Collaboration), Class. Quant. Grav. **31**, 165013 (2014), 1401.6066.
- [37] C. Cahillane, J. Betzwieser, D. Brown, E. Goetz, E. D. Hall, K. Izumi, S. Kandhasamy, S. Karki, J. S. Kissel, G. Mendell, et al. (2017), 1708.03023.

- [38] K. Cannon et al., Astrophys. J. **748**, 136 (2012), 1107.2665.
- [39] C. Messick et al., Phys. Rev. **95**, 042001 (2017), 1604.04324.
- [40] A. H. Nitz, I. W. Harry, J. L. Willis, C. M. Biwer, D. A. Brown, L. P. Pekowsky, T. Dal Canton, A. R. Williamson, T. Dent, C. D. Capano, et al., *PyCBC Software*, https://github.com/ligo-cbc/pycbc (2017).
- [41] A. H. Nitz, T. Dent, T. Dal Canton, S. Fairhurst, and D. A. Brown (2017), 1705.01513.
- [42] A. H. Nitz (2017), 1709.08974.
- [43] S. A. Usman et al., Class. Quant. Grav. 33, 215004 (2016), 1508.02357.
- [44] T. Dal Canton et al., Phys. Rev. **D90**, 082004 (2014), 1405.6731.
- [45] B. S. Sathyaprakash and S. V. Dhurandhar, Phys. Rev. D 44, 3819 (1991).
- [46] B. J. Owen, Phys. Rev. D 53, 6749 (1996), gr-qc/9511032.
- [47] B. J. Owen and B. S. Sathyaprakash, Phys. Rev. D 60, 022002 (1999), 9808076.
- [48] D. A. Brown, I. Harry, A. Lundgren, and A. H. Nitz, Phys.Rev. D 86, 084017 (2012), 1207.6406.
- [49] I. W. Harry, B. Allen, and B. Sathyaprakash, Phys. Rev. D 80, 104014 (2009), 0908.2090.
- [50] P. Ajith, N. Fotopoulos, S. Privitera, A. Neunzert, and A. J. Weinstein, Phys. Rev. D 89, 084041 (2014), 1210.6666.
- [51] L. Blanchet, T. Damour, G. Esposito-Farèse, and B. R. Iyer, Phys. Rev. Lett. 93, 091101 (2004), 0406012.
- [52] L. Blanchet, Living Rev. Rel. 17, 2 (2014), 1310.1528.
- [53] A. Bohé et al., Phys. Rev. D 95, 044028 (2017), 1611.03703.
- http://link.aps.org/supplemental/10.1103/PhysRevLett.118.2211**Q**34] L. S. Collaboration and V. Collaboration, GCN **21474**, 1 B. P. Abbott et al. (LIGO Scientific Collaboration, (2017).
 - [55] S. Klimenko et al., Phys. Rev. D 93, 042004 (2016), 1511.05999.
 - [56] B. Allen, W. G. Anderson, P. R. Brady, D. A. Brown, and J. D. E. Creighton, Phys. Rev. D 85, 122006 (2012), 0509116.
 - [57] B. P. Abbott et al. (LIGO Scientific, Virgo, ASKAP, BOOTES, DES, DEC, Fermi GBM, Fermi-LAT, GRAWITA, INTEGRAL, iPTF, InterPlanetary Network, J-GEM, La Silla-QUEST Survey, Liverpool Telescope, LOFAR, MASTER, MAXI, MWA, Pan-STARRS, PESSTO, Pi of the Sky, SkyMapper, Swift, C2PU, TOROS, VISTA), Astrophys. J. Lett. 826, L13 (2016), 1602.08492.
 - [58] B. P. Abbott et al. (LIGO Scientific, Virgo, ASKAP, BOOTES, DES, DEC, Fermi GBM, Fermi-LAT, GRAWITA, INTEGRAL, iPTF, InterPlanetary Network, J-GEM, La Silla-QUEST Survey, Liverpool Telescope, LOFAR, MASTER, MAXI, MWA, Pan-STARRS, PESSTO, Pi of the Sky, SkyMapper, Swift, C2PU, TOROS, VISTA), Astrophys. J. Suppl. 225, 8 (2016), 1604.07864.
 - [59] S. Adrián-Martínez et al. (LIGO Scientific Collaboration, Virgo Collaboration, IceCube, ANTARES), Phys. Rev. D 93, 122010 (2016), 1602.05411.
 - [60] R. L. Forward, Phys. Rev. D 17, 379 (1978).
 - [61] K. Thorne, in *Three Hundred Years of Gravitation*, edited by S. W. Hawking and W. Israel (Cambridge University Press, 1987), pp. 330–458, ISBN 9780521343121.

- [62] S. Fairhurst, New J. Phys. **11**, 123006 (2009), [Erratum: New J. Phys.13,069602(2011)], 0908.2356.
- [63] S. Fairhurst, Class. Quantum Grav. 28, 105021 (2011), 1010.6192.
- [64] P. Ajith and S. Bose, Phys. Rev. D 79, 084032 (2009), 0901.4936.
- [65] L. P. Singer and L. Price, Phys. Rev. D 93, 024013 (2016), 1508.03634.
- [66] J. Veitch et al., Phys. Rev. D 91, 042003 (2015), 1409.7215.
- [67] L. P. Singer et al., Astrophys. J. 829, L15 (2016), 1603.07333.
- [68] L. P. Singer et al., Astrophys. J. Suppl. 226, 10 (2016), 1605.04242.
- [69] H.-Y. Chen and D. E. Holz (2016), 1612.01471.
- [70] S. Countryman, GCN 21475, 1 (2017), IceCube circular.
- [71] D. Dornic, GCN **21479**, 1 (2017), ANTARES circular.
- [72] J. Alvarez-Muniz, GCN 21523, 1 (2017), Pierre Auger circular.
- [73] A. Pozanenko, GCN 21476, 1 (2017), INTEGRAL SPI-ACS circular.
- [74] F. Verrecchia, GCN 21477, 1 (2017), AGILE MCAL circular.
- [75] V. Savchenko, GCN 21478, 1 (2017), INTEGRAL SPI-ACS and MCAL circular.
- [76] D. Kocevski, GCN 21481, 1 (2017), Fermi LAT circular.
- [77] F. Verrecchia, GCN **21482**, 1 (2017), *AGILE* circular.
- [78] A. Goldstein, GCN 21484, 1 (2017), Fermi GBM circular.
- [79] D. Svinkin, GCN 21500, 1 (2017), Konus/WIND circular.
- [80] F. Schussler, GCN 21673, 1 (2017), H.E.S.S. circular.
- [81] V. Savchenko, GCN 21695, 1 (2017), INTEGRAL circular.
- [82] A. Lien, GCN 21483, 1 (2017), Swift BAT circular.
- [83] S. Xiong, GCN 21486, 1 (2017), Insight-HXMT circular.
- [84] S. Sugita, GCN **21494**, 1 (2017), MAXI GSC circular.
- [85] P. Evans, GCN **21503**, 1 (2017), Swift XRT circular.
- [86] S. J. Smartt, GCN 21488, 1 (2017), Pan-STARRS circular.
- [87] P. D'Avanzo, GCN **21495**, 1 (2017), REM circular.
- [88] Y. Utsumi, GCN 21497, 1 (2017), Subaru Hyper Suprime-Cam circular.
- [89] G. Greco, GCN 21498, 1 (2017), ESO-VST circular.
- [90] V. Lipunov, GCN **21499**, 1 (2017), MASTER circular.
- [91] C. Copperwheat, GCN 21512, 1 (2017), Liverpool Telescope circular.
- [92] G. P. Smith, GCN 21692, 1 (2017), Gemini and VLT circular.
- [93] M. Soares-Santos, GCN 21789, 1 (2017), DECam circular.
- [94] G. Hosseinzadeh, GCN 21860, 1 (2017), LCOGT circular.
- [95] S. Yang, GCN 21878, 1 (2017), DLT40 circular.
- [96] M. Im, GCN 21885, 1 (2017), LSGT circular.
- [97] D. Roberts, GCN **21888**, 1 (2017), PIRATE circular.
- [98] N. Tanvir, GCN 21890, 1 (2017), VISTA circular.
- [99] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. Lett. 116, 241102 (2016), 1602.03840.
- [100] F. Pretorius, Phys. Rev. Lett. 95, 121101 (2005), gr-qc/0507014.
- [101] M. Campanelli, C. O. Lousto, P. Marronetti, and Y. Zlochower, Phys. Rev. Lett. 96, 111101 (2006), gr-qc/0511048.
- [102] J. G. Baker, J. Centrella, D.-I. Choi, M. Koppitz, and J. van Meter, Phys. Rev. Lett. 96, 111102 (2006), gr-qc/0511103.

- [103] M. Pürrer, Phys. Rev. D 93, 064041 (2016), 1512.02248.
- [104] S. Husa, S. Khan, M. Hannam, M. Pürrer, F. Ohme, X. J. Forteza, and A. Bohé, Phys. Rev. D 93, 044006 (2016), 1508.07250.
- [105] S. Khan, S. Husa, M. Hannam, F. Ohme, M. Pürrer, X. J. Forteza, and A. Bohé, Phys. Rev. D 93, 044007 (2016), 1508.07253.
- [106] M. Hannam, P. Schmidt, A. Bohé, L. Haegel, S. Husa, F. Ohme, G. Pratten, and M. Pürrer, Phys. Rev. Lett. 113, 151101 (2014), 1308.3271.
- [107] R. Smith, S. E. Field, K. Blackburn, C.-J. Haster, M. Pürrer, V. Raymond, and P. Schmidt, Phys. Rev. D 94, 044031 (2016), 1604.08253.
- [108] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. D 94, 064035 (2016), 1606.01262.
- [109] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Classical Quantum Gravity 34, 104002 (2017), 1611.07531.
- [110] T. D. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Phys. Rev. X 6, 041014 (2016), 1606.01210.
- [111] G. Lovelace et al., Class. Quant. Grav. 33, 244002 (2016), 1607.05377.
- [112] T. B. Littenberg and N. J. Cornish, Phys.Rev. D 91, 084034 (2015), 1410.3852.
- [113] W. M. Farr, B. Farr, and T. Littenberg, Tech. Rep. LIGO-T1400682, LIGO Project (2015), URL https://dcc.ligo.org/LIGO-T1400682/public.
- [114] A. Pai, S. Dhurandhar, and S. Bose, Phys. Rev. D 64, 042004 (2001).
- [115] F. Hofmann, E. Barausse, and L. Rezzolla, Astrophys. J. 825, L19 (2016), 1605.01938.
- [116] X. Jiménez-Forteza, D. Keitel, S. Husa, M. Hannam, S. Khan, and M. Pürrer, Phys. Rev. D 95, 064024 (2017), 1611.00332.
- [117] J. Healy and C. O. Lousto, Phys. Rev. D 95, 024037 (2017), 1610.09713.
- [118] N. K. Johnson-McDaniel et al., Tech. Rep. LIGO-T1600168, LIGO Scientific Collaboration and Virgo Collaboration (2016), URL https://dcc.ligo.org/LIGO-T1600168/public.
- [119] D. Keitel et al., Phys. Rev. D **96**, 024006 (2017), 1612.09566.
- [120] A. Krolak and B. F. Schutz, General Relativity and Gravitation 19, 1163 (1987).
- [121] D. E. Holz and S. A. Hughes, Astrophys. J. 629, 15 (2005), astro-ph/0504616.
- [122] P. A. R. Ade et al. (Planck), Astron. Astrophys. 594, A13 (2016), 1502.01589.
- [123] S. Kullback and R. A. Leibler, Ann. Math. Statist. **22**, 79 (1951).
- [124] S. Vitale, D. Gerosa, C.-J. Haster, K. Chatziioannou, and A. Zimmerman (2017), 1707.04637.
- [125] L. S. Finn and D. F. Chernoff, Phys. Rev. D 47, 2198 (1993), 9301003.
- [126] C. Cutler and E. E. Flanagan, Phys. Rev. D 49, 2658 (1994), 9402014.
- [127] L. Blanchet, T. Damour, B. R. Iyer, C. M. Will, and A. Wiseman, Phys. Rev. Lett. 74, 3515 (1995), gr-qc/9501027.

- [128] J. Veitch, M. Pürrer, and I. Mandel, Phys. Rev. Lett. 115, 141101 (2015), 1503.05953.
- [129] P. B. Graff, A. Buonanno, and B. Sathyaprakash, Phys. Rev. **D92**, 022002 (2015), 1504.04766.
- [130] C.-J. Haster, Z. Wang, C. P. L. Berry, S. Stevenson, J. Veitch, and I. Mandel, Mon. Not. Roy. Astron. Soc. 457, 4499 (2016), 1511.01431.
- [131] A. Ghosh, W. Del Pozzo, and P. Ajith, Phys. Rev. D 94, 104070 (2016), 1505.05607.
- [132] E. Racine, Phys. Rev. **D78**, 044021 (2008), 0803.1820.
- [133] P. Ajith et al., Phys. Rev. Lett. **106**, 241101 (2011), 0909.2867.
- [134] C. O. Lousto and Y. Zlochower, Phys. Rev. D 89, 021501 (2014), 1307.6237.
- [135] D. Gerosa, M. Kesden, U. Sperhake, E. Berti, and R. O'Shaughnessy, Phys. Rev. **D92**, 064016 (2015), 1506.03492.
- [136] T. A. Apostolatos, C. Cutler, G. J. Sussman, and K. S. Thorne, Phys. Rev. **D49**, 6274 (1994).
- [137] P. Schmidt, F. Ohme, and M. Hannam, Phys. Rev. D 91, 024043 (2015), 1408.1810.
- [138] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Astrophys. J. Lett. 833, L1 (2016),
- [139] B. P. Abbott et al. (LIGO Scientific Collaboration, Virgo Collaboration), Astrophys. J. Suppl. 227, 14 (2016),
- [140] L. Blanchet and B. S. Sathyaprakash, Phys. Rev. Lett. 74, 1067 (1995).
- [141] C. K. Mishra, K. G. Arun, B. R. Iyer, and B. S.

- Sathyaprakash, Phys. Rev. **D82**, 064010 (2010), 1005.0304.
- [142] T. G. F. Li, W. Del Pozzo, S. Vitale, C. Van Den Broeck, M. Agathos, J. Veitch, K. Grover, T. Sidery, R. Sturani, and A. Vecchio, Phys. Rev. **D85**, 082003 (2012), 1110.0530.
- [143] C. M. Will, Living Rev. Rel. 17, 4 (2014), 1403.7377.
- [144] C. Will, Theory and Experiment in Grav-**Physics** (Cambridge University itational 1993), ISBN 9780521439732, Press. URL https://books.google.com/books?id=BhnUITA7sDIC.
- [145] J. M. Weisberg, D. J. Nice, and J. H. Taylor, Astrophys. J. **722**, 1030 (2010), 1011.0718.
- [146] P. C. C. Freire, N. Wex, G. Esposito-Farese, J. P. W. Verbiest, M. Bailes, B. A. Jacoby, M. Kramer, I. H. Stairs, J. Antoniadis, and G. H. Janssen, Mon. Not. Roy. Astron. Soc. 423, 3328 (2012), 1205.1450.
- [147] I. H. Stairs, Living Rev. Rel. 6, 5 (2003), astroph/0307536.
- [148] N. Wex (2014), arXiv:1402.5594.
- [149] K. Chatziioannou, N. Yunes, and N. Cornish, Phys. Rev. D 86, 022004 (2012), [Erratum: Phys. Rev.D95,no.12,129901(2017)], 1204.2585.
- [150] M. Isi, M. Pitkin, and A. J. Weinstein, Phys. Rev. D 96, 042001 (2017), 1703.07530.
- [151] T. Callister, A. S. Biscoveanu, N. Christensen, M. Isi, A. Matas, O. Minazzoli, T. Regimbau, M. Sakellariadou, J. Tasson, and E. Thrane (2017), 1704.08373.
- [152] M. Isi, Tech. Note, LIGO-P1700276 (2017), URL https://dcc.ligo.org/P1700276/public.

Authors

T. D. Creighton, ¹⁰³ J. Cripe, ² S. G. Crowder, ¹⁰⁴ T. J. Cullen, ^{29,2} A. Cumming, ⁴⁶ L. Cunningham, ⁴⁶ E. Cuoco, ³⁰ T. Dal Canton, ⁵⁰ G. Dálya, ⁵⁶ S. L. Danilishin, ^{22,10} S. D'Antonio, ³³ K. Danzmann, ^{22,10} A. Dasgupta, ¹⁰⁵ C. F. Da Silva Costa, ⁵ V. Dattilo, ³⁰ I. Dave, ⁶¹ M. Davier, ²⁸ D. Davis, ⁴⁴ E. J. Daw, ¹⁰⁶ B. Day, ⁷⁷ S. De, ⁴⁴ D. DeBra, ⁵² J. Degallaix, ²⁶ M. De Laurentis, ^{17,4} S. Deléglise, ⁷¹ W. Del Pozzo, ^{59,23,24} N. Demos, ¹⁵ T. Denker, ¹⁰ T. Dent, ¹⁰ R. De Pietri, ^{107,108} V. Dergachev, ³⁸ R. De Rosa, ^{79,4} R. T. DeRosa, ⁷ C. De Rossi, ^{26,30} R. DeSalvo, ¹⁰⁹ O. de Varona, ¹⁰ J. Devenson, ²⁷ S. Dhurandhar, ¹⁹ M. C. Díaz, ¹⁰³ L. Di Fiore, ⁴ M. Di Giovanni, ^{110,95} T. Di Girolamo, ^{51,79,4} A. Di Lieto, ^{23,24} S. Di Pace, ^{97,35} I. Di Palma, ^{97,35} F. Di Renzo, ^{23,24} Z. Doctor, ⁹¹ V. Dolique, ²⁶ F. Donovan, ¹⁵ K. L. Dooley, ¹¹ S. Doravari, ¹⁰ I. Dorrington, ³⁶ R. Douglas, ⁴⁶ M. Dovale Álvarez, ⁵⁹ T. P. Downes, ²¹ M. Drago, ¹⁰ C. Dreissigacker, ¹⁰ J. C. Driggers, ⁴⁷ Z. Du, ⁸² M. Ducrot, ⁸ P. Dupej, ⁴⁶ S. E. Dwyer, ⁴⁷ T. B. Edo, ¹⁰⁶ M. C. Edwards, ⁷² A. Effler, ⁷ H.-B. Eggenstein, ^{38,10} P. Ehrens, ¹ J. Eichholz, ¹ S. S. Eikenberry, ⁵ R. A. Eisenstein, ¹⁵ R. C. Essick, ¹⁵ D. Estevez, ⁸ S. Farinon, ⁶⁰ B. Farr, ⁹¹ W. M. Farr, ⁵⁹ E. J. Fauchon-Jones, ³⁶ M. Favata, ¹¹¹ M. Fays, ³⁶ C. Fee, ⁸⁴ H. Fehrmann, ¹⁰ J. Feicht, ¹ M. M. Fejer, ⁵² A. Fernandez-Galiana, ¹⁵ I. Ferrante, ^{23,24} E. C. Ferreira, ¹⁶ F. Ferrini, ³⁰ F. Fidecaro, ^{23,24} D. Finstad, ⁴¹ I. Fiori, ³⁰ D. Fiorucci, ³⁹ M. Fishbach, ⁹¹ R. P. Fisher, ⁴⁴ M. Fitz-Axen, ⁴⁵ R. Flaminio, ^{26,112} M. Fletcher, ⁴⁶ H. Fong, ⁹⁰ J. A. Font, ^{85,113} P. W. F. Forsyth, ⁵⁷ S. S. Forsyth, ⁷⁷ J.-D. Fournier, ⁶⁷ S. Frasca, ^{97,35} F. Frasconi, ²⁴ Z. Frei, ⁵⁶ A. Freise, ⁵⁹ R. Frey, ⁷⁰ V. Frey, ²⁸ E. M. Fries, ¹ P. Fritschel, ¹⁵ V. V. Frolov, ⁷ P. Fulda, ⁵ M. Gennai, ¹ L. Glover, ¹⁰⁹ E. Goetz, ¹¹⁸ R. Goetz, ⁵ S. Gomes, ³⁶ B. Goncharov, ⁶ G. González, ² J. M. Gonzalez Castro, ^{23,24} A. Gopakumar, ¹¹⁹ M. L. Gorodetsky, ⁶² S. E. Gossan, ¹ M. Gosselin, ³⁰ R. Gouaty, ⁸ A. Grado, ¹²⁰ A. C. Graet, ⁴⁶ M. Granata, ²⁶ A. Grant, ⁴⁶ S. Gras, ¹⁵ C. Gray, ⁴⁷ G. Greco, ^{121,122} X. Guo, ⁸² A. Gupta, ⁶⁴ M. K. Gupta, ¹⁰⁵ K. E. Gushwa, ¹ E. K. Gustafson, ¹ R. Gustafson, ¹¹⁸ O. Halim, ^{18,17} B. R. Hall, ⁶⁹ E. D. Hall, ¹⁵ E. Z. Hamilton, ³⁶ G. Hammond, ⁴⁶ M. Haney, ¹²³ M. M. Harke, ¹⁰ J. Hansson, ⁷¹ T. Hardwick, ² J. Harrs, ^{17,18} G. M. Harry, ¹²⁴ I. W. Harry, ³⁸ M. J. Hart, ⁴⁶ C.-J. Haster, ⁹⁰ K. Haughian, ⁴⁶ J. Heanly, ⁴⁸ A. W. Heptonstall, ¹ M. Heurs, ^{10,22} S. Hild, ⁴⁶ T. Heitmann, ⁶⁷ P. Hello, ²⁸ G. Hemming, ³⁰ M. Hendry, ⁴⁶ I. S. Heng, ⁴⁶ J. Henning, ⁴⁶ A. W. Heptonstall, ¹ M. Heurs, ^{10,22} S. Hild, ⁴⁶ T. Hinderer, ⁵⁶ D. Hoak, ³⁰ D. Hofman, ²⁶ K. Holt, ⁷ D. E. Holtz, ⁵⁹ J. Houghia, ³⁶ C. Horst, ²¹ J. Hough, ⁴⁶ E. A. Houston, ⁴⁶ E. J. Howell, ⁵⁵ Y. M. Hu, ¹⁰ E. A. Huerta, ¹² D. Huet, ²⁸ B. Hughey, ³⁷ S. Husa, ¹⁰² S. H. Huttner, ⁴⁶ T. Huynh-Dinh, ⁷ N. Indik, ¹⁰ R. Inta, ³⁶ G. Intini, ^{97,35} H. N. Isa, ⁴⁶ J.-M. Isac, ⁷¹ M. Isi, ¹ B. R. Iyer, ²⁰ K. Izumi, ⁴⁷ T. Jacqmin, ⁷¹ K. Jani, ⁷⁷ P. Jaranowski, ¹²⁵ S. Jawahar, ⁶³ F. Jiménez-Forteza, ¹⁰² W. W. Johnson, ²⁸ B. Kamai, ¹ S. Kandhasamy, ⁷ G. Kang, ⁴⁰ J. B. Kanner, ¹⁸ S. J. Kapadia, ²¹ S. Karki, ⁷⁰ K. S. Karvinen, ¹⁰ M. Kasprzack, ²⁰ M. Katolik, ¹² E. Kasavounidis, ¹⁵ W. Katzman, ⁷ S. Kaufer, ²² K. Kawabe, ⁴⁷ F. Kéfélian, ⁶⁷ D. Keitel, ⁴⁶ A. J. Kemball, ¹² R. Kennedy, ¹⁶⁶ C. Kent, ³⁶ J. S. Kin, ¹³⁰ N. Kim, ³⁰ W. Kim, ¹³¹ Y.-M. Kim, ⁹² S. J. Kimbrell, ⁷⁷ E. J. King, ⁷³ P. J. King, ⁴⁷ M. Kinley-Hanlon, ¹²⁴ R. Kirchhoff, ¹⁰ J. S. Kissel, ⁴⁷ L. Kleybolte, ³⁴ S. Klimenko, ⁵ T. D. Knowles, ⁴¹ P. Koch, ¹⁰ S. M. Koehlenb D. E. McClelland, ²⁵ S. McCormick, ⁷ L. McCuller, ¹⁵ S. C. McGuire, ¹³⁹ G. McIntyre, ¹ J. McIver, ¹ D. J. McManus, ²⁵ L. McNeill, ⁶ T. McRae, ²⁵ S. T. McWilliams, ⁴¹ D. Meacher, ⁶⁴ G. D. Meadors, ^{38,10} M. Mehmet, ¹⁰ J. Meidam, ¹⁴ E. Mejuto-Villa, ⁹ A. Melatos, ⁹⁶ T. McRae, ²⁵ S. T. McWilliams, ⁴¹ D. Meacher, ⁶⁴ G. D. Meadors, ^{38,10} M. Mehmet, ¹⁰ J. Meidam, ¹⁴ E. Mejuto-Villa, ⁹ A. Melatos, ⁹⁶ G. Mendell, ⁴⁷ R. A. Mercer, ²¹ E. L. Merilh, ⁴⁷ M. Merzougui, ⁶⁷ S. Meshkov, ¹ C. Messenger, ⁴⁶ C. Messick, ⁶⁴ R. Metzdorff, ⁷¹ P. M. Meyers, ⁴⁵ H. Miao, ⁵⁹ C. Michel, ²⁶ H. Middleton, ⁵⁹ E. E. Mikhailov, ¹⁴⁰ L. Milano, ^{79,4} A. L. Miller, ^{5,97,35} B. B. Miller, ⁸⁹ J. Miller, ¹⁵ M. Millhouse, ¹⁰¹ M. C. Milovich-Goff, ¹⁰⁹ O. Minazzoli, ^{67,141} Y. Minenkov, ³³ J. Ming, ³⁸ C. Mishra, ¹⁴² S. Mitra, ¹⁹ V. P. Mitrofanov, ⁶² G. Mitselmakher, ⁵ R. Mittleman, ¹⁵ D. Moffa, ⁸⁴ A. Moggi, ²⁴ K. Mogushi, ¹¹ M. Mohan, ³⁰ S. R. P. Mohapatra, ¹⁵ M. Montani, ^{121,122} C. J. Moore, ¹³ D. Moraru, ⁴⁷ G. Moreno, ⁴⁷ S. Morisaki, ⁸¹ S. R. Morriss, ¹⁰³ B. Mours, ⁸ C. M. Mow-Lowry, ⁵⁹ G. Mueller, ⁵ A. W. Muir, ³⁶ Arunava Mukherjee, ¹⁰ D. Mukherjee, ²¹ S. Mukherjee, ¹⁰³ N. Mukund, ¹⁹ A. Mullavey, ⁷ J. Munch, ⁷³ E. A. Muñiz, ⁴⁴ M. Muratore, ³⁷ P. G. Murray, ⁴⁶ K. Napier, ⁷⁷ I. Nardecchia, ^{32,33} L. Naticchioni, ^{97,35} R. K. Nayak, ¹⁴³ J. Neilson, ¹⁰⁹ G. Nelemans, ^{66,14} T. J. N. Nelson, ⁷ M. Nery, ¹⁰ A. Neunzert, ¹¹⁸ L. Nevin, ¹ J. M. Newport, ¹²⁴ G. Newton[‡], ⁴⁶ K. K. Y. Ng, ⁹³ T. T. Nguyen, ²⁵ D. Nichols, ⁶⁶ A. B. Nielsen, ¹⁰ S. Nissanke, ^{66,14} A. Nitz, ¹⁰ A. Noack, ¹⁰ F. Nocera, ³⁰ D. Nolting, ⁷ C. North, ³⁶ L. K. Nuttall, ³⁶ J. Oberling, ⁴⁷ G. D. O'Dea, ¹⁰⁹ G. H. Ogin, ¹⁴⁴ J. J. Oh, ¹³¹ S. H. Oh, ¹³¹ F. Ohme, ¹⁰ M. A. Okada, ¹⁶ M. Oliver, ¹⁰² P. Oppermann, ¹⁰ Richard J. Oram, ⁷ B. O'Reilly, ⁷ R. Ormiston, ⁴⁵ L. F. Ortega, ⁵ R. O'Shaughnessy, ⁵⁸ S. Ossokine, ³⁸ D. J. Ottaway, ⁷³

```
H. Overmier, P. B. J. Owen, S. A. E. Pace, S. J. Page, S. Page, S. A. Pai, S. Pai, Pai, S. Pai, S.
```

(LIGO Scientific Collaboration and Virgo Collaboration)

[†]Deceased, February 2017. [‡]Deceased, December 2016.

```
<sup>1</sup>LIGO, California Institute of Technology, Pasadena, CA 91125, USA
```

²Louisiana State University, Baton Rouge, LA 70803, USA

³Università di Salerno, Fisciano, I-84084 Salerno, Italy

⁴INFN, Sezione di Napoli, Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy

⁵University of Florida, Gainesville, FL 32611, USA

⁶OzGrav, School of Physics & Astronomy, Monash University, Clayton 3800, Victoria, Australia

⁷LIGO Livingston Observatory, Livingston, LA 70754, USA

⁸Laboratoire d'Annecy-le-Vieux de Physique des Particules (LAPP), Université Savoie Mont Blanc, CNRS/IN2P3, F-74941 Annecy,

⁹University of Sannio at Benevento, I-82100 Benevento, Italy and INFN, Sezione di Napoli, I-80100 Napoli, Italy

¹⁰Albert-Einstein-Institut, Max-Planck-Institut f
ür Gravitationsphysik, D-30167 Hannover, Germany

¹¹The University of Mississippi, University, MS 38677, USA

¹²NCSA, University of Illinois at Urbana-Champaign, Urbana, IL 61801, USA

¹³University of Cambridge, Cambridge CB2 1TN, United Kingdom

- ¹⁴Nikhef, Science Park, 1098 XG Amsterdam, The Netherlands ¹⁵LIGO, Massachusetts Institute of Technology, Cambridge, MA 02139, USA
 ¹⁶Instituto Nacional de Pesquisas Espaciais, 12227-010 São José dos Campos, São Paulo, Brazil ¹⁷Gran Sasso Science Institute (GSSI), I-67100 L'Aquila, Italy ¹⁸INFN, Laboratori Nazionali del Gran Sasso, I-67100 Assergi, Italy ¹⁹Inter-University Centre for Astronomy and Astrophysics, Pune 411007, India ²⁰International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bengaluru 560089, India ²¹University of Wisconsin-Milwaukee, Milwaukee, WI 53201, USA ²²Leibniz Universität Hannover, D-30167 Hannover, Germany ²³Università di Pisa, I-56127 Pisa, Italy ²⁴INFN, Sezione di Pisa, I-56127 Pisa, Italy ²⁵OzGrav, Australian National University, Canberra, Australian Capital Territory 0200, Australia ²⁶Laboratoire des Matériaux Avancés (LMA), CNRS/IN2P3, F-69622 Villeurbanne, France SUPA, University of the West of Scotland, Paisley PA1 2BE, United Kingdom
 LAL, Univ. Paris-Sud, CNRS/IN2P3, Université Paris-Saclay, F-91898 Orsay, France ²⁹California State University Fullerton, Fullerton, CA 92831, USA ³⁰European Gravitational Observatory (EGO), I-56021 Cascina, Pisa, Italy ³¹Chennai Mathematical Institute, Chennai 603103, India ³²Università di Roma Tor Vergata, I-00133 Roma, Italy ³³INFN, Sezione di Roma Tor Vergata, I-00133 Roma, Italy ³⁴Universität Hamburg, D-22761 Hamburg, Germany ³⁵INFN, Sezione di Roma, I-00185 Roma, Italy ³⁶Cardiff University, Cardiff CF24 3AA, United Kingdom ³⁷Embry-Riddle Aeronautical University, Prescott, AZ 86301, USA ³⁸ Albert-Einstein-Institut, Max-Planck-Institut für Gravitationsphysik, D-14476 Potsdam-Golm, Germany ³⁹APC, AstroParticule et Cosmologie, Université Paris Diderot, CNRS/IN2P3, CEA/Irfu, Observatoire de Paris, Sorbonne Paris Cité, F-75205 Paris Cedex 13, France ⁴⁰Korea Institute of Science and Technology Information, Daejeon 34141, Korea ⁴¹West Virginia University, Morgantown, WV 26506, USA ⁴²Università di Perugia, I-06123 Perugia, Italy ⁴³INFN, Sezione di Perugia, I-06123 Perugia, Italy ⁴⁴Syracuse University, Syracuse, NY 13244, USA ⁴⁵University of Minnesota, Minneapolis, MN 55455, USA ⁴⁶SUPA, University of Glasgow, Glasgow G12 8QQ, United Kingdom ⁴⁷LIGO Hanford Observatory, Richland, WA 99352, USA ⁴⁸Caltech CaRT, Pasadena, CA 91125, USA ⁴⁹Wigner RCP, RMKI, H-1121 Budapest, Konkoly Thege Miklós út 29-33, Hungary ⁵⁰NASA Goddard Space Flight Center, Greenbelt, MD 20771, USA ⁵¹Columbia University, New York, NY 10027, USA ⁵²Stanford University, Stanford, CA 94305, USA ⁵³Università di Camerino, Dipartimento di Fisica, I-62032 Camerino, Italy ⁵⁴Università di Padova, Dipartimento di Fisica e Astronomia, I-35131 Padova, Italy ⁵⁵INFN, Sezione di Padova, I-35131 Padova, Italy ⁵⁶Institute of Physics, Eötvös University, Pázmány P. s. 1/A, Budapest 1117, Hungary ⁵⁷Nicolaus Copernicus Astronomical Center, Polish Academy of Sciences, 00-716, Warsaw, Poland ⁵⁸Rochester Institute of Technology, Rochester, NY 14623, USA ⁵⁹University of Birmingham, Birmingham B15 2TT, United Kingdom ⁶⁰INFN, Sezione di Genova, I-16146 Genova, Italy ⁶¹RRCAT, Indore MP 452013, India
- ⁶²Faculty of Physics, Lomonosov Moscow State University, Moscow 119991, Russia
- ⁶³SUPA, University of Strathclyde, Glasgow G1 1XQ, United Kingdom
- ⁶⁴The Pennsylvania State University, University Park, PA 16802, USA
- ⁶⁵OzGrav, University of Western Australia, Crawley, Western Australia 6009, Australia
- ⁶⁶Department of Astrophysics/IMAPP, Radboud University Nijmegen, P.O. Box 9010, 6500 GL Nijmegen, The Netherlands
- ⁶⁷ Artemis, Université Côte d'Azur, Observatoire Côte d'Azur, CNRS, CS 34229, F-06304 Nice Cedex 4, France
- ⁶⁸Institut FOTON, CNRS, Université de Rennes 1, F-35042 Rennes, France
- ⁶⁹Washington State University, Pullman, WA 99164, USA
- ⁷⁰University of Oregon, Eugene, OR 97403, USA
- ⁷¹Laboratoire Kastler Brossel, UPMC-Sorbonne Universités, CNRS, ENS-PSL Research University, Collège de France, F-75005 Paris,
- ⁷²Carleton College, Northfield, MN 55057, USA

- ⁷³OzGrav, University of Adelaide, Adelaide, South Australia 5005, Australia
- ⁷⁴ Astronomical Observatory Warsaw University, 00-478 Warsaw, Poland
- ⁷⁵VU University Amsterdam, 1081 HV Amsterdam, The Netherlands
- ⁷⁶University of Maryland, College Park, MD 20742, USA
- ⁷⁷Center for Relativistic Astrophysics, Georgia Institute of Technology, Atlanta, GA 30332, USA
- ⁷⁸Université Claude Bernard Lyon 1, F-69622 Villeurbanne, France
- ⁷⁹Università di Napoli 'Federico II,' Complesso Universitario di Monte S.Angelo, I-80126 Napoli, Italy
- ⁸⁰Dipartimento di Fisica, Università degli Studi di Genova, I-16146 Genova, Italy
- ⁸¹RESCEU, University of Tokyo, Tokyo, 113-0033, Japan.
- ⁸²Tsinghua University, Beijing 100084, China
- ⁸³Texas Tech University, Lubbock, TX 79409, USA
- ⁸⁴Kenyon College, Gambier, OH 43022, USA
- ⁸⁵Departamento de Astronomía y Astrofísica, Universitat de València, E-46100 Burjassot, València, Spain
- ⁸⁶Museo Storico della Fisica e Centro Studi e Ricerche Enrico Fermi, I-00184 Roma, Italy
- ⁸⁷National Tsing Hua University, Hsinchu City, 30013 Taiwan, Republic of China
- ⁸⁸Charles Sturt University, Wagga Wagga, New South Wales 2678, Australia
- ⁸⁹Center for Interdisciplinary Exploration & Research in Astrophysics (CIERA), Northwestern University, Evanston, IL 60208, USA
- ⁹⁰Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S 3H8, Canada
- ⁹¹University of Chicago, Chicago, IL 60637, USA
- ⁹²Pusan National University, Busan 46241, Korea
- ⁹³The Chinese University of Hong Kong, Shatin, NT, Hong Kong
- ⁹⁴INAF, Osservatorio Astronomico di Padova, I-35122 Padova, Italy
- ⁹⁵INFN, Trento Institute for Fundamental Physics and Applications, I-38123 Povo, Trento, Italy
- ⁹⁶OzGrav, University of Melbourne, Parkville, Victoria 3010, Australia
- 97 Università di Roma 'La Sapienza,' I-00185 Roma, Italy
- ⁹⁸Université Libre de Bruxelles, Brussels 1050, Belgium
- ⁹⁹Sonoma State University, Rohnert Park, CA 94928, USA
- ¹⁰⁰Departamento de Matemáticas, Universitat de València, E-46100 Burjassot, València, Spain
- ¹⁰¹Montana State University, Bozeman, MT 59717, USA
- ¹⁰²Universitat de les Illes Balears, IAC3—IEEC, E-07122 Palma de Mallorca, Spain
- ¹⁰³The University of Texas Rio Grande Valley, Brownsville, TX 78520, USA
- ¹⁰⁴Bellevue College, Bellevue, WA 98007, USA
- ¹⁰⁵Institute for Plasma Research, Bhat, Gandhinagar 382428, India
- ¹⁰⁶The University of Sheffield, Sheffield S10 2TN, United Kingdom
- ¹⁰⁷Dipartimento di Scienze Matematiche, Fisiche e Informatiche, Università di Parma, I-43124 Parma, Italy
- ¹⁰⁸INFN, Sezione di Milano Bicocca, Gruppo Collegato di Parma, I-43124 Parma, Italy
- ¹⁰⁹California State University, Los Angeles, 5151 State University Dr, Los Angeles, CA 90032, USA
- ¹¹⁰Università di Trento, Dipartimento di Fisica, I-38123 Povo, Trento, Italy
- ¹¹¹Montclair State University, Montclair, NJ 07043, USA
- ¹¹²National Astronomical Observatory of Japan, 2-21-1 Osawa, Mitaka, Tokyo 181-8588, Japan
- ¹¹³Observatori Astronòmic, Universitat de València, E-46980 Paterna, València, Spain
- ¹¹⁴School of Mathematics, University of Edinburgh, Edinburgh EH9 3FD, United Kingdom
- ¹¹⁵University and Institute of Advanced Research, Koba Institutional Area, Gandhinagar Gujarat 382007, India
- ¹¹⁶IISER-TVM, CET Campus, Trivandrum Kerala 695016, India
- ¹¹⁷University of Szeged, Dóm tér 9, Szeged 6720, Hungary
- ¹¹⁸University of Michigan, Ann Arbor, MI 48109, USA
- ¹¹⁹Tata Institute of Fundamental Research, Mumbai 400005, India
- ¹²⁰INAF, Osservatorio Astronomico di Capodimonte, I-80131, Napoli, Italy
- ¹²¹Università degli Studi di Urbino 'Carlo Bo,' I-61029 Urbino, Italy
- ¹²²INFN, Sezione di Firenze, I-50019 Sesto Fiorentino, Firenze, Italy
- ¹²³Physik-Institut, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland
- ¹²⁴American University, Washington, D.C. 20016, USA
- ¹²⁵University of Białystok, 15-424 Białystok, Poland
- ¹²⁶University of Southampton, Southampton SO17 1BJ, United Kingdom
- ¹²⁷University of Washington Bothell, 18115 Campus Way NE, Bothell, WA 98011, USA
- ¹²⁸Institute of Applied Physics, Nizhny Novgorod, 603950, Russia
- ¹²⁹Korea Astronomy and Space Science Institute, Daejeon 34055, Korea
- ¹³⁰Inje University Gimhae, South Gyeongsang 50834, Korea
- ¹³¹National Institute for Mathematical Sciences, Daejeon 34047, Korea
- ¹³²NCBJ, 05-400 Świerk-Otwock, Poland
- ¹³³Institute of Mathematics, Polish Academy of Sciences, 00656 Warsaw, Poland

- ¹³⁴Hillsdale College, Hillsdale, MI 49242, USA
- ¹³⁵Hanyang University, Seoul 04763, Korea
- ¹³⁶Seoul National University, Seoul 08826, Korea
- ¹³⁷NASA Marshall Space Flight Center, Huntsville, AL 35811, USA
- ¹³⁸ESPCI, CNRS, F-75005 Paris, France
- ¹³⁹Southern University and A&M College, Baton Rouge, LA 70813, USA
- ¹⁴⁰College of William and Mary, Williamsburg, VA 23187, USA
- ¹⁴¹Centre Scientifique de Monaco, 8 quai Antoine Ier, MC-98000, Monaco
- ¹⁴²Indian Institute of Technology Madras, Chennai 600036, India
- ¹⁴³IISER-Kolkata, Mohanpur, West Bengal 741252, India
- ¹⁴⁴Whitman College, 345 Boyer Avenue, Walla Walla, WA 99362 USA
- ¹⁴⁵Indian Institute of Technology Bombay, Powai, Mumbai, Maharashtra 400076, India
- ¹⁴⁶Scuola Normale Superiore, Piazza dei Cavalieri 7, I-56126 Pisa, Italy
- ¹⁴⁷Université de Lyon, F-69361 Lyon, France
- ¹⁴⁸Hobart and William Smith Colleges, Geneva, NY 14456, USA
- ¹⁴⁹OzGrav, Swinburne University of Technology, Hawthorn VIC 3122, Australia
- ¹⁵⁰Janusz Gil Institute of Astronomy, University of Zielona Góra, 65-265 Zielona Góra, Poland
- ¹⁵¹University of Washington, Seattle, WA 98195, USA
- ¹⁵²King's College London, University of London, London WC2R 2LS, United Kingdom
- ¹⁵³Indian Institute of Technology, Gandhinagar Ahmedabad Gujarat 382424, India
- ¹⁵⁴Indian Institute of Technology Hyderabad, Sangareddy, Khandi, Telangana 502285, India
- ¹⁵⁵International Institute of Physics, Universidade Federal do Rio Grande do Norte, Natal RN 59078-970, Brazil
- ¹⁵⁶Andrews University, Berrien Springs, MI 49104, USA
- ¹⁵⁷Università di Siena, I-53100 Siena, Italy
- ¹⁵⁸Trinity University, San Antonio, TX 78212, USA
- ¹⁵⁹Abilene Christian University, Abilene, TX 79699, USA
- ¹⁶⁰Colorado State University, Fort Collins, CO 80523, USA

PACS numbers: