



Find solutions for your homework

Search

[home](#) / [study](#) / [math](#) / [statistics and probability](#) / [statistics and probability solutions manuals](#) / [the statistical sleuth](#) / [3rd edition](#) / [chapter 4](#) / [problem 31dp](#)

# The Statistical Sleuth | (3rd Edition)

[See this solution in the app](#)

## Problem

Effect of Group Therapy on Survival of Breast Cancer Patients. Researchers randomly assigned metastatic breast cancer patients to either a control group or a group that received weekly 90-minute sessions of group therapy and self-hypnosis, to see whether the latter treatment improved the patients' quality of life. The group therapy involved discussion and support for coping with the disease, but the patients were not led to believe that the therapy would affect the progression of their disease. Surprisingly, it was noticed in a follow-up 10 years later that the group therapy patients appeared to have lived longer. The data on number of months of survival after beginning of the study are shown in Display 4.17. (Data from a graph in D. Spiegel, J. R. Bloom, H. C. Kraemer, and E. Gottheil, "Effect of Psychosocial Treatment on Survival of Patients with Metastatic Breast Cancer," *Lancet* (October 14, 1989): 888–91.) Notice that three of the women in the treatment group were still alive at the time of the follow-up, so their survival times are only known to be larger than 122 months. Is there indeed evidence of an effect of the group therapy treatment on survival time and, if so, how much more time can a breast cancer patient expect to live if she receives this therapy? Analyze the data as best as possible and write a brief report of the findings.

### DISPLAY 4.17 Months of survival after beginning of study for 58 breast cancer patients

#### Control Patients ( $n = 24$ )

2, 6, 8, 10, 12, 12, 14, 14, 14, 16, 16, 16, 18, 18, 18, 20, 22, 22, 26, 34, 36, 38, 40, 48

#### Patients Given Group Therapy for One Year ( $n = 34$ )

2, 2, 4, 4, 4, 6, 6, 8, 10, 10, 12, 14, 16, 16, 16, 18, 20, 22, 32, 36, 46, 46, 48, 48, 58, 58, 66, 72, 72, 82, 122, 122\*, 122\*, 122\*

\*These three patients were still alive at the end of the 122-month study period.

## Step-by-step solution

### Step 1 of 4

There is information about the effect of group therapy on survival of breast cancer patients.

The objective is to test whether there is indeed evidence of an effect of the group therapy treatment on survival time.

For this testing, need to give hypothesis, rank-sum (Mann-Whitney) test statistics and find  $p$ -value then take decision basis on  $p$ -value.

[Comment](#)

### Step 2 of 4

The null and alternative hypothesis can be defined as follows:

$H_0$  : There is no effect of the group therapy treatment on survival time.

$H_a$  : There is an effect of the group therapy treatment on survival time.


The Mann-Whitney test statistic is calculated as follows:

## Post a question

Answers from our experts for your tough homework questions

[Continue to post](#)

20 questions remaining



**Snap a photo from your phone to post a question**  
We'll send you a one-time download link

888-888-8888

[Text me](#)

By providing your phone number, you agree to receive a one-time automated text message with a link to the app. Standard messaging rates may apply.

## My Textbook Solutions



The Statistical  
Sleuth  
3rd Edition



Business  
Statistics...  
2nd Edition



Computer...  
5th Edition

[View all solutions](#)

## Statistics Chegg tutors who can help right now



Aditya  
NIT Bhopal

77%



Adrian  
National University...

22%



Mario  
Universidad Centro...

199%

[Find me a tutor](#)



Chapter 4, Problem 31DP

2 Bookmarks

Show all steps: **ON**4) Click **Define Groups** and enter the group value.5) Click **Continue**.6) Select **Mann-Whitney U** in **Test Type**.7) Click **Ok**.
[Comment](#)
**Step 3 of 4**

SPSS Output:

Ranks				
	CancerGroup	N	Mean Rank	Sum of Ranks
SurvivalMonth	Control	24	26.54	637.00
	Therapy	34	31.59	1074.00
	Total	58		

Test Statistics <sup>a</sup>	
	SurvivalMonth
Mann-Whitney U	337.000
Wilcoxon W	637.000
Z	-1.123
Asymp. Sig. (2-tailed)	.262

a. Grouping Variable: CancerGroup

[Comment](#)
**Step 4 of 4**

From the SPSS output, the Mann-Whitney test statistic is -1.123 and the one-sided  $p$ -value is 0.262.

Decision:

The  $p$ -value is greater than 0.05 which is 0.262, so do not reject the null hypothesis. There is insufficient evidence that there is an effect of the group therapy treatment on survival time, so it is not possible to give time for breast cancer patient expect to live if she receives this therapy.

[Comments \(1\)](#)
Isn't the  $p$ -value two sided here?


Was this solution helpful?

Chapter 4, Problem 31DP

2 Bookmarks

Show all steps: ON

Salvage Logging. When wildfires ravage forests, the timber industry argues that logging the burned trees enhances forest...  
[See solution](#)

Motivation and Creativity. Find a 95% confidence interval for the treatment effect (poem creativity score after intrinsic motivation questionnaire minus poem creativity score after extrinsic motivation questionnaire, from Section 1.1.1 (file...  
[See solution](#)

[See more problems in subjects you study](#)

COMPANY	▼
LEGAL & POLICIES	▼
CHEGG PRODUCTS AND SERVICES	▼
CHEGG NETWORK	▼
CUSTOMER SERVICE	▼



© 2003-2020 Chegg Inc. All rights reserved.