

```

GET DATA /TYPE=XLSX
  /FILE='C:\Users\Jayaraj David\Downloads\project_survey_data.xlsx'
  /SHEET=name 'Sheet1'
  /CELLRANGE=full
  /READNAMES=on
  /ASSUMEDSTRWIDTH=32767.
EXECUTE.
DATASET NAME DataSet1 WINDOW=FRONT.
GET DATA /TYPE=XLSX
  /FILE='C:\Users\Jayaraj David\Downloads\project_survey_data.xlsx'
  /SHEET=name 'Sheet1'
  /CELLRANGE=full
  /READNAMES=on
  /ASSUMEDSTRWIDTH=32767.
EXECUTE.
DATASET NAME DataSet2 WINDOW=FRONT.
DATASET CLOSE DataSet1.
FREQUENCIES VARIABLES=Age Gender Education_level
  /ORDER=ANALYSIS.

```

Frequencies

Statistics

		Age	Gender	Education_level
N	Valid	209	209	209
	Missing	0	0	0

Frequency Table

Age

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	100	47.8	47.8	47.8
<18	7	3.3	3.3	51.2
18 - 21	64	30.6	30.6	81.8
22 - 25	38	18.2	18.2	100.0
Total	209	100.0	100.0	

Gender

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	100	47.8	47.8	47.8
Female	54	25.8	25.8	73.7
Male	55	26.3	26.3	100.0
Total	209	100.0	100.0	

Education_level

	Frequency	Percent	Valid Percent	Cumulative Percent
Valid	100	47.8	47.8	47.8
High School / Diploma	8	3.8	3.8	51.7
Postgraduate	71	34.0	34.0	85.6
Undergraduate	30	14.4	14.4	100.0
Total	209	100.0	100.0	

```
SAVE OUTFILE='C:\Users\Jayaraj David\OneDrive\Desktop\MSC AMS\Semester 3\AMS506PR Research Methodology and Multivariate Statistics\Mini_Project\project_dataset.sav'
/COMPRESSED.
REGRESSION
/MISSING LISTWISE
/STATISTICS COEFF OUTS R ANOVA
/CRITERIA=PIN(.05) POUT(.10)
/NOORIGIN
/DEPENDENT stresslevel
/METHOD=ENTER acad_wl peer_pressure Social_isol finance health fam_expect lack_sup.
```

Regression

```
[DataSet2] C:\Users\Jayaraj David\OneDrive\Desktop\MSC AMS\Semester 3\AMS506PR Research Methodology and Multivariate Statistics\Mini_Project\project_dataset.sav
```

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	lack_sup, health, Social_isol, acad_wl, fam_expect, peer_pressure, finance ^b	.	Enter

a. Dependent Variable: stress level

b. All requested variables entered.

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.799 ^a	.639	.614	.8073

a. Predictors: (Constant), lack_sup, health, Social_isol, acad_wl, fam_expect, peer_pressure, finance

ANOVA^a

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	116.492	7	16.642	25.537	.000 ^b
	Residual	65.819	101	.652		
	Total	182.312	108			

a. Dependent Variable: stress level

b. Predictors: (Constant), lack_sup, health, Social_isol, acad_wl, fam_expect, peer_pressure, finance

Coefficients^a

Model	Unstandardized Coefficients		Standardized Coefficients	t	Sig.
	B	Std. Error	Beta		
1 (Constant)	-.236	.279		-.845	.400
acad_wl	.352	.081	.323	4.340	.000
peer_pressure	.172	.082	.160	2.095	.039
Social_isol	.032	.079	.033	.410	.683
finance	.328	.080	.345	4.088	.000
health	.081	.066	.087	1.227	.223
fam_expect	.171	.076	.167	2.240	.027
lack_sup	.006	.067	.006	.086	.931

a. Dependent Variable: stress level

REGRESSION

```

/MISSING LISTWISE
/STATISTICS COEFF OUTS R ANOVA
/CRITERIA=PIN(.05) POUT(.10)
/NOORIGIN
/DEPENDENT stresslevel
/METHOD=ENTER Coping_freq.

```

Regression

Variables Entered/Removed^a

Model	Variables Entered	Variables Removed	Method
1	Coping_freq ^b	.	Enter

a. Dependent Variable: stress level

b. All requested variables entered.

Model Summary

Model	R	R Square	Adjusted R Square	Std. Error of the Estimate
1	.616 ^a	.379	.373	1.0287

a. Predictors: (Constant), Coping_freq

ANOVA^a

Model		Sum of Squares	df	Mean Square	F	Sig.
1	Regression	69.090	1	69.090	65.293	.000 ^b
	Residual	113.222	107	1.058		
	Total	182.312	108			

a. Dependent Variable: stress level

b. Predictors: (Constant), Coping_freq

Coefficients^a

Model		Unstandardized Coefficients		Standardized Coefficients	t	Sig.
		B	Std. Error	Beta		
1	(Constant)	5.247	.266		19.702	.000
	Coping_freq	-.795	.098	-.616	-8.080	.000

a. Dependent Variable: stress level