

AI-Driven Knowledge Graph and Retrieval-Augmented Clinical Decision Support for Rare Genetic Diseases

Project Title: AI-Driven Knowledge Graph and Retrieval-Augmented Clinical Decision Support for Rare Genetic Diseases

Submitted by: Shradha Gajelli, Jayashanmuhesh Rajasekar, Shrikesh Chaudhari

1. INTRODUCTION

Rare genetic diseases affect approximately 30 million Americans, with an estimated 80% having genetic origins.¹ Among these, muscular dystrophies represent a clinically heterogeneous group of inherited disorders characterized by progressive muscle weakness and degeneration, encompassing over 30 genetic subtypes.² The diagnostic journey for patients with rare genetic diseases often spans 5-7 years from symptom onset to genetic confirmation, with muscular dystrophies typically requiring 2-5 years for definitive diagnosis.³ This diagnostic odyssey results from multiple challenges: clinicians must interpret genetic variants across more than 200 genes associated with neuromuscular disorders, synthesize data fragmented across disparate databases (ClinVar, OMIM, GeneReviews, PubMed, LOVD), and navigate complex genotype-phenotype correlations where the same gene can produce vastly different phenotypes.

The complexity of variant interpretation is compounded by the prevalence of Variants of Uncertain Significance (VUS), which account for 40-50% of genetic test results and require extensive literature review, database searches, and expert consultation.⁴ For clinicians managing rare genetic diseases, this translates to 30-60 minutes spent per variant researching across multiple platforms—time that delays therapeutic decisions and potentially life-altering interventions. The emergence of mutation-specific therapies, such as exon-skipping drugs for Duchenne Muscular Dystrophy (DMD), has elevated the urgency of accurate and rapid genetic diagnosis, as therapeutic efficacy is contingent upon precise genotype matching.⁵

Knowledge Graphs and Retrieval-Augmented Generation as Clinical Innovation

Knowledge graphs offer a structured approach to representing biomedical entities (genes, variants, diseases, phenotypes, treatments) and their complex interrelationships through a network of nodes and edges. When integrated with Retrieval-Augmented Generation (RAG)—a technique that enhances large language models by incorporating external, curated data sources—knowledge graphs enable real-time, evidence-based recommendations at the point of care.^{6,7} RAG addresses fundamental limitations of standalone AI models, including the tendency to generate inaccurate ("hallucinated") content and lack of access to current medical literature, by grounding responses in vetted, domain-specific knowledge bases.⁸

This integration bridges the gap between genomic data generation and clinical interpretation, transforming variant calls into actionable clinical insights. By synthesizing information across variant pathogenicity databases, clinical management guidelines, pharmacological databases, and active clinical trial registries, a RAG-enabled knowledge graph can support clinicians in variant interpretation, differential diagnosis, treatment selection, and clinical trial matching—addressing critical decision points that currently rely on time-intensive manual research.

Thesis Statement

This project proposes building an AI-driven knowledge graph integrated with a RAG pipeline to enhance evidence-based decision support for four genetically and phenotypically distinct muscular dystrophies: Duchenne Muscular Dystrophy (DMD gene), Becker Muscular Dystrophy (DMD gene), Limb-Girdle Muscular Dystrophy Type R1/LGMD2A (CAPN3 gene), and LAMA2-related Congenital Muscular Dystrophy (LAMA2 gene). This system will reduce diagnostic delays, improve treatment selection accuracy, and provide actionable clinical insights by integrating genomic data with evidence-based medicine, ultimately advancing precision medicine for rare disease populations.

2. PROBLEM STATEMENT & OPPORTUNITY

Current State: Data Fragmentation and Clinical Workflow Challenges

The genomic medicine landscape is characterized by profound information silos that impede clinical decision-making. ClinVar, maintained by the National Institutes of Health, contains over 2.5 million variant entries with pathogenicity classifications, yet cross-referencing these with phenotypic databases (OMIM's 16,000+ gene-disease relationships) and therapeutic guidelines (GeneReviews' expert-authored summaries) requires manual navigation of disparate interfaces.^{9,10} Locus-specific databases such as the UMD-DMD database, which catalogs over 10,000 DMD variants, provide granular mutation data but exist independently from clinical management resources.¹¹ PubMed's 35+ million articles represent an overwhelming corpus that, while comprehensive, lacks structured accessibility for time-constrained clinicians.

Specific Clinical Gaps Addressed by This Project

1. Variant Interpretation Delays

The clinical significance of genetic variants falls along a spectrum defined by the American College of Medical Genetics and Genomics (ACMG) guidelines: Pathogenic, Likely Pathogenic, Uncertain Significance, Likely Benign, and Benign.¹² Variants of Uncertain Significance (VUS) constitute 40-50% of genetic test results and require evidence synthesis from population databases, functional studies, computational predictions, and familial segregation data.⁴ For muscular dystrophies, this process is particularly time-sensitive given the availability of mutation-specific therapies such as eteplirsen (exon 51 skipping, FDA approved 2016),

golodirsen and viltolarsen (exon 53 skipping, approved 2019-2020), and casimersen (exon 45 skipping, approved 2021) for DMD patients with specific deletion patterns.^{13,14} Each day of delayed interpretation potentially defers access to disease-modifying therapy.

2. Differential Diagnosis Complexity

Muscular dystrophies present significant diagnostic challenges due to phenotypic overlap. LGMD2A (calpainopathy), BMD, and other LGMD subtypes share clinical features including proximal muscle weakness, elevated creatine kinase levels, and variable age of onset.¹⁵ Distinguishing these conditions requires pattern recognition across genetic findings, muscle biopsy results, imaging characteristics, and cardiac involvement—information that is rarely synthesized in a unified format. Misdiagnosis leads to inappropriate management: for instance, LGMD2A typically lacks cardiac involvement, obviating the intensive cardiac surveillance protocols required for DMD/BMD.¹⁶

3. Treatment Selection Challenges

The advent of mutation-specific therapies has introduced a new paradigm in DMD management but has also created complexity in matching patients to appropriate treatments. Each of the four FDA-approved exon-skipping antisense oligonucleotides targets specific deletion patterns, with eteplirsen applicable to approximately 13% of DMD patients (those amenable to exon 51 skipping), and golodirsen/viltolarsen and casimersen each applicable to approximately 8% (exons 53 and 45, respectively).^{13,17} Determining eligibility requires computational analysis of whether a patient's deletion pattern would restore the reading frame following exon skipping—a determination that is non-trivial for clinicians without bioinformatics support. Additionally, the clinical trial landscape for neuromuscular diseases includes over 150 active trials globally, yet matching patients to appropriate studies requires awareness of eligibility criteria that include specific genetic findings.¹⁸

4. Genotype-Phenotype Correlation Complexity

The DMD gene exemplifies the challenge of variable phenotypes from a single genetic locus. Out-of-frame deletions and nonsense mutations typically result in complete dystrophin loss and the severe DMD phenotype (wheelchair dependence by age 12, life expectancy 20s-30s), while in-frame deletions produce partially functional dystrophin and the milder BMD phenotype (ambulation into 30s-50s, near-normal life expectancy with cardiac management).^{19,20} The "reading frame rule," which predicts phenotype based on whether a deletion maintains the translational reading frame, has approximately 90% predictive accuracy but requires molecular genetics expertise to apply.²¹ Similarly, LAMA2 mutations demonstrate allelic heterogeneity: complete laminin- α 2 deficiency causes severe congenital muscular dystrophy (MDC1A) with no independent ambulation, while partial deficiency allows for milder, later-onset presentations that may achieve walking.²²

Opportunity Statement

A unified, AI-enhanced knowledge framework can provide actionable insights at the point of care, accelerating precision medicine for rare disease patients. By integrating structured genomic knowledge with retrieval-augmented generation, clinicians can access real-time, evidence-based recommendations for variant interpretation, differential diagnosis, and treatment selection—reducing diagnostic delays from years to days and enabling timely access to life-altering therapies. This system democratizes access to subspecialty expertise, particularly benefiting community hospitals and rural healthcare settings where genetic specialists may be unavailable.

Alignment with Clinical Quality Measures

This project aligns with multiple domains of clinical quality improvement:

- **Care Coordination:** Integrates multidisciplinary evidence spanning genetics, cardiology, pulmonology, and neurology into unified recommendations
- **Diagnostic Accuracy:** Reduces misdiagnosis through comprehensive genotype-phenotype correlation and differential diagnosis support
- **Treatment Appropriateness:** Matches patients to eligible mutation-specific therapies and clinical trials
- **Patient Safety:** Prevents inappropriate treatments based on genetic findings (e.g., avoiding unnecessary cardiac interventions in LGMD2A patients)
- **Health Equity:** Democratizes access to expert-level rare disease knowledge regardless of geographic location or institutional resources

Quantifiable Clinical Impact

Evidence supports the significant clinical impact of timely genetic diagnosis in muscular dystrophies. Corticosteroid initiation in DMD patients prolongs ambulation by 2-3 years, but this benefit is maximized with early treatment.²³ Appropriate exon-skipping therapy selection can slow disease progression in eligible patients.²⁴ Early diagnosis of LAMA2-CMD enables proactive respiratory management, improving outcomes and reducing acute hospitalizations.²⁵ Conversely, LGMD2A genetic confirmation prevents unnecessary cardiac interventions and guides appropriate counseling regarding prognosis, which differs substantially from DMD/BMD.²⁶

3. CLINICAL RATIONALE

3.1 Duchenne Muscular Dystrophy (DMD gene)

Epidemiology and Clinical Significance

Duchenne Muscular Dystrophy (DMD) represents the most common fatal childhood genetic disease, with a global birth prevalence of 19.8 per 100,000 live male births, corresponding to approximately 1 in 3,500-5,000 male births.²⁷ As an X-linked recessive disorder, DMD predominantly affects males, though 8-10% of female carriers may experience mild symptoms due to skewed X-inactivation.²⁸ The clinical course is characterized by progressive proximal muscle weakness beginning at ages 2-5 years, with classic early signs including delayed motor milestones, frequent falls, difficulty climbing stairs, and Gowers' maneuver (using hands to push up from the floor). Calf pseudohypertrophy, a pathognomonic finding resulting from muscle fiber replacement with fat and connective tissue, is often present.²⁹

Disease progression follows a predictable trajectory: loss of ambulation typically occurs by age 12, followed by progressive scoliosis, respiratory insufficiency requiring non-invasive ventilation (NIV) by the late teens, and dilated cardiomyopathy which develops in nearly all patients.³⁰ Historically, death occurred in the late teens to early twenties from cardiopulmonary failure. However, contemporary multidisciplinary management—including corticosteroids, cardiac surveillance, and respiratory support—has extended median survival to the late twenties to early thirties in high-resource settings, with some patients living into their forties.³¹

Genetic Mechanism

The DMD gene, located on chromosome Xp21, is the largest known human gene, spanning 2.2 million base pairs and comprising 79 exons.³² Out-of-frame deletions (60-65% of cases), duplications (5-10%), or nonsense mutations (10-15%) result in complete dystrophin absence. The dystrophin protein, normally expressed at the muscle fiber sarcolemma, provides structural integrity by linking the intracellular cytoskeleton to the extracellular matrix through the dystrophin-associated glycoprotein complex.³³ Dystrophin deficiency renders muscle membranes susceptible to contraction-induced damage, triggering cycles of degeneration and regeneration that ultimately lead to muscle fiber replacement with fibrofatty tissue.

Evidence-Based Management Guidelines

The landmark 2018 DMD Care Considerations, published in three parts in The Lancet Neurology, represents the gold standard for comprehensive DMD management.^{23,34,35} These guidelines, developed by an international expert working group, provide evidence-graded recommendations across the patient lifespan:

Corticosteroid Therapy:

- **Indication:** All boys with DMD capable of walking
- **Agents:** Prednisone (0.75 mg/kg/day) or deflazacort (0.9 mg/kg/day)
- **Evidence Level:** Level A (high-quality evidence from multiple studies)
- **Efficacy:** Prolongs ambulation by 2-3 years, delays scoliosis, preserves respiratory and cardiac function

- **Adverse Effects:** Require monitoring for weight gain, bone health (bisphosphonates for fracture prevention), behavioral changes, cushingoid features

Cardiac Management:

- **Surveillance:** Cardiac MRI or echocardiography annually starting age 10
- **Prophylactic ACE inhibitors:** Initiate even if asymptomatic, based on evidence of delaying cardiomyopathy onset
- **Evidence:** Multiple studies demonstrate reduced mortality with early cardiac intervention³⁶

Respiratory Management:

- **Surveillance:** Pulmonary function tests (PFTs) every 6 months after age 12 or loss of ambulation
- **NIV Initiation:** When forced vital capacity (FVC) <50% predicted or symptoms of hypoventilation
- **Cough Assistance:** Mechanical insufflation-exsufflation for peak cough flow <270 L/min
- **Evidence:** NIV prolongs survival by years and improves quality of life³⁷

Genetic Therapies Approved 2016-2023

The therapeutic landscape for DMD has been revolutionized by mutation-specific approaches:

Exon-Skipping Antisense Oligonucleotides (AONs):

- **Eteplirsen (Exondys 51):** FDA approved September 2016 for exon 51 skipping (~13% of DMD patients)
- **Golodirsen (Vyondys 53):** FDA approved December 2019 for exon 53 skipping (~8% of patients)
- **Viltolarsen (Viltepso):** FDA approved August 2020 for exon 53 skipping (~8% of patients)
- **Casimersen (Amondys 45):** FDA approved February 2021 for exon 45 skipping (~8% of patients)
^{13,14,38}

Mechanism: These phosphorodiamidate morpholino oligomers (PMOs) bind to exon-specific sequences in pre-mRNA, causing the spliceosome to skip the targeted exon during processing. For patients with out-of-frame deletions, skipping an adjacent exon can restore the reading frame, resulting in production of internally truncated but partially functional dystrophin (similar to BMD). For example, a patient with a deletion of exons 45-52 (out-of-frame) would be amenable to exon 53 skipping with golodirsen or viltolarsen, restoring the frame and converting the molecular phenotype to BMD-like.

Ataluren (Translarna): Approved in Europe (not FDA approved in the US) for nonsense mutations, which constitute 10-15% of DMD cases. Mechanism involves readthrough of premature stop codons.³⁹

Gene Therapy:

- **Elevidys (delandistrogene moxeparvovec):** FDA approved June 2023 for patients aged 4-5 years. Uses AAV vector to deliver a shortened (micro-dystrophin) gene. First gene therapy approved for DMD.

How Knowledge Graph + RAG Supports Clinical Decisions

Example Clinical Scenario:

- **Patient:** 5-year-old male with elevated CK (15,000 U/L), proximal weakness, positive Gowers' sign
- **Genetic Testing:** DMD gene deletion of exons 48-52

Knowledge Graph Retrieval:

1. Query deletion pattern: exons 48-52 → Out-of-frame deletion (frameshift)
2. Retrieve DMD phenotype information (severe course)
3. Reading frame analysis: Deletion of exon 53 would restore frame (exons 48-52 deleted + skip 53 = in-frame 48-53 deletion)
4. Retrieve eligible drugs: Golodirsen (Vyondys 53), Viltolarsen (Viltepso)
5. Retrieve DMD management guidelines: Corticosteroids, cardiac surveillance, respiratory monitoring
6. Retrieve active clinical trials: Query ClinicalTrials.gov for DMD trials matching deletion pattern

RAG-Generated Clinical Summary: "Genetic findings confirm Duchenne Muscular Dystrophy (DMD) with out-of-frame deletion of exons 48-52. Patient is eligible for exon 53 skipping therapy:

Treatment Recommendations:

1. **Exon-Skipping Therapy:** Golodirsen (Vyondys 53, 30 mg/kg IV weekly) or Viltolarsen (Viltepso, 80 mg/kg IV weekly). Evidence: FDA-approved based on dystrophin expression increases; monitor renal function (nephrotoxicity risk).
2. **Corticosteroids:** Initiate deflazacort 0.9 mg/kg/day. Evidence: Prolongs ambulation 2-3 years (Level A).
3. **Cardiac Surveillance:** Baseline cardiac MRI; initiate ACE inhibitor at age 10 regardless of symptoms (prophylactic benefit demonstrated).
4. **Respiratory Monitoring:** PFTs every 6 months after age 12.
5. **Clinical Trials:** [List NCT identifiers for active DMD trials]

Prognosis: With contemporary management including mutation-specific therapy, anticipated wheelchair dependence age 13-15, life expectancy 30s-40s with cardiac/respiratory support.

Genetic Counseling: X-linked recessive; mother is obligate carrier. 50% risk for male siblings. Offer carrier testing for female relatives."

3.2 Becker Muscular Dystrophy (BMD gene - same as DMD)

Epidemiology and Clinical Distinction

Becker Muscular Dystrophy (BMD), with an incidence of 1 in 18,000-30,000 male births, represents an allelic variant of DMD—caused by mutations in the same DMD gene but resulting in a milder phenotype.

⁴¹ This fundamental genetic paradox—identical locus, vastly different clinical courses—presents a critical challenge for clinical decision support systems and underscores the importance of genotype-phenotype correlation. BMD typically manifests between ages 5-15 years with slower progressive proximal muscle weakness, preserved ambulation into the third to fifth decade or beyond, and near-normal life expectancy with appropriate cardiac management.⁴² However, phenotypic variability is substantial; some individuals remain nearly asymptomatic into adulthood, while others experience more rapid progression approaching the DMD trajectory.

Genetic Mechanism: The Reading Frame Rule

The distinction between DMD and BMD phenotypes is predominantly explained by the "reading frame rule," formulated by Monaco et al. in their seminal 1988 Science publication.²¹ In-frame deletions or duplications preserve the triplet codon reading frame, allowing translation of a shortened but partially functional dystrophin protein with intact N-terminal (actin-binding) and C-terminal (dystroglycan-binding) domains. This internally truncated dystrophin provides some structural support to the muscle membrane, resulting in the milder BMD phenotype. Conversely, out-of-frame mutations introduce premature stop codons, resulting in complete dystrophin absence and the severe DMD phenotype.

The reading frame rule demonstrates approximately 90% predictive accuracy, with exceptions typically involving:

- Mutations affecting critical functional domains
- Mosaic expression of dystrophin
- Alternative splicing events
- Mutations in regulatory regions affecting expression levels⁴³

Example:

- DMD: Deletion of exons 45-51 → Out-of-frame (frameshift) → No dystrophin → DMD
- BMD: Deletion of exons 45-50 → In-frame deletion → Truncated dystrophin → BMD

Clinical Management Differences from DMD

Corticosteroid Approach: BMD management is less standardized than DMD, with corticosteroid use determined on an individual basis rather than universal recommendation. When used, dosing is typically more conservative than DMD protocols.⁴⁴

Cardiac Surveillance - Critical Importance: A key clinical insight from Melacini et al. (1996) is that cardiomyopathy in BMD can occur independently of skeletal muscle severity.⁴⁵ Patients with minimal or no muscle weakness may develop life-threatening dilated cardiomyopathy, necessitating:

- Annual cardiac MRI or echocardiography regardless of ambulatory status
- Prophylactic ACE inhibitor therapy upon detection of cardiac involvement
- Beta-blockers for heart failure management
- Consideration for implantable cardioverter-defibrillator (ICD) for arrhythmias

Exercise Recommendations: Unlike DMD, where vigorous exercise may accelerate muscle damage, moderate aerobic exercise is beneficial in BMD for maintaining cardiovascular fitness and muscle function.⁴⁶

CDS Clinical Decision Points

1. Phenotype Prediction:

- **Input:** DMD gene deletion pattern
- **Output:** Prediction of DMD vs. BMD based on reading frame analysis
- **Clinical Utility:** Appropriate prognostic counseling, life planning, treatment intensity decisions

2. Prognostic Counseling:

- **DMD:** Wheelchair by age 12, life expectancy 20s-30s
- **BMD:** Ambulatory into 30s-50s+, near-normal life expectancy with cardiac management
- Profound implications for patient and family expectations, educational planning, career decisions

3. Differential Diagnosis: BMD phenotypically overlaps with LGMD, requiring genetic confirmation for accurate diagnosis. Both present with:

- Later childhood/adolescent onset proximal weakness
- Elevated CK levels
- Preserved ambulation into adulthood

- Distinguishing features: BMD has X-linked inheritance, more consistent cardiac involvement than most LGMDs

Knowledge Graph Application Example

Clinical Scenario:

- **Patient:** 10-year-old male, difficulty running, CK 8,000 U/L, positive family history (maternal uncle had "muscle disease")
- **Genetic Testing:** DMD gene deletion of exons 45-50

Knowledge Graph Retrieval:

1. Deletion pattern: exons 45-50 → Frame analysis → In-frame deletion
2. Reading frame rule application → BMD phenotype predicted
3. Retrieve BMD clinical course data
4. Retrieve BMD-specific management recommendations
5. Query family history implications (X-linked inheritance)

RAG-Generated Clinical Summary: "In-frame deletion of exons 45-50 in the DMD gene is consistent with Becker Muscular Dystrophy (BMD) phenotype.

Predicted Clinical Course:

- Milder phenotype than DMD
- Expected ambulation into adulthood (30s-50s or beyond)
- Variable progression; some patients remain minimally symptomatic
- Late-onset cardiomyopathy risk (can occur independently of skeletal muscle severity)

Management Recommendations:

1. **Cardiac Surveillance (CRITICAL):** Annual cardiac MRI starting now; early ACE inhibitor if cardiac dysfunction detected (Evidence: Melacini 1996, Bushby 1993)
2. **Corticosteroids:** Not routinely recommended; consider on individual basis if rapid progression
3. **Exercise:** Encourage moderate aerobic exercise (beneficial in BMD, unlike DMD)
4. **Physical Therapy:** Maintain range of motion, prevent contractures
5. **NOT Eligible for Exon-Skipping Therapies:** In-frame deletion produces functional dystrophin; exon-skipping would not provide benefit

Prognosis: Near-normal life expectancy with cardiac management. Most maintain ambulation beyond age 40.

Genetic Counseling: X-linked inheritance; mother is carrier (confirmed by family history). 50% risk for male siblings, 50% risk daughters are carriers. Offer cascade genetic testing for at-risk family members."

3.3 Limb-Girdle Muscular Dystrophy Type R1 (LGMD2A) - CAPN3 Gene

Epidemiology and Clinical Presentation

Limb-Girdle Muscular Dystrophy Type R1 (LGMDR1), historically termed LGMD2A, represents the most common autosomal recessive LGMD subtype, accounting for 10-30% of all LGMD cases and an estimated 30-47% of LGMD cases in some populations.^{47,48} The updated nomenclature (LGMDR1, where "R" denotes recessive and "1" indicates it was the first recessive form identified) reflects contemporary classification systems.⁴⁹ Autosomal recessive inheritance distinguishes LGMDR1 from DMD/BMD, with equal gender distribution and 25% recurrence risk for siblings of affected individuals. Founder mutations in specific populations (Basque Country, Réunion Island, Amish communities) result in locally elevated prevalence.⁵⁰

Clinical Features:

- **Age of Onset:** Typically 8-15 years, with range from 2-40 years⁵¹
- **Distribution of Weakness:**
 - Pelvic girdle > shoulder girdle initially
 - **Scapular winging** prominent and often an early, distinguishing feature
 - Progression generally symmetric and proximal
- **Gait Abnormalities:** Waddling gait, tendency to walk on tiptoes, difficulty with stairs
- **Muscle Hypertrophy:** Calf pseudohypertrophy common (similar to DMD/BMD)
- **Contractures:** Achilles tendon contractures, laxity of abdominal muscles
- **CK Elevation:** 5-30 times normal (less than DMD but significantly elevated)
- **Cardiac Involvement:** **Typically absent** (key distinguishing feature from DMD/BMD)⁵²
- **Cognition:** Normal intelligence; no CNS involvement
- **Ambulation Loss:** Variable, typically 10-20 years after symptom onset (20s-40s); milder phenotypes may retain ambulation beyond age 60

Genetic Mechanism

The CAPN3 gene, mapped to chromosome 15q15.1, encodes calpain-3, a muscle-specific calcium-activated neutral protease (94 kDa protein, hence alternative name p94).⁵³ Calpain-3 serves dual functions:

1. **Proteolytic Function:** Cleaves sarcomeric and cytoskeletal proteins (titin, filamin, talin), designating them for ubiquitin-proteasome degradation—a critical step in muscle remodeling
2. **Structural Function:** Anchors to the sarcomere, contributing to muscle fiber integrity

Over 500 CAPN3 mutations have been identified, demonstrating considerable allelic heterogeneity.

⁵⁴ Loss-of-function mutations result in impaired muscle membrane repair, altered calcium homeostasis, and dysregulation of the ubiquitin-proteasome pathway. Intriguingly, approximately 20% of LGMDR1 patients have normal calpain-3 protein levels on Western blot but lack autocatalytic activity—the protein's self-cleavage that is essential for function—underscoring the importance of genetic over solely protein-based diagnosis.⁵⁵

Diagnostic Challenges: Overlapping Phenotypes

LGMDR1 presents significant differential diagnostic challenges due to phenotypic overlap with:

- **Becker Muscular Dystrophy:** Both have later onset, proximal weakness, preserved ambulation into adulthood
- **Other LGMD Subtypes:** LGMDR2 (dysferlinopathy), LGMDR9 (FKRP-related), sarcoglycanopathies
- **Facioscapulohumeral Muscular Dystrophy (FSHD):** Both feature scapular winging

Distinguishing Features Favoring LGMDR1:

- Autosomal recessive inheritance (vs. X-linked for BMD)
- Prominent scapular winging (early and consistent)
- **Absence of cardiac involvement** (unlike DMD/BMD)
- Specific muscle MRI patterns (preferential involvement of posterior thigh compartment)⁵⁶

Evidence-Based Management

The comprehensive GeneReviews chapter by Vissing et al. (2016) provides authoritative clinical guidance for calpainopathy management.⁵⁷

Current Management (No Disease-Modifying Therapy Available):

1. Physical Therapy:

- Maintain functional capacity and range of motion
- Prevent contractures (especially Achilles tendons)

- Strengthen unaffected muscle groups
2. **Assistive Devices:** Ankle-foot orthoses (AFOs), wheelchairs, adaptive equipment as disease progresses
 3. **Respiratory Monitoring:** Less common than DMD but can occur; annual PFTs recommended
 4. **Baseline Cardiac Screening:** Although typically spared, baseline ECG/echocardiogram recommended; routine surveillance not required if normal
 5. **Orthopedic Management:** Scoliosis surveillance, contracture management
 6. **Genetic Counseling:**
 - 25% recurrence risk for siblings
 - Carrier testing for relatives
 - Prenatal diagnosis options for future pregnancies

Emerging Therapies:

- **Gene Therapy:** CAPN3 gene replacement strategies in preclinical and early clinical development⁵⁸
- **Clinical Trial Matching:** Critical for patients to be aware of enrollment opportunities

CDS Value for Differential Diagnosis

Example Clinical Scenario:

- **Patient:** 12-year-old female with difficulty running, shoulder weakness
- **Examination:** Prominent bilateral scapular winging, proximal muscle weakness (4/5 hip flexors, 4+/5 shoulder abductors), normal cardiac examination
- **Laboratory:** CK 4,500 U/L
- **Family History:** First cousin with similar symptoms (suggests autosomal recessive)

Knowledge Graph Retrieval:

1. Clinical features: Scapular winging + proximal weakness + adolescent onset + normal cardiac → High probability LGMDR1
2. CK level: Consistent with multiple myopathies
3. Autosomal recessive inheritance pattern (affected cousin)
4. Differential diagnosis database query:
 - LGMDR1 (CAPN3): Matches all features

- BMD (DMD): Less likely (female gender, autosomal recessive pattern, prominent scapular winging)
- FSHD: Possible (scapular winging) but typically asymmetric facial involvement
- Other LGMD subtypes: Possible

RAG-Generated Clinical Summary: "Clinical presentation most consistent with Limb-Girdle Muscular Dystrophy Type R1 (LGMDR1, calpainopathy).

Differential Diagnosis:

1. LGMDR1 (CAPN3) - Most Likely

- Rationale: Scapular winging + proximal weakness + adolescent onset + autosomal recessive pattern + normal cardiac + CK elevation

2. BMD (DMD) - Less Likely

- Against: Female gender, prominent scapular winging, family history suggests AR inheritance

3. FSHD - Possible

- Against: Symmetric involvement, no facial weakness

Recommended Genetic Testing: Neuromuscular gene panel including: CAPN3 (LGMDR1), DMD, FKRP (LGMDR9), DYSF (LGMDR2), ANO5 (LGMDR12), sarcoglycan genes (LGMDR3-6)

If CAPN3 Mutations Confirmed:

1. Prognosis: Variable progression; ambulation loss typically 20s-40s; wheelchair dependence expected

2. Management:

- Physical therapy (maintain function, prevent contractures)
- Respiratory monitoring (annual PFTs)
- **Cardiac surveillance NOT routinely required** (spared in LGMDR1, unlike DMD/BMD)
- Orthopedic follow-up (scoliosis, contractures)

3. Clinical Trial Screening: Check for gene therapy trials (CAPN3 gene replacement)

4. Genetic Counseling:

- Autosomal recessive: 25% recurrence risk for siblings
- Offer carrier testing to family members
- Prenatal diagnosis options for future pregnancies (PGT, CVS, amniocentesis)

Prognosis: Life expectancy typically normal; quality of life impacted by progressive weakness and eventual wheelchair dependence. Cardiac and respiratory complications less common than DMD."

3.4 LAMA2-Related Congenital Muscular Dystrophy (LAMA2 gene)

Epidemiology and Clinical Significance

LAMA2-related muscular dystrophy, also termed merosin-deficient congenital muscular dystrophy type 1A (MDC1A), represents one of the most common forms of congenital muscular dystrophy worldwide, accounting for approximately one-third of CMD cases.⁵⁹ Prevalence is estimated at 0.6-0.7 per 100,000 in the UK and Italy.⁶⁰ As an autosomal recessive disorder, LAMA2-related dystrophies affect both genders equally and carry a 25% recurrence risk for siblings.

The clinical spectrum encompasses two primary phenotypes determined by mutation severity:

1. Severe, Early-Onset Form (Classic MDC1A):

- **Complete Laminin- α 2 Deficiency:** Typically due to nonsense or frameshift mutations resulting in absent protein
- **Clinical Features:**
 - **Neonatal Presentation:** Severe hypotonia evident at birth ("floppy infant"), poor spontaneous movements
 - **Feeding Difficulties:** Poor suck, failure to thrive, gastroesophageal reflux, aspiration risk
 - **Respiratory Insufficiency:** Early onset, often requiring ventilatory support in infancy
 - **Contractures:** Develop early (hips, knees, ankles, elbows)
 - **Ambulation:** Most patients never achieve independent walking
 - **Scoliosis:** Progressive, often severe
 - **White Matter Abnormalities:** Characteristic T2 hyperintensities on brain MRI (diagnostic clue, present in >90% of cases)⁶¹
 - **Cognition:** Typically **normal** (key differentiating feature from other CMDs with brain involvement)
 - **Seizures:** Occur in 5-30% of patients, often with focal features and autonomic symptoms; more common with extensive cortical malformations⁶²
 - **Cardiac:** Generally **spared** (no cardiomyopathy, unlike DMD/BMD)
 - **CK Elevation:** 5-20 times normal
 - **Life Expectancy:** Teens to 30s with supportive care (reduced compared to general population)

2. Milder, Later-Onset Form (Partial LAMA2 Deficiency):

- **Partial Laminin- α 2 Deficiency:** Typically due to missense mutations allowing residual protein expression
- **Clinical Features:**
 - Onset in childhood or adolescence
 - Limb-girdle pattern of weakness
 - Some patients achieve ambulation
 - Variable severity, milder contractures
 - White matter changes still present on MRI⁶³

Genetic Mechanism

The LAMA2 gene, located on chromosome 6q22-q23, encodes the laminin- α 2 chain (also called merosin).⁶⁴ This subunit combines with laminin- β 1 and laminin- γ 1 chains to form the heterotrimeric laminin-211 protein (also called laminin-2 or merosin), a critical component of the basement membrane in skeletal muscle, peripheral nerves, and brain. Laminin-211 bridges the sarcolemma to the extracellular matrix, binding intracellularly to α -dystroglycan (part of the dystrophin-glycoprotein complex) and extracellularly to collagen and other matrix proteins.

Pathophysiology: Complete or partial absence of laminin- α 2 disrupts this structural linkage, causing:

- Muscle fiber membrane instability and susceptibility to contraction-induced damage
- Defective muscle regeneration and increased apoptosis
- Peripheral nerve dysmyelination (causing delayed motor nerve conduction velocities)
- Brain white matter abnormalities (mechanism incompletely understood, possibly related to blood-brain barrier dysfunction)

Over 300 LAMA2 mutations have been reported, with genotype-phenotype correlations:⁶⁵

- Nonsense, frameshift, splice-site mutations → Complete deficiency → Severe MDC1A
- Missense mutations → Partial deficiency → Milder phenotypes

Distinctive Diagnostic Features

Brain MRI - Pathognomonic White Matter Changes: The presence of diffuse, symmetric T2 hyperintensities in the cerebral white matter is a hallmark diagnostic feature of LAMA2-CMD, present in >90% of patients with complete deficiency.⁶⁶ This finding, combined with elevated CK and early-onset hypotonia, strongly suggests LAMA2-CMD even before genetic confirmation. Importantly, these white matter changes do

not correlate with cognitive impairment; most patients have normal intelligence despite extensive white matter signal abnormality.

Muscle Biopsy:

- Immunohistochemistry: Absent or reduced laminin- α 2 staining (merosin deficiency)
- Dystrophic changes: Fiber size variation, necrosis, regeneration, fibrosis
- Note: Secondary laminin- α 2 deficiency can occur in dystroglycanopathies, necessitating genetic confirmation

Evidence-Based Management

The GeneReviews chapter on LAMA2 Muscular Dystrophy provides comprehensive clinical guidance.
⁶⁷

Current Management (No Curative Therapy):

1. Respiratory Support:

- **Critical intervention:** Respiratory insufficiency is life-threatening
- Non-invasive ventilation (NIV): Often required in infancy or early childhood
- Tracheostomy: May be necessary for severe cases
- Cough assistance: Mechanical insufflation-exsufflation for secretion clearance
- Surveillance: Regular PFTs, sleep studies to detect hypoventilation

2. Nutritional Support:

- Nasogastric or gastrostomy tube feeding for safe nutrition
- Prevents aspiration, ensures adequate caloric intake
- Multidisciplinary approach with speech pathology, gastroenterology

3. Orthopedic Management:

- Contracture prevention: Physical therapy, stretching, night splints
- Scoliosis surveillance: Bracing or surgical intervention if severe
- Hip dislocation monitoring

4. Neurological Management:

- Seizure monitoring (5-30% incidence)
- Antiepileptic drugs as needed

5. Supportive Equipment:

- Wheelchairs, standers, adaptive equipment
- Environmental modifications

Emerging Therapies:

- **Omigapil:** Anti-apoptotic compound showed muscle histology improvement in Phase 2 trial but failed to demonstrate functional benefit in Phase 3⁶⁸
- **Antisense Oligonucleotides:** Strategies to upregulate LAMA1 (laminin- α 1) as compensatory mechanism in preclinical development
- **Gene Therapy:** AAV-mediated LAMA2 gene delivery in preclinical studies⁶⁹

CDS Critical Role: Early Diagnosis & Proactive Management

Example Clinical Scenario:

- **Patient:** 3-month-old infant with severe hypotonia, poor feeding, weak cry
- **Examination:** Generalized hypotonia, minimal antigravity movements, normal alertness and social interaction
- **Laboratory:** CK 3,500 U/L
- **Brain MRI:** Diffuse, symmetric T2 hyperintensities in cerebral white matter

Knowledge Graph Retrieval:

1. Clinical features: Early-onset hypotonia + elevated CK + white matter changes → High probability LAMA2-CMD
2. White matter changes database: Characteristic finding in LAMA2-CMD (>90% sensitive)
3. Retrieve LAMA2 mutation types and predicted severity
4. Retrieve LAMA2-CMD management protocols
5. Query prognosis data based on phenotype

RAG-Generated Clinical Summary: "Clinical presentation and MRI findings highly suggestive of LAMA2-related Congenital Muscular Dystrophy (MDC1A).

Diagnostic Findings:

- Diffuse cerebral white matter T2 hyperintensities: Pathognomonic for LAMA2-CMD (present in >90% of complete deficiency cases)

- Early-onset severe hypotonia + elevated CK: Consistent with congenital muscular dystrophy
- Normal cognition expected despite white matter changes (distinctive feature)

Recommended Immediate Actions:

1. **Genetic Testing:** LAMA2 gene sequencing (confirm diagnosis; nonsense/frameshift mutations predict complete deficiency and severe phenotype)
2. **Pulmonology Referral:** URGENT
 - Assess respiratory function (PFTs if feasible, sleep study)
 - Anticipate need for non-invasive ventilation (NIV) or tracheostomy
 - Respiratory failure is leading cause of morbidity/mortality
3. **Gastroenterology/Nutrition:** Evaluate feeding safety (swallow study); consider G-tube placement to prevent aspiration and ensure growth
4. **Neurology:** Seizure risk 5-30%; monitor for focal seizures (may have visual/autonomic features)

If LAMA2 Mutations Confirmed:

Prognosis:

- Complete laminin- α 2 deficiency (if nonsense/frameshift mutations): Severe phenotype
 - No independent ambulation expected
 - Respiratory support likely required (NIV or tracheostomy)
 - Progressive contractures, scoliosis
 - **Normal cognition** (reassuring for family)
 - Life expectancy: Teens to 30s with supportive care (variable, depends on respiratory management)
- Partial deficiency (if missense mutations): Milder phenotype possible; some achieve ambulation

Management Plan:

1. Respiratory support (NIV/tracheostomy as needed)
2. Nutritional support (G-tube feeding)
3. Physical therapy (maintain range of motion, prevent contractures)
4. Orthopedic surveillance (scoliosis, hip dislocation)
5. Neurology follow-up (seizure monitoring)
6. Supportive equipment (wheelchair, standers, adaptive devices)

Emerging Therapies:

- Gene therapy trials in preclinical phase (LAMA2 gene delivery)
- LAMA1 upregulation strategies under investigation
- Recommend enrollment in natural history studies

Genetic Counseling:

- Autosomal recessive: 25% recurrence risk for future pregnancies
- Offer carrier testing to parents (confirm diagnosis)
- Prenatal diagnosis options: Chorionic villus sampling (CVS) or amniocentesis with LAMA2 sequencing
- Preimplantation genetic testing (PGT) for future pregnancies

Prognosis Summary: While LAMA2-CMD is a severe, life-limiting condition, proactive management—particularly respiratory support and nutritional optimization—improves quality of life and extends survival. Normal cognition allows for meaningful engagement and development despite physical limitations."

4. INTEGRATION OF EVIDENCE-BASED MEDICINE

Defining Evidence-Based Medicine

David Sackett, widely regarded as a pioneer of evidence-based medicine (EBM), defined it in his seminal 1996 publication as "the conscientious, explicit and judicious use of current best evidence in making decisions about the care of individual patients." Sackett emphasized that EBM requires integrating three fundamental pillars:

1. **Best Available Research Evidence:** Systematic, methodologically rigorous clinical research
2. **Individual Clinical Expertise:** The proficiency and judgment clinicians develop through experience
3. **Patient Values and Preferences:** Individual circumstances, concerns, and expectations

This definition underscores that evidence alone is insufficient; clinical decision-making must synthesize research findings with contextual factors unique to each patient.

The Challenge of EBM in Rare Diseases

Rare genetic diseases present a fundamental paradox for evidence-based medicine: they are precisely the conditions that would benefit most from precision medicine approaches, yet they suffer from the sparsest evidence base. The hallmarks of high-quality evidence—large randomized controlled trials (RCTs), meta-

analyses, and systematic reviews—are often logistically and economically infeasible for diseases affecting hundreds to thousands of patients globally. Consequently, rare disease evidence derives from:

1. **Patient Registries:** Natural history data from systematically collected cohorts
 - Example: Treat-NMD DMD Global Database (>10,000 patients across 40 countries)⁷¹
 - Provides longitudinal outcome data, genotype-phenotype correlations, survival statistics
2. **Case Series and Observational Studies:** Retrospective or prospective cohort studies
 - Example: Fanin et al. (2004) characterized 238 LGMDR1 patients, establishing genotype-phenotype correlations⁷²
 - Provides clinically actionable data despite lack of randomized controls
3. **Expert Consensus Guidelines:** Systematic reviews of available evidence synthesized by multidisciplinary panels
 - Example: Birnkrant et al. (2018) DMD Care Considerations—graded recommendations based on evidence quality^{23,34,35}
 - Fills gaps where formal trial evidence is unavailable
4. **Genetic Databases with Pathogenicity Assertions:** Curated variant-disease associations
 - Example: ClinVar submissions undergo expert review, with evidence codes (ACMG criteria) supporting classifications¹²
 - Provides mutation-specific outcome predictions
5. **Approval of Therapies Based on Surrogate Endpoints:** For rapidly progressive, life-threatening conditions, regulatory agencies (FDA, EMA) may approve therapies based on biochemical outcomes rather than clinical efficacy
 - Example: Exon-skipping drugs approved based on dystrophin expression increases, not functional outcomes¹³
 - Confirmatory trials required post-approval to validate clinical benefit

How RAG + Knowledge Graph Enables EBM for Rare Diseases

Retrieval-Augmented Generation addresses the "data poverty" paradox by systematically aggregating, structuring, and synthesizing fragmented evidence sources into clinically actionable formats.

1. Structured Evidence Aggregation: Data Sources Integrated

A. Variant Pathogenicity Databases:

- **ClinVar (NCBI):** >2.5 million variant entries with expert-reviewed pathogenicity classifications (Pathogenic, Likely Pathogenic, VUS, Likely Benign, Benign) based on ACMG criteria^{>9,12}</sup>
- **LOVD (Leiden Open Variation Database):** Locus-specific variant databases curated by disease experts (e.g., UMD-DMD database for DMD variants)^{>11}</sup>
- **Human Gene Mutation Database (HGMD):** Comprehensive repository of disease-causing mutations

B. Gene-Disease Association Databases:

- **OMIM (Online Mendelian Inheritance in Man):** Authoritative compendium of human genes and genetic phenotypes with >16,000 entries^{>10}</sup>
- **Orphanet:** Rare disease classifications, prevalence data, natural history information

C. Clinical Management Guidelines:

- **GeneReviews (UW/NCBI):** Expert-authored, peer-reviewed disease summaries with evidence-based management recommendations; updated regularly^{>57,67}</sup>
- **Professional Society Guidelines:** Birnkrant DMD Care Considerations (Lancet Neurology 2018), ACMG variant interpretation standards (2015)^{>12,23,34,35}</sup>

D. Pharmacological Databases:

- **DrugBank:** Comprehensive drug information including mechanisms, targets, FDA approval status, dosing
- **Clinical Trial Registries (ClinicalTrials.gov):** Active trials with eligibility criteria, contact information for enrollment

E. Biomedical Literature:

- **PubMed/PubMed Central:** >35 million citations with abstracts; full-text articles when available
- **Natural Language Processing (NLP):** Extract structured information from unstructured text (treatment outcomes, genotype-phenotype correlations)

F. Disease-Specific Resources:

- **Treat-NMD:** Standards of care for neuromuscular diseases, patient registry data, outcome measures

2. Knowledge Graph Structure: Entities and Relationships

Entities (Nodes):

- **Genes:** DMD, CAPN3, LAMA2 (with gene location, function, expression patterns)
- **Variants:** Deletions (e.g., DMD exons 45-50), point mutations (e.g., CAPN3 c.550delA), duplications

- **Properties:** Genomic coordinates, transcript consequence, ACMG classification, population frequency
- **Diseases:** DMD, BMD, LGMDR1, LAMA2-CMD
 - **Properties:** Inheritance pattern, age of onset, clinical features, prognosis
- **Phenotypes:** Muscle weakness (specific distribution), cardiomyopathy, white matter changes, scapular winging, respiratory insufficiency
 - Encoded using Human Phenotype Ontology (HPO) terms for standardization
- **Drugs:** Deflazacort, eteplirsen, golodirsen, viltolarsen, casimersen, elevidys, ataluren
 - **Properties:** Mechanism, indication, dosing, adverse effects, FDA approval date
- **Clinical Features:** CK level ranges, cardiac ejection fraction, FVC percentages, ambulation age
- **Guidelines:** DMD Care Considerations 2018, ACMG Variant Standards 2015
- **Clinical Trials:** NCT identifiers, eligibility criteria, trial phase, enrollment status, contact information

Relationships (Edges):

- Gene --[causes]--> Disease (with inheritance pattern: X-linked, autosomal recessive)
- Variant --[pathogenic_for]--> Disease (with ACMG classification: Pathogenic, Likely Pathogenic, VUS)
- Variant --[reading_frame_status]--> In-frame / Out-of-frame (critical for DMD/BMD distinction)
- Disease --[presents_with]--> Phenotype (with frequency: Always/Usually/Sometimes/Rarely)
- Disease --[managed_by]--> Intervention (with evidence level: A/B/C based on guideline grading)
- Variant --[eligible_for]--> Drug (mutation-specific eligibility: exon deletion pattern → exon-skipping drug)
- Disease --[surveillance_includes]--> Monitoring protocol (e.g., DMD → annual cardiac MRI)
- Disease --[prognosis]--> Outcome measure (ambulation age, life expectancy)
- Variant --[enrolled_in]--> Clinical trial (active trials for specific genotypes)

3. RAG Component: Evidence-Linked Clinical Summaries

Retrieval Module:

- **Query Processing:** User input (patient variant, clinical question) → Parsed into structured query
- **Graph Traversal:** Navigate relationships to gather relevant entities
 - Example: Variant node → Gene node → Disease node → Phenotype nodes → Treatment nodes → Guideline nodes → Trial nodes

- **Context Assembly:** Compile retrieved information into structured context (variant details, disease characteristics, management recommendations, trial options)
- **Relevance Ranking:** Prioritize most pertinent information based on evidence level (Level A guidelines prioritized over Level C), recency (recent publications weighted higher), and specificity (mutation-specific data prioritized)

Generation Module:

- **Large Language Model Integration:** GPT-4, Claude, or other advanced LLMs for natural language synthesis
- **Prompt Engineering:** Structured prompts ensuring:
 - Evidence-based reasoning (cite sources, indicate evidence levels)
 - Clinical appropriateness (recommendations align with established guidelines)
 - Actionable format (specific next steps, dosing information, referral recommendations)
 - Source transparency (citations to guidelines, databases, publications)
- **Output Components:**
 - **Diagnosis/Phenotype Prediction:** Based on genotype (reading frame analysis for DMD/BMD)
 - **Treatment Recommendations:** Evidence-graded (Level A/B/C)
 - **Surveillance Protocols:** Cardiac monitoring frequency, PFT schedule, orthopedic follow-up
 - **Clinical Trial Matching:** Eligible trials with NCT identifiers, enrollment contacts
 - **Genetic Counseling Points:** Inheritance pattern, recurrence risk, cascade testing recommendations
 - **Prognosis:** Evidence-based expectations for disease trajectory, life expectancy

Example RAG Workflow:

User Query: "8-year-old with DMD, exon 45-52 deletion. Treatment options?"

Retrieval Process:

1. **Identify Variant:** Exon 45-52 deletion in DMD gene
2. **Frame Analysis:** Deletion spans 8 exons → Calculate reading frame
 - Exons 45-52: Total nucleotides deleted not divisible by 3 → Out-of-frame deletion
3. **Phenotype Prediction:** Out-of-frame → DMD phenotype (not BMD)
4. **Retrieve Eligible Drugs:**
 - Deletion of exon 53 would restore frame (exons 45-53 deletion would be in-frame)

- Drugs targeting exon 53: Golodirsen (Vyondys 53), Viltolarsen (Viltepso)

5. **Retrieve Evidence:** Clinical trial data, FDA approval information, safety profiles

6. **Retrieve Guidelines:** Birnkrant 2018 recommendations (corticosteroids, cardiac surveillance, respiratory monitoring)

7. **Retrieve Clinical Trials:** Query ClinicalTrials.gov for active DMD trials accepting patients with exon 45-52 deletions

RAG-Generated Output:

"Patient with DMD exon 45-52 deletion has Duchenne Muscular Dystrophy (out-of-frame deletion predicts severe phenotype). Deletion pattern is amenable to exon 53 skipping therapy.

FDA-Approved Treatment Options:

1. Exon 53 Skipping:

- **Golodirsen (Vyondys 53):** 30 mg/kg IV weekly
- **Viltolarsen (Viltepso):** 80 mg/kg IV weekly
- **Evidence:** FDA accelerated approval (2019, 2020) based on dystrophin expression increases in muscle biopsies; confirmatory trials ongoing
- **Efficacy:** Aims to slow disease progression by restoring reading frame
- **Monitoring:** Renal function (nephrotoxicity risk reported with PMO drugs); urinalysis and serum creatinine every 3 months

2. Corticosteroids (Standard of Care):

- **Deflazacort:** 0.9 mg/kg/day OR **Prednisone:** 0.75 mg/kg/day
- **Evidence:** Level A (prolonged ambulation by 2-3 years in multiple RCTs; Cochrane review)²³
- **Monitoring:** Weight, blood pressure, bone density (bisphosphonates if low BMD), behavioral changes, cataracts

3. Gene Therapy:

- **Elevidys (delandistrogene moxeparvovec):** Consider if age 4-5 years (FDA approved June 2023)
- One-time AAV-mediated micro-dystrophin gene delivery

Surveillance (Birnkrant 2018 Guidelines):

- **Cardiac:** Baseline cardiac MRI now; initiate ACE inhibitor at age 10 even if asymptomatic (prophylactic benefit)
- **Respiratory:** PFTs every 6 months starting age 12 or loss of ambulation
- **Orthopedic:** Scoliosis screening; anticipate bracing if curves >20 degrees
- **Physical Therapy:** Maintain range of motion, prevent contractures

Clinical Trials: [List NCT identifiers for active DMD trials enrolling patients with exon 45-52 deletions]

Prognosis with Contemporary Management:

- Ambulation loss expected age 13-15
- Life expectancy 30s-40s with comprehensive cardiac/respiratory care
- Exon-skipping therapy may modestly delay progression (ongoing confirmatory trials)

Sources:

- FDA Drug Approval: Golodirsen (December 2019), Viltolarsen (August 2020)
- Clinical Guideline: Birnkrant et al., Lancet Neurol 2018 (Parts 1-3)
- Variant Database: ClinVar, LOVD-UMD DMD"

Alignment with CDS Five Rights Framework⁷³

1. **Right Information:** Evidence-graded recommendations sourced from curated databases and peer-reviewed guidelines
2. **Right Person:** Delivered to clinician managing patient with specific genetic findings
3. **Right Format:** Concise, actionable summary with clear next steps (not overwhelming database dumps)
4. **Right Channel:** Integrated into clinical workflow (EHR via SMART on FHIR, genomic report interfaces, consult dashboards)
5. **Right Time:** At point of care when genetic results return or clinical questions arise

ACMG Standards Integration

The ACMG/AMP variant interpretation guidelines (Richards et al., 2015) provide a standardized framework for classifying sequence variants using 28 evidence criteria with defined strength levels (Very Strong, Strong, Moderate, Supporting) for pathogenicity (P) or benign (B) interpretation.¹² The knowledge graph encodes these criteria:

ACMG Evidence Codes:

- **PVS1 (Very Strong Pathogenic):** Null variant (nonsense, frameshift, splice site, exonic deletion) in gene where loss-of-function is disease mechanism
 - Example: LAMA2 nonsense mutation → Complete protein loss → MDC1A
- **PS1 (Strong Pathogenic):** Same amino acid change as established pathogenic variant
- **PM2 (Moderate Pathogenic):** Absent or extremely rare in population databases (ExAC, gnomAD)
- **PP3 (Supporting Pathogenic):** Multiple computational tools predict deleterious effect
- **BP4 (Supporting Benign):** Multiple computational tools predict no impact

RAG Integration: When queried about a variant, the system retrieves ACMG evidence codes supporting the classification and generates an explanation:

"Variant LAMA2 c.2043C>A (p.Tyr681*) classified as **Pathogenic** based on:

- PVS1: Nonsense variant introduces premature stop codon, predicting loss of functional laminin- α 2 (null mechanism established for LAMA2-CMD)
- PM2: Absent in gnomAD population database (not a benign polymorphism)
- Consistent with other reported nonsense mutations causing severe MDC1A phenotype
- ACMG Classification: **Pathogenic** (PVS1 + PM2 meets criteria per Richards 2015)
- Clinical Correlation: Predict complete laminin- α 2 deficiency → Severe early-onset CMD phenotype"

Quality of Evidence Indicators

Each clinical recommendation is tagged with evidence level to enable informed decision-making:

- **Level A:** High-quality evidence (RCTs, systematic reviews, validated clinical practice guidelines)
 - Example: "Corticosteroids prolong ambulation in DMD (Level A evidence, Cochrane review)"
- **Level B:** Moderate-quality evidence (cohort studies, case series with consistent results)
 - Example: "Cardiac MRI surveillance in BMD detects cardiomyopathy (Level B evidence, Melacini 1996 cohort)"
- **Level C:** Expert opinion, case reports, mechanistic rationale
 - Example: "Physical therapy for LGMDR1 maintains function (Level C evidence, expert consensus)"

This grading allows clinicians to weigh the strength of recommendations when making shared decisions with patients.

5. SYSTEM CONCEPT & TECHNICAL OVERVIEW

System Architecture Overview

The proposed clinical decision support system integrates three core components: (1) a knowledge graph structuring biomedical entities and relationships, (2) a retrieval-augmented generation pipeline combining information retrieval with large language model synthesis, and (3) a user interface enabling clinician interaction. This architecture is designed to transform fragmented genomic data into actionable clinical insights while maintaining transparency and evidence traceability.

Component 1: Knowledge Graph Construction

Graph Database Technology:

- **Neo4j or AWS Neptune:** Graph databases optimized for relationship traversal
- **RDF Triple Stores (Apache Jena, Blazegraph):** Alternative approach using semantic web standards
- **Property Graph Model:** Nodes (entities) with properties; edges (relationships) with directional types and properties

Entity Types (Nodes):

1. Genes:

- Properties: Gene symbol (DMD, CAPN3, LAMA2), HGNC ID, chromosomal location, protein product, expression patterns, function
- Example: {symbol: "DMD", location: "Xp21", protein: "Dystrophin", size: "79 exons"}

2. Variants:

- Properties: HGVS nomenclature (c.5899C>T, p.Arg1967*), genomic coordinates (GRCh38), variant type (deletion, duplication, SNV), ACMG classification, ClinVar ID, population frequency (gnomAD)
- Example: {hgvs: "c.5899C>T", consequence: "nonsense", clinvar_id: "RCV000012345", classification: "Pathogenic"}

3. Diseases:

- Properties: Disease name, OMIM ID, inheritance pattern, age of onset range, life expectancy, cardiac involvement, respiratory involvement
- Example: {name: "Duchenne Muscular Dystrophy", omim: "310200", inheritance: "X-linked recessive", onset: "2-5 years"}

4. Phenotypes (HPO Terms):

- Properties: HPO ID, phenotype description, body system, severity
- Example: {hpo_id: "HP:0003560", term: "Muscular dystrophy", system: "Musculoskeletal"}

5. Drugs:

- Properties: Generic name, brand name, mechanism, indication, dosing, route, FDA approval date, adverse effects, cost
- Example: {generic: "Golodirsen", brand: "Vyondys 53", mechanism: "Exon 53 skipping", dose: "30 mg/kg IV weekly", fda_approval: "2019-12-12"}

6. Clinical Features:

- Properties: Feature type (CK level, cardiac EF, FVC), normal range, abnormal threshold, units
- Example: {type: "CK", units: "U/L", normal: "< 200", DMD_range: "5000-20000"}

7. Guidelines:

- Properties: Title, authoring body, publication year, evidence grade, recommendation text, DOI
- Example: {title: "DMD Care Considerations Part 1", authors: "Birnkrant et al.", year: 2018, journal: "Lancet Neurol"}

8. Clinical Trials:

- Properties: NCT identifier, title, phase, status, eligibility criteria, contact information, locations
- Example: {nct: "NCT03375164", phase: "3", status: "Recruiting", indication: "DMD exon 53 amenable"}

Relationship Types (Edges):

1. Gene --[CAUSES]--> Disease

- Properties: Inheritance pattern, penetrance
- Example: (DMD) --[CAUSES {pattern: "X-linked recessive"}]--> (Duchenne MD)

2. Variant --[PATHOGENIC_FOR]--> Disease

- Properties: ACMG classification, evidence codes, confidence
- Example: (DMD exon 45-52 deletion) --[PATHOGENIC_FOR {acmg: "Pathogenic", evidence: ["PVS1"]}]-> (DMD)

3. Variant --[READING_FRAME]--> Frame Status

- Properties: Frame status (in-frame, out-of-frame)
- Example: (DMD exon 45-50 deletion) --[READING_FRAME {status: "in-frame"}]--> (BMD phenotype)

4. Disease --[PRESENTS_WITH]--> Phenotype

- Properties: Frequency (always, usually, sometimes, rarely), age of onset

- Example: (DMD) --[PRESENTS_WITH {frequency: "always", onset: "childhood"}]--> (Proximal muscle weakness)

5. Disease --[MANAGED_BY]--> Intervention

- Properties: Evidence level (A/B/C), guideline source, dosing
- Example: (DMD) --[MANAGED_BY {level: "A", source: "Birnkrant 2018"}]--> (Deflazacort 0.9mg/kg/day)

6. Variant --[ELIGIBLE_FOR]--> Drug

- Properties: Mechanism explanation, FDA approval status
- Example: (DMD exon 45-52 deletion) --[ELIGIBLE_FOR {mechanism: "Exon 53 skip restores frame"}]--> (Golodirsen)

7. Disease --[SURVEILLANCE_INCLUDES]--> Monitoring Protocol

- Properties: Frequency, starting age, stopping criteria
- Example: (DMD) --[SURVEILLANCE_INCLUDES {frequency: "annual", start_age: 10}]--> (Cardiac MRI)

Data Sources & Integration Pipeline:

1. ClinVar (Variant-Disease Associations):

- **Access:** FTP download or Entrez E-utilities API
- **Extraction:** Parse XML/VCF files for variant-disease assertions, ACMG classifications, submitter interpretations
- **Update Frequency:** Weekly incremental updates

2. OMIM (Gene-Disease Relationships):

- **Access:** API with academic license
- **Extraction:** Parse phenotype descriptions, gene functions, inheritance patterns
- **Update Frequency:** Monthly

3. GeneReviews (Clinical Management):

- **Access:** Web scraping or API (if available)
- **Extraction:** NLP to extract diagnostic criteria, management recommendations, prognosis
- **Update Frequency:** Quarterly (as reviews are updated)

4. LOVD (Locus-Specific Variants):

- **Access:** API or database dumps
- **Extraction:** Variant-phenotype correlations, founder mutations
- **Update Frequency:** Monthly

5. PubMed/PubMed Central (Literature):

- **Access:** Entrez E-utilities API
- **Extraction:** NLP/text mining for genotype-phenotype correlations, treatment outcomes, clinical trial results
- **Tools:** MetaMap (UMLS concept extraction), PubTator (entity recognition)
- **Update Frequency:** Continuous (query recent publications monthly)

6. ClinicalTrials.gov (Active Trials):

- **Access:** API
- **Extraction:** Trial eligibility criteria (genetic eligibility parsed), contact information, enrollment status
- **Update Frequency:** Weekly

7. DrugBank (Pharmacological Data):

- **Access:** API or database download
- **Extraction:** Drug mechanisms, indications, FDA approval dates, dosing, adverse effects
- **Update Frequency:** Quarterly

8. Treat-NMD (Standards of Care):

- **Access:** Manual curation from published care standards
- **Extraction:** Surveillance protocols, outcome measures
- **Update Frequency:** Annually

Data Curation Workflow:

1. **Automated Extraction:** APIs, database dumps, scheduled scraping
2. **Transformation:** Convert to standardized formats (FHIR Genomics, GA4GH standards where applicable)
3. **Quality Control:** Validate data integrity (missing fields, duplicate entries)
4. **Expert Review:** Clinical geneticist reviews high-impact assertions (e.g., novel pathogenic variants)

5. **Graph Loading:** Batch insert into graph database with relationship creation

6. **Versioning:** Timestamp updates; maintain historical versions for auditing

Component 2: RAG (Retrieval-Augmented Generation) Pipeline

Retrieval Module:

1. Query Processing:

- **Input:** Natural language query or structured input (variant HGVS, patient age, clinical features)
- **Parsing:** Extract key entities (genes, variants, diseases, phenotypes) using Named Entity Recognition (NER)
- **Intent Classification:** Determine query type (variant interpretation, treatment recommendation, differential diagnosis, prognosis)

2. Graph Traversal:

- **Starting Node:** Identified variant or disease
- **Path Exploration:**
 - Variant → Gene → Disease → Phenotypes → Treatments → Guidelines → Trials
 - Multi-hop reasoning (e.g., 3-hop: Variant → Disease → Phenotype, Variant → Drug eligibility)
- **Relationship Filtering:** Prioritize high-confidence relationships (e.g., "Pathogenic" over "VUS")
- **Context Assembly:** Compile all retrieved nodes and relationships into structured JSON context

Example Query: "DMD patient, exon 45-52 deletion, treatment options"

Graph Traversal Steps:

1. Identify Variant Node: DMD exon 45-52 deletion
2. Traverse: Variant --[PATHOGENIC_FOR]--> DMD (disease node)
3. Traverse: DMD --[PRESENTS_WITH]--> Muscle weakness, Cardiomyopathy (phenotype nodes)
4. Traverse: DMD --[MANAGED_BY]--> Deflazacort, Golodirsen, Cardiac MRI (treatment/surveillance nodes)
5. Traverse: Variant --[ELIGIBLE_FOR]--> Golodirsen, Viltolarsen (drug nodes)
6. Traverse: DMD --[GUIDELINE]--> Birnkrant 2018 (guideline node)
7. Retrieve Trial: DMD --[CLINICAL_TRIAL]--> NCT numbers

Retrieved Context (Structured JSON):

```
json

{
  "variant": {
    "deletion": "exons 45-52",
    "gene": "DMD",
    "frame_status": "out-of-frame",
    "phenotype_prediction": "Duchenne MD"
  },
  "disease": {
    "name": "Duchenne Muscular Dystrophy",
    "onset": "2-5 years",
    "progression": "Wheelchair by age 12",
    "life_expectancy": "20s-30s with care"
  },
  "treatments": [
    {
      "drug": "Golodirsen",
      "indication": "Exon 53 skipping",
      "dose": "30 mg/kg IV weekly",
      "evidence": "FDA approved 2019",
      "eligibility": "Exon 45-52 deletion amenable"
    },
    {
      "drug": "Deflazacort",
      "dose": "0.9 mg/kg/day",
      "evidence": "Level A (prolongs ambulation 2-3 years)"
    }
  ],
  "surveillance": [
    { "test": "Cardiac MRI", "frequency": "Annual", "start_age": 10 },
    { "test": "PFTs", "frequency": "Every 6 months", "start_age": 12 }
  ],
  "guidelines": {
    "source": "Birnkrant et al., Lancet Neurol 2018",
    "doi": "10.1016/S1474-4422(18)30024-3"
  },
  "trials": ["NCT03375164", "NCT02500381"]
}
```

3. Relevance Ranking:

- **Evidence Level Weighting:** Level A guidelines prioritized over Level C
 - **Recency:** Recent publications (within 5 years) weighted higher for rapidly evolving fields
 - **Specificity:** Mutation-specific data prioritized over general disease information
 - **Source Authority:** Peer-reviewed guidelines > individual case reports
-

Generation Module:

1. Large Language Model Selection:

- **Options:** GPT-4 (OpenAI), Claude (Anthropic), Med-PaLM (Google), Meditron (open-source)
- **Considerations:**
 - Context window (need to include extensive retrieved data)
 - Medical domain fine-tuning (Med-PaLM preferable for clinical accuracy)
 - Hallucination mitigation (grounding in retrieved context critical)

2. Prompt Engineering:

System Prompt:

You are a clinical genetics expert assistant. Generate evidence-based clinical recommendations for genetic variants in muscular dystrophies (DMD, BMD, LGMDR1, LAMA2-CMD).

INSTRUCTIONS:

1. Base all recommendations ONLY on the provided knowledge graph context (do not hallucinate information)
2. Cite evidence sources explicitly (guidelines, databases, publications)
3. Grade evidence quality (Level A/B/C)
4. Provide actionable next steps (specific dosing, referrals, surveillance)
5. Include genetic counseling points (inheritance, recurrence risk)
6. Identify clinical trial opportunities when relevant
7. Format output for clinician readability (structured sections, bullet points)
8. Avoid medical jargon where possible; explain technical terms

CRITICAL SAFETY RULES:

- Never recommend off-label treatments without explicit evidence
- Acknowledge uncertainty (use "may," "suggest" rather than definitive when evidence weak)
- Highlight contraindications and monitoring requirements
- Defer to human clinical judgment for complex cases

User Prompt:

PATIENT CASE:

- Age: 8 years
- Gender: Male
- Genetic Finding: DMD gene, deletion of exons 45-52
- Clinical Question: What treatment options are available?

KNOWLEDGE GRAPH CONTEXT:

{Retrieved JSON context from graph}

TASK:

Generate a comprehensive clinical summary including:

1. Phenotype prediction (DMD vs BMD)
2. Treatment recommendations (FDA-approved, evidence-based)
3. Surveillance protocols
4. Clinical trial eligibility
5. Genetic counseling points
6. Prognosis

Cite all sources explicitly (guidelines, FDA approvals, etc.).

3. LLM Processing:

- **Input:** System prompt + User prompt + Retrieved context
- **Generation Parameters:**
 - Temperature: 0.3 (low to minimize creativity/hallucination)
 - Max tokens: 2000
 - Top-p: 0.9 (nucleus sampling)
- **Post-Processing:**
 - Validate citations (ensure all cited sources exist in knowledge graph)
 - Check for hallucinations (flag any claims not grounded in retrieved context)
 - Format output (add structured headings, bullet points)

4. Output Structure:

Generated Clinical Summary:

PATIENT: 8-year-old male with DMD exon 45-52 deletion

GENETIC ANALYSIS:

- Deletion Pattern: Exons 45-52 (out-of-frame)
- Predicted Phenotype: Duchenne Muscular Dystrophy (severe)
- Reading Frame Restoration: Exon 53 skipping would restore frame

TREATMENT RECOMMENDATIONS:

1. EXON-SKIPPING THERAPY (Mutation-Specific):

- Golodirsen (Vyondys 53): 30 mg/kg IV weekly
Evidence: FDA approved December 2019 for exon 53 skipping
- Viltolarsen (Viltepso): 80 mg/kg IV weekly
Evidence: FDA approved August 2020 for exon 53 skipping
Mechanism: Skipping exon 53 restores reading frame
Monitoring: Renal function (urinalysis, creatinine every 3 months)

2. CORTICOSTEROIDS (Standard of Care):

- Deflazacort: 0.9 mg/kg/day OR Prednisone: 0.75 mg/kg/day
Evidence: Level A (Birnkrant 2018, Lancet Neurol)
Efficacy: Prolongs ambulation 2-3 years
Monitoring: Weight, bone density, blood pressure, behavior

SURVEILLANCE PROTOCOLS (Birnkrant 2018):

- Cardiac: Baseline MRI now; ACE inhibitor at age 10; annual cardiac MRI
- Respiratory: PFTs every 6 months starting age 12
- Orthopedic: Scoliosis screening annually

CLINICAL TRIALS:

- NCT03375164 (Golodirsen extension study)
- NCT02500381 (Casimersen/Golodirsen long-term outcomes)

GENETIC COUNSELING:

- Inheritance: X-linked recessive
- Mother: Obligate carrier (50% risk for male siblings)
- Female relatives: Offer carrier testing

PROGNOSIS:

- Wheelchair dependence expected age 13-15
- Life expectancy 30s-40s with comprehensive care
- Exon-skipping therapy aims to slow progression

SOURCES:

[1] FDA Drug Approval: Golodirsen (December 2019), Viltolarsen (August 2020)

[2] Clinical Guideline: Birnkrant DJ et al. Lancet Neurol 2018;17(3):251-267

[3] Variant Database: ClinVar Accession RCV000012345

Component 3: User Interface & Integration

Dashboard Features:

1. Variant Input Interface:

- **Upload Genetic Report:** Parse VCF, HGVS nomenclature from lab reports
- **Manual Entry:** Dropdown menus for gene selection, variant type, exon numbers
- **Validation:** Check variant format, flag novel variants not in ClinVar

2. Query Interface:

- **Natural Language:** "What treatments are available for this patient?"
- **Structured Forms:** Checkboxes for query type (Diagnosis, Treatment, Prognosis, Trials)
- **Patient Context:** Age, gender, symptom onset, family history (optional inputs to refine recommendations)

3. Results Display:

A. Phenotype Prediction:

- **Visual:** Traffic light system (Green: Benign, Yellow: VUS, Red: Pathogenic)
- **Text:** "This deletion predicts Duchenne Muscular Dystrophy (DMD) phenotype based on out-of-frame reading."
- **Confidence:** "Reading frame rule has 90% predictive accuracy (Monaco 1988)"

B. Treatment Recommendations:

- **Tabular format:** Drug | Dose | Route | Evidence Level | FDA Status
- **Expandable details:** Mechanism, adverse effects, monitoring requirements
- **Visual:** Evidence pyramid (Level A at top, C at bottom)

C. Clinical Trial Matches:

- **List:** NCT identifiers with trial names
- **Eligibility:** "Patient meets genetic eligibility criterion"

- **Contact:** Principal investigator email, enrollment phone number
- **Link:** Direct link to ClinicalTrials.gov page

D. Knowledge Graph Visualization:

- **Interactive graph:** Central variant node connected to disease, drugs, phenotypes
- **Hoverable edges:** Display relationship properties (e.g., "ELIGIBLE_FOR: Exon 53 skipping restores frame")
- **Filtering:** Toggle node types (show/hide phenotypes, trials, etc.)

4. Evidence Transparency:

- **Clickable citations:** Hyperlinks to PubMed, guideline PDFs, ClinVar entries
- **Source credibility indicators:** Peer-reviewed guideline (high), case report (low)
- **Evidence trail:** Show which knowledge graph nodes contributed to each recommendation

5. Export Function:

- **PDF Report:** Generate structured clinical summary for EHR documentation
- **HL7 FHIR Format:** Export recommendations as structured data for EHR integration
- **Patient-Friendly Summary:** Plain language version for shared decision-making

Integration Points:

1. MANA AI Platform:

- **Embed RAG-CDS module** within existing health informatics infrastructure
- **Single sign-on (SSO)** for clinician authentication
- **Role-based access control** (geneticists, neurologists, genetic counselors)

2. EHR Integration (Future):

- **SMART on FHIR App:** Embed as application within Epic, Cerner, Allscripts
- **CDS Hooks:** Trigger recommendations when genetic test results entered
- **FHIR Genomics Resources:** Consume genetic data in standardized format (Observation-genetics, DiagnosticReport)

3. Genetic Lab Interfaces:


- **API:** Receive variant calls directly from sequencing platforms (Illumina, PacBio)

- **Automated Workflow:** Variant uploaded → RAG analysis triggered → Report generated

4. Clinical Trial Registries:

- **Auto-Update:** Scheduled queries to ClinicalTrials.gov API for new trials
- **Patient Alerts:** Notify when new eligible trial opens

Simple Workflow Diagram:



6. EXPECTED CLINICAL & ORGANIZATIONAL IMPACT

Patient-Level Impact

1. Reduced Diagnostic Delays:

Current State Burden: The average diagnostic odyssey for rare diseases spans 5-7 years, with patients consulting 7-8 physicians before receiving accurate diagnosis.³ For muscular dystrophies specifically, delays of 2-5 years from symptom onset to genetic confirmation are common.⁷⁴ This delay has profound consequences:

- Missed therapeutic windows (corticosteroids most effective when initiated before age 5 in DMD)
- Unnecessary diagnostic procedures (invasive muscle biopsies, multiple specialist consultations)
- Psychological burden on families (uncertainty, lack of prognosis)
- Financial costs (\$5,000-\$10,000 in wasteful testing before diagnosis)⁷⁵

With RAG-CDS:

- Rapid variant interpretation reduces genetic confirmation to days-weeks rather than months-years
- Reading frame analysis automated (DMD vs. BMD distinction instant)
- Differential diagnosis support accelerates workup (targeted genetic testing panels rather than broad sequencing)
- Early diagnosis enables timely intervention

Clinical Significance:

- **DMD:** Corticosteroid initiation before age 5 prolongs ambulation by 2-3 years; delay beyond age 7 reduces benefit²³
- **LAMA2-CMD:** Early respiratory support planning prevents acute crises, reduces hospitalizations
- **BMD:** Timely distinction from DMD provides accurate prognostic counseling, avoids inappropriate treatment intensity

2. Access to Precision Therapies:

Current Barrier: Mutation-specific therapies revolutionize treatment but introduce complexity in patient-therapy matching. For DMD:

- Eteplirsen (exon 51 skipping): Applicable to ~13% of patients
- Golodirsen/Viltolarsen (exon 53 skipping): ~8% each
- Casimersen (exon 45 skipping): ~8%
- Eligibility determination requires bioinformatics expertise to predict reading frame restoration—clinicians may not recognize eligible patients without computational support.^{13,76}

With RAG-CDS:

- Automatic eligibility matching: Deletion pattern → Reading frame analysis → Eligible drugs identified
- Reduces "therapeutic nihilism": Clinicians aware of all treatment options
- Prevents missed opportunities: Patient doesn't miss eligible therapy due to lack of awareness

Example Impact:

- Patient with DMD exon 45-52 deletion identified as eligible for golodirsen/viltolarsen
- Exon-skipping therapy initiated → Slowed disease progression (ongoing confirmatory trials assessing magnitude of benefit)
- Without CDS: Risk of clinician unaware of eligibility → patient receives only standard corticosteroids

3. Clinical Trial Enrollment:

Current State: <5% of rare disease patients enroll in clinical trials despite >7,000 rare diseases with >10,000 active trials globally.⁷⁷ Barriers include:

- Clinician unawareness of relevant trials
- Complex eligibility criteria (genetic specifications buried in lengthy protocols)
- Lack of patient-trial matching infrastructure

With RAG-CDS:

- Real-time trial matching based on genotype
- NCT identifiers, contact information, eligibility summaries presented at point of care
- Increased enrollment → Accelerated therapeutic development

Example Impact:

- LGMDR1 patient identified as eligible for CAPN3 gene therapy Phase 1/2 trial
- Patient enrolled → Potential access to experimental therapy; contributes to research
- Broader Impact: Faster trial completion → Earlier regulatory approval

4. Improved Prognostic Counseling:

Genetic Heterogeneity Challenge: Same gene (DMD) produces vastly different phenotypes (DMD vs. BMD):

- DMD: Wheelchair by age 12, life expectancy 20s-30s
- BMD: Ambulatory into 30s-50s, near-normal life expectancy with cardiac management Accurate phenotype prediction critical for:
- Educational planning (mainstream vs. adapted education)
- Career counseling (physical demands of occupation)
- Life planning (independent living feasibility, family planning)

- Psychological preparation

With RAG-CDS:

- Reading frame analysis → Accurate DMD vs. BMD prediction
- LAMA2 mutation type → Complete vs. partial deficiency → Severity prediction
- Evidence-based prognosis with confidence intervals

Example Impact:

- Family learns child has BMD (not DMD) → Appropriate expectations, long-term planning
 - Avoids unnecessary psychological burden of DMD prognosis when BMD is correct diagnosis
-

Provider-Level Impact

1. Reduced Cognitive Load:

Current State Burden: Clinicians managing rare diseases face:

- Information overload: >2.5 million ClinVar variants, 16,000+ OMIM entries, 35+ million PubMed articles
- Time scarcity: 30-60+ minutes per variant spent manually cross-referencing databases
- Burnout: Tedious, repetitive knowledge retrieval tasks

With RAG-CDS:

- Integrated evidence retrieval in seconds
- Synthesized recommendations rather than raw database dumps
- More time for patient interaction, counseling, complex decision-making

Quantifiable Impact:

- Time savings: 30-60 minutes → <5 minutes per variant interpretation
- Burnout reduction: Automation of repetitive tasks improves job satisfaction⁷⁸

2. Enhanced Diagnostic Accuracy:

Current Diagnostic Errors:

- Misdiagnosis rates in rare diseases: 20-40% initially incorrect or delayed diagnosis⁷⁹
- LGMDR1 frequently misdiagnosed as BMD due to phenotypic overlap

- VUS misinterpretation: Variants classified as VUS may be reclassified as pathogenic with additional evidence

With RAG-CDS:

- Differential diagnosis support: System flags overlapping conditions, suggests distinguishing tests
- Variant reclassification alerts: ClinVar updates trigger re-analysis
- Pattern recognition: Combination of features (scapular winging + normal cardiac + elevated CK + autosomal recessive) → LGMDR1 flagged as most likely

Quantifiable Impact:

- Diagnostic accuracy improvement: 20-30% reduction in misdiagnosis (extrapolated from CDS studies in other domains)^{>80}
- Faster definitive diagnosis: Targeted genetic panels rather than broad exome sequencing

3. Evidence Access at Point of Care:

Current State:

- Guidelines exist but not readily accessible during patient encounters
- Clinicians rely on memory, outdated knowledge, or time-consuming literature searches
- Variation in care: Practice patterns differ between institutions, specialists

With RAG-CDS:

- Real-time guideline recommendations embedded in workflow
- Evidence-graded (Level A/B/C) so clinicians understand strength of recommendation
- Standardized care: All providers receive same evidence-based guidance

Quantifiable Impact:

- Guideline adherence: Improvement from 30-50% to 70-90% (observed in other CDS implementations)^{>81}
- Quality improvement: Appropriate corticosteroid initiation, cardiac surveillance compliance

4. Education & Expertise Democratization:

Current Expertise Gap:

- Genetic specialists concentrated in academic medical centers (limited rural access)

- Primary care and general neurology have limited rare disease training
- Knowledge explosion: New therapies, trials, variants published continuously (impossible for individual clinician to stay current)

With RAG-CDS:

- Community hospitals access subspecialty-level recommendations
- Non-specialists manage rare diseases with expert-level guidance
- Continuous knowledge updates: System reflects latest evidence (provider doesn't need to manually update knowledge)

Quantifiable Impact:

- Geographic equity: Reduced disparity in care between urban academic centers and rural/community settings
- Provider confidence: Surveys show increased comfort managing rare diseases with CDS support⁸²

Healthcare System / Organizational Impact

1. Cost Reduction:

Diagnostic Odyssey Costs:

- Average cost of diagnostic odyssey before rare disease diagnosis: \$5,000-\$10,000 in unnecessary testing⁷⁵
- Includes: Muscle biopsies (\$1,000-\$3,000), EMG/NCS (\$500-\$1,500), MRI (\$1,000-\$3,000), multiple specialist consultations
- Multiply by 30 million Americans with rare diseases → Substantial societal cost

With RAG-CDS:

- Faster diagnosis → Fewer unnecessary tests
- Targeted genetic testing: Mutation-specific panels (lower cost) rather than whole exome sequencing (\$3,000-\$5,000)
- Early intervention prevents complications: LAMA2-CMD respiratory support reduces ICU admissions

Quantifiable Savings:

- Estimated 20-30% reduction in pre-diagnosis testing costs
- Avoided acute complications (e.g., respiratory failure in LAMA2-CMD) save \$10,000-\$50,000 per hospitalization

2. Precision Medicine Infrastructure:

Current Fragmentation:

- Each institution maintains separate databases, workflows
- Data silos impede research, quality improvement
- Lack of interoperability between genetic labs, EHRs, research databases

With RAG-CDS as Foundation:

- Standardized knowledge representation (FHIR Genomics, GA4GH standards)
- Interoperable data exchange between institutions
- Scalable to other disease domains: Framework extends beyond muscular dystrophies to metabolic disorders (PKU, MCAD), connective tissue disorders (Marfan, Ehlers-Danlos), hematologic conditions (hemophilia, sickle cell)

Organizational Benefit:

- Research acceleration: Aggregated genomic-phenotypic data supports outcome studies
- Participation in consortia: Interoperable data enables multi-institution collaborations (Treat-NMD, EURO-NMD)

3. Quality Metrics & Accreditation:

Meaningful Use / MIPS:

- Demonstrates integration of genomic data into clinical decision-making (Meaningful Use criteria)
- CDS implementation contributes to Merit-based Incentive Payment System (MIPS) quality score

Patient-Centered Outcomes Research (PCOR):

- Rapid diagnosis improves patient experience, satisfaction scores
- Shared decision-making facilitated by evidence-based prognosis

Specialty Certification:

- Supports Neuromuscular Center of Excellence designation

- Quality improvement data: Adherence to DMD Care Considerations 2018 guidelines

4. Population Health:

Carrier Screening:

- Identify at-risk families before symptom onset
- LGMDR1, LAMA2-CMD: Autosomal recessive → 25% recurrence risk for siblings
- Cascade testing: First-degree relatives (50% carrier risk for X-linked; 50% carrier risk for AR if parent carrier)

Newborn Screening Integration:

- Early DMD detection (elevated CK on newborn screen) → Pre-symptomatic corticosteroid trial (research ongoing)
- CDS flags newborn with elevated CK + family history → Expedited genetic testing

Registry Linkage:

- Connect patients to natural history studies: Treat-NMD DMD Global Database
- Long-term outcome tracking: Real-world evidence for therapy effectiveness

Quantifiable Impact:

- Family planning: Genetic counseling reduces recurrence in subsequent pregnancies (prenatal diagnosis, preimplantation genetic testing)
- Population-level disease burden reduction

Scalability & Future Extensions

Disease Expansion:

Phase 1 (Current Proposal): 4 muscular dystrophies (DMD, BMD, LGMDR1, LAMA2-CMD)

- Proof-of-concept demonstrating feasibility, clinical utility
- Foundation: Knowledge graph infrastructure, RAG pipeline, user interface

Phase 2 (Year 2-3): Expand to all muscular dystrophies

- Additional LGMD subtypes: LGMDR2 (dysferlinopathy, DYSF), LGMDR9 (FKRP), sarcoglycanopathies (SGCA, SGCB, SGCG, SGCD)

- Other muscular dystrophies: Facioscapulohumeral MD (FSHD, DUX4), Myotonic Dystrophy (DM1, DMPK; DM2, CNBP), Emery-Dreifuss MD (EMD, LMNA)
- Congenital myopathies: Central core disease (RYR1), nemaline myopathy (NEB, ACTA1)

Phase 3 (Year 4-5): Broader neuromuscular disorders

- Charcot-Marie-Tooth disease (PMP22, MPZ, GJB1, MFN2)
- Spinal Muscular Atrophy (SMN1)
- Metabolic myopathies (CPT2, PYGM, GAA)

Phase 4 (Year 5+): Other rare disease categories

- Metabolic disorders: Phenylketonuria (PAH), MCAD deficiency (ACADM)
- Connective tissue: Marfan syndrome (FBN1), Ehlers-Danlos (COL5A1, COL5A2)
- Hematologic: Hemophilia A (F8), Sickle cell disease (HBB)
- Immunodeficiency: SCID subtypes (IL2RG, ADA)

Technical Scalability:

- Knowledge graph: Neo4j supports billions of nodes/edges (sufficient for >7,000 rare diseases)
- RAG pipeline: Modular design allows parallel processing of multiple queries
- Cloud deployment (AWS, Azure, GCP): Auto-scaling based on demand

EHR Integration:

HL7 FHIR Standards:

- **FHIR Genomics Implementation Guide:** Standardized representation of genetic variants (Observation-genetics resource)
- **DiagnosticReport:** Genetic test results in structured format
- **MedicationRequest:** Treatment recommendations as actionable FHIR resources

SMART on FHIR App:

- Embeddable within Epic (largest US EHR), Cerner, Allscripts
- Single sign-on (OAuth 2.0): Clinician authentication via EHR credentials
- Context-aware: Automatically pull patient demographics, prior genetic tests from EHR

CDS Hooks:

- Event-driven triggers: When genetic test result entered → RAG-CDS analysis triggered automatically
- Card-based recommendations: Appear within EHR workflow (similar to drug interaction alerts)

FHIR Resources Generated by CDS:

- **ServiceRequest:** Referral to genetic counselor, cardiac surveillance order
- **CarePlan:** Structured management plan (corticosteroids, PFTs, MRI surveillance)
- **RiskAssessment:** Prognosis, life expectancy estimates

Clinical Genomics Ecosystem:

Genetic Lab Integration:

- API connections: Receive variant calls from Illumina BaseSpace, Pacific Biosciences SMRT Link, Oxford Nanopore EPI2ME
- VCF/HGVS parsing: Automated extraction of variants from sequencing reports
- Bi-directional: CDS sends interpretations back to lab report (annotated VCF)

Pharmacogenomics Expansion:

- Integrate CPIC (Clinical Pharmacogenomics Implementation Consortium) guidelines
- Drug-gene interactions: CYP2D6 status → Codeine metabolism predictions
- Dosing recommendations: DPYD → Fluoropyrimidine (5-FU, capecitabine) dosing adjustments

Multi-Omic Integration (Research Use):

- Link genomic variants with transcriptomics (RNA-seq: aberrant splicing)
- Proteomics: Dystrophin quantification from muscle biopsy
- Metabolomics: Biomarker correlations (CK levels, urinary metabolites)
- Research Platform: Integrated multi-omic view for genotype-phenotype research

Global Collaboration:

International Registries:

- Link to Treat-NMD Global Database (DMD/BMD patients across 40+ countries)
- European neuromuscular registries (EURO-NMD, AFM-Téléthon registries)
- Data sharing: Anonymized variant-phenotype data contributes to global knowledge

Multi-Language Support:

- Translate guidelines, drug information into multiple languages
- Cultural adaptation: Dosing adjustments for population-specific factors (pharmacogenomics)

Low-Resource Settings:

- Cloud-based deployment: No on-premise infrastructure required
 - Regions lacking genetic subspecialists benefit most from democratized expertise
 - Telemedicine integration: Remote genetic counseling enabled by CDS support
-

Measurable Outcomes for Evaluation

Primary Outcomes:

1. **Time to Diagnosis:** Reduce from 2-5 years to <6 months
 - Metric: Days from symptom onset to genetic confirmation
 - Goal: 50% reduction in diagnostic delay
2. **Treatment Appropriateness:** Increase eligible patients receiving mutation-specific therapy
 - Metric: % of DMD patients with amenable deletions receiving exon-skipping drugs
 - Goal: From <50% to >90%

Secondary Outcomes:

1. **Clinician Satisfaction:**
 - Metric: Survey-based (5-point Likert scale) on time saved, confidence in decisions
 - Goal: >80% report positive impact
2. **Clinical Trial Enrollment:**
 - Metric: % of eligible patients enrolled in trials
 - Goal: Increase from <5% to >20%
3. **Diagnostic Accuracy:**
 - Metric: Rate of final diagnosis concordant with CDS suggestion
 - Goal: >85% concordance
4. **Guideline Adherence:**
 - Metric: % of DMD patients receiving guideline-recommended surveillance (cardiac MRI, PFTs)

- Goal: Increase from 60% to >85%
-

7. CONCLUSION

This project addresses a fundamental challenge in modern healthcare: translating the exponential growth of genomic knowledge into actionable clinical decisions that improve patient outcomes. Muscular dystrophies—with their extreme genetic heterogeneity, overlapping phenotypes, and rapidly evolving therapeutic landscape—exemplify the complexity clinicians face in genomic medicine. By constructing a knowledge graph integrating genetic variants, diseases, phenotypes, treatments, and clinical guidelines, and augmenting it with retrieval-augmented generation, we create a system that not only retrieves information but synthesizes it into evidence-based clinical insights.

Innovation Highlights

The proposed system addresses critical clinical challenges unique to rare genetic diseases:

1. Same Gene, Profoundly Different Diseases: The DMD gene paradigm—where out-of-frame mutations cause devastating Duchenne phenotype while in-frame mutations produce milder Becker phenotype—demonstrates the necessity of computational genotype-phenotype correlation. Our system automates reading frame analysis, instantly predicting phenotype and guiding appropriate prognostic counseling. This distinction has profound implications: a family learning their child has BMD rather than DMD fundamentally changes their expectations, life planning, and treatment intensity.

2. Mutation-Specific Precision Therapies: The FDA approval of four exon-skipping drugs for DMD (2016-2021) revolutionized treatment but introduced matching complexity. Each drug targets specific deletion patterns: eteplirsen for ~13% of patients, golodirsen/viltolarsen for ~8% each, and casimersen for ~8%. Determining eligibility requires predicting whether skipping a specific exon restores the reading frame—a computational task beyond most clinicians' capacity. Our RAG-enabled system automatically matches patient deletion patterns to eligible therapies, preventing missed treatment opportunities due to clinician unawareness.

3. Differential Diagnosis Through Pattern Recognition: LGMDR1 (calpainopathy) presents diagnostic challenges due to phenotypic overlap with BMD and other LGMD subtypes. The knowledge graph encodes distinguishing features: prominent scapular winging, autosomal recessive inheritance, absence of cardiac involvement. When queried with a clinical scenario, the RAG system retrieves these patterns, assigns likelihood scores based on feature combinations, and suggests targeted genetic testing—accelerating diagnosis from years to weeks.

4. Early-Onset Disease Management: LAMA2-related congenital muscular dystrophy requires proactive respiratory management from infancy to prevent life-threatening crises. The pathognomonic brain MRI finding—diffuse white matter T2 hyperintensities—combined with severe hypotonia and elevated CK, enables early

genetic diagnosis. Our system flags this pattern, triggers urgent pulmonology referral, and provides prognostic information (normal cognition despite white matter changes) that is critical for family counseling.

Bridging Genomic Data and Clinical Evidence

The RAG-powered knowledge graph fundamentally addresses the challenge at the heart of precision medicine: translating genetic information into clinical action. Current workflows require clinicians to:

1. Query ClinVar for variant pathogenicity
2. Search OMIM for disease characteristics
3. Review GeneReviews for management guidelines
4. Search PubMed for treatment evidence
5. Query ClinicalTrials.gov for trial eligibility
6. Manually synthesize across sources

This fragmented process, requiring 30-60+ minutes per variant, is unsustainable as genetic testing becomes routine. Our system performs these queries in seconds, synthesizes findings, and presents evidence-based recommendations with transparent citations—providing what individual clinicians cannot: comprehensive, up-to-date, synthesized knowledge at the point of care.

Transforming Rare Disease Care: Immediate and Tangible Impact

The consequences of this system are not theoretical but immediate:

- A 3-year-old diagnosed with DMD at symptom onset rather than age 7 gains 4 years of corticosteroid benefit—translating to years of additional ambulation, preserved respiratory/cardiac function, and improved quality of life.
- A patient with DMD exon 45-52 deletion receives golodirsen exon-skipping therapy rather than corticosteroids alone—accessing mutation-specific treatment proven to increase dystrophin expression.
- A family learns their 10-year-old has BMD (not DMD) through accurate reading frame analysis—avoiding years of unnecessary psychological burden and enabling appropriate long-term planning for a near-normal life expectancy.
- An infant with LAMA2-CMD receives respiratory support before the first crisis through early genetic diagnosis facilitated by brain MRI pattern recognition—preventing ICU admission, improving outcomes.
- A 12-year-old with LGMDR1 avoids unnecessary cardiac interventions through accurate diagnosis distinguishing calpainopathy (cardiac sparing) from DMD/BMD (cardiac involvement)—saving costs, preventing patient anxiety.

Broader Implications for Precision Medicine

With 7,000 rare diseases affecting 30 million Americans, and 80% having genetic origins, this framework has scalability far beyond muscular dystrophies. The knowledge graph architecture generalizes to any monogenic disorder: metabolic diseases (PKU, MCAD deficiency), connective tissue disorders (Marfan, Ehlers-Danlos), hematologic conditions (hemophilia, sickle cell disease), immunodeficiencies (SCID variants), and beyond. As whole genome sequencing costs approach \$100, the bottleneck shifts from data generation to data interpretation—precisely where AI-augmented clinical decision support excels.

Vision Forward: Democratizing Genomic Medicine

This project lays groundwork for a future where genomic medicine is not confined to academic medical centers with genetic subspecialists, but accessible to any clinician caring for any patient, anywhere. A future where:

- **Diagnostic odysseys are measured in weeks, not years:** Automated variant interpretation, differential diagnosis support, and targeted genetic testing panels accelerate definitive diagnosis.
- **Precision therapies reach every eligible patient:** Real-time eligibility matching ensures no patient misses mutation-specific treatment due to clinician unawareness.
- **Evidence-based medicine adapts in real-time:** As new therapies are approved, guidelines updated, and genetic discoveries published, the knowledge graph reflects current best evidence within days—not the years required for knowledge translation via continuing medical education.
- **Health equity is advanced:** Rural hospitals, community practices, and low-resource settings access subspecialty-level rare disease expertise, reducing geographic and socioeconomic disparities in care quality.
- **Clinician cognitive load is reduced:** Automation of tedious knowledge retrieval tasks allows clinicians to focus on complex clinical reasoning, patient counseling, and compassionate care—reducing burnout and improving job satisfaction.

Final Statement

By combining structured knowledge representation, AI-driven synthesis, and clinical expertise grounded in evidence-based medicine, we can close the gap between genomic possibility and clinical reality. This is not incremental improvement but transformative change: from reactive to proactive, from fragmented to integrated, from delayed to immediate. This is the promise of retrieval-augmented generation for clinical decision support—turning genetic data into clinical wisdom, at the speed and scale that rare disease patients deserve.

REFERENCES

1. National Institutes of Health. Rare Diseases. Available at: <https://rarediseases.info.nih.gov/>

2. Theadom T, Rodrigues M, Roxburgh R, et al. Prevalence of muscular dystrophies: a systematic literature review. *Neuroepidemiology*. 2014;43(3-4):259-268.
3. Rare Disease Day. The diagnostic odyssey. Available at: <https://www.rarediseaseday.org/>
4. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
5. Aartsma-Rus A, Ginjaar IB, Bushby K. The importance of genetic diagnosis for Duchenne muscular dystrophy. *J Med Genet*. 2016;53(3):145-151.
6. Lewis P, Perez E, Piktus A, et al. Retrieval-augmented generation for knowledge-intensive NLP tasks. *Adv Neural Inf Process Syst*. 2020;33:9459-9474.
7. Miao J, Thongprayoon C, Suppadungsuk S, et al. Integrating retrieval-augmented generation with large language models in nephrology: advancing practical applications. *Medicina (Kaunas)*. 2024;60(3):445.
8. Khanna R, Nguyen A, Samala R. Enhancing medical AI with retrieval-augmented generation: a mini narrative review. *Digit Health*. 2025;11:20552076241311433.
9. Landrum MJ, Lee JM, Benson M, et al. ClinVar: improving access to variant interpretations and supporting evidence. *Nucleic Acids Res*. 2018;46(D1):D1062-D1067.
10. Amberger JS, Bocchini CA, Schiettecatte F, Scott AF, Hamosh A. OMIM.org: Online Mendelian Inheritance in Man (OMIM®), an online catalog of human genes and genetic disorders. *Nucleic Acids Res*. 2015;43(Database issue):D789-798.
11. Aartsma-Rus A, Van Deutekom JC, Fokkema IF, Van Ommen GJ, Den Dunnen JT. Entries in the Leiden Duchenne muscular dystrophy mutation database: an overview of mutation types and paradoxical cases that confirm the reading-frame rule. *Muscle Nerve*. 2006;34(2):135-144.
12. Richards S, Aziz N, Bale S, et al. Standards and guidelines for the interpretation of sequence variants: a joint consensus recommendation of the American College of Medical Genetics and Genomics and the Association for Molecular Pathology. *Genet Med*. 2015;17(5):405-424.
13. Shirley M. Casimersen: First Approval. *Drugs*. 2021;81(7):875-879.
14. Dhillon S. Viltolarsen: First Approval. *Drugs*. 2020;80(10):1027-1031.
15. Vissing J, Barresi R, Witting N, et al. A heterozygous 21-bp deletion in CAPN3 causes dominantly inherited limb girdle muscular dystrophy. *Brain*. 2016;139(Pt 8):2154-2163.
16. Fanin M, Nascimbeni AC, Fulizio L, Angelini C. The frequency of limb girdle muscular dystrophy 2A in northeastern Italy. *Neuromuscul Disord*. 2005;15(3):218-224.

17. Frank DE, Schnell FJ, Akana C, et al. Increased dystrophin production with golodirsen in patients with Duchenne muscular dystrophy. *Neurology*. 2020;94(21):e2270-e2282.
18. ClinicalTrials.gov. Search: Muscular Dystrophy. Available at: <https://clinicaltrials.gov/>
19. Monaco AP, Bertelson CJ, Liechti-Gallati S, Moser H, Kunkel LM. An explanation for the phenotypic differences between patients bearing partial deletions of the DMD locus. *Genomics*. 1988;2(1):90-95.
20. Bushby KM, Gardner-Medwin D. The clinical, genetic and dystrophin characteristics of Becker muscular dystrophy. I. Natural history. *J Neurol*. 1993;240(2):98-104.
21. Monaco AP, Bertelson CJ, Liechti-Gallati S, Moser H, Kunkel LM. An explanation for the phenotypic differences between patients bearing partial deletions of the DMD locus. *Genomics*. 1988;2(1):90-95.
22. Geranmayeh F, Clement E, Feng LH, et al. Genotype-phenotype correlation in a large population of muscular dystrophy patients with LAMA2 mutations. *Neuromuscul Disord*. 2010;20(4):241-250.
23. Birnkrant DJ, Bushby K, Bann CM, et al. Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and neuromuscular, rehabilitation, endocrine, and gastrointestinal and nutritional management. *Lancet Neurol*. 2018;17(3):251-267.
24. Clemens PR, Rao VK, Connolly AM, et al. Safety, tolerability, and efficacy of viltolarsen in boys with Duchenne muscular dystrophy amenable to exon 53 skipping: a phase 2 randomized clinical trial. *JAMA Neurol*. 2020;77(8):982-991.
25. Philpot J, Bagnall A, King C, Dubowitz V, Muntoni F. Feeding problems in merosin deficient congenital muscular dystrophy. *Arch Dis Child*. 1999;80(6):542-547.
26. Richard I, Broux O, Allamand V, et al. Mutations in the proteolytic enzyme calpain 3 cause limb-girdle muscular dystrophy type 2A. *Cell*. 1995;81(1):27-40.
27. Mah JK, Korngut L, Dykeman J, Day L, Pringsheim T, Jette N. A systematic review and meta-analysis on the epidemiology of Duchenne and Becker muscular dystrophy. *Neuromuscul Disord*. 2014;24(6):482-491.
28. Bushby K, Finkel R, Birnkrant DJ, et al. Diagnosis and management of Duchenne muscular dystrophy, part 1: diagnosis, and pharmacological and psychosocial management. *Lancet Neurol*. 2010;9(1):77-93.
29. Mercuri E, Bönnemann CG, Muntoni F. Muscular dystrophies. *Lancet*. 2019;394(10213):2025-2038.
30. Birnkrant DJ, Bushby K, Bann CM, et al. Diagnosis and management of Duchenne muscular dystrophy, part 2: respiratory, cardiac, bone health, and orthopaedic management. *Lancet Neurol*. 2018;17(4):347-361.
31. Passamano L, Taglia A, Palladino A, et al. Improvement of survival in Duchenne Muscular Dystrophy: retrospective analysis of 835 patients. *Acta Myol*. 2012;31(2):121-125.
32. Hoffman EP, Brown RH Jr, Kunkel LM. Dystrophin: the protein product of the Duchenne muscular dystrophy locus. *Cell*. 1987;51(6):919-928.

33. Ervasti JM, Campbell KP. Membrane organization of the dystrophin-glycoprotein complex. *Cell*. 1991;66(6):1121-1131.
34. Birnkrant DJ, Bushby K, Bann CM, et al. Diagnosis and management of Duchenne muscular dystrophy, part 2: respiratory, cardiac, bone health, and orthopaedic management. *Lancet Neurol*. 2018;17(4):347-361.
35. Birnkrant DJ, Bushby K, Bann CM, et al. Diagnosis and management of Duchenne muscular dystrophy, part 3: primary care, emergency management, psychosocial care, and transitions of care across the lifespan. *Lancet Neurol*. 2018;17(5):445-455.
36. Duboc D, Meune C, Lerebours G, Devaux JY, Vaksman G, Bécane HM. Effect of perindopril on the onset and progression of left ventricular dysfunction in Duchenne muscular dystrophy. *J Am Coll Cardiol*. 2005;45(6):855-857.
37. Simonds AK, Muntoni F, Heather S, Fielding S. Impact of nasal ventilation on survival in hypercapnic Duchenne muscular dystrophy. *Thorax*. 1998;53(11):949-952.
38. Heo YA. Golodirsen: First Approval. *Drugs*. 2020;80(3):329-333.
39. Finkel RS, Flanigan KM, Wong B, et al. Phase 2a study of ataluren-mediated dystrophin production in patients with nonsense mutation Duchenne muscular dystrophy. *PLoS One*. 2013;8(12):e81302.
40. U.S. Food and Drug Administration. FDA approves first gene therapy for treatment of certain patients with Duchenne muscular dystrophy. June 22, 2023.
41. Mah JK, Korngut L, Dykeman J, Day L, Pringsheim T, Jette N. A systematic review and meta-analysis on the epidemiology of Duchenne and Becker muscular dystrophy. *Neuromuscul Disord*. 2014;24(6):482-491.
42. Bushby KM, Gardner-Medwin D. The clinical, genetic and dystrophin characteristics of Becker muscular dystrophy. I. Natural history. *J Neurol*. 1993;240(2):98-104.
43. Beggs AH, Koenig M, Boyce FM, Kunkel LM. Detection of 98% of DMD/BMD gene deletions by polymerase chain reaction. *Hum Genet*. 1990;86(1):45-48.
44. Griggs RC, Miller JP, Greenberg CR, et al. Efficacy and safety of deflazacort vs prednisone and placebo for Duchenne muscular dystrophy. *Neurology*. 2016;87(20):2123-2131.
45. Melacini P, Fanin M, Danieli GA, et al. Cardiac and respiratory involvement in advanced stage Duchenne and Becker muscular dystrophies. *Neuromuscul Disord*. 1996;6(5):367-376.
46. Gumerson JD, Michele DE. The dystrophin-glycoprotein complex in the prevention of muscle damage. *J Biomed Biotechnol*. 2011;2011:210797.
47. Beckmann JS, Richard I, Hillaire D, et al. A gene for limb-girdle muscular dystrophy maps to chromosome 15 by linkage. *C R Acad Sci III*. 1991;312(3):141-148.

48. Fanin M, Nascimbeni AC, Aurino S, et al. Frequency of LGMD gene mutations in Italian patients with distinct clinical phenotypes. *Neurology*. 2009;72(16):1432-1435.
49. Straub V, Murphy A, Udd B; LGMD workshop study group. 229th ENMC international workshop: Limb girdle muscular dystrophies - Nomenclature and reformed classification Naarden, the Netherlands, 17-19 March 2017. *Neuromuscul Disord*. 2018;28(8):702-710.
50. Pantoja-Melendez CA, Miranda-Duarte A, Roque-Ramirez B, Zenteno JC. Epidemiological and molecular characterization of a Mexican population isolate with high prevalence of limb-girdle muscular dystrophy type 2A due to a novel calpain-3 mutation. *PLoS One*. 2017;12(1):e0170280.
51. Fanin M, Fulizio L, Nascimbeni AC, et al. Molecular diagnosis in LGMD2A: mutation analysis or protein testing? *Hum Mutat*. 2004;24(1):52-62.
52. Vissing J, Barresi R, Witting N, et al. A heterozygous 21-bp deletion in CAPN3 causes dominantly inherited limb girdle muscular dystrophy. *Brain*. 2016;139(Pt 8):2154-2163.
53. Richard I, Broux O, Allamand V, et al. Mutations in the proteolytic enzyme calpain 3 cause limb-girdle muscular dystrophy type 2A. *Cell*. 1995;81(1):27-40.
54. Kramerova I, Kudryashova E, Tidball JG, Spencer MJ. Null mutation of calpain 3 (p94) in mice causes abnormal sarcomere formation in vivo and in vitro. *Hum Mol Genet*. 2004;13(13):1373-1388.
55. Fanin M, Nascimbeni AC, Fulizio L, Trevisan CP, Meznaric-Petrusa M, Angelini C. Loss of calpain-3 autocatalytic activity in LGMD2A patients with normal protein expression. *Am J Pathol*. 2003;163(5):1929-1936.
56. Mercuri E, Bushby K, Ricci E, et al. Muscle MRI findings in patients with limb girdle muscular dystrophy with calpain 3 deficiency (LGMD2A) and early contractures. *Neuromuscul Disord*. 2005;15(2):164-171.
57. Vissing J, Barresi R, Witting N, et al. Calpainopathy. In: Adam MP, Feldman J, Mirzaa GM, et al., editors. *GeneReviews* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2025. Updated 2016 May 5.
58. Bartoli M, Poupiot J, Goyenvallé A, et al. Noninvasive monitoring of therapeutic gene transfer in animal models of muscular dystrophies. *Gene Ther*. 2006;13(1):20-28.
59. Allamand V, Guicheney P. Merosin-deficient congenital muscular dystrophy, autosomal recessive (MDC1A, MIM  #156225), LAMA2 gene coding for $\alpha 2$ chain of laminin). *Eur J Hum Genet*. 2002;10(2):91-94.
60. Mostacciuolo ML, Miorin M, Martinello F, Angelini C, Perini P, Trevisan CP. Genetic epidemiology of congenital muscular dystrophy in a sample from north-east Italy. *Hum Genet*. 1996;97(3):277-279.
61. Philpot J, Sewry C, Pennock J, Dubowitz V. Clinical phenotype in congenital muscular dystrophy: correlation with expression of merosin in skeletal muscle. *Neuromuscul Disord*. 1995;5(4):301-305.

62. Natera-de Benito D, Töpf A, Vilchez JJ, et al. Molecular characterization of congenital muscular dystrophies in Spain. *Ann Neurol*. 2020;88(2):293-307.
63. Tan E, Topaloglu H, Sewry C, et al. Late onset muscular dystrophy with cerebral white matter changes due to partial merosin deficiency. *Neuromuscul Disord*. 1997;7(2):85-89.
64. Helbling-Leclerc A, Zhang X, Topaloglu H, et al. Mutations in the laminin alpha 2-chain gene (LAMA2) cause merosin-deficient congenital muscular dystrophy. *Nat Genet*. 1995;11(2):216-218.
65. Oliveira J, Santos R, Soares-Silva I, et al. LAMA2 gene mutation update: toward a more comprehensive picture of the laminin- α 2 variome and its related phenotypes. *Hum Mutat*. 2018;39(10):1314-1337.
66. Philpot J, Cowan F, Pennock J, et al. Merosin-deficient congenital muscular dystrophy: the spectrum of brain involvement on magnetic resonance imaging. *Neuromuscul Disord*. 1999;9(2):81-85.
67. Yonekawa T, Nishino I. LAMA2 Muscular Dystrophy. In: Adam MP, Feldman J, Mirzaa GM, et al., editors. *GeneReviews* [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2025. Updated 2020 Sep 17.
68. Girgenrath M, Dominov JA, Kostek CA, Miller JB. Inhibition of apoptosis improves outcome in a model of congenital muscular dystrophy. *J Clin Invest*. 2004;114(11):1635-1639.
69. Goudenege S, Lebel C, Huot NB, et al. Myoblasts derived from normal hESCs and dystrophic hiPSCs efficiently fuse with existing muscle fibers following transplantation. *Mol Ther*. 2012;20(11):2153-2167.
70. Sackett DL, Rosenberg WM, Gray JA, Haynes RB, Richardson WS. Evidence based medicine: what it is and what it isn't. *BMJ*. 1996;312(7023):71-72.
71. Bladen CL, Salgado D, Monges S, et al. The TREAT-NMD DMD Global Database: analysis of more than 7,000 Duchenne muscular dystrophy mutations. *Hum Mutat*. 2015;36(4):395-402.
72. Fanin M, Nascimbeni AC, Aurino S, et al. Frequency of LGMD gene mutations in Italian patients with distinct clinical phenotypes. *Neurology*. 2009;72(16):1432-1435.
73. Osheroff JA, Teich JM, Middleton B, Steen EB, Wright A, Detmer DE. A roadmap for national action on clinical decision support. *J Am Med Inform Assoc*. 2007;14(2):141-145.
74. Rare Disease Impact Report. Global Genes. 2013.
75. Zuryski Y, Deverell M, Dalkeith T, et al. Australian children living with rare diseases: experiences of diagnosis and perceived consequences of diagnostic delays. *Orphanet J Rare Dis*. 2017;12(1):68.
76. Aartsma-Rus A, Fokkema I, Verschuuren J, et al. Theoretic applicability of antisense-mediated exon skipping for Duchenne muscular dystrophy mutations. *Hum Mutat*. 2009;30(3):293-299.
77. Hwang TJ, Tomasi PA, Bourgeois FT. Delays in completion and results reporting of clinical trials under the Paediatric Regulation in the European Union: A cohort study. *PLoS Med*. 2018;15(3):e1002520.

78. Shanafelt TD, Dyrbye LN, Sinsky C, et al. Relationship between clerical burden and characteristics of the electronic environment with physician burnout and professional satisfaction. *Mayo Clin Proc.* 2016;91(7):836-848.
79. Anderson M, Elliott EJ, Zurynski YA. Australian families living with rare disease: experiences of diagnosis, health services use and needs for psychosocial support. *Orphanet J Rare Dis.* 2013;8:22.
80. Sutton RT, Pincock D, Baumgart DC, Sadowski DC, Fedorak RN, Kroeker KI. An overview of clinical decision support systems: benefits, risks, and strategies for success. *NPJ Digit Med.* 2020;3:17.
81. Bright TJ, Wong A, Dhurjati R, et al. Effect of clinical decision-support systems: a systematic review. *Ann Intern Med.* 2012;157(1):29-43.
82. Kawamoto K, Houlihan CA, Balas EA, Lobach DF. Improving clinical practice using clinical decision support systems: a systematic review of trials to identify features critical to success. *BMJ.* 2005;330(7494):765.