

# Bayesian Optimisation for Likelihood Free Inference

Make model parameterisation go brrr

Jacob Cumming

University of Melbourne

April 2024



# Notation

- ▶ Model is considered a (random) function  $f(\boldsymbol{\theta})$  that maps  $\boldsymbol{\theta}$  (a vector of parameters) to a model output, that can be transformed into  $\mathbf{X}$ , that has the same shape as:
- ▶  $\mathbf{X}_{\text{obs}}$ , a vector of outputs given to us usually in the forms of summary statistics (incidence, prevalence, hospitalisations etc).

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$
- ▶ Or even  $\mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}})) := \Pr(S(\mathbf{X}_{\text{obs}})|\theta)$ , where  $S(\mathbf{X}_{\text{obs}})$  is a (vector of) summary statistic(s)

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$
- ▶ Or even  $\mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}})) := \Pr(S(\mathbf{X}_{\text{obs}})|\theta)$ , where  $S(\mathbf{X}_{\text{obs}})$  is a (vector of) summary statistic(s)
- ▶  $\hat{\theta} = \arg \max_{\theta} \mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}}))$

# Parameter inference would become easy if we had...

- ▶ An explicit form for the likelihood:  $\mathcal{L}(\theta|\mathbf{X}_{\text{obs}}) := \Pr(\mathbf{X}_{\text{obs}}|\theta)$
- ▶ Or even  $\mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}})) := \Pr(S(\mathbf{X}_{\text{obs}})|\theta)$ , where  $S(\mathbf{X}_{\text{obs}})$  is a (vector of) summary statistic(s)
- ▶  $\hat{\theta} = \arg \max_{\theta} \mathcal{L}(\theta|S(\mathbf{X}_{\text{obs}}))$
- ▶  $\Pr(\theta|S(\mathbf{X}_{\text{obs}})) \propto \Pr(S(\mathbf{X}_{\text{obs}})|\theta) \Pr(\theta)$

# The Sad Truth

- ▶ As models become more complicated, explicit likelihoods don't exist (think agent based models).

# A Standard Bayesian Solution

- ▶ Approximate Bayesian Computation (ABC)
  1. Sample from prior
  2. Run model
  3. Accept or reject parameters run based on how well  $\mathbf{X}$  'matches'  $\mathbf{X}_{\text{obs}}$ .



# What is 'matches'

- ▶ Discrepancy function  $D : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$

- ▶ Can be a norm such as

$$\|S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})\|_p := (\sum_{i=1}^d |S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})|^p)^{1/p}$$


# What is 'matches'

- ▶ Discrepancy function  $D : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ 
  - ▶ Can be a norm such as
$$\|S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})\|_p := (\sum_{i=1}^d |S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})|^p)^{1/p}$$
  - ▶ Care should be taken to rescale  $S(\mathbf{X}_{\text{obs}})$  and  $S(\mathbf{X})$  appropriately (ie via a covariance matrix).


# What is 'matches'

- ▶ Discrepancy function  $D : \mathcal{X} \times \mathcal{X} \rightarrow \mathbb{R}$ 
  - ▶ Can be a norm such as
$$\|S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})\|_p := (\sum_{i=1}^d |S(\mathbf{X}) - S(\mathbf{X}_{\text{obs}})|^p)^{1/p}$$
  - ▶ Care should be taken to rescale  $S(\mathbf{X}_{\text{obs}})$  and  $S(\mathbf{X})$  appropriately (ie via a covariance matrix).
- ▶  $D(S(\mathbf{X}), S(\mathbf{X}_{\text{obs}}))$ , gives acceptance probability of  $\theta$ .

# Acceptance Probability



## Attempt 2



# Overall Idea of my Research

- ▶ What if we could 'predict' discrepancy values we hadn't seen before?

# Overall Idea of my Research


- ▶ What if we could 'predict' discrepancy values we hadn't seen before?
- ▶ For parameters 'close' to parameters we've already tried it should be easy.

# Gaussian Processes


- ▶ A class of random functions
- ▶ Common examples - Brownian motion, Ornstein Uhlenbeck process
- ▶ Model the mean discrepancy using one of these (kriging)




Function estimation with  $\ell = 0.5$



Function estimation with  $\ell = 1$



Function estimation with  $\ell = 2$



# Sample frame title